Efficient Transamidation of Primary Carboxamides by in Situ Activation with *N,N*-Dialkylformamide Dimethyl Acetals

Thomas A. Dineen, Matthew A. Zajac, Andrew G. Myers*

Supporting Information

Index:

General Experimental Procedures	S-1
Materials	S-2
Instrumentation	S-2
Preparation of <i>N,N</i> -Diisopropylformamide Dimethyl Acetal	S-3
Preparation of <i>N'</i> -Acyl <i>N,N</i> -Diisopropylformamides	S-3
Transamidation of <i>N'</i> -Acyl <i>N,N</i> -Diisopropylformamides using Scandium Triflate as Catalyst	S-5
Preparation of a 0.7 M Solution of Zirconium Chloride in Acetonitrile	S-8
Transamidation using <i>N,N</i> -Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter	S-8
Transamidation using <i>N,N</i> -Dimethylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter	S-18

General Experimental Procedures: All reactions were performed in flame-dried glassware fitted with rubber septa under a positive pressure of argon, unless otherwise noted. Air- and moisture-sensitive liquids were transferred via syringe or stainless steel cannula. Organic solutions were concentrated by rotary evaporation below 35 °C at 40 mmHg. Analytical and preparative thin-layer chromatography (TLC) was performed using glass plates pre-coated with a 0.25-mm layer of silica gel impregnated with a fluorescent indicator (254 nm). TLC plates were visualized by exposure to ultraviolet light and/or submersion in aqueous ceric ammonium molybdate solution (CAM) or solutions of ninhydrin in *n*-butanol, followed by

brief heating on a hot plate (215 °C, 10-15 s). Flash-column chromatography was performed as described by Still et al.,¹ employing silica gel (60 Å, standard grade) purchased from Sorbent Technologies.

Materials: Commercial solvents and reagents were used as received with the following exceptions. Dichloromethane and ether were purified by the method of Pangborn et al.² Benzylamine, pyrrolidine, *N*-benzylmethylamine, and *s*-butylamine were distilled from calcium hydride at 760 mmHg under an atmosphere of argon and the distilled solution was stored under argon. Free-base forms of α -amino ester substrates were prepared from commercial hydrochloride salts by dissolution of the latter in a solution of aqueous sodium carbonate, extraction of the aqueous solution with ether, drying of the ethereal solutions over anhydrous sodium sulfate, and concentration of the dried solutions to provide the free-base form of the amino acid. 5-A Molecular sieves were activated by washing the commercial pellets with dichloromethane, crushing the washed sieves in a mortar and pestle, and heating the resulting powder at 170 °C under vacuum (0.1 mmHg) for 24 h. The crushed, activated sieves were stored in an oven at 150 °C until needed.

Instrumentation: Proton nuclear magnetic resonance spectra (¹H NMR) were recorded on Varian Mercury 400 (400 MHz) or Varian INOVA 500 (500 Mhz) at 23 °C. Chemical shifts are expressed in parts per million (ppm, δ scale) downfield from tetramethylsilane and are referenced to residual protium in the NMR solvent (CHCl₃, δ 7.26). Data are represented as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, sep = septet, m = multiplet and/or multiple resonances, br = broad), coupling constant in Hertz, integration and assignment. Carbon nuclear magnetic resonance spectra (¹³C NMR) were recorded on a Varian Mercury 400 (100 MHz) at 23 °C. Chemical shifts are reported in parts per million downfield from tetramethylsilane and are referenced to the carbon resonances of the solvent (CDCl₃, δ 77.0). Samples prepared infrared (IR) spectroscopy were prepared as neat films by evaporation of dichloromethane solutions; infrared spectra were recorded using a Perkin-Elmer FT-IR spectrometer referenced to a polystyrene standard. Data are represented as follows: frequency of absorption (cm⁻¹), intensity of absorption (vs = very strong, s = strong, m = medium, w = weak, br = broad). High-resolution mass spectra were obtained at the Harvard University Mass Spectrometry Facility.

¹ Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923.

² Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, *15*, 1518.

***N,N*-Diisopropylformamide dimethyl acetal (DIF-DMA).** The following is a slight modification of the procedure of Bredereck et al.³ *N,N*-Diisopropylformamide (Aldrich Co., 28.1 mL, 25.0 g, 193 mmol) and dimethyl sulfate (Aldrich Co., 18.8 mL, 25.0 g, 199 mmol) were combined in a 250-mL, round-bottom flask fitted with a Teflon-coated stir-bar. The resulting clear, yellow solution was heated at 85 °C, with stirring, for 4 d. The resulting brown reaction mixture was then cooled to 23 °C, ether (20 mL) was added to the cooled solution, and the resulting biphasic mixture was stirred vigorously for 5 min. The upper, ethereal layer was removed with a cannula (18-ga), and an additional portion of ether (20 mL) was added to the residue. The upper, ethereal layer was removed with a cannula (18-ga). The remaining thick, brown mixture was poured into a solution of sodium methoxide in methanol (prepared by the addition of 10.0 g (435 mmol) of sodium metal to 100 mL dry methanol) in a 250-mL, round-bottom flask fitted with a Teflon-coated stir bar. The resulting brown mixture was stirred for 14 h at 23 °C, then was extracted with hexane (5 × 100 mL). The hexane extracts were combined, and the combined solution was dried over anhydrous sodium sulfate. The dried solution was concentrated to afford a clear, colorless liquid containing a white, suspended precipitate. The mixture was distilled (bath 105 °C) to give 22 g of DIF-DMA as a clear, colorless liquid (65%, 88–89 °C, 40 mmHg), contaminated with ~3% *N,N*-diisopropylformamide: ¹H NMR (400 MHz, CDCl₃), δ 4.61 (s, 1 H, (CH₃O)₂CHN), 3.34 (sep, J = 7.0 Hz, 2 H, NCH(CH₃)₂), 3.25 (s, 6 H, OCH₃), 1.06 (d, J = 7.0 Hz, 12 H, CH(CH₃)₂).

Preparation of *N*'-Benzoyl *N,N*-Diisopropylformamidine from Benzamide and *N,N*-Diisopropylformamide Dimethyl Acetal. A 50-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (2.5 g) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry

³ Bredereck, H.; Simchem, G.; Rebsdatt, S.; Kantlehn, W.; Horn, P.; Wahl, R.; Hoffmann, H.; Grieshab, P. *Chem. Ber.* **1968**, *101*, 41.

the molecular sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. Benzamide (300 mg, 2.48 mmol) and dichloromethane (25 mL) were added in sequence. The resulting suspension was stirred for 5 min at 23 °C, and *N,N*-diisopropylformamide dimethyl acetal (DIF-DMA, 530 µL, 2.97 mmol, 1.25 equiv) was added via syringe. After stirring for 3 h at 23 °C, the reaction mixture was filtered through Celite, and the filter cake was washed with dichloromethane (3 × 10 mL). The filtrates were combined, and the combined solution was evaporated. The residue was purified by flash-column chromatography on silica gel (50% ether-hexane) providing 558 mg of *N'*-benzoyl *N,N*-diisopropylformamidine as a white, amorphous solid (97%): ¹H NMR (400 MHz, CDCl₃) δ 8.83 (s, 1 H, NCHN), 8.30-8.25 (m, 2 H, Ar-H), 7.46-7.51 (m, 1 H, Ar-H), 7.45-7.39 (m, 2 H, Ar-H), 4.82 (sep, J = 6.8 Hz, 1 H, isopropyl CH(CH₃)₂), 3.75 (sep, J = 6.8 Hz, 1 H, isopropyl CH(CH₃)₂), 1.38 (d, J = 6.8 Hz, 6 H, isopropyl CH(CH₃)₂), 1.37 (d, J = 6.8 Hz, 6 H, CH(CH₃)₂); ¹³C NMR (100 MHz, CDCl₃) δ 177.4, 157.9, 137.0, 131.4, 129.5, 127.6, 48.7, 47.5, 23.3, 19.6; FTIR (neat) 3066 (vw), 2975 (m), 2937 (w), 1640 (s), 1582 (s), 1561 (vs), 1499 (s) cm⁻¹; HRMS-ES (*m/z*): [M + H]⁺ calcd. for C₁₄H₂₁N₂O 233.1654; found 233.1655.

Preparation of *N'*-(*N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninyl) *N,N*-Diisopropylformamidine from *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide and *N,N*-Diisopropylformamidine Dimethyl Acetal.

N,N-diisopropylformamide dimethyl acetal (DIF-DMA, 140 µL, 1.40 mmol, 1.5 equiv) was added to a stirring solution of *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide (120 mg, 0.45 mmol) in dichloromethane (2.5 mL) at 23 °C. After stirring for 2 h at 23 °C, the reaction mixture was filtered through Celite, and the filter cake was washed with dichloromethane (3 × 10 mL). The filtrates were combined, and the combined solution was evaporated. The residue was purified by flash-column chromatography on silica gel (80% ether-hexane) to give 161.2 mg of *N'*-(*N*^α-(*tert*-butoxycarbonyl)-L-phenylalaninyl) *N,N*-diisopropylformamidine as a white solid (95%): ¹H NMR (500 MHz, C₆D₆) δ 8.52 (s, 1 H, NCHN), 7.34-

7.23 (m, 2 H, Ar-**H**), 7.16-6.80 (m, 3 H, Ar-**H**), 5.84 (d, 1 H, $J = 7.0$ Hz, CONHCH₂), 5.08 (dt, 1 H, $J = 7.0$, 5.5 Hz, NHCHCO), 4.17 (sep, 1 H, $J = 6.8$ Hz, isopropyl CH(CH₃)₂), 3.50 (ABq [$J = 13.2$ Hz] of d [$J = 5.5$ Hz], 2H, CHCH₂Ph), 2.73 (sep, 1 H, $J = 6.8$ Hz, isopropyl CH(CH₃)₂), 1.43 (s, 10 H, *tert*-butoxy OC(CH₃)₃), 0.84-0.79 (m, 6 H, isopropyl CH(CH₃)₂), 0.60-0.54 (m, 6 H, isopropyl CH(CH₃)₂).

Preparation of *N'*-(2,3:4,6-*O*-Isopropylidene-2-keto-gulonanyl) *N,N*-Diisopropylformamidine from 2,3:4,6-*O*-Isopropylidene-2-keto-gulonamide and *N,N*-Diisopropylformamide Dimethyl Acetal. *N,N*-diisopropylformamide dimethyl acetal (DIF-DMA, 320 μ L, 1.60 mmol, 2.0 equiv) was added to a stirring solution of 2,3:4,6-*O*-isopropylidene-2-keto-gulonamide (218 mg, 0.80 mmol) in dichloromethane (2.0 mL) at 23 °C. After stirring for 26 h at 23 °C, the reaction mixture was filtered through Celite, and the filter cake was washed with dichloromethane (3×10 mL). The filtrates were combined, and the combined solution was evaporated. The residue was purified by flash-column chromatography on silica gel (50% ether-ethyl acetate) to give 305 mg of *N'*-(2,3:4,6-*O*-isopropylidene-2-keto-gulonanyl) *N,N*-diisopropylformamidine as a clear, colorless oil (99%): ¹H NMR (400 MHz, CDCl₃) δ 8.68 (s, 1 H, NCHN), 5.02 (s, 1 H, OCHCH), 4.80 (sep, 1 H, $J = 6.8$ Hz, isopropyl CH(CH₃)₂), 4.27 (d, 1 H, $J = 2.4$ Hz OCHCH), 4.18 (q, 1 H, $J = 2.4$ Hz,), 4.15-4.04 (m, 2 H, OCH₂CH), 3.69 (sep, 1 H, $J = 6.8$ Hz, isopropyl CH(CH₃)₂), 1.52 (s, 3 H, O₂-CCH₃), 1.42 (s, 6 H, O₂C(CH₃)₂), 1.33 (s, 3 H, O₂CCH₃), 1.33 (d, 6 H, $J = 6.8$ Hz, isopropyl CH(CH₃)₂), 1.26 (d, 6 H, $J = 6.8$ Hz, isopropyl CH(CH₃)₂).

Transamidation of *N'*-Benzoyl *N,N*-Diisopropylformamidine with Benzylamine to form *N*-Benzyl Benzamide using Scandium Triflate as Catalyst. Scandium triflate (25 mg, 0.05 mmol, 0.1 equiv) was added to a stirring solution of benzylamine (164 μ L, 1.50 mmol, 3 equiv) and *N'*-benzoyl *N,N*-

diisopropylformamidine (116 mg, 0.500 mmol) in ether (2.5 mL) at 23 °C. The mixture was stirred for 5 h at 23 °C, then was poured into 0.1 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether, and the ethereal layers were combined. The combined layers were washed with saturated aqueous sodium chloride solution (10 mL), the washed solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (60% ether-hexane) to afford 84.1 mg of *N*-benzyl benzamide as a white solid (95% yield, 92% over two steps).

Transamidation of *N'*-Benzoyl *N,N*-Diisopropylformamidine with Pyrrolidine to form *N*-Benzoyl Pyrrolidine using Scandium Triflate as Catalyst. Scandium triflate (25 mg, 0.05 mmol, 0.1 equiv) was added to a stirring solution of pyrrolidine (130 μ L, 1.50 mmol, 3 equiv) and *N'*-benzoyl *N,N*-diisopropylformamidine (116 mg, 0.500 mmol) in ether (2.5 mL) at 23 °C. The mixture was stirred for 5 h at 23 °C, then was poured into 0.1 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether, and the ethereal layers were combined. The combined layers were washed with saturated aqueous sodium chloride solution (10 mL), the washed solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel eluting with ether-hexane (25 $\rightarrow$ 100%) to afford 83.4 mg of *N*-benzoylpyrrolidine⁴ as a clear, colorless oil (95% yield, 92% over two steps): ¹H NMR (500 MHz, CDCl₃), δ 7.53-7.51 (m, 2 H, Ar-**H**), 7.42-7.37 (m, 3 H, Ar-**H**), 3.66 (t, *J* = 7.0 Hz, 2 H, CONCH₂), 3.43 (t, *J* = 6.5 Hz, 2 H, CONCH₂), 1.97 (p, *J* = 7.0 Hz, 2 H, NCH₂CH₂), 1.88 (p, *J* = 7.0 Hz, 2 H, NCH₂CH₂).

⁴ Alonso, E.; Ramón, D. J.; Yus, M. *Tetrahedron* **1998**, 54, 13629.

Transamidation of *N'*-(*N'*-(*tert*-Butoxycarbonyl)-L-phenylalaninyl) *N,N*-Diisopropylformamidine with Pyrrolidine using Scandium Triflate as Catalyst. Scandium triflate (14.9 mg, 0.03 mmol, 0.1 equiv) was added to a stirring solution of pyrrolidine (62 μ L, 0.75 mmol, 2.5 equiv) and *N'*-(*N'*-(*tert*-butoxycarbonyl)-L-phenylalaninyl) *N,N*-diisopropylformamidine (112 mg, 0.30 mmol) in ether (1.5 mL) at 23 $^{\circ}$ C. The mixture was stirred for 5 h at 23 $^{\circ}$ C, then was poured into 0.1 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether, and the ethereal layers were combined. The combined layers were washed with saturated aqueous sodium chloride solution (10 mL), the washed solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (35% ethyl acetate-hexane) to afford 94.0 mg of *N'*-(*tert*-butoxycarbonyl)-L-phenylalanine-*N*-pyrrolidinamide⁵ as a clear, colorless oil (98%): ^1H NMR (500 MHz, C_6D_6) δ 7.28-7.20 (m, 5 H, Ar-**H**), 5.40 (d, 1 H, $J = 8.0$ Hz, CONHCH₂), 4.59 (dd, 1 H, $J = 14.8, 8.0$ Hz, NHCHCO), 3.45-3.28 (m, 3 H, pyrrolidinyl NCH₂), 3.02-2.92 (m, 2 H, CHCH₂Ph), 2.57-2.53 (m, 1 H, pyrrolidinyl NCH), 1.76-1.53 (m, 4 H, pyrrolidinyl NCH₂CH₂), 1.42 (s, 10 H, *tert*-butoxy OC(CH₃)₃).

Transamidation of *N'*-(2,3:4,6-*O*-Isopropylidene-2-keto-gulonany) *N,N*-Diisopropylformamidine with Benzylamine to form 2,3:4,6-*O*-Isopropylidene-2-keto-*N*-benzyl-L-gulonamide using Scandium Triflate as Catalyst. Scandium triflate (11.0 mg, 0.022 mmol, 0.1 equiv) was added to a stirring solution of benzylamine (60 μ L, 0.55 mmol, 2.5 equiv) and *N'*-(2,3:4,6-*O*-isopropylidene-2-keto-gulonany) *N,N*-diisopropylformamidine (85 mg, 0.22 mmol) in ether (1.1 mL) at 23 $^{\circ}$ C. The mixture was stirred for 5 h at 23 $^{\circ}$ C, then was poured into 0.1 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated

⁵ Richmond, M. L.; Sprout, C. M.; Seto, C. T. *J. Org. Chem.* **2005**, 70, 8835.

and extracted with two 5-mL portions of ether, and the ethereal layers were combined. The combined layers were washed with saturated aqueous sodium chloride solution (10 mL), the washed solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (60% ether-hexane) to afford 78.1 mg of 2,3:4,6-*O*-isopropylidene-2-keto-*N*-benzyl-L-gulonamide as a clear, colorless gum (98%): ^1H NMR (400 MHz, CDCl_3), δ 7.44 (br s, 1 H, CONHCH₂), 7.35-7.20 (m, 5 H, Ar-H), 4.63 (s, 1 H, OCHCH), 4.56-4.46 (m, 2 H, NHCH₂Ph), 4.32 (s, 1 H, OCHCH), 4.17 (s, 1 H, OCHCH), 4.07 (ABq, 2 H, $J = 14.7$ Hz, OCH₂CH), 1.58 (s, 3 H, O₂CCH₃), 1.55 (s, 3 H, O₂CCH₃), 1.40 (s, 3 H, O₂CCH₃), 1.09 (s, 3 H, O₂CCH₃); ^{13}C NMR (100 MHz, CDCl_3) δ 167.3, 137.7, 128.6, 127.7, 127.4, 114.6, 110.6, 97.5, 88.0, 73.4, 72.5, 60.0, 43.5, 28.7, 27.1, 25.7, 18.4. FTIR (neat) 2985 (m), 1688 (s), 1521 (m) cm^{-1} ; HRMS-ES (m/z): $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{19}\text{H}_{26}\text{NO}_6$ 364.1760; found 364.1766.

Preparation of a 0.7 M Solution of Zirconium Chloride in Acetonitrile. Acetonitrile (28 mL, freshly distilled from calcium hydride) was added via syringe over 15 min to a 50-mL round-bottom flask – cooled to $-78\text{ }^\circ\text{C}$ in a Dry Ice-acetone bath – fitted with a Teflon-coated stir bar and rubber septum and containing solid zirconium chloride (4.60 g, 19.7 mmol). A white suspension formed. The reaction flask was removed from the cooling bath. The solid began dissolving as the mixture warmed (dissolution appeared to begin at $\sim 0\text{ }^\circ\text{C}$) and the mixture quickly warmed to $40\text{--}50\text{ }^\circ\text{C}$. The mixture was maintained at this temperature by gentle heating with a heat gun until the solids had dissolved ($\sim 5\text{--}10$ min). The resulting clear, colorless solution was allowed to cool, and the cooled solution was stirred at $23\text{ }^\circ\text{C}$ for 30 min. The septum-capped flask containing the clear, colorless solution of zirconium chloride in acetonitrile was stored in a dessicator containing Drierite. Over a 4-month time period, the solution yellowed, but this did not appear to have any effect in transamidation reactions.

Transamidation of Benzamide with Benzylamine to form *N*-Benzyl Benzamide using *N,N*-Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 5-mL, round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (170 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. Benzamide (20.0 mg, 0.165 mmol) and dichloromethane (1.7 mL) were added in sequence. The resulting suspension was stirred for 5 min at 23 °C, then *N,N*-diisopropylformamide dimethyl acetal (DIF-DMA, 43.0 μL, 0.21 mmol, 1.25 equiv) was added via syringe. After stirring for 2 h at 23 °C, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (30% ethyl acetate-hexane, *R_f* = 0.12 and 0.48 for benzamide and the acyl amidine, respectively). Benzylamine (36 μL, 0.33 mmol, 2.0 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 120 μL, 0.08 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (3 mL),⁶ and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 × 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (60% ether–hexane) providing 32.3 mg of *N*-benzyl benzamide⁷ as a white solid (93%): ¹H NMR (400 MHz, CDCl₃) δ 7.80-7.78 (m, 2 H, Ar-**H**), 7.52-7.28 (m, 8 H, Ar-**H**), 6.55 (br s, 1 H, CONH), 4.64 (d, *J* = 5.5 Hz, 2 H, CH₂Ph).

⁶ Addition of ether aids in the subsequent filtration of the reaction mixture.

⁷ Matsuda, F.; Itoh, S.; Hattori, N.; Yanagiya, M.; Matsumoto, T. *Tetrahedron* **1985**, *41*, 3625.

Transamidation of Benzamide with *sec*-Butylamine to form *N-sec*-Butyl Benzamide using *N,N*-Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and 5-A MS (330 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. Benzamide (40.0 mg, 0.330 mmol) and dichloromethane (3.3 mL) were added in sequence. The resulting suspension was stirred for 5 min, and *N,N*-diisopropylformamide dimethyl acetal (DIF-DMA, 84.0 μ L, 0.413 mmol, 1.25 equiv) was added via syringe. After stirring for 2 h at 23 °C, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (30% ethyl acetate-hexane, R_f = 0.12 and 0.48 for benzamide and the acyl amidine, respectively). *sec*-Butyl amine (67.0 μ L, 0.66 mmol, 2.0 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 190 μ L, 0.132 mmol, 0.4 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 $\times$ 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (80% ether-hexane) to afford 56.7 mg of *N-sec*-butyl benzamide⁸ as a white solid (97%): ¹H NMR (500 MHz, CDCl₃) δ 7.76-7.74 (m, 2 H, Ar-**H**), 7.46-7.39 (m, 3 H, Ar-**H**), 6.08 (br s, 1 H, CONH**H**), 4.10 (sep, J = 7 Hz, 1 H, NHCH(CH₃)CH₂), 1.55 (sextet, J = 7 Hz, 2 H, CHCH₂-CH₃), 1.21 (d, J = 6.6 Hz, 3 H, CHCH₃), 0.95 (t, J = 7.3 Hz, 3 H, CH₂CH₃).

⁸ Tillack, A.; Rudloff, I.; Beller, M. *Eur. J. Org. Chem.* **2001**, 523.

Transamidation of Benzamide with *N*-Benzylmethylamine to form *N*-Benzyl-*N*-methyl Benzamide using *N,N*-Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and 5-Å MS (330 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. Benzamide (40.0 mg, 0.330 mmol) and dichloromethane (3.3 mL) were added in sequence. The resulting suspension was stirred for 5 min, and *N,N*-diisopropylformamide dimethyl acetal (DIF-DMA, 84.0 µL, 0.413 mmol, 1.25 equiv) was added via syringe. After stirring for 2 h at 23 °C, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (30% ethyl acetate-hexane, R_f = 0.12 and 0.48 for benzamide and the acyl amidine, respectively). *N*-Benzylmethylamine (106 µL, 0.825 mmol, 2.5 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 235 µL, 0.165 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 × 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (50% ether-hexane) to afford 70.0 mg of *N*-benzyl-*N*-methyl benzamide⁹ as a clear, colorless oil (94%): ¹H NMR (400 MHz, CDCl₃) δ 7.47-7.30 (m, 9 H, Ar-**H**), 7.19-7.16 (m, 1 H, Ar-**H**), 4.77 (br s, 0.5 × 2 H, NCH₂Ph), 4.52 (br s, 0.5 × 2 H, NCH₂Ph), 3.03 (br s, 0.5 × 3 H, NCH₃), 2.87 (br s, 0.5 × 3 H, NCH₃).

⁹ Leggio, A.; Liguori, A.; Napoli, A.; Siciliano, C.; Sindona, G. *J. Org. Chem.* **2001**, 66, 2246.

Transamidation of Benzamide with *N,O*-Dimethylhydroxylamine to form *N*-Methoxy-*N*-methyl Benzamide using *N,N*-Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-Å MS (350 mg) and evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. Benzamide (40.0 mg, 0.330 mmol) and dichloromethane (3.3 mL) were added in sequence. The resulting suspension was stirred for 5 min, and *N,N*-diisopropylformamide dimethyl acetal (DIF-DMA, 84.0 µL, 0.413 mmol, 1.25 equiv) was added via syringe. After stirring for 2 h at 23 °C, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (30% ethyl acetate-hexane, R_f = 0.12 and 0.48 for benzamide and the acyl amidine, respectively). *N,O*-Dimethylhydroxylamine (72 µL, 0.99 mmol, 3.0 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 235 µL, 0.165 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 × 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (50% ether-hexane) to afford 47.4 mg of *N*-methoxy-*N*-methyl benzamide¹⁰ as a clear, colorless oil (87%): ¹H NMR (500 MHz, CDCl₃) δ 7.65 (app d, J = 7.8 Hz, 2 H, Ar-**H**), 7.45-7.37 (m, 3 H, Ar-**H**), 3.54 (s, 3 H, NCH₃), 3.34 (s, 3 H, OCH₃).

¹⁰ Leggio, A.; Liguori, A.; Napoli, A.; Siciliano, C.; Sindona, G. *J. Org. Chem.* **2001**, 66, 2246.

Transamidation of Benzamide with Morpholine to form *N*-Benzoyl Morpholine using *N,N*-Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and 5-Å MS (330 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. Benzamide (40.0 mg, 0.330 mmol) and dichloromethane (3.3 mL) were added in sequence. The resulting suspension was stirred for 5 min, and *N,N*-Diisopropylformamide dimethyl acetal (DIF-DMA, 84.0 µL, 0.413 mmol, 1.25 equiv) was added via syringe. After stirring for 2 h at 23 °C, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (30% ethyl acetate-hexane, R_f = 0.12 and 0.48 for benzamide and the acyl amidine, respectively). Morpholine (58.0 µL, 0.66 mmol) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 190 µL, 0.132 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 × 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (80% ether-hexane) to afford 54.5 mg of *N*-benzoyl morpholine¹¹ as a white solid (87%): ¹H NMR (500 MHz, CDCl₃) δ 7.41-7.38 (m, 5 H, Ar-**H**), 3.43-3.75 (m, 8 H, **CH**₂).

¹¹ Kondo, K.; Sekimoto, E.; Nakao, J.; Murakami, Y. *Tetrahedron* **2000**, 56, 5843.

Transamidation of Pivalamide with Benzylamine to form *N*-Benzyl Pivalamide using *N,N*-Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (400 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. Pivalamide (40.0 mg, 0.395 mmol) and dichloromethane (4.0 mL) were added in sequence. The resulting suspension was stirred for 5 min, and *N,N*-Diisopropylformamide dimethyl acetal (DIF-DMA, 84.0 μ L, 0.413 mmol, 1.25 equiv) was added via syringe. The mixture was stirred for 24 h at 23 °C.¹² Benzylamine (86 μ L, 0.79 mmol, 2 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 226 μ L, 0.158 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 $\times$ 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (50% ether-hexane) to afford 68.5 mg of *N*-benzyl pivalamide¹³ as a white solid (91%): ¹H NMR (500 MHz, CDCl₃) δ 7.36-7.27 (m, 5 H, Ar-**H**), 5.89 (br s, 1 H, **NH**), 4.44 (d, *J* = 5.6 Hz, 2 H, **NHCH**₂Ph), 1.23 (s, 9 H **CH**(**CH**₃)₃).

¹² The reaction was stirred for 24 h to ensure complete formation of the acyl amidine.

¹³ Kondo, K.; Iida, T.; Fujita, H.; Suzuki, T.; Yamaguchi, K.; Murakami, Y. *Tetrahedron* **2000**, 56, 8883.

Transamidation of *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide with *N*-Benzylmethylamine to form *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalanine-*N*-benzyl-*N*-methylamide using *N,N*-Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter.

A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (190 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide (50.0 mg, 0.189 mmol) and dichloromethane (1.9 mL) were added in sequence. The resulting suspension was stirred for 5 min, and *N,N*-diisopropylformamide dimethyl acetal (DIF-DMA, 48.0 μL, 0.234 mmol, 1.25 equiv) was added via syringe. After stirring for 4 h at 23 °C, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (50% ethyl acetate-hexane, *R*_f = 0.18 and 0.42 for the primary amide and the acyl amidine, respectively). *N*-Benzylmethylamine (61 μL, 0.47 mmol, 2.5 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 135 μL, 0.09 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 × 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (40% ether-hexane) to afford 65.3 mg of *N*^α-(*tert*-butoxycarbonyl)-L-phenylalanine-*N*-benzyl-*N*-methylamide¹⁴ as a white solid (94%): ¹H NMR (500 MHz, CDCl₃) δ 7.28-6.98 (m, 10 H, Ar-**H**), 5.46 (d, *J* = 8.8 Hz, 0.75 × 1 H, Boc**NH**), 5.39 (d, *J* = 8.8 Hz, 0.25 × 1 H, Boc**NH**), 4.88 (dt, *J* = 7.3 Hz, 1 H, NH**CH**CO), 4.55 (A of ABX, *J* = 14.6

¹⁴ Hagiwara, D.; Miyake, H.; Morimoto, H.; Murai, M.; Fujii, T.; Matsuo, M. *J. Med. Chem.* **1992**, 35, 3184.

Hz, 0.75×1 H, CHCH_2Ph), 4.44 (B of ABX, $J = 14.6$ Hz, 0.75×1 H, CHCH_2Ph), 4.36 (A of ABX, $J = 16.6$ Hz, 0.25×1 H, CHCH_2Ph), 4.12 (B of ABX, $J = 16.6$ Hz, 0.25×1 H, CHCH_2Ph), 3.10 (d, $J = 7.3$ Hz, 2H, CH_2Ph), 2.83 (s, 0.25×3 H, NCH_3), 2.62 (s, 0.75×3 H, NCH_3), 1.43 (s, 0.75×9 H, $\text{CH}(\text{CH}_3)_4$), 1.39 (s, 0.25×9 H, $\text{CH}(\text{CH}_3)_3$).

Transamidation of N^α -(*tert*-Butoxycarbonyl)-L-phenylalaninamide with L-Phenylalanine-*tert*-butyl ester to form Boc-L-Phe-L-Phe-Ot-Bu using N,N -Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (190 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. N^α -(*tert*-Butoxycarbonyl)-L-phenylalaninamide (50.0 mg, 0.189 mmol) and dichloromethane (1.9 mL) were added in sequence, The resulting suspension was stirred for 5 min, and N,N -diisopropylformamide dimethyl acetal (DIF-DMA, 48.0 μL , 0.234 mmol, 1.25 equiv) was added via syringe. After stirring for 4 h at 23 °C, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (50% ethyl acetate-hexane, $R_f = 0.18$ and 0.42 for the primary amide and the acyl amidine, respectively). L-Phenylalanine-*tert*-butyl ester (83 mg, 0.38 mmol, 2.0 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 135 μL , 0.09 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3×3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal

solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (40% ether-hexane) to afford 81.5 mg of Boc-L-Phe-L-Phe-O*t*-Bu as a white solid (92%): ^1H NMR (400 MHz, CDCl_3) δ 7.29-7.18 (m, 8 H, Ar-**H**), 7.06-7.04 (m, 2 H, Ar-**H**), 6.34 (br d, $J = 4.8$ Hz, 1 H, CONH**CH**), 4.98 (br s, 1 H, BocNH**H**), 4.64 (app d, $J = 5.6$ Hz, 1 H, NH**CH**CO), 4.35 (br s, 1 H, NH**CH**CO), 3.05-3.01 (m, 4 H, C**CH**₂Ph), 1.40 (s, 9 H, CH(**CH**₃)₃), 1.35 (s, 9 H, CH(**CH**₃)₃); ^{13}C NMR (100 MHz, CDCl_3) δ 170.5, 169.9, 155.2, 136.5, 136.0, 129.5, 129.3, 128.6, 128.3, 126.9, 82.3, 80.6, 55.7, 53.6, 38.3, 38.1, 28.2, 27.8; IR (neat) 3421 (vw), 3323 (vw), 2983 (w), 1721 (m), 1671 (m), 1497 (m), 1368 (m), 1156 (s), 905 (vs) cm^{-1} ; HRMS-ES (m/z): $[\text{M} + \text{H}]^+$ calcd. for $\text{C}_{27}\text{H}_{37}\text{N}_2\text{O}_5$ 469.2602; found 469.2681.

Transamidation of (*S*)-Mandelamide with Benzylamine to form *N*-Benzyl (*S*)-Mandelamide using *N,N*-Diisopropylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and 5-Å MS (330 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. (*S*)-Mandelamide (50.0 mg, 0.331 mmol) and dichloromethane (3.3 mL) were added in sequence. The resulting suspension was stirred for 5 min, *N,N*-diisopropylformamide dimethyl acetal (DIFDMA, 83.0 μL , 0.414 mmol, 1.25 equiv) was added via syringe. After stirring for 2 h at 23 °C, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (80% ethyl acetate-hexane, $R_f = 0.27$ and 0.64 for (*S*)-mandelamide and the acyl amidine, respectively). Benzylamine (90.0 μL , 0.827 mmol, 2.5 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 235 μL , 0.165 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3×3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was

separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (70% ether-hexane) to afford 61.0 mg of *N*-benzyl (*S*)-mandelamide¹⁵ as a white solid (76%): ¹H NMR (500 MHz, CDCl₃) δ 7.39-7.23 (m, 8 H, Ar-**H**), 7.18-7.16 (m, 2 H, Ar-**H**), 6.88 (br s, 1 H, CONH), 5.01 (s, 1 H, HOCH**CO**), 4.38 (dd, *J* = 5.9, 2.9 Hz, 2 H, CH₂Ph).

Transamidation of *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide with Benzylamine to form *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalanine-*N*-benzylamide using *N,N*-Dimethylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (190 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide (50.0 mg, 0.189 mmol) and dichloromethane (1.9 mL) were added in sequence. After stirring for 5 min at 23 °C, *N,N*-dimethylformamide dimethyl acetal (DMF-DMA, 33.0 μL, 0.25 mmol, 1.3 equiv) was added via syringe, and the resulting mixture was heated to reflux. After stirring at reflux for 4 h, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (80% ethyl acetate-hexane, *R_f* = 0.44 and 0.38 for the primary amide and the acyl amidine, respectively), and the mixture was allowed to cool to 23 °C. Benzylamine (41 μL, 0.38 mmol, 2.0 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 135 μL, 0.095 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 × 3 mL), the filtrates were combined, and the combined solution was

¹⁵ Kolasa, T.; Miller, M. J. *J. Org. Chem.* **1987**, 52, 4978.

washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (60% ether–hexane) to afford 63.0 mg of *N*^α-(*tert*-butoxycarbonyl)-L-phenylalanine-*N*-benzylamide¹⁶ as a white solid (94%): ¹H NMR (400 MHz, CDCl₃) δ 7.28-7.17 (m, 8H, Ar-**H**), 7.08 (br s, 2 H, Ar-**H**), 6.31 (br t, *J* = 5.6 Hz, 1 H, CONHCH₂), 4.38-3.28 (m, 3 H, NHCHCO and NHCH₂Ph), 3.07 (d, *J* = 6.4 Hz, 2 H, CHCH₂Ph), 1.38 (s, 9 H, CH(CH₃)₃).

Transamidation of *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide with *N*-Benzylmethylamine to form *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalanine-*N*-benzyl-*N*-methylamide using *N,N*-Dimethylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (220 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide (52.4 mg, 0.198 mmol) and dichloromethane (2.0 mL) were added in sequence. After stirring for 5 min at 23 °C, and *N,N*-dimethylformamide dimethyl acetal (DMF-DMA, 34.0 μL, 0.256 mmol, 1.3 equiv) was added via syringe, and the resulting mixture was heated to reflux. After stirring at reflux for 4 h, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (80% ethyl acetate-hexane, *R*_f = 0.44 and 0.38 for the primary amide and the acyl amidine, respectively), and the mixture was allowed to cool to 23 °C. *N*-Benzylmethylamine (76 μL, 0.59 mmol, 3.0 equiv) was added to the reaction mixture at this point

¹⁶ Fujita, Y.; Tsuda, Y.; Li, T.; Motoyama, T.; Takahashi, M.; Shimizu, Y.; Yokoi, T.; Sasaki, Y.; Ambo, A.; Kita, A.; Jinsmaa, Y.; Bryant, S. D.; Lazarus, L. H.; Okada, Y. *J. Med. Chem.* **2004**, *47*, 3591.

followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 140 μ L, 0.098 mmol, 0.5 equiv). After stirring for 1 h at 23 $^{\circ}$ C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 $\times$ 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (30% ethyl acetate-hexane) to afford 66.1 mg of *N $^{\alpha}$ -(tert-butoxycarbonyl)-L-phenylalanine-N-benzyl-N-methylamide*¹¹ as a clear oil (91%).

Transamidation of *N $^{\alpha}$ -(tert-Butoxycarbonyl)-L-phenylalaninamide* with *N,O*-Dimethylhydroxylamine to form *N $^{\alpha}$ -(tert-Butoxycarbonyl)-L-phenylalanine-N-methoxy-N-methylamide* using *N,N*-Dimethylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (215 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 $^{\circ}$ C, and an argon atmosphere was introduced. *N $^{\alpha}$ -(tert-Butoxycarbonyl)-L-phenylalaninamide* (55.0 mg, 0.208 mmol) and dichloromethane (2.1 mL) were added in sequence. After stirring for 5 min at 23 $^{\circ}$ C, *N,N*-dimethylformamide dimethyl acetal (DMF-DMA, 36.0 μ L, 0.271 mmol, 1.3 equiv) was added via syringe, and the resulting mixture was heated to reflux. After stirring at reflux for 4 h, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (80% ethyl acetate-hexane, R_f = 0.44 and 0.38 for the primary amide and the acyl amidine, respectively), and the mixture was allowed to cool to 23 $^{\circ}$ C. *N,O*-Dimethylhydroxylamine (46 μ L, 0.62 mmol, 3.0 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 150

μL , 0.104 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 $\times$ 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (30% ethyl acetate-hexane) to afford 55.6 mg of *N* ^{α} -(*tert*-butoxycarbonyl)-L-phenylalanine-*N*-methoxy-*N*-methylamide¹⁷ as a clear oil (87%): ¹H NMR (500 MHz, CDCl₃) δ 7.29-7.16 (m, 5 H, Ar-**H**), 5.16 (br d, *J* = 7.3 Hz, 1 H, BocNHCH), 4.95 (br d, *J* = 5.9 Hz, 1 H, NHCHCO), 3.65 (s, 3 H, NCH₃), 3.16 (s, 3 H, OCH₃), 3.04 (dd, *J* = 13.2, 5.9 Hz, 1 H, CHCH₂Ph), 2.87 (dd, *J* = 13.0, 7.3 Hz, 1 H, CHCH₂Ph), 1.38 (s, 9 H, CH(CH₃)₃).

Transamidation of *N* ^{α} -(*tert*-Butoxycarbonyl)-L-phenylalaninamide with Morpholine to form *N* ^{α} -(*tert*-Butoxycarbonyl)-L-phenylalanine-*N*-morpholinamide using *N,N*-Dimethylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (210 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. *N* ^{α} -(*tert*-Butoxycarbonyl)-L-phenylalaninamide (52.0 mg, 0.196 mmol) and dichloromethane (2.0 mL) were added in sequence. After stirring for 5 min at 23 °C, *N,N*-dimethylformamide dimethyl acetal (DMF-DMA, 34.0 μL , 0.256 mmol, 1.3 equiv) was added via syringe, and the resulting mixture was heated to reflux. After stirring at reflux for 4 h, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (80% ethyl acetate-hexane, *R_f* = 0.44 and 0.38 for the primary amide and the acyl

¹⁷ Sawamura, M.; Nakayama, Y.; Kato, T.; Ito, Y. *J. Org. Chem.* **1995**, 60, 1727.

amidine, respectively), and the mixture was allowed to cool to 23 °C. Morpholine (43 µL, 0.49 mmol, 2.5 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 140 µL, 0.098 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 × 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (40% ethyl acetate-hexane) to afford 56.7 mg of *N*^α-(*tert*-butoxycarbonyl)-L-phenylalanine-*N*-morpholinamide¹⁸ as a clear oil (87%): ¹H NMR (500 MHz, CDCl₃) δ 7.29-7.18 (m, 5 H, Ar-**H**), 5.44 (d, J = 8.8 Hz, 1 H, BocNHCH), 4.78 (dt, J = 8.8, 5.9 Hz, 1 H, NHCHCO), 2.84-3.03 (m, 4 H, morpholine C-**H**), 3.24-3.27 (m, 1 H, morpholine C-**H**), 3.38-3.58 (m, 5 H, morpholine C-**H** and CHCH₂Ph), 1.41 (s, 9 H, CH(CH₃)₃).

Transamidation of *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide with Aniline to form *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalanine-*N*-phenylamide using *N,N*-Dimethylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and of crushed 5-A MS (225 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide (55.2 mg, 0.208 mmol) and dichloromethane (2.1 mL) were added in sequence. After stirring for 5 min at 23 °C, *N,N*-

¹⁸ (a) Katritzky, A. R.; He, H.-Y.; Jiang, R.; Long, Q. *Tetrahedron: Asymmetry* **2001**, 12, 2427; Douat, C.; Heitz, A.; Martinez, J.; Fehrentz, J.-A. *Tetrahedron Lett.* **2000**, 41, 37.

dimethylformamide dimethyl acetal (DMF-DMA, 36.0 μ L, 0.271 mmol, 1.3 equiv) was added via syringe, and the resulting mixture was heated to reflux. After stirring at reflux for 4 h, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (80% ethyl acetate-hexane, R_f = 0.44 and 0.38 for the primary amide and the acyl amidine, respectively), and the mixture was allowed to cool to 23 $^{\circ}$ C. Aniline (38 μ L, 0.42 mmol, 2.0 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 200 μ L, 0.141 mmol, 0.5 equiv). After stirring for 1 h at 23 $^{\circ}$ C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3×3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (30% ethyl acetate-hexane) to afford 56.0 mg of *N* $^{\alpha}$ -(*tert*-butoxycarbonyl)-L-phenylalanine-*N*-phenylamide¹⁹ as a white solid (79%): ^1H NMR (500 MHz, CDCl_3) δ 8.09 (br s, CONHPh), 7.37-7.25 (m, 9 H, Ar-H), 7.14-7.06 (m, 1 H, Ar-H), 5.35 (br s, 1 H, BocNHCH), 4.56 (br s, 1 H, NHCHCO), 3.19-3.07 (m, 2 H, CHCH₂Ph), 1.41 (s, 9 H, CH(CH₃)₃).

Transamidation of *N* $^{\alpha}$ -(*tert*-Butoxycarbonyl)-L-phenylalaninamide with L-Phenylalanine Methyl Ester to form Boc-L-Phe-L-Phe-OMe using *N,N*-Dimethylformamide Dimethyl Acetal as Activator and Zirconium Chloride as Promoter. A 10-mL round-bottom flask containing a Teflon-coated stir bar and crushed 5-A MS (210 mg) was evacuated (0.1 mmHg), and a good flame was applied for ~2 min to dry

¹⁹ Obrecht, D.; Bohdal, U.; Broger, C.; Bur, D.; Lehmann, C.; Ruffieux, R.; Schönholzer, P.; Spiegler, C.; Müller, K. *Helv. Chim. Acta* **1995**, 78, 563.

the sieves. The flame was removed, the flask was allowed to cool to 23 °C, and an argon atmosphere was introduced. *N*^α-(*tert*-Butoxycarbonyl)-L-phenylalaninamide (50.0 mg, 0.189 mmol) and dichloromethane (1.9 mL) were added in sequence. After stirring for 5 min at 23 °C, *N,N*-dimethylformamide dimethyl acetal (DMF-DMA, 33.0 μL, 0.246 mmol, 1.3 equiv) was added via syringe, and the resulting mixture was heated to reflux. After stirring at reflux for 4 h, analysis of the reaction mixture by thin-layer chromatography showed that complete conversion of the starting material to the acyl amidine had occurred (80% ethyl acetate-hexane, *R*_f = 0.44 and 0.38 for the primary amide and the acyl amidine, respectively), and the mixture was allowed to cool to 23 °C. L-Phenylalanine methyl ester (68 mg, 0.38 mmol, 2.0 equiv) was added to the reaction mixture at this point followed 3 min later by a solution of zirconium chloride in acetonitrile (0.7 M, 135 μL, 0.095 mmol, 0.5 equiv). After stirring for 1 h at 23 °C, the reaction mixture was diluted with ether (2 mL), and the diluted solution was filtered through a pad of Celite. The Celite pad was washed with ether (3 × 3 mL), the filtrates were combined, and the combined solution was washed with 0.2 N aqueous sulfuric acid solution (10 mL). The aqueous layer was separated and extracted with two 5-mL portions of ether. The ethereal layers were combined, and the combined layers were washed sequentially with saturated aqueous sodium bicarbonate solution (5 mL) and saturated aqueous sodium chloride solution (10 mL). After washing, the ethereal solution was dried over anhydrous magnesium sulfate, and the dried solution was concentrated. The residue was purified by flash column chromatography on silica gel (30% ethyl acetate-hexane) to afford 76.6 mg of Boc-L-Phe-L-Phe-OMe²⁰ as a white solid (95%): ¹H NMR (400 MHz, CDCl₃) δ 7.30-7.18 (m, 8 H, Ar-**H**), 6.99 (m, 2 H, Ar-**H**), 6.30 (d, *J* = 7.0 Hz, 1 H, CON**H**C), 4.95 (s, 1 H, BocN**H**C), 4.78 (app d, *J* = 6.6 Hz, 1 H, NH**C**HCO), 4.34-4.33 (m, 1 H, BocNH**C**HCO), 3.66 (s, 3 H, COO**CH**₃), 3.06-3.02 (m, 4 H, **CH****CH**₂Ph), 1.40 (s, 9 H, **CH**(**CH**₃)₃).

²⁰ Rzepecki, P.; Gallmeier, H.; Geib, N.; Cernovska, K.; König, B.; Schrader, T. *J. Org. Chem.* **2004**, 69, 5168.