Efforts Towards the Total Synthesis of (–)-Kendomycin

David R. Williams and Khalida Shamim*

Department of Chemistry

Indiana University

Bloomington

Indiana IN 47405

Supporting Information

The experimental details and characterization data along with ^1H NMR and ^{13}C NMR spectra for compounds **2**, **7**, **10**, **11**, **13**, **14**, **15**, **20**, **21**, **22** and **23** are provided.

7-Benzyloxy-4S,6S-dimethyl-1-heptyne (7).

Aldehyde (50 mg, 213 μmol) was combined with K_2CO_3 (58 mg, 426 μmol) and MeOH (3.5 mL) at room temperature. After 5 min, dimethyl-1-diazo-2-oxopropylphosphonate (49 mg, 256 μmol) was added and the reaction mixture was stirred for a period of 12 h. The reaction mixture was then diluted with pentane (10 mL) and brine (10 mL) resulting in a white precipitate in the aqueous layer while the organic layer was clear and colorless. The aqueous layer was separated and extracted with pentane (2 x 15 mL). The organic phases were combined, washed with saturated aqueous NaHCO_3 (2 x 30 mL), dried over MgSO_4 , filtered and concentrated to provide a yellow oil. The product was purified by flash column chromatography (15 g SiO_2 , 5% EtOAc in hexanes) to afford 47 mg (96%) of alkyne **7** as a clear colorless oil: R_f 0.5 (10% EtOAc in hexanes); $[\alpha]_D^{21} -9.4$ (c 1.19, CHCl_3); IR (neat) 3308, 2958, 2925, 2853, 1454, 1379, 1362, 1097, 1029 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.37 – 7.29 (m, 5H), 4.52 (s, 2H), 3.33 (A of ABX, $J_{AB} = 9.0$ Hz, $J_{AX} = 5.83$ Hz, 1H), 3.27 (B of ABX, $J_{AB} = 9.0$ Hz, $J_{BX} = 6.8$ Hz, 1H), 2.21 – 2.06 (m, 2H), 1.98 – 1.97 (m, 1H), 1.92 – 1.75 (m, 2H), 1.36 – 1.20 (m, 2H), 0.99 (d, $J = 6.6$ Hz, 3H), 0.95 (d, $J = 6.6$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ 139.0, 128.6, 127.8, 127.7, 83.6, 76.7, 73.3, 69.4, 40.4, 31.2, 30.1, 26.9, 19.3, 17.2; MS (CI, CH_4) m/e (relative intensity) 181 (10), 157 (14), 117 (13), 109 (16), 99 (48), 91 (100); HRMS m/e calcd. for $\text{C}_{16}\text{H}_{23}\text{O}$ ($\text{M}+\text{H}$) $^+$ 231.1745 found 231.1749.

(R)-3-((E)-(3S,7S,9S)-10-Benzyloxy-3,5,7,9-tetramethyl-dec-4-en-1-yn-1-yl)-4-phenyl-oxazolidin-2-one (10).

In a schlenk flask a suspension of Cp_2ZrCl_2 (28 mg, 100 μmol) in dry 1,2-dichloro ethane (1.5 mL) was treated at 0 °C with a 2M solution of Me_3Al in toluene (0.7 mL) followed by addition of a solution of 1-alkyne **7** (108 mg, 470 μmol) in 1,2-dichloro ethane (0.5 mL). The reaction mixture was stirred at room temperature for 3 h, the solvent was then removed *in vacuo*, and dry ether (2 mL) was added. The solution of alkenyl alane was added at –23 °C to a mixture of anhydrous copper cyanide (45 mg, 510 μmol) in THF(4 mL) and a 0.5 M solution of 1-hexynyllithium in 5:1 THF/hexane (2.2 mL). After stirring at –23 °C for 5 min, a solution of oxazolidinone **9** (92 mg, 400 μmol) in THF (1 mL) was added dropwise. Stirring at –23 °C was continued for an additional 20 min and the mixture was then quenched by cannulation into a cold solution of saturated $\text{NH}_4\text{Cl}/\text{NH}_4\text{OH}$ (20 mL (9:1)) with subsequent stirring overnight. The mixture was then extracted three times with ether (3 x 50 mL). The combined organic phases were dried (MgSO_4), filtered and concentrated to obtain an oil which contained a 7:1 ratio of product diastereomers. The major product was purified by flash column chromatography (30 g SiO_2 , 30% Et_2O in hexanes) to afford 157 mg (70%) of pure conjugate addition product **10** as a clear colorless oil: R_f 0.3 (25% EtOAc in hexanes); $[\alpha]_D^{21}$ –29.9 (c 0.87, CHCl_3); IR (neat) 3066, 3031, 2957, 2923, 2870, 2845, 1782, 1707, 1443, 1383, 1315, 1200 cm^{-1} ;

^1H NMR (400 MHz, CDCl_3) δ 7.40 – 7.28 (m, 10 H), 5.42 (dd, $J = 8.6, 3.5$ Hz, 1H), 4.93 (d, $J = 8.3$ Hz, 1H), 4.68 (t, $J = 8.8$ Hz, 1H), 4.51 (s, 2H), 4.27 (dd, $J = 8.9, 3.5$ Hz, 1H), 3.31 (A of ABX, $J_{\text{AB}} = 9.0$ Hz, $J_{\text{AX}} = 5.9$ Hz, 1H), 3.22 (B of ABX, $J_{\text{AB}} = 9.0$ Hz, $J_{\text{BX}} = 7.1$ Hz, 1H), 3.02 – 2.92 (m, 2H), 2.88 – 2.83 (m, 1H), 1.90 – 1.82 (m, 2H), 1.74 – 1.60 (m, 2H), 1.50 (s, 3H), 1.14 – 0.99 (m, 2H), 0.95 (d, $J = 6.3$ Hz, 3H), 0.90 (d, $J = 6.7$ Hz, 3H), 0.71 (d, $J = 6.3$ Hz, 3H); ^{13}C NMR (101MHz, CDCl_3) δ 171.7, 153.7, 139.1, 138.8, 133.6, 130.5, 129.1, 128.6, 128.3, 127.5, 127.4, 125.9, 72.9, 69.8, 57.6, 48.4, 42.6, 40.7, 30.9, 29.7, 29.2, 27.7, 20.9, 19.2, 16.9, 15.9; MS (CI, CH_4) m/e (relative intensity) 386 (66), 370 (27), 232 (33), 223 (38), 205 (40), 164 (36), 123 (51), 99 (100), 91 (61), 83 (28); HRMS m/e calcd. for $\text{C}_{30}\text{H}_{40}\text{NO}_4$ ($\text{M}+\text{H}$) $^+$ 478.2957 found 478.2971.

(E)-(4S,8S,10S)-11-Benzyloxy-4,6,8,10-tetramethyl-undec-5-enal (11).

A solution of starting nitrile (30 mg, 92 μmol) in dry toluene (2.3 mL) was cooled to -78°C and treated with DIBAL (0.1 mL, 1M in hexanes). After 20 min, H_2O (0.01 mL) was added and the solution was stirred at room temperature for 0.5 h. Then 4N NaOH (0.01 mL) was added dropwise. After 1 h, MgSO_4 was added and stirring was continued for another 0.5 h. The reaction mixture was then filtered and concentrated to provide a colorless oil. The crude product was purified by flash column chromatography (10 g SiO_2 , 8% EtOAc in hexanes) to afford 27 mg (89%) of aldehyde **11** as a clear colorless oil: R_f 0.3 (10% EtOAc in hexanes): ^1H NMR (400 MHz, CDCl_3) δ 9.70 (t, $J = 1.8$ Hz,

1H), 7.32 – 7.31 (m, 4H), 7.27 – 7.25 (m, 1H), 4.80 (d, $J = 9.5$ Hz, 1H), 4.48 (s, 2H), 3.28 (A of ABX, $J_{AB} = 9.0$ Hz, $J_{AX} = 5.9$ Hz, 1H), 3.20 (B of ABX, $J_{AB} = 9.0$ Hz, $J_{BX} = 6.9$ Hz, 1H), 2.38 – 2.22 (m, 3H), 1.93 – 1.81 (m, 2H), 1.77 – 1.61 (m, 3H), 1.50 (s, 3H), 1.49 – 1.40 (m, 1H), 1.15 – 0.99 (m, 2H), 0.93 (d, $J = 6.6$ Hz, 3H), 0.87 (d, $J = 6.6$ Hz, 3H), 0.76 (d, $J = 6.4$ Hz, 3H). ^{13}C NMR (101MHz, CDCl_3) δ 202.9, 138.7, 133.8, 131.4, 128.2, 127.4, 127.3, 76.6, 72.9, 48.6, 42.2, 40.8, 31.9, 30.8, 29.7, 27.6, 21.3, 19.2, 16.8, 16.1.

(E)-(2S,4S,5R,8S,12S,14S)-15-Benzyloxy-5-hydroxy-1-(4-methoxy-benzyloxy)-2,4,8,10,12,14-hexamethyl-pentadec-9-en-3-one (13).

Preparation of (–)-(Ipc)₂BOTf :- Trifluoromethane sulfonic acid (0.3 mL, 2.96 mmol) was added to a suspension of (–)-(Ipc)₂BH (0.85 g, 2.96 mmol) in dry hexanes (2 mL) at 0 °C. After 15 min, the mixture was warmed to room temperature and stirred gently for 2 h. Hydrogen was evolved in the process and the evolution subsided after the 2 h interval at room temperature. The upper colorless solution was transferred via cannula into a dry flask. This stock solution of boron triflate (assuming 60% conversion) was used in the aldol reactions within 24 h.

To a stirred solution of (–)-(Ipc)₂BOTf (0.17 mL, 0.8 M) in CH₂Cl₂ (0.5 mL) at 0 °C was added Et₃N (0.04 mL, 272 μmol), followed by ketone **12** (21 mg, 91 μmol) in 0.5 mL of CH₂Cl₂. After stirring for 3 h at 0 °C, aldehyde **11** (15 mg, 45 μmol) in CH₂Cl₂ (0.5 mL) was added, and the reaction was stirred at that temperature for another 1 h. The reaction was then maintained with stirring at –15 °C for a period of 12 h. The mixture was then partitioned between Et₂O (3x 25 mL) and a pH 7 buffer solution (20 mL) and the combined organic extracts were concentrated *in vacuo*. The residue was suspended in MeOH (1 mL) and pH7 buffer (0.3 mL) and cooled to 0 °C. H₂O₂ solution (0.5 mL, 30% aqueous solution) was added dropwise and stirring continued at room temperature for 1 to 2 h. The mixture was then poured into distilled water (10 mL) and extracted with CH₂Cl₂ (3x 10 mL). The combined organic extracts were washed sequentially with saturated aqueous NaHCO₃ solution (15 mL) and brine (15 mL), and then dried (MgSO₄) and concentrated *in vacuo* to afford an oily residue which was a 10:1 ratio of product diastereomers. The major compound was then separated by performing flash column chromatography (30 g SiO₂, 12% EtOAc in hexanes) to afford 19 mg (72%) of pure aldol product **13** as a clear colorless oil: R_f 0.08 (10% EtOAc in hexanes); [α]_D²¹ +3.1 (c 0.90, CHCl₃); IR (neat) 3474, 2954, 2925, 2866, 2359, 2328, 1704, 1610, 1514, 1456, 1374, 1248, 1097, 1036 cm^{–1}; ¹H NMR (400 MHz, CDCl₃) δ 7.37 – 7.26 (m, 5H), 7.22 (d, *J* = 8.6 Hz, 2H), 6.88 (d, *J* = 8.6 Hz, 2H), 4.87 (d, *J* = 9.3 Hz, 1H), 4.52 (s, 2H), 4.44 (A of AB, *J*_{AB} = 11.5 Hz, 1H), 4.40 (B of AB, *J*_{AB} = 11.5 Hz, 1H), 3.88 – 3.77 (m, 4H), 3.64 (A of ABX, *J*_{AB} = 8.8 Hz, *J*_{AX} = 8.8 Hz, 1H), 3.42 (B of ABX, *J*_{AB} = 8.8 Hz, *J*_{BX} = 5.1 Hz, 1H), 3.32 (A of ABX *J*_{AB} = 8.9 Hz, *J*_{AX} = 5.6 Hz, 1H), 3.23 (B of ABX, *J*_{AB} = 8.9 Hz, *J*_{BX} = 7.2 Hz, 1H), 3.13 – 3.04 (m, 1H), 2.87 (d, *J* = 3.0 Hz, 1H), 2.66 (dq, *J* = 7.2, 2.9

Hz, 1H), 2.38 – 2.28 (m, 1H), 1.98 – 1.82 (m, 2H), 1.80 – 1.62 (m, 2H), 1.60 – 1.48 (m, 4H), 1.40 – 1.20 (m, 3H), 1.20 – 1.00 (m, 8H), 0.98 – 0.80 (m, 6H), 0.78 – 0.70 (d, J = 6.4 Hz, 3H); ^{13}C NMR (101MHz, CDCl_3) δ 218.5, 159.2, 138.8, 132.6, 132.5, 129.9, 129.1, 128.2, 127.4, 127.3, 113.7, 76.6, 73.0, 72.9, 72.0, 71.3, 55.2, 50.0, 48.6, 45.1, 40.9, 33.9, 32.2, 32.1, 30.9, 27.7, 21.4, 19.2, 16.9, 16.0, 13.8, 8.8; MS (CI, CH_4) m/e (relative intensity) 427 (7), 241 (9), 199 (6), 122 (9), 121 (100), 99 (6); HRMS m/e calcd. for $\text{C}_{36}\text{H}_{55}\text{O}_5$ ($\text{M}+\text{H}$) $^+$ 567.4044 found 567.4037.

Formation of Acetonide (**14**).

To a solution of starting 1,3-diol (88 mg, 155 μmol) in dry CH_2Cl_2 (1 mL) and 2,2-dimethoxy propane (1 mL) at room temperature was added a few crystals of pyridinium *p*-toluenesulfonic acid. After stirring for 3 h, the solvent was removed *in vacuo* and the reaction was subjected to flash column chromatography (20 g SiO_2 , 7% EtOAc in hexanes) and to afford 78 mg (83%) of acetonide **14** as a clear colorless oil: R_f 0.6 (20% EtOAc in hexanes); $[\alpha]_D^{21}$ -2.0 (c 0.90, CHCl_3); IR (neat) 2955, 2915, 2869, 1613, 1514 1455, 1380, 1362, 1248, 1226, 1172, 1097 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.32 – 7.31 (m, 4H), 7.27– 7.22 (m, 3H), 6.84 (d, J = 8.6 Hz, 2H), 4.84 (d, J = 9.4 Hz, 1H), 4.48 (s, 2H), 4.40 (s, 2H), 3.78 (s, 3H), 3.68 – 3.64 (m, 1H), 3.41 – 3.36 (m, 2H), 3.30 – 3.27

(m, 2H), 3.21 – 3.17 (m, 1H), 2.36 – 2.28 (m, 1H), 1.93 – 1.79 (m, 3H), 1.76 – 1.70 (m, 2H), 1.69 – 1.62 (m, 1H), 1.51 (s, 3H), 1.44 – 1.34 (m, 1H), 1.29 – 1.20 (m, 8H), 1.14 – 1.00 (m, 2H), 0.90 – 0.87 (m, 10H), 0.78 – 0.76 (m, 6H); ^{13}C NMR (101MHz, CDCl_3) δ 159.0, 138.8, 132.8, 132.2, 130.8, 129.1, 128.2, 127.4, 127.3, 113.7, 100.1, 76.6, 74.0, 72.9, 72.7, 72.6, 69.5, 55.2, 48.6, 40.8, 36.4, 36.3, 33.8, 32.2, 30.9, 28.5, 27.7, 25.1, 23.6, 21.3, 19.1, 16.9, 16.1, 11.8, 11.2; MS (EI) m/e (relative intensity) 550 (8), 220 (13), 137 (19), 122 (35), 121 (100), 99 (27), 91 (78); HRMS m/e calcd. for $\text{C}_{38}\text{H}_{57}\text{O}_5$ ($\text{M}-\text{CH}_3$) $^+$ 593.4201 found 593.4195.

Formation of Benzothiazole (15).

To a stirred solution of the starting primary alcohol (23 mg, 45 μmol) in dry THF (0.3 mL) at room temperature was added 2-mercapto-1,3-benzothiazole (9 mg, 54 μmol) and triphenylphosphine (14 mg, 54 μmol). The resulting solution was cooled to 0 $^{\circ}\text{C}$ and diisopropylazodicarboxylate (11 μl , 58 μmol) was added dropwise. The cooling bath was then removed and the mixture was allowed to warm to room temperature with stirring for 4 h, after which time, it was concentrated *in vacuo* and subjected to flash column chromatography (10 g SiO_2 , 2% EtOAc in hexanes) to afford 26 mg (87%) of pure

thioether **15** as a clear colorless oil: R_f 0.14 (5% EtOAc in hexanes); $[\alpha]_D^{21} -2.9$ (c 0.51, CHCl_3); IR (neat) 3066, 2955, 2914, 2859, 1620, 1513, 1455, 1428, 1379, 1247, 1226, 1163, 1089, 994 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.86 (d, $J = 8.1$ Hz, 1H), 7.75 (d, $J = 8.1$ Hz, 1H), 7.41 (t, $J = 7.7$ Hz, 1H), 7.31 – 7.26 (m, 3H), 6.89 (d, $J = 8.1$ Hz, 2H), 4.91 (d, $J = 9.3$ Hz, 1H), 4.44 (s, 2H), 3.82 (s, 3H), 3.73 – 3.68 (m, 1H), 3.45 – 3.39 (m, 3H), 3.32 (A of ABX, $J_{AB} = 8.7$ Hz, $J_{AX} = 6.3$ Hz, 1H), 3.19 (B of ABX, $J_{AB} = 8.7$ Hz, $J_{BX} = 7.4$ Hz, 1H), 2.42 – 2.32 (m, 1H), 2.10 – 1.96 (m, 2H), 1.90 – 1.70 (m, 4H), 1.67 – 1.58 (m, 2H), 1.48 – 1.38 (m, 2H), 1.36 – 1.24 (m, 11H), 1.06 (d, $J = 6.6$ Hz, 3H), 1.00 – 0.90 (m, 6H), 0.88 – 0.78 (m, 6H); ^{13}C NMR (101MHz, CDCl_3) δ 167.6, 159.0, 153.3, 135.1, 133.1, 132.0, 130.8, 129.1, 125.9, 124.0, 121.4, 120.8, 113.7, 100.1, 74.0, 72.8, 72.7, 69.5, 55.2, 48.4, 43.6, 41.4, 36.4, 36.3, 33.8, 32.3, 30.8, 28.5, 28.1, 25.1, 23.6, 21.3, 19.2, 19.1, 16.1, 11.8, 11.2; MS (CI, CH_4) m/e (relative intensity) 668 (29), 611 (37), 610 (84), 490 (28), 472 (26), 390 (14), 137 (17), 121 (100); HRMS m/e calcd. for $\text{C}_{39}\text{H}_{58}\text{NO}_4\text{S}_2$ ($\text{M}+\text{H}$) $^+$ 668.3802 found 668.3804.

Formation of **2** by Parikh-Doering Oxidation.

To a solution of starting primary alcohol (23 mg, 42 μmol) in dry CH_2Cl_2 (0.6 mL) at 0 $^\circ\text{C}$ was added SO_3 .pyridine complex (27 mg, 169 μmol) and then diisopropylethylamine

(0.05 mL, 297 μ mol). After 5 min, dry DMSO (0.04 mL, 593 μ mol) was added. After stirring for 5 min, the reaction was quenched with brine (10 mL) and diluted with Et₂O (10 mL). The organic phase was separated and washed with saturated aqueous NaHCO₃ (1 x 15 mL). The organic layer was dried over anhydrous MgSO₄, filtered, concentrated *in vacuo* and then subjected to flash column chromatography (10 g SiO₂, 10% EtOAc in hexanes) and to afford 20 mg (85%) of pure aldehyde **2** as a clear colorless oil: R_f 0.37 (20% EtOAc in hexanes); IR (neat) 2820, 2720, 1715cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 9.71 (s, 1H), 7.86 (d, *J* = 8.1 Hz, 1H), 7.76 (d, *J* = 8.1 Hz, 1H), 7.42 (t, *J* = 7.2 Hz, 1H), 7.30 (t, *J* = 7.3 Hz, 1H), 4.90 (d, *J* = 9.3 Hz, 1H), 3.79 – 3.72 (m, 2H), 3.41 (A of ABX, *J*_{AB} = 12.6 Hz, *J*_{AX} = 5.7 Hz, 1H), 3.20 (B of ABX, *J*_{AB} = 12.6 Hz, *J*_{BX} = 7.5 Hz, 1H), 2.46 – 2.34 (m, 2H), 2.08 – 1.98 (m, 2H), 1.87 – 1.70 (m, 3H), 1.57 (s, 3H), 1.48 – 1.42 (m, 2H), 1.33 -1.26 (m, 10H), 1.15 (d, *J* = 7.0 Hz, 3H), 1.06 (d, *J* = 6.6 Hz, 3H), 0.95 (d, *J* = 6.6 Hz, 3H), 0.86 (d, *J* = 6.8 Hz, 3H), 0.82 (d, *J* = 6.2 Hz, 3H).

1-Bromo-2-dimethoxymethyl-3,5-dimethoxy-4-methyl-benzene (19).

Trimethoxymethane (132 mg, 1.24 mmol) was added dropwise to a stirred solution of aldehyde **22** (50 mg, 0.19 mmol) containing few crystals of *p*-toluenesulfonic acid.

Molecular sieves (4A°) (80 mg) were also added. The mixture was then warmed to 70 °C and stirring was continued for 12 h. The reaction mixture was then concentrated *in vacuo*

and subjected to flash column chromatography (50 g, SiO₂, 15 % EtOAc in hexanes) to afford 53 mg of dimethyl acetal **19** (90%) as clear colorless oil: *R_f* 0.15 (10% EtOAc in hexanes); IR (neat) 2992, 2935, 2827, 1589, 1466, 1449, 1355, 1293, 1232, 1121, 1075, 990 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.86 (s, 1H), 5.62 (s, 1H), 3.83 (s, 3H), 3.78 (s, 3H), 3.48 (s, 6H), 2.11 (s, 3H); ¹³C NMR (101MHz, CDCl₃) δ 158.9, 158.8, 123.0, 120.3, 119.7, 111.6, 105.2, 62.0, 55.8, 55.7, 9.0; MS (LoRes ESI) *m/e* (relative intensity) 289 (12), 287 (12), 261 (93), 259 (100), 251 (20), 249 (11); HRMS *m/e* calcd. for C₁₂H₁₇BrO₄Na (M+Na)⁺ 327.0202 found 327.0203.

(S)-2-[(4R,5S,6R)-6-[(E)-(3S,7S,9S)-10-(Benzothiazol-2-ylsulfanyl)-3,5,7,9-tetramethyl-dec-4-enyl]-2,2,5-trimethyl-[1,3]dioxan-4-yl]-1-[2-(1,1-dimethoxyethyl)-3,5-dimethoxy-4-methyl-phenyl]-propan-1-ol (20).

To a stirred solution of aryl bromide **19** (15 mg, 51 μmol) in dry THF (0.3 mL) at –78 °C was added nBuLi (0.03 mL, 2.5 M in Hexanes), and the reaction mixture was stirred for 45 min at that temperature. Aldehyde **2** (23 mg, 42 μmol) dissolved in THF (0.3 mL) was then added dropwise. The reaction was maintained at –78 °C for an additional 0.5 h. and

was then quenched by addition of saturated aqueous NH_4Cl (10 mL). After extraction with Et_2O (2 x 20 mL), the combined Et_2O phases were dried (Na_2SO_4), filtered, and concentrated *in vacuo* to provide an oily residue which proved to be a 2:1 mixture of product diastereomers. The two products were then purified by flash column chromatography (20 g SiO_2 , 15% EtOAc in Hexanes) to afford 22 mg of alcohol **20** (68%). The major product diastereomer is fully characterized as follows: R_f 0.2 (20% EtOAc in hexanes); $[\alpha]_D^{21}$ -18.0 (c 0.40, CHCl_3); IR (neat) 3513, 2955, 2923, 1596, 1456, 1453, 1429, 1374, 1222, 1124, 1070, 993 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.82 (d, $J = 8.0$ Hz, 1H), 7.72 (d, $J = 7.8$ Hz, 1H), 7.40 – 7.34 (m, 1H), 7.28 – 7.20 (m, 1H), 7.03 (s, 1H), 5.64 (s, 1H), 5.55 (s, 1H), 4.86 (d, $J = 9.0$ Hz, 1H), 3.83 (s, 3H), 3.71 – 3.63 (m, 5H), 3.54 (s, 1H), 3.43 (s, 3H), 3.37 (A of ABX, $J_{AB} = 12.7$ Hz, $J_{AX} = 5.7$ Hz, 1H), 3.29 (s, 3H), 3.15 (B of ABX, $J_{AB} = 12.7$ Hz, $J_{BX} = 7.4$ Hz, 1H), 2.36 – 2.29 (m, 1H), 2.16 – 2.08 (m, 4H), 2.03 – 1.94 (m, 2H), 1.80 – 1.64 (m, 3H), 1.52 (s, 3H), 1.44 – 1.36 (m, 4H), 1.34 – 1.18 (m, 7H), 1.01 (d, $J = 6.6$ Hz, 3H), 0.90 (d, $J = 6.6$ Hz, 3H), 0.86 (d, $J = 7.2$ Hz, 3H), 0.81 (d, $J = 6.8$ Hz, 3H), 0.78 (d, $J = 6.4$ Hz, 3H); ^{13}C NMR (101MHz, CDCl_3) δ 167.7, 158.0, 157.6, 153.3, 141.8, 135.1, 133.0, 132.1, 126.0, 124.1, 121.4, 120.9, 120.2, 117.1, 105.9, 103.5, 100.6, 80.1, 74.3, 69.6, 61.6, 56.2, 55.5, 55.2, 48.4, 43.6, 41.4, 39.9, 36.5, 33.8, 32.3, 30.9, 29.7, 28.5, 28.1, 25.0, 24.0, 21.4, 19.2, 19.1, 16.1, 11.6, 9.0, 5.5; MS (LoRes ESI) m/e (relative intensity) 762 (69), 709 (100), 708 (47), 650 (54), 632 (24), 470 (75), 430 (59), 418 (12); HRMS m/e calcd. for $\text{C}_{43}\text{H}_{65}\text{NO}_7\text{S}_2\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 794.4099 found 794.6005.

2-((2S,4S,8S,E)-10-((4R,5S,6R)-6-((S)-1-(4-Methoxybenzyloxy)propan-2-yl)-2,2,5-trimethyl-1,3-dioxan-4-yl)-2,4,6,8-tetramethyldec-6-enylsulfonyl)benzo[d]thiazole (21).

To a solution of sulfide **15** (15 mg, 0.02 mmol) in EtOH (0.7 mL) at 0 °C was added the oxidant [20 µL of a premixed solution of 320 mg of ammonium molybdate in 50% H₂O₂ (360 µL)] followed by NaOAc (15 mg, 0.18 mmol). After stirring 4 h at 0 °C, the reaction mixture was diluted with EtOAc (5 mL), washed with H₂O (5 mL) and then brine (5 mL). The organic layer was separated, dried (MgSO₄), filtered and concentrated providing a yellow oil. The crude product was purified by flash column chromatography (5 mL SiO₂, 15% EtOAc : Hexanes) to afford 13 mg (85%) of sulfone **21** as a clear colorless oil: *R_f* 0.35 (20% EtOAc in Hexanes); [α]_D²¹ -2.9 (c 0.82, CHCl₃); IR (neat) 3061, 2958, 2930, 2869, 1615, 1513, 1458, 1380, 1327, 1247, 1226, 1149, 1089, 1024 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.23 (d, *J* = 8.6 Hz, 1H), 8.04 (d, *J* = 7.7 Hz, 1H), 7.67 – 7.60 (m, 2H), 7.27 (d, *J* = 7.7 Hz, 2H), 6.89 (d, *J* = 8.6 Hz, 2H), 4.85 (d, *J* = 9.1 Hz, 1H), 4.44 (s, 2H), 3.82 (s, 3H), 3.72 – 3.66 (m, 1H), 3.54 – 3.49 (m, 1H), 3.45 – 3.30 (m, 4H), 2.44 – 2.28 (m, 2H), 1.90 – 1.82 (m, 2H), 1.80 – 1.72 (m, 2H), 1.69 – 1.61 (m, 4H), 1.52 (s, 3H), 1.32 – 1.22 (m, 12H), 1.13 (d, *J* = 6.6 Hz, 3H), 0.94 – 0.91 (m, 6H),

0.81 – 0.79 (m, 6H); ^{13}C NMR (101MHz, CDCl_3) δ . 167.1, 159.3, 153.0, 136.9, 133.5, 131.9, 131.0, 129.4, 128.2, 127.9, 125.7, 122.6, 114.0, 100.4, 74.3, 73.0, 72.9, 69.8, 61.8, 55.5, 48.4, 44.2, 36.7, 36.6, 34.0, 32.5, 28.8, 28.0, 26.5, 25.3, 23.9, 21.6, 19.8, 19.2, 16.3, 12.1, 11.4; MS (ESI) m/e (relative intensity) 695 (8), 681 (10), 680 (28), 638 (10), 637 (25), 623 (24), 622 (100), 616 (5); HRMS m/e calcd. for $\text{C}_{39}\text{H}_{57}\text{NO}_6\text{S}_2\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 722.3525 found 722.3520.

(4R,5S,6R)-4-((3S,4E,7S,9S,10E)-11-(6-bromo-2,4-dimethoxy-3-methylphenyl)-3,5,7,9-tetramethylundeca-4,10-dienyl)-6-((S)-1-(4-methoxybenzyloxy)propan-2-yl)-2,2,5-trimethyl-1,3-dioxane (23).

To a solution of sulfone **21** (13 mg, 0.02 mmol) in THF (0.2 mL), an LDA solution (0.05 mL, 0.5 M) was added dropwise at $-78\text{ }^\circ\text{C}$ and the reaction mixture was stirred for 25 min. A solution of aldehyde **22** (10.3 mg, 0.04 mmol) in THF (0.15 mL) was then added dropwise, and the mixture was immediately warmed to room temperature. After stirring for 1 h, the reaction was quenched with saturated aqueous NH_4Cl (5 mL), and extracted with EtOAc (2 X 5 mL). The organic phases were combined, dried (MgSO_4), filtered and concentrated to exclusively provide *E*-olefin. Product was purified by flash column

chromatography (5 mL SiO₂, 5% Et₂O in Hexanes) to afford 12 mg (80%) of **23** as clear colorless oil: R_f 0.45 (20% EtOAc in hexanes); [α]_D²¹ +8.4 (c 0.77, CHCl₃); IR (neat) 2954, 2925, 2852, 1612, 1589, 1513, 1453, 1380, 1247, 1226, 1123 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.23 (d, *J* = 8.0 Hz, 2H), 6.84 (d, *J* = 8.0 Hz, 2H), 6.83 (s, 1H), 6.30 (d, *J* = 16.0 Hz, 1H), 6.14 (dd, *J* = 16.0, 8.2 Hz, 1H), 4.86 (d, *J* = 9.4 Hz, 1H), 4.40 (s, 2H), 3.78 (s, 3H), 3.77 (s, 3H), 3.69 – 3.64 (m, 1H), 3.58 (s, 3H), 3.41 – 3.36 (m, 2H), 3.29 – 3.26 (m, 1H), 2.44 – 2.31 (m, 2H), 2.09 – 2.07 (m, 4H), 1.85 – 1.78 (m, 1H), 1.73 – 1.63 (m, 3H), 1.53 (s, 3H), 1.41 – 1.20 (m, 12H), 1.04 (d, *J* = 6.6 Hz, 3H), 0.90 (d, *J* = 5.6 Hz, 3H), 0.88 (d, *J* = 6.6 Hz, 3H), 0.79 (d, *J* = 6.2 Hz, 3H), 0.76 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (101MHz, CDCl₃) δ 159.3, 157.7, 157.4, 142.9, 133.0, 132.8, 131.1, 129.4, 124.6, 122.6, 121.3, 120.1, 114.0, 111.2, 100.4, 74.3, 73.1, 73.0, 69.7, 59.7, 56.0, 55.5, 48.0, 44.8, 36.7, 36.6, 36.0, 34.1, 32.5, 28.7, 28.5, 25.3, 23.9, 21.6, 20.9, 20.2, 16.4, 12.1, 11.4, 9.2; MS (CI, CH₄) *m/e* (relative intensity) 744 (34), 743 (32), 742 (30), 730 (36), 729 (94), 727 (100), 725 (27), 713 (12), 709 (10); HRMS *m/e* calcd. for C₄₂H₆₃BrO₆ 742.3808, found 742.3803.

OBSERVE H1

STANDARD 1H OBSERVE

Pulse Sequence: s2pul

File: ks-3-l44

Date: Apr 27 2004

VXR400

PULSE SEQUENCE: standard

OBSERVE H1

Pulse Sequence: s2pul

STANDARD 1H OBSERVE

Archive directory: /i400/dob/vnmrsys/data
Sample directory:

Pulse Sequence: s2pul

User: 1-12-87

File: Ks-4-64

Date: Mar 4 2005

I500

PULSE SEQUENCE: standard

OBSERVE H1

¹³C OBSERVE

Archive directory: /i400/ksh/vnmrsys/data
Sample directory:
File: CARBON

Pulse Sequence: s2pul

Date: Jun 1 2004

i400

PULSE SEQUENCE: standard

OBSERVE C13

DECOUPLE H1

FORM 1041-101
OBSERVE H1

```
Archive directory: /i400/ksh/vnmrsys/data
Sample directory:
File: CARBON
```

Date: Jun 7 2004
I400

PULSE SEQUENCE: standard
OBSERVE C13
DECOUPLE H1

13

Archive directory: /400/dob/hymn/systa

Sample directory:
File: PROTON

Pulse Sequence: s2pul

User: 1-12-87

Date: Mar 1 2005

1400

PULSE SEQUENCE: standard

OBSERVE H1

13C OBSERVE

Archive directory: /MOOfish/mmmsys/data

Sample directory:

File: CARBON

Pulse Sequence: s2pul

Date: Jun 10 2004

DOM

PULSE SEQUENCE: standard

OBSERVE C13

DECOUPLE H1

```
Archive directory: /vvr400/dob/vmrsys/data
Sample directory:
```

Pulse Sequence: s2pul
File: ks-4-112
Date: Apr 28 2005
VXR400.
PULSE SEQUENCE: standard
OBSERVE H1

13C OBSERVE

```
Archive directory: /vvr400/dob/vnmr/sys/data
Sample directory:
File: CARBON
```

Pulse Sequence: s2pul

Date: Apr 28 2005

VXR400

PULSE SEQUENCE: standard

OBSERVE C13

DECOUPLE H1

Active directory: hmc02bdcvrennyss\data
Sample directory:
File: PROTON
Pulse Sequence: s2pf1
Date: Apr 14 2005
V079400
PULSE SEQUENCE: standard
OBSERVE: 1H

2

STANDARD 1H OBSERVE

Archive directory: /i400/dob/vnmrsys/data

Sample directory:

Pulse Sequence: sbpul

User: 1-12-87

File: ks-4-109-upper

Date: Apr 15 2005

1500

PULSE SEQUENCE: standard

OBSERVE H1

¹³C OBSERVE

Archive directory: /vnr400/dob/vnmrSYS/data
Sample directory:
File: CARBON

Pulse Sequence: s2pul

Date: Apr 16 2005

VXR400

PULSE SEQUENCE: standard

OBSERVE C13

DECOUPLE H1

Archive directory: /A000/bdb/vnmr/sys/data
Sample directory:
File: PROTON

User: 1-12-87

1400

PULSE SEQUENCE: standard
OBSERVE H1

STANDARD CARBON PARAMETERS

Archive directory: /nmr/i500/jas/vnmrsys/data
Sample directory:

Pulse Sequence: s2pul

User: 1-14-87

File: ks-c.001

Date: Jun 15 2005

I500

PULSE SEQUENCE: standard

OBSERVE C13

DECOUPLE H1

23