This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Eicosapentaenoic acid is more effective than docosahexaenoic acid in inhibiting proinflammatory mediator production and transcription from LPS-induced human asthmatic alveolar macrophage cells

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1016/j.clnu.2008.10.012>

PUBLISHER

© Elsevier Ltd and European Society for Clinical Nutrition and Metabolism

VERSION

AM (Accepted Manuscript)

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Mickleborough, Timothy D., Sandra L. Tecklenburg, Gregory S. Montgomery, and Martin R. Lindley. 2019. "Eicosapentaenoic Acid Is More Effective Than Docosahexaenoic Acid in Inhibiting Proinflammatory Mediator Production and Transcription from Lps-induced Human Asthmatic Alveolar Macrophage Cells". figshare. <https://hdl.handle.net/2134/11341>.

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.

For the full text of this licence, please go to:
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

Eicosapentaenoic acid is more effective than docosahexaenoic acid in
inhibiting pro-inflammatory mediator production and transcription from LPS-
induced human asthmatic alveolar macrophage cells

¹Timothy D. Mickleborough Ph.D, ¹Sandra L. Tecklenburg M.S, ²Gregory S Montgomery, M.D.
and ³Martin R. Lindley PhD.

¹Exercise Biochemistry Laboratory, Department of Kinesiology, Indiana University, Bloomington,
Indiana. ²Section of Pediatric Pulmonology, Indiana University School of Medicine and Riley Hospital
for Children, Indianapolis, Indiana. ³Human Cellular and Molecular Research Laboratory, Loughborough
University, Loughborough, LE11 3TU, United Kingdom

Correspondence:

Timothy D. Mickleborough, Ph.D.
Department of Kinesiology
Indiana University
1025. E. 7th Street, HPER 112
Bloomington, IN 47401

Phone: 812-855-0753
Fax: 812-855-3193
Email: tmickleb@indiana.edu

Running title: Fish oil and proinflammatory mediator generation

Abstract

Background & aims: The purpose of the study was to determine which of the active constituents of fish oil, eicosapentaenoic acid (EPA) or docosahexaenoic acid (DHA), is most effective in suppressing proinflammatory mediator generation and cytokine expression from LPS-stimulated human asthmatic alveolar macrophages (AM ϕ).

Methods: The AM ϕ were obtained from twenty one asthmatic adults using fiberoptic bronchoscopy. Cells were pretreated with DMEM, pure EPA, an EPA-rich media (45% EPA/10% DHA), pure DHA, a DHA-rich media (10% EPA/50% DHA) or Lipovenos^R (*n*-6 PUFA), and then exposed to **Dulbecco's Modified Eagle's Medium** (DMEM) (-) or LPS (+). Supernatants were analyzed for leukotriene (LT)B₄, prostaglandin (PG)D₂, tumor necrosis factor (TNF)- α and interleukin (IL)-1 β production. Detection of TNF- α and IL-1 β mRNA expression levels were quantified by reverse transcriptase polymerase chain reaction.

Results: 120 μ M pure EPA and EPA-rich media significantly ($p < 0.05$) suppressed TNF- α and IL-1 β mRNA expression and the production of LTB₄, PGD₂ and TNF- α and IL-1 β in LPS-stimulated primary AM ϕ cells obtained from asthmatic patients to a much greater extent than 120 μ M pure DHA and DHA-rich media respectively.

Conclusions: This study has shown for the first time that EPA is a more potent inhibitor than DHA of inflammatory responses in human asthmatic AM ϕ cells.

Keywords: eicosapentaenoic acid, docosahexaenoic acid, fish oil, asthma, proinflammatory mediators

1 Introduction

2
3 Over the past three decades there has been significant interest in the therapeutic potential of fish
4 oils for various inflammatory conditions such as rheumatoid arthritis, inflammatory bowel diseases, and
5 asthma. Fish oil, rich in omega-3 (*n*-3) polyunsaturated fatty acids (PUFA), such as eicosapentaenoic acid
6 (EPA) and docosahexaenoic acid (DHA), appear to have additional antiphlogistic properties primarily
7 through their effects on the neutrophil and macrophage (M ϕ) component of the inflammatory response ¹⁻⁷.

8 Eicosapentaenoic acid can compete with arachidonic acid (AA), as a substrate for
9 cyclooxygenase (COX)-2 and 5-lipoxygenase (5-LO) enzymes and be converted to less inflammatory
10 eicosanoids ^{6, 8}. At present the mechanism(s) underpinning the anti-inflammatory effects of DHA are
11 unclear, but may be related to altered gene transcription and translation via direct or indirect actions on
12 intracellular signaling pathways ⁹⁻¹¹. In addition, *n*-3 PUFA-derived mediators such as lipoxin,
13 docosatrienes resolvins and neuroprotectins may also have anti-inflammatory, pro-resolving and
14 protective properties ¹².

15 The observational evidence on fish oil effects has been relatively consistent in demonstrating
16 protection against asthma and/or allergy in relation to a high intake, and ecological and other cross-
17 sectional data support the hypothesis that *n*-6 PUFA may increase and *n*-3 PUFA may decrease asthma
18 risk ¹³. While the clinical data on the effect of fish oil supplementation in asthma has been equivocal
19 ¹⁴ supplementing the diet with fish oil in individuals with exercise-induced asthma (EIA) has yielded
20 promising results ^{15, 16}. Our laboratory has shown that 3 weeks of fish oil supplementation reduced the
21 severity of EIA, airway inflammation and bronchodilator use, and improved asthma symptoms scores in
22 elite athletes ¹⁵ and asthmatic individuals ¹⁶.

23 The majority of studies investigating the effects of *n*-3 PUFA on asthma/EIA have either
24 employed fish oils rich in EPA or oils which contain a heterogeneous blend of EPA and DHA. Only a
25 few studies have examined the effects of supplementing asthmatic patients with pure EPA and/or DHA,
26 with conflicting results ¹⁴. Data is therefore insufficient to make recommendations for intake of specific
27 *n*-3 PUFA in asthma, e.g. EPA versus DHA versus EPA + DHA combined ¹⁷. Although many studies

1 have investigated the effects of EPA and DHA on macrophage function in animal models and cell lines,
2 there is little evidence about the effects of these lipids on primary human macrophages obtained from
3 asthmatic patients.

4 Therefore, the main aim of this study was to compare the individual effects of EPA and DHA,
5 and a variety of heterogeneous blends of EPA and DHA, on eicosanoid and cytokine generation from
6 LPS-stimulated human asthmatic alveolar macrophages (AM ϕ). In addition, the effects of EPA and DHA
7 on cytokine mRNA expression were investigated in the LPS-stimulated AM ϕ .

8 9 **Methods**

10
11 Twenty non-smoking adults with asthma were recruited to this study. Asthma was diagnosed by a
12 history of recurrent wheezing and chest tightness and a previous physician diagnosis. All subjects had
13 clinically treated mild-to- moderate persistent asthma, with an FEV₁ greater than 70% of predicted ¹⁸.
14 Inhaled corticosteroids, 5-lipoxygenase inhibitors and leukotriene receptor antagonists were withheld for
15 4 weeks prior to fiberoptic bronchoscopy. Subjects were also excluded if they had a history of taking *n*-3
16 PUFA supplements or consumed more than one fish meal per week. A group of nonasthmatic (control)
17 subjects was not included in the present study as it has been shown that fish oil supplementation does not
18 alter pulmonary function or inflammatory mediator generation in this population ¹⁵. The local Institutional
19 Research Ethics Committee approved the study protocol.

20 21 **Fiberoptic Bronchoscopy.**

22 Fiberoptic bronchoscopy was used to obtain BALF from each subject. Using local anesthesia with
23 lidocaine (2% wt/vol) to the upper airways and larynx, a fiberoptic bronchoscope was passed through the
24 nasal passages into the trachea. The bronchoscope was wedged in the right middle lobe and 4 × 60-ml
25 aliquots of prewarmed sterile 0.9% NaCl solution were instilled. This solution was aspirated through the
26 bronchoscope, collected in prechilled glass bottles, and stored on ice and processed within 30 min.

1 Separation of AM ϕ from BALF.

2 AM ϕ cells were separated from BALF using previously described methods, with slight
 3 modifications ¹⁹. Briefly, the BALF was filtered through a single layer of coarse sterile gauze to remove
 4 mucus clumps and then centrifuged at 1,000 g for 10 min at 4° C to obtain a cell pellet. The cell pellet was
 5 washed once in 50 ml of Ca²⁺/ Mg²⁺ free Hanks' balanced salt solution (HBSS). The cells were counted on
 6 a hemocytometer slide using a Kimura counterstain and viability assessed by the trypan blue exclusion
 7 test. Cytospins were performed, using 10⁴ cells per slide, and stained with May-Grunwald-Giemsa in
 8 order to obtain differential cell counts (Table 1). The remaining cells were resuspended at a concentration
 9 of 2 × 10⁶ AM ϕ per milliliter in RPMI 1640 medium supplemented with 10% (vol/vol) fetal calf serum,
 10 2 mM L-glutamine, 100 U/ml penicillin, and 100 µg/ml streptomycin. 2 × 10⁶ AM ϕ /well were plated onto
 11 6-well plates and allowed to adhere for 90 min in a humidified incubator in 95% air, 5% CO₂ (vol/vol), at
 12 37° C. Nonadherent cells were removed by washing three times with RPMI 1640 medium, leaving the
 13 adherent macrophages. The resulting AM ϕ population was > 95% pure, as assessed by staining and
 14 morphologic analysis. The AM ϕ from each individual were harvested with a cell scraper and combined
 15 into one aliquot.

17 Experimental Design.

18 Cells were divided into six treatment groups, pure EPA [*cis*-5, 8, 11, 14, 17-eicosapentanoic acid
 19 (Sigma-Aldrich, St. Louis, MO)], an EPA-rich media [(EPAX 4510 TG (45% EPA/10% DHA) (Pronova
 20 Biocare, Lysaker, Norway)], pure DHA [*cis*-4, 7, 10, 13, 16, 19-docosahexanoic acid (Sigma-Aldrich, St.
 21 Louis, MO)], a DHA-rich media [(EPAX 1050 TG (10% EPA/50% DHA) (Pronova Biocare, Lyasker,
 22 Norway)], Lipovenos® (Fresenius-Kabi, Bad-Homburg, Germany: an *n*-6 PUFA) or Dulbecco's
 23 Modified Eagle's Medium (DMEM) (control media).

24 All fatty acids were dissolved in distilled H₂O, aliquoted, and stored under an N₂ stream, and
 25 stored at -80°C for no longer than 1 week prior to use. EPAX 4510 TG is a triglyceridic oil containing
 26 approximately 45% EPA and 10% DHA, with saturated fatty acids and monounsaturated fatty acids

comprising 9-12% and 20-24% respectively. EPAX 1050 TG is triglyceridic oil containing approximately 10% EPA and 50% DHA, with saturated fatty acids and monounsaturated fatty acids comprising 2-10% and 5-15% respectively.

AM ϕ cells were suspended in Dulbecco's modified Eagle's medium (DMEM) (Mediatech, Herndon, VA), supplemented with 10% heat inactivated endotoxin-free fetal bovine serum (FBS, Interger), 100 U/ml penicillin, and 100 U/ml streptomycin (GIBCO-BRL) and incubated at 37°C in 5% CO₂. In all experiments, cells were plated in 24-well plates at a density of 1 x 10⁶ cells/well and allowed to adhere for 2 hr and then incubated with either a high-dose (120 μ M) or low dose (25 μ M) (of a 100mM stock solution) of pure EPA, pure DHA, EPAX 4510, EPAX 1050 or Lipovenos^R (*n*-6 PUFA) for 4 hr².²⁰ The medium was then aspirated and the cells rinsed with 10 ml of sterile PBS twice. After washing the pure EPA, pure DHA, EPAX 4510 TG, EPAX 1050 TG or Lipovenos^R media from the plates, fresh DMEM supplemented with 10% FBS and L-glutamine was used to carry out all subsequent incubations with or without LPS. The cells were then stimulated with LPS (1 μ g/ml) (BD Diagnostics, Sparks, MD) or incubated with medium alone for 3 hr²⁰. The supernatant was gently aspirated and stored at -70°C for subsequent competitive enzyme immunoassay (EIA) analysis of leukotriene (LT) B₄ (Cayman Chemical, Ann Arbor, MI), prostaglandin (PG) D₂- methoxime (MOX) (Cayman Chemical, Ann Arbor, MI), tumor necrosis factor (TNF)- α (Cayman Chemical, Ann Arbor, MI), and interleukin (IL)-1 β (Cayman Chemical, Ann Arbor, MI) using previously described methods¹⁶. Because PGD₂ is a relatively unstable compound, we measured PGD₂-MOX), a stable derivative of PGD₂.

Evaluation of cytokine mRNA expression

Total RNA was isolated from macrophage monolayers using a commercially available kit (Qiagen, Valencia, CA) following the protocol provided. Detection of mRNA by real-time polymerase chain reaction was performed on an ABI-PRISM[®] 7700 Sequence Detector (Perkin-Elmer Applied Biosystems, Foster City, CA, USA) as previously described¹⁰. The TaqMan real-time PCR was performed on the cytokines TNF- α and IL-1 β using pre-developed assay reagent kits. All reagents

necessary for running a *TaqMan* real time PCR assay were purchased from Perkin-Elmer Applied Biosystems. Each 25- μ l reaction contained forward and reverse primers, 20 ng of cDNA, 2x *TaqMan* Universal PCR Mastermix and *TaqMan* probe. The primers (Life Technologies, Grand Island, NY) and probes (Applied Biosystems, Foster City, CA) used in the TNF- α assay were: forward primer, 5'-TGATCCGAGACGTGGAA -3'; reverse primer, 5'-ACCGCCTGGAGTTCTGGAA -3'; and for the *TaqMan* Probe (5' 6-FAM, 3' TAMRA labeled), TGGCAGAAGAGGCACTCCCCCAA. For the IL-1 β assay: forward primer, 5'-CTGATGGCCCTAAACAGATGAAG – 3'; reverse, 5'-GGTCGGAGATTCGTAGCAGCTGGAT – 3'; and for the *TaqMan* Probe (5' 6-FAM, 3' TAMRA labeled), ATGAACAACAAAAATGCCTCGTGCTGTCTG. All reactions were performed in triplicate under the following conditions: 50°C for 2 min and 95°C for 10 min, followed by 40 cycles of 95°C for 15 s and 60°C for 1 min. Relative expression levels of IL-1 β and TNF- α mRNA were calculated using the $2^{-\Delta\Delta CT}$ method²¹ after confirmation that the efficiency of the real-time PCR reaction was similar for the 2 target genes over a range of template concentrations²². The fold change for each target gene, normalized to GAPDH, was calculated for each sample using the equation $x \text{ (amount of target)} = 2^{-\Delta\Delta CT}$. The mean ($\pm$ SD) fold change in gene expression was then determined from the triplicate samples, and expressed relative to vehicle control (DMEM).

Statistical Analysis

Data were analyzed using the SPSS version 15 statistical software (SPSS Inc., Chicago, IL, USA). A one-way ANOVA was used to analyze the data. Where a significant *F*-ratio was found, Fisher's protected least-square difference post-hoc test was used to detect differences in group means. All reported p-values were considered significant at the 0.05 level.

Results

The fluid recovered, total cell count and percent differential airway cell counts recovered from bronchial samples are presented in Table 1. The unstimulated AM ϕ cells did not demonstrate any significant differences ($p>0.05$) in LTB₄, PGD₂, TNF- α or IL-1 β production among the high dose (120 μ M) treatment groups (Figure 1, panel A-D). DMEM pretreated controls demonstrated a significant increase ($p<0.05$) in LTB₄, PGD₂, TNF- α and IL-1 β production in response to LPS stimulation (Figure 1, panel A-D). The inhibitory effect of 120 μ M pure EPA on LPS-stimulated LTB₄, PGD₂, TNF- α and IL-1 β production was significantly ($p<0.05$) greater than that of 120 μ M pure DHA, EPAX4510, EPAX1050 and Lipovenos^R. EPA and EPAX 4510 pretreatment significantly reduced ($p<0.05$) LPS-stimulated LTB₄, PGD₂, TNF- α and IL-1 β production by 84.6% and 66.3%, 81.7% and 60.5%, 90.2% and 72.9%, and 88.2% and 73.6% respectively compared to DMEM pretreatment (Figure 1, panel A-D). Similarly, 120 μ M pure DHA and EPAX 1050 pretreatment significantly decreased ($p<0.05$) LPS-stimulated LTB₄, PGD₂, TNF- α and IL-1 β production by 50.3%, 39.7%, 18.3%, 51.7% and 31.5%, 51.6% and 33.2% respectively compared to DMEM pretreatment (Figure 1, panel A-D). In contrast Lipovenos[®] had no significant ($p>0.05$) effect on LPS-induction of pro-inflammatory mediators compared to control cells (Figure 1, panel A-D).

Since differences in LPS-stimulated eicosanoid and cytokine production were observed at the high dose (120 μ M) we also examined whether these differences would be noticeable at a lower dose of *n*-3 PUFA (25 μ M). Figure 2 (panel A-B) presents the effects of LPS-stimulated LTB₄ and TNF- α production in AM ϕ cells pretreated with 25 μ M pure EPA, pure DHA, EPAX4510, EPAX1050 and Lipovenos^R. The anti-inflammatory response at the 25 μ M dose followed a similar pattern as the 120 μ M *n*-3 PUFA dose, but had significantly ($p<0.05$) less of an inhibitory effect on LPS-stimulated AM ϕ LTB₄ and TNF- α production (Figure 2, panel A-B). A comparable effect was seen for LPS-stimulated PGD₂ and IL-1 β production (data not shown).

The effect of the high dose (120 μ M) and low dose (25 μ M) *n*-3PUFA was also examined at the transcriptional level. We investigated the effects of 25 μ M and 120 μ M pure DHA, EPAX4510,

EPAX1050 and Lipovenos^R on LPS-stimulated AM ϕ TNF- α and IL-1 β mRNA expression (Figure 3, panel A-B). The inhibitory effect of 120 μ M pure-EPA on LPS-stimulated TNF- α and IL-1 β mRNA expression was significantly greater ($p < 0.05$) than that of 120 μ M pure-DHA, EPAX4510, EPAX1050 and Lipovenos^R (Figure 3, panel A). Interestingly, the effect of 25 μ M *n*-3 PUFA on LPS-stimulated TNF- α and IL-1 β mRNA expression followed a similar pattern to the 120 μ M pretreated cells (Figure 3, panel A) but had significantly less ($p < 0.05$) of an inhibitory effect (Figure 3, panel B).

Discussion

This study has demonstrated for the first time that pure EPA media **reduced** TNF- α and IL-1 β mRNA expression and the production of LTB₄, PGD₂ and TNF- α and IL-1 β from LPS-stimulated primary AM ϕ cells obtained from asthmatic patients to a much greater extent than pure DHA. Interestingly, the EPA-rich media (EPAX 4510) significantly reduced cytokine mRNA expression and generation of eicosanoids and cytokines from LPS-stimulated AM ϕ cells to a much greater degree than both the pure DHA and the DHA-rich media (EPAX 1050), suggesting that the greater the EPA content of an *n*-3 PUFA formulation the greater the eicosanoid and cytokine **reduction**. Whilst these experiments were conducted at a comparatively high *n*-3 PUFA dose (120 μ M) we also observed a similar pattern of LPS-stimulated AM ϕ cytokine and eicosanoid production and cytokine mRNA expression inhibition on a much lower *n*-3 PUFA concentration (25 μ M). Although our findings agree with a number of human studies that have shown that fish oil suppressed cytokine production in LPS-stimulated mononuclear cells^{4, 23, 24}, other studies have not shown this response²⁵⁻²⁷.

While there are a few studies that have assessed the efficacy of *n*-3 PUFA on cytokine production using human THP-1 monocytes^{10, 28, 29} and murine macrophages^{2, 7, 20}, our study is the first to assess the efficacy of a variety of *n*-3 PUFA blends on eicosanoid and cytokine production from AM ϕ obtained directly from human asthmatic patients. Our data, for the most part, agree and expand the few studies that have examined either the individual or combined effects of EPA and DHA on macrophage function in vitro. Lo et al.⁷ observed a reduction in TNF- α production and mRNA expression in LPS-stimulated

1 murine RAW macrophage incubated in the presence of 114 μ M EPA for 24 h. Similarly Zhao et al.³⁰ has
 2 shown that LPS-stimulated human THP-1 monocytes pretreated with 60 μ M EPA for 24 h significantly
 3 decreased TNF- α production and mRNA expression. In contrast Zhao et al.³¹ has shown in a follow-up
 4 study that human THP-1 monocytes pretreated with 100 μ M DHA for 24 h significantly reduced LPS
 5 (1 μ g/ml)-stimulated IL-6, IL-1 β and TNF- α production and mRNA expression. Chu and coworkers²⁸
 6 have reported using low doses of EPA (10 μ M) and DHA (10 μ M) significantly reduced TNF- α and IL- β
 7 generation from LPS-stimulated human THP-1 monocytes. It has been demonstrated that murine RAW
 8 264.7 macrophages pretreated for 4 h with Omegaven®, a high purity emulsion containing 1.25 - 2.82 g
 9 EPA and 1.44 - 3.09 g DHA, resulted in a significant suppression of LPS (1 μ g/ml)-stimulated TNF- α
 10 production by 48%² and TNF- α mRNA expression by 47%²⁰ compared to control media, while
 11 Lipovenos® (an *n*-6 PUFA emulsion) did not alter cytokine production compared to control medium
 12 alone.

13 Recently Weldon et al.¹⁰ sought to investigate the differential effects of pure EPA and DHA on
 14 cytokine expression from activated human THP-1 monocyte-derived macrophages in vitro. Equivalent
 15 doses of EPA and DHA significantly decreased LPS-stimulated THP-1 monocyte TNF- α , IL-1 β and
 16 TNF- α production and mRNA expression compared to control cells. However, whether at a relatively
 17 high dose (100 μ M) or low dose (25 μ M), DHA had a much greater inhibitory effect on cytokine
 18 production and mRNA expression than the equivalent EPA dose. This is in contrast with the findings
 19 from the present study which found that EPA at a high and low dose had a significantly greater inhibitory
 20 effect on LPS-stimulated human AM ϕ LTB₄, PGD₂, TNF- α - and IL- β generation and TNF- α - and IL- β
 21 mRNA expression compared to a comparable DHA dose. The divergent findings between the present
 22 study and the Weldon et al.¹⁰ study are difficult to reconcile, but may in part be related to the use of
 23 different cell types used. For example, the present study used primary human AM ϕ taken directly from
 24 the airways of asthmatic patients, whereas the cell line used in the experiments by Weldon et al.¹⁰ were
 25 human THP-1 monocytes, which although may behave like native monocyte –derived macrophages in

1 comparison to other human myeloid cell lines, may express a different physiologic response compared to
2 primary human AM ϕ when exposed to a *n*-3 PUFA.

3 The present study has demonstrated that *n*-3 PUFA inhibits LTB₄ and PGD₂ from LPS-stimulated
4 human AM ϕ . Our results concur with several studies in which dietary supplementation with EPA and
5 DHA **reduced** inflammatory eicosanoids such as products generated via the 5-lipoxygenase pathway of
6 neutrophils and monocytes and an attenuation of LTB₄-mediated chemotaxis⁶, and decreased PGE₂
7 production in LPS-stimulated murine RAW 264.7 cells pretreated with Omegaven® compared to control
8 cells³.

9 In the present study we evaluated the release of particular proinflammatory eicosanoids, derived
10 from both the cyclooxygenase (COX) and 5-lipoxygenase (5-LO) pathway such as PGD₂ and LTB₄, and a
11 few key proinflammatory cytokines (IL-1 β and TNF- α) that have been directly implicated in the
12 pathogenesis of asthma. LTB₄ is a potent neutrophil chemoattractant factor in the airways³², while PGD₂
13 is a potent bronchoconstrictor and is thought to play a role in pathogenesis of asthma, in particular during
14 the early asthmatic response to allergen³³. IL-1 β induces airway neutrophilia, and increased expression of
15 IL-1 β in asthmatic airway epithelium has been reported, together with an increased an increased number
16 of AM ϕ expressing IL-1 β ³⁴. TNF- α is also released from AM ϕ from asthmatic patients after allergen
17 challenge³⁵, and may have an important amplifying effect in asthmatic inflammation³⁶. Since, both IL-
18 1 β and TNF- α both activate and are activated by nuclear factor-kappaB (NF- κ B), this positive regulatory
19 loop may amplify and perpetuate the asthmatic inflammatory response³⁷.

20 Our findings indicate that that EPA is a more potent inhibitor than DHA of LPS-stimulated
21 eicosanoid and cytokine generation from human asthmatic AM ϕ . At present the data from the few studies
22 assessing the comparative effects of EPA and DHA on in vitro inflammatory mediator generation is
23 equivocal. Khalfoun et al.³⁸ have demonstrated a more potent inhibition of IL-6 from LPS-stimulated
24 lymphocytes on EPA compared to DHA, while Weldon et al.¹⁰ more recently demonstrated a more potent
25 inhibition on DHA compared with EPA in inhibiting LPS-stimulated human THP-1 monocyte cytokine
26 production. On the other hand Moon et al.³⁹ observed no difference between EPA and DHA on IL-6

secretion from murine macrophages, and Chu et al.²⁸ observed no difference between EPA and DHA on TNF- α and IL-1 β production from human THP-1 monocytes.

The biological mechanisms underpinning the more potent anti-inflammatory effects of EPA compared to DHA in the present study may be related to diverse mechanisms of action. EPA can cause dual inhibition of cyclooxygenase (COX)-2 and 5-lipoxygenase (LOX) pathways. EPA is a much less preferred substrate compared with AA for both pathways, and generally by substrate competition inhibits release of AA derived eicosanoids, thus reducing the generation of proinflammatory 'tetraene' 4-series leukotrienes (LTs) and 2-series prostanoids, and production of cytokines from inflammatory cells^{6,8}. EPA-derived metabolites have lower biological activity compared to their analogous AA-derivatives⁴⁰. We have previously shown that a fish oil diet decreased LTB₄ and increased LTB₅ generation from activated polymorphonuclear leukocytes obtained from asthmatic patients¹⁶. In addition, the anti-inflammatory effects of EPA may occur by modulating intracellular signal pathways which, in turn, influence gene activation and cytokine production. Lo and coworkers⁷ showed that RAW macrophages incubated in EPA-rich media altered NF- κ B activity (suppression of p65/p50 dimer), while Zhao et al.³⁰ demonstrated that EPA inhibited LPS-induced NF- κ B activation in human THP-1 monocytes. Lipopolysaccharide stimulation of monocytes activates several intracellular pathways, including I κ B kinase and mitogen-activated kinase (MAPK) pathways (ERK, JNK and p38). These signaling pathways activate a variety of transcription factors such as NF- κ B and activator-protein 1 (AP-1), leading to the activation of cytokine gene expression⁴¹. It has been reported that LPS-stimulated CD14 expression and toll-like receptor (TLR)-4-induced signaling pathways are down-regulated by *n*-3 PUFA⁴², providing mechanisms through which EPA may exert its effects on both AP-1 and NF- κ B activation.

Although DHA may have similar anti-inflammatory effects as EPA, it does not act by direct competition with AA. DHA can decrease the release of AA from membrane phospholipids by decreasing phospholipase A2 activity, and decreasing the responsiveness of TLR-4 to LPS, thereby suppressing NF- κ B activation and subsequent inflammatory gene transcription¹¹. Komatsu et al.⁴³ has shown that 60 μ M of DHA inhibited NF- κ B activity in interferon- γ and LPS-stimulated RAW

macrophages, while Weldon et al. recently demonstrated that DHA was more effective than EPA in inhibiting p65 expression and increased I κ B α expression in LPS-stimulated THP-1 monocytes. Interestingly, Novak and coworkers²⁰ have reported reduced LPS-stimulated RAW macrophage NF- κ B activity pre-treated with an *n*-3 PUFA lipid emulsion (Omegaven®). The inhibition of NF- κ B activity by either EPA or DHA is important especially since AM ϕ and bronchial epithelial cells from stable asthmatics exhibit increased NF- κ B activity compared with cells from healthy individuals³⁷, and therefore it has been suggested that NF- κ B plays a pivotal role in the pathogenesis of asthma^{37, 44}.

Interestingly, Li and coworkers⁴⁵ recently demonstrated that EPA and DHA down-regulate LPS-induced activation of NF- κ B via a peroxisome proliferator-activated receptor (PPAR)- γ -dependent pathway in human kidney cells. These results suggest that PPAR- γ activation by EPA and DHA may be one of the underlying mechanisms for the beneficial effects of fish oil. Due to the fact that a new class of mediator families derived from fish oil, the EPA-and DHA-derived resolvins (RvE1 and RvD1) and the DHA-derived protectin (PD1), which act locally, and possess potent anti-inflammatory novel bioactions, suggest potentially novel therapeutic treatment strategies for asthma¹². In addition, since we have previously shown that the amount of LTB₅ generated from activated PMNLs was markedly increased following fish oil supplementation in asthmatic patients¹⁶, further studies should investigate the effect of fish oil on EPA and DHA- derived (anti-inflammatory) mediators generated from human asthmatic AM ϕ ⁴⁶.

In conclusion the present study has shown for the first time that EPA is a more potent inhibitor than DHA of inflammatory responses in human asthmatic AM ϕ cells. In addition, the present study has shown that the greater the EPA content of a fish oil formulation the greater the inhibition of the inflammatory response. Elucidating the mechanism of this modulation could help us to understand how dietary *n*-3 PUFA achieves their specific effects on airway inflammation in asthmatic individuals. The clinical relevance of the present study, along with previous work from our group^{15, 16}, suggest that EPA-rich fish oil may provide a therapeutic option for adults with asthma. In view of the clinical consequences,

these findings point towards prophylactic and acute therapeutic effects of fish oil supplementation in inflammatory diseases such as asthma. It is possible that pharmacotherapy could be decreased in some patients with asthma in concert with increased fish-oil ingestion if both the drug and *n*-3 PUFA are exerting their therapeutic effects through the same molecular actions.

Conflict of Interest

The authors report no conflict of interest.

Acknowledgments

The authors would like to thank the participating asthmatic adults for volunteering to this study.

References

1. Babcock TA, Helton WS, Anwar KN, *et al.* Synergistic anti-inflammatory activity of omega-3 lipid and rofecoxib pretreatment on macrophage proinflammatory cytokine production occurs via divergent NF-kappaB activation. *JPEN J Parenter Enteral Nutr* 2004;**28**(4):232-9; discussion 239-40.
2. Babcock TA, Helton WS, Hong D, *et al.* Omega-3 fatty acid lipid emulsion reduces LPS-stimulated macrophage TNF-alpha production. *Surg Infect (Larchmt)* 2002;**3**(2):145-9.
3. Babcock TA, Novak T, Ong E, *et al.* Modulation of lipopolysaccharide-stimulated macrophage tumor necrosis factor-alpha production by omega-3 fatty acid is associated with differential cyclooxygenase-2 protein expression and is independent of interleukin-10. *J Surg Res* 2002;**107**(1):135-9.
4. Endres S, Ghorbani R, Kelley VE, *et al.* The effect of dietary supplementation with n-3 polyunsaturated fatty acids on the synthesis of interleukin-1 and tumor necrosis factor by mononuclear cells. *N Engl J Med* 1989;**320**(5):265-71.
5. Endres S, Meydani SN, Ghorbani R, *et al.* Dietary supplementation with n-3 fatty acids suppresses interleukin-2 production and mononuclear cell proliferation. *J Leukoc Biol* 1993;**54**(6):599-603.
6. Lee TH, Hoover RL, Williams JD, *et al.* Effect of dietary enrichment with eicosapentaenoic and docosahexaenoic acids on in vitro neutrophil and monocyte leukotriene generation and neutrophil function. *N Engl J Med* 1985;**312**(19):1217-24.
7. Lo CJ, Chiu KC, Fu M, *et al.* Fish oil decreases macrophage tumor necrosis factor gene transcription by altering the NF kappa B activity. *J Surg Res* 1999;**82**(2):216-21.

- 1 8. Thien FC, Hallsworth MP, Soh C, *et al.* Effects of exogenous eicosapentaenoic acid on
2 generation of leukotriene C4 and leukotriene C5 by calcium ionophore-activated human
3 eosinophils in vitro. *J Immunol* 1993;**150**(8 Pt 1):3546-52.
- 4 9. Gorjao R, Verlengia R, Lima TM, *et al.* Effect of docosahexaenoic acid-rich fish oil
5 supplementation on human leukocyte function. *Clin Nutr* 2006;**25**(6):923-38.
- 6 10. Weldon SM, Mullen AC, Loscher CE, *et al.* Docosahexaenoic acid induces an anti-
7 inflammatory profile in lipopolysaccharide-stimulated human THP-1 macrophages more
8 effectively than eicosapentaenoic acid. *J Nutr Biochem* 2007;**18**(4):250-8.
- 9 11. Lee JY, Zhao L, Youn HS, *et al.* Saturated fatty acid activates but polyunsaturated fatty
10 acid inhibits Toll-like receptor 2 dimerized with Toll-like receptor 6 or 1. *J Biol Chem*
11 2004;**279**(17):16971-9.
- 12 12. Serhan CN. Resolution phase of inflammation: novel endogenous anti-inflammatory and
13 proresolving lipid mediators and pathways. *Annu Rev Immunol* 2007;**25**:101-37.
- 14 13. Mickleborough TD, Rundell KW. Dietary polyunsaturated fatty acids in asthma- and
15 exercise-induced bronchoconstriction. *Eur J Clin Nutr* 2005;**59**(12):1335-46.
- 16 14. Reisman J, Schachter HM, Dales RE, *et al.* Treating asthma with omega-3 fatty acids:
17 where is the evidence? A systematic review. *BMC Complement Altern Med* 2006;**6**:26.
- 18 15. Mickleborough TD, Murray RL, Ionescu AA, *et al.* Fish Oil Supplementation Reduces
19 Severity of Exercise-induced Bronchoconstriction in Elite Athletes. *Am J Respir Crit*
20 *Care Med* 2003;**168**(10):1181-9.
- 21 16. Mickleborough TD, Lindley MR, Ionescu AA, *et al.* Protective effect of fish oil
22 supplementation on exercise-induced bronchoconstriction in asthma. *Chest*
23 2006;**29**(1):39-49.

- 1 17. Akabas SR, Deckelbaum RJ. Summary of a workshop on n-3 fatty acids: current status of
2 recommendations and future directions. *Am J Clin Nutr* 2006;**83**(6 Suppl):1536S-1538S.
- 3 18. Hallstrand TS, Moody MW, Aitken ML, *et al.* Airway immunopathology of asthma with
4 exercise-induced bronchoconstriction. *J Allergy Clin Immunol* 2005;**116**(3):586-93.
- 5 19. Davies JQ, Gordon S. Isolation and culture of human macrophages. *Methods Mol Biol*
6 2005;**290**:105-16.
- 7 20. Novak TE, Babcock TA, Jho DH, *et al.* NF-kappa B inhibition by omega -3 fatty acids
8 modulates LPS-stimulated macrophage TNF-alpha transcription. *Am J Physiol Lung Cell*
9 *Mol Physiol* 2003;**284**(1):L84-9.
- 10 21. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
11 quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001;**25**(4):402-8.
- 12 22. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR.
13 *Nucleic Acids Res* 2001;**29**(9):e45.
- 14 23. Caughey GE, Mantzioris E, Gibson RA, *et al.* The effect on human tumor necrosis factor
15 alpha and interleukin 1 beta production of diets enriched in n-3 fatty acids from vegetable
16 oil or fish oil. *Am J Clin Nutr* 1996;**63**(1):116-22.
- 17 24. Gallai V, Sarchielli P, Trequattrini A, *et al.* Cytokine secretion and eicosanoid production
18 in the peripheral blood mononuclear cells of MS patients undergoing dietary
19 supplementation with n-3 polyunsaturated fatty acids. *J Neuroimmunol* 1995;**56**(2):143-
20 53.
- 21 25. Blok WL, Deslypere JP, Demacker PN, *et al.* Pro- and anti-inflammatory cytokines in
22 healthy volunteers fed various doses of fish oil for 1 year. *Eur J Clin Invest*
23 1997;**27**(12):1003-8.

- 1 26. Kew S, Mesa MD, Tricon S, *et al.* Effects of oils rich in eicosapentaenoic and
2 docosahexaenoic acids on immune cell composition and function in healthy humans. *Am*
3 *J Clin Nutr* 2004;**79**(4):674-81.
- 4 27. Schmidt EB, Varming K, Moller JM, *et al.* No effect of a very low dose of n-3 fatty acids
5 on monocyte function in healthy humans. *Scand J Clin Lab Invest* 1996;**56**(1):87-92.
- 6 28. Chu AJ, Walton MA, Prasad JK, *et al.* Blockade by polyunsaturated n-3 fatty acids of
7 endotoxin-induced monocytic tissue factor activation is mediated by the depressed
8 receptor expression in THP-1 cells. *J Surg Res* 1999;**87**(2):217-24.
- 9 29. Zhao Y, Chen LH. Eicosapentaenoic acid prevents lipopolysaccharide-stimulated DNA
10 binding of activator protein-1 and c-Jun N-terminal kinase activity. *J Nutr Biochem*
11 2005;**16**(2):78-84.
- 12 30. Zhao Y, Joshi-Barve S, Barve S, *et al.* Eicosapentaenoic acid prevents LPS-induced
13 TNF-alpha expression by preventing NF-kappaB activation. *J Am Coll Nutr*
14 2004;**23**(1):71-8.
- 15 31. Zhao G, Etherton TD, Martin KR, *et al.* Anti-inflammatory effects of polyunsaturated
16 fatty acids in THP-1 cells. *Biochem Biophys Res Commun* 2005;**336**(3):909-17.
- 17 32. Alam R. Chemokines in allergic inflammation. *J Allergy Clin Immunol* 1997;**99**(3):273-
18 7.
- 19 33. Johnston SL, Freezer NJ, Ritter W, *et al.* Prostaglandin D2-induced bronchoconstriction
20 is mediated only in part by the thromboxane prostanoid receptor. *Eur Respir J*
21 1995;**8**(3):411-5.

- 1 34. Sousa AR, Lane SJ, Nakhosteen JA, *et al.* Expression of interleukin-1 beta (IL-1beta) and
2 interleukin-1 receptor antagonist (IL-1ra) on asthmatic bronchial epithelium. *Am J Respir*
3 *Crit Care Med* 1996;**154**(4 Pt 1):1061-6.
- 4 35. Gosset P, Tsicopoulos A, Wallaert B, *et al.* Increased secretion of tumor necrosis factor
5 alpha and interleukin-6 by alveolar macrophages consecutive to the development of the
6 late asthmatic reaction. *J Allergy Clin Immunol* 1991;**88**(4):561-71.
- 7 36. Shah A, Church MK, Holgate ST. Tumour necrosis factor alpha: a potential mediator of
8 asthma. *Clin Exp Allergy* 1995;**25**(11):1038-44.
- 9 37. Hart LA, Krishnan VL, Adcock IM, *et al.* Activation and Localization of Transcription
10 Factor, Nuclear Factor-kappa B, in Asthma. *Am. J. Respir. Crit. Care Med.*
11 1998;**158**(5):1585-1592.
- 12 38. Khalfoun B, Thibault F, Watier H, *et al.* Docosahexaenoic and eicosapentaenoic acids
13 inhibit in vitro human endothelial cell production of interleukin-6. *Adv Exp Med Biol*
14 1997;**400B**:589-97.
- 15 39. Moon Y, Pestka JJ. Deoxynivalenol-induced mitogen-activated protein kinase
16 phosphorylation and IL-6 expression in mice suppressed by fish oil. *J Nutr Biochem*
17 2003;**14**(12):717-26.
- 18 40. Goldman DW, Pickett WC, Goetzl EJ. Human neutrophil chemotactic and degranulating
19 activities of leukotriene B5 (LTB5) derived from eicosapentaenoic acid. *Biochem*
20 *Biophys Res Commun* 1983;**117**(1):282-8.
- 21 41. Pelaia G, Cuda G, Vatrella A, *et al.* Mitogen-activated protein kinases and asthma. *J Cell*
22 *Physiol* 2005;**202**(3):642-53.

42. Lee JY, Plakidas A, Lee WH, *et al.* Differential modulation of Toll-like receptors by fatty acids: preferential inhibition by n-3 polyunsaturated fatty acids. *J. Lipid Res.* 2003;**44**(3):479-486.
43. Komatsu W, Ishihara K, Murata M, *et al.* Docosahexaenoic acid suppresses nitric oxide production and inducible nitric oxide synthase expression in interferon-gamma plus lipopolysaccharide-stimulated murine macrophages by inhibiting the oxidative stress. *Free Radic Biol Med* 2003;**34**(8):1006-16.
44. Gagliardo R, Chanez P, Mathieu M, *et al.* Persistent activation of nuclear factor-kappaB signaling pathway in severe uncontrolled asthma. *Am J Respir Crit Care Med* 2003;**168**(10):1190-8.
45. Li H, Ruan XZ, Powis SH, *et al.* EPA and DHA reduce LPS-induced inflammation responses in HK-2 cells: evidence for a PPAR-gamma-dependent mechanism. *Kidney Int* 2005;**67**(3):867-74.
46. Miles EA, Allen E, Calder PC. In vitro effects of eicosanoids derived from different 20-carbon Fatty acids on production of monocyte-derived cytokines in human whole blood cultures. *Cytokine* 2002;**20**(5):215-23.

Figure Legends

Figure 1. Effect 120 μ M EPA-rich media (EPAX 4510), pure EPA, pure DHA, DHA-rich media (EPAX 1050) and LipovenosR (*n*-6 PUFA) on LTB₄ (panel A), PGD₂ (panel B), TNF- α (panel C) and IL-1 β (panel D) production by LPS-stimulated human AM ϕ . The data represent mean $\pm$ SD from six separate experiments, each done in triplicate. * $p < 0.05$ vs. DMEM + LPS. Ψ $p < 0.05$ vs. 25 μ M treatment within each inflammatory mediator. A difference in letter (a through e) designates significant differences ($p < 0.05$) between treatments (EPAX 4510, EPA, DHA, EPAX 1050 and Lipovenos[®]).

Figure 2. Effect 25 μ M EPA-rich media (EPAX 4510), pure EPA, pure DHA, DHA-rich media (EPAX 1050) and LipovenosR (*n*-6 PUFA) on LTB₄ (panel A) and TNF- α (panel B) production by LPS-stimulated human AM ϕ . The data represent mean $\pm$ SD from six separate experiments, each done in triplicate. * $p < 0.05$ vs. DMEM + LPS. A difference in letter (a through e) designates significant differences ($p < 0.05$) between treatments (EPAX 4510, EPA, DHA, EPAX 1050 and Lipovenos[®]).

Figure 3. Effect 120 μ M (panel A) and 25 μ M (panel B) EPA-rich media (EPAX 4510), pure EPA, pure DHA, DHA-rich media (EPAX 1050) and LipovenosR (*n*-6 PUFA) on TNF- α and IL-1 β mRNA levels in LPS-stimulated human alveolar macrophages. TNF- and IL-1 mRNA levels were normalized to GAPDH and expressed relative to DMEM. The data represent mean $\pm$ SD. * $p < 0.05$ relative to DMEM. Ψ $p < 0.05$ vs. 25 μ M treatment within each cytokine. A difference in letter (a through e) designates a significant difference ($p < 0.05$) between each treatment (EPAX 4510, EPA, DHA, EPAX 1050 and Lipovenos[®]).