This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Electron transfer reactions in ternary systems on silica gel surfaces: evidence for radical cation diffusion

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1039/c0pp00049c>

PUBLISHER

© Royal Society of Chemistry and Owner Societies

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at:
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Williams, Sian L., Iain Kirkpatrick, and David R. Worrall. 2019. "Electron Transfer Reactions in Ternary Systems on Silica Gel Surfaces: Evidence for Radical Cation Diffusion". figshare.
<https://hdl.handle.net/2134/16134>.

Electron Transfer Reactions in Ternary Systems on Silica Gel Surfaces: Evidence for Radical Cation Diffusion

Sian L. Williams,* Iain Kirkpatrick, and David R. Worrall

Received (in XXX, XXX) Xth XXXXXXXXX 200X, Accepted Xth XXXXXXXXX 200X

First published on the web Xth XXXXXXXXX 200X

DOI: 10.1039/b000000x

Electron transfer reactions have been studied between 9-anthracenecarboxylic acid co-adsorbed with perylene on silica gel surfaces employing azulene as a molecular shuttle in order to facilitate hole transfer. In this paper we present for the first time a ternary system that unambiguously demonstrates an appreciable mobility of radical cations on the silica gel surface. Rates of hole transfer from the 9-anthracenecarboxylic acid radical cation to perylene via azulene have been studied using diffuse reflectance laser flash photolysis spectroscopy. Azulene has been shown to enhance the rate of electron transfer in the ternary system, demonstrating significant mobility of the azulene and its radical cation species on silica gel surfaces. The data shows that the azulene radical cation can diffuse at an appreciable rate on the silica gel surface.

Introduction

The photochemistry and photophysics of molecules adsorbed to oxide surfaces and contained within zeolites has been the subject of a number of studies¹⁻⁴². The rapid mobility of molecules on oxide surfaces has been previously reported^{2,6-11} and studied through both energy and electron transfer reactions. The latter have also been studied on other porous materials such as clays^{2,43-46}. Silica gel, which was employed in this study, is widely used in industry as a catalytic support and a greater understanding of reaction kinetics on these materials is significant in optimising these applications.

Multi-photon ionisation of arenes and the subsequent electron transfer reactions of their radical cations on silica gel have been previously reported^{3,4,13,14} and the influence of the nature and loading of the electron donor and electron acceptor compounds on the reaction rate investigated. The kinetics of decay of radical ions on silica gel is not straightforward since the surface is heterogeneous in nature and rates rarely conform to simple exponential kinetics. We have used the model described by Alberly *et al.*^{4,12-15,50} to characterise our data sets, ensuring a comprehensive exploration of the parameter space in order to obtain a global optimum value for the rate constants^{13,14}.

We have previously shown that energy and electron transfer reactions in these systems are dependent upon the rate of diffusion^{13,14}, and in the case of azulene as energy acceptor / electron donor, rates of reaction are largely determined by the rate of diffusion of this relatively more mobile species. We have demonstrated also that energetics play a role in determining electron transfer rates, and have shown a Marcus-type dependence of rate on the free energy for electron transfer^{4,14}. In this study we also found evidence for some steric influence on reaction rates.

In previous work we studied the activation energy of back electron transfer of some derivatised anthracenes^{12,15}. There was a correlation between the activation energy and the size of the radical cation which suggested that movement of the radical cation and not of the electron dominated the observed kinetics. In this paper we present for the first time a ternary

system that unambiguously demonstrates the mobility of radical cations on the silica gel surface.

Experimental

Samples were prepared as follows: Silica gel (Davisil grade 635, 60-100 mesh, 6 nm pore size, surface area 480 m² g⁻¹; Aldrich Chemical Co.) was dried at a temperature of 125 °C under a vacuum of 5 x 10⁻⁵ mbar for 8 hours, and the vessel re-pressurised with dry nitrogen. 9-anthracenecarboxylic acid (minimum 99%, Aldrich Chemical Co.), azulene (99%, Aldrich Chemical Co.) and perylene (99%, Aldrich Chemical Co.) were dissolved in acetonitrile (spectrophotometric grade, Aldrich Chemical Co.) and a known weight of the resulting solution added to the dried silica gel. The mixture was allowed to equilibrate for a period of one hour, with periodic agitation. The solvent was then removed under vacuum to a pressure of 5 x 10⁻⁵ mbar. The resulting sample was then sealed under vacuum into a cylindrical glass cuvette (22 mm diameter x 10 mm path length). Sample loadings were determined from the mass of solution added to the silica gel. The loadings were varied between 0.1 μmol g⁻¹ and 2.5 μmol g⁻¹ for all three components. These concentrations correspond to low surface coverages, with the highest being less than 0.1% of a homogenous 2D surface monolayer.

Ground state diffuse reflectance spectra were recorded using a Perkin-Elmer Lambda Bio 40 spectrophotometer equipped with a Spectralon integrating sphere using barium sulphate as a reference.

The nanosecond diffuse reflectance laser flash photolysis apparatus has been described previously in references 16 and 27. In the flash photolysis experiments, excitation of the samples was with the third (355 nm, 110 mJ/pulse) harmonic of a Surelite I Nd:YAG laser (Continuum). The pulse energy was attenuated using solutions of sodium nitrite in water, to ensure that transient reflectance changes were kept below 10%, where the change in reflectance is directly proportional to the concentration of transient species⁴⁷⁻⁴⁹. Diffusely reflected analysing light from a 275 W xenon arc lamp (Oriel) was collected and focussed onto the entrance slit of an f/3.4

grating monochromator (Applied Photophysics) and detected with a side-on photomultiplier tube (Hamamatsu R928). Signal capture was by an LT364 Waverunner digitising oscilloscope (LeCroy), interfaced to an IBM-compatible PC.

Transient decay data has been analysed using a model which assumes a log Gaussian distribution of rate constants described previously^{13,14,50} according to equation 1

$$\frac{C}{C_0} = \frac{\int_{-\infty}^{\infty} \exp(-t^2) \exp\left[-\bar{k} t \exp(\gamma t)\right] dt}{\int_{-\infty}^{\infty} \exp(-t^2) dt} \quad (1)$$

which was transformed to have finite integrals as described in the appendix of reference⁵⁰. Here C and C_0 are transient concentrations at times $t = t$ and $t = 0$ after the laser pulse, γ is the width of the distribution and $\bar{k}$ is the mean rate constant. At low sample loadings and small reflectance changes (less than 10%), C and C_0 can be replaced by ΔR and ΔR_0 , the reflectance changes at times $t = t$ and $t = 0$ (relative to the laser pulse) respectively.

Results and Discussion

Transient absorption spectra

Transient absorption spectra were obtained using 355 nm laser excitation of the respective samples. Fig. 1 shows the transient difference spectrum obtained from photolysis of 9-anthracenecarboxylic acid co-adsorbed with perylene on silica gel following laser excitation at 355 nm. Anthracene triplet (420 nm) and radical cation (400 nm and 710 nm) absorption bands were assigned based upon previous measurements^{3,4,13,14}.

Fig. 1 Transient difference spectra of 9-anthracenecarboxylic acid ($2.5 \mu\text{mol g}^{-1}$) co-adsorbed with perylene ($0.25 \mu\text{mol g}^{-1}$) on silica gel at 2 ms (solid line), 10 ms (dashed line) and 50 ms (dotted line) following laser excitation at 355 nm.

The perylene radical cation (530 nm) absorption band was assigned by comparison to literature data of measurements

carried out on surfaces and in solution⁵¹⁻⁵⁴. The 9-anthracenecarboxylic acid triplet (420 nm) decays quite rapidly by both unimolecular decay and by energy transfer to perylene.

Note that in this study 9-carboxylic acid anthracene is used since it has been demonstrated to be immobile on the silica gel surface¹³ as evidenced by the lack of delayed fluorescence following laser excitation. Hence any dynamic quenching observed is the result of the diffusional characteristics of the co-adsorbed species. The perylene radical cation (530 nm) appears concomitantly with the anthracene radical cation (710 nm) decay. Kinetic analysis of these species shows that, as expected, the rate of appearance of the perylene radical cation and disappearance of the anthracene radical cation are the same to within experimental error (see Table 1).

Co-adsorption of azulene into the 9-anthracenecarboxylic acid / perylene system changes the observed electron transfer kinetics (*vide infra*) but does not affect the appearance of the transient difference spectra owing to both the azulene triplet and radical cation absorptions being outside the spectral range studied.

Electron Transfer Kinetics

The kinetics of electron transfer for 9-anthracenecarboxylic acid / perylene systems in both the presence and absence of varying loadings of azulene on silica gel have been measured following laser excitation at 355 nm. The kinetic data was analysed as in previous publications^{13,14} using the dispersive kinetic model of Albery et al⁵⁰ which provides analytical parameters with which to characterise the decay or (in the case of perylene) rise rate constant as a function of concentration.

The extracted parameters are then analysed to yield a bimolecular rate constant characteristic of the process according to equation 2:

$$\bar{k} = \bar{k}_0 + k_q [\text{quencher}] \quad (2)$$

where $\bar{k}_0$ is the unquenched decay rate constant, and k_q is the bimolecular quenching constant. Fig. 2 shows a plot of fitted $\bar{k}$ versus quencher concentration for electron transfer in the 9-anthracenecarboxylic acid / azulene / perylene system. The addition of azulene to the system decouples the decay of the anthracene radical cation and the rise of the perylene radical cation. The rate of electron transfer from the azulene to the 9-anthracene carboxylic acid radical is very rapid due to the large diffusion coefficient for the azulene on silica gel. As we have shown in a previous publication¹⁴, molecular size plays a role in determining diffusion coefficients on silica gel, and hence the rate of diffusion of the azulene is significantly faster than that of the perylene. Given the 9-anthracenecarboxylic acid is, as previously demonstrated, hydrogen bonded and therefore fixed to the silica gel surface, the rate of electron transfer is governed in part by how quickly the azulene diffuses across the silica gel surface, and as shown previously¹⁴ there are also energetic considerations (*vide infra*).

In the absence of azulene the bimolecular rate constant for both perylene radical cation formation and 9-anthracenecarboxylic acid radical cation decay are the same to within experimental error, and the electron transfer rate in this case is governed by the rate of diffusion of the perylene. The introduction of azulene to the system causes the bimolecular rate constants for both the formation of perylene radical cation and the decay of the anthracene radical cation species to increase, but not to the same extent as would be the case if the mobility of the azulene radical were the same as for free azulene and this alone was the determining factor in the

Fig. 2 Electron transfer plots for azulene to 9-anthracenecarboxylic acid (CAn) radical cation (●) and perylene (Pe) to azulene radical cation (○) following laser excitation at 355 nm as a function of azulene concentration.

electron transfer rate. However, were the azulene radical to be immobile on the surface there may be expected to be no enhancement to the perylene rise kinetics on addition of azulene based on purely diffusional arguments. As previously mentioned, energetic factors have been shown to influence electron transfer rates and a Marcus dependence was shown for a range of electron donors^{4,14}. In this instance the oxidation potential of the perylene (0.92 V vs Ag/AgCl⁵⁵, relative to 1.26 V for anthracene and 1.04 V for azulene versus the same reference electrode¹⁴) is such that the free energy for electron transfer between the azulene radical cation and perylene is lower than that between the anthracene radical cation and perylene, which would suggest that on energetic grounds electron transfer between the former two should be slower than between the latter (see figure 8 in reference 14).

Additionally the similarity in free energy for electron transfer in the 9-anthracenecarboxylic acid / azulene and 9-anthracenecarboxylic acid / perylene couples shows that the slow rate of electron transfer in the anthracene / perylene system is a result of slow diffusion by the perylene rather than being dominated by energetic considerations. Therefore the observed *increased* rate of electron transfer to perylene observed in the presence of azulene must be ascribable to

considerable mobility of the azulene radical cation, where diffusion of the azulene radical to the perylene is faster than diffusion of the perylene to the azulene radical. Even allowing for energetic effects, however, the rate of diffusion of the azulene radical is less than that of the free azulene, since an order of magnitude reduction in the electron transfer rate would not be expected on energetic grounds alone. However, it clearly has a larger diffusion coefficient than perylene demonstrating the complex interaction of parameters which determine diffusion rates on silica gel. A summary of the measured bimolecular rate constants are given in Table 1.

The table shows that the introduction of azulene in the perylene / 9-anthracenecarboxylic acid system greatly increases the rates of electron transfer in this ternary system, demonstrating the effectiveness of a small molecule as a shuttle of, in this instance, a hole between two relatively immobile species separated by a relatively large distance. This constitutes to our knowledge the first unambiguous demonstration of the mobility of radical cations on the silica gel surface, and also the first demonstration of a molecular shuttle of this type in an amorphous system.

Table 1 Bimolecular rate constants for electron transfer between azulene (Az) and the 9-anthracenecarboxylic acid (An-9-CA) radical cation and perylene (Pe) and the azulene radical cation, where *x* denotes the fact that the concentration of that species is changing.

System	$k_{\text{et}} / \text{g mol}^{-1} \text{s}^{-1}$	
	Perylene ^{•+} rise 530 nm	Anthracene ^{•+} decay 710 nm
1 An-9-CA : <i>x</i> Pe	$8.3 \pm 0.8 \times 10^7$	$6.5 \pm 0.8 \times 10^7$
1 An-9-CA : <i>x</i> Az : 1 Pe	$1.1 \pm 0.2 \times 10^9$	$1.4 \pm 0.3 \times 10^{10}$

Conclusions

We have demonstrated electron transfer in this ternary system using azulene for hole transfer between 9-anthracenecarboxylic acid radical cation as an electron acceptor and perylene as an electron donor. The radical cation species of both perylene and 9-anthracenecarboxylic acid can be seen in the transient absorption spectra. The absorption bands are well separated from one another as well as that of the triplet excited states of the parent species and hence their kinetics can be independently measured.

The presence of an azulene molecular hole shuttle has been shown to substantially enhance the rate of electron transfer. From comparison of bimolecular rate constants and consideration of possible energetic influences, mobility of the azulene radical cation is clearly demonstrated, with its rate of diffusion being still rapid relative to perylene even though it is shown to be slower than for the neutral azulene species.

This study shows the potential of such shuttles as electron transfer catalysts and offers a means of controlling charge separation times in these systems.

Notes and references

1. J.K. Thomas and E.H. Ellison, Various aspects of the constraints imposed on the photochemistry of systems in porous silica, *Adv. Colloid Interface Sci.*, 2001, **89**, 195-238.
2. J.K. Thomas, Physical Aspects of Radiation-Induced Processes on SiO₂, g-Al₂O₃, Zeolites, and Clays, *Chem. Rev.*, 2005, **105**, 1683-1734.
3. F. Wilkinson, D.R. Worrall and S.L. Williams, Primary Photochemical Processes of Anthracene Adsorbed on Silica-Gel, *J. Phys. Chem.*, 1995, **99**, 6689-6696.
4. D.R. Worrall, S.L. Williams and F. Wilkinson, Electron transfer reactions of anthracene adsorbed on silica gel, *J. Phys. Chem. B*, 1997, **101**, 4709-4716.
5. D. Oelkrug, S. Reich, F. Wilkinson and P.A. Leicester, Photoionization on Insulator Surfaces - Diffuse Reflectance Laser Flash-Photolysis of Distyrylbenzenes Adsorbed on Silica and Alumina, *J. Phys. Chem.*, 1991, **95**, 269-274.
6. D. Oelkrug, S. Uhl, F. Wilkinson and C.J. Willsher, Bonding and mobility of acridine on some metal-oxide surfaces as studied by delayed fluorescence and transient absorption-spectroscopy, *J. Phys. Chem.*, 1989, **93**, 4551-4556.
7. R.W. Kessler and F. Wilkinson, Diffuse reflectance triplet-triplet absorption-spectroscopy of aromatic-hydrocarbons chemisorbed on γ -alumina, *J. Chem. Soc., Faraday Trans. 1*, 1981, **77**, 309-320.
8. I.K. Lednev, N. Mathivanan and L.J. Johnston, Photochemistry of Stilbene Adsorbed on Silica-Gel and Nax Zeolite - a Diffuse-Reflectance Laser Flash-Photolysis Study, *J. Phys. Chem.*, 1994, **98**, 11444-11451.
9. G.J. Kavarnos and N.J. Turro, Photosensitization by reversible electron-transfer - theories, experimental-evidence, and examples, *Chem. Rev.*, 1986, **86**, 401-449.
10. A. Liu, M.C. Sauer, D.M. Loffredo and A.D. Trifunac, Transient absorption spectra of aromatic radical cations in hydrocarbon solutions, *J. Photochem. Photobiol. A:Chem.*, 1992, **67**, 197-208.
11. D.R. Worrall, S.L. Williams and F. Wilkinson, Electron transfer on insulator surfaces: Exciplex emission and the role of electron diffusion in determining radical deactivation rates, *J. Phys. Chem. A*, 1998, **102**, 5484-5490.
12. D.R. Worrall, S.L. Williams, F. Wilkinson, J.E. Crossley, H. Bouas-Laurent and J.P. Desvergne, Spectroscopy and ion-electron recombination kinetics of radical ions of anthracenes and substituted anilines on silica gel, *J. Phys. Chem. B*, 1999, **103**, 9255-9261.
13. D.R. Worrall, I. Kirkpatrick and S.L. Williams, Controlling factors in electron and energy transfer reactions on silica gel surfaces, *Photochem. Photobiol. Sci.*, 2002, **1**, 896-901.
14. D.R. Worrall, I. Kirkpatrick and S.L. Williams, Bimolecular processes on silica gel surfaces: energetic factors in determining electron-transfer rates, *Photochem. Photobiol. Sci.* 2004, **3**, 63-70.
15. D.R. Worrall, S.L. Williams and T. Ganguly, Ion-electron recombination on silica gel surfaces: experiment and modelling, *Photochem. Photobiol. Sci.* 2006, **5**, 844-849.
16. D.R. Worrall and S.L. Williams, Diffuse-Reflectance Laser Flash Photolysis, *Encyclopedia of Modern Optics Volume 1*, Guenther, R.D., Steel, D.G and Bayvel, L. (eds), Elsevier, Oxford, 2004, pp 31-38, ISBN 0-12-227600-0.
17. S. Pankasem and J.K. Thomas, Pyrene, Pyrene Derivatives, and 1,1'-Binaphthyl as Luminescent Probes for Photophysical Studies of Alumina Surfaces, *J. Phys. Chem.*, 1991, **95**, 7385-7393.
18. S. Pankasem and J.K. Thomas, Reflectance Spectroscopic Studies of the Cation Radical and the Triplet of Pyrene on Alumina, *J. Phys. Chem.*, 1991, **95**, 6990-6996.
19. G. Beck and J.K. Thomas, Photochemical Processes on Aluminum-Oxide - a Pulsed Laser Study of Photoinduced Reactions, *Chem. Phys. Lett.*, 1983, **94**, 553-557.
20. D. Oelkrug, G. Krabichler, W. Honnen, F. Wilkinson and C.J. Willsher, Photophysical Behavior of Diphenylpolyenes Adsorbed on Alumina by Diffuse Reflectance Laser Flash-Photolysis, *J. Phys. Chem.*, 1988, **92**, 3589-3594.
21. Y. Mao and J.K. Thomas, Photochemical-Reactions of Biphenyl and Its Chlorinated Derivatives on the Surfaces of Gamma-Alumina and Silica-Alumina, *J. Chem. Soc.-Faraday Trans.*, 1992, **88**, 3079-3086.
22. Y. Mao and J.K. Thomas, Photochemical-Reactions of Pyrene on Surfaces of Gamma-Alumina and Silica-Alumina, *Langmuir*, 1992, **8**, 2501-2508.
23. K.K. Iu and J.K. Thomas, Photophysical Properties of Pyrene in Zeolites .2. Effects of Coadsorbed Water, *Langmuir*, 1990, **6**, 471-478.
24. L.J. Johnston, J.C. Scaiano, J.L. Shi, W. Siebrand and F. Zerbetto, Observation and Modeling of the Recombination Kinetics of Diphenylmethyl Radicals in the Cavities of Na-X Zeolite, *J. Phys. Chem.*, 1991, **95**, 10018-10024.
25. S. Hashimoto, N. Fukazawa, H. Fukumura and H. Masuhara, Observation and Characterization of Excimer Emission from Anthracene Included in Nax Zeolite, *Chem. Phys. Lett.*, 1994, **219**, 445-451.
26. X.S. Liu, K.K. Iu, J.K. Thomas, H.Y. He and J. Klinowski, Spectroscopic Studies of Protonated Aromatic Species and Radical Cations in H⁺-Zeolites, *J. Am. Chem. Soc.*, 1994, **116**, 11811-11818.
27. F. Wilkinson and R. Beer, Diffuse reflectance laser photolysis of adsorbed molecules, in *Photochemical processes in organised molecular systems*, K. Honda, Editor, 1991, Elsevier Science: Amsterdam. p. 377-398.
28. G.H. Zhang, J.K. Thomas, A. Eremenko, T. Kikteva and F. Wilkinson, Photoinduced charge-transfer reaction between pyrene and N,N'-dimethylaniline on silica gel surfaces, *J. Phys. Chem. B*, 1997, **101**, 8569-8577.
29. D. Oelkrug, M. Gregor and S. Reich, Translational Mobility in the Adsorbed State - Bimolecular Deactivation Kinetics of Laser-Excited Azaaromatics on Microporous Silica, *Photochem. Photobiol.*, 1991, **54**, 539-546.
30. D. Oelkrug, S. Uhl, F. Wilkinson and C.J. Willsher, Bonding and Mobility of Acridine on Some Metal-Oxide Surfaces as Studied by Delayed Fluorescence and Transient Absorption- Spectroscopy, *J. Phys. Chem.*, 1989, **93**, 4551-4556.
31. R. Krasnansky, K. Koike and J.K. Thomas, Gaussian Approximation to the Unique Heterogeneous Langmuir- Hinshelwood Type Fluorescence Quenching at the Silica-Gel Gas- Solid Interface - Pyrene and 9,10-Diphenylanthracene Singlet Quenching by Oxygen, *J. Phys. Chem.*, 1990, **94**, 4521-4528.

32. S.A. Ruetten and J.K. Thomas, Fluorescence and triplet quantum yields of arenes on surfaces, *J. Phys. Chem. B*, 1998, **102**, 598-606.
33. R. Dabestani, J. Higgin, D. Stephenson, I.N. Ivanov and M.E. Sigman, Photophysical and photochemical processes of 2-methyl, 2-ethyl, and 2-tert-butylanthracenes on silica gel, A substituent effect study. *J. Phys. Chem. B*, 2000, **104**, 10235-10241.
34. H. Garcia and H.D. Roth, Generation and reactions of organic radical cations in zeolites, *Chem. Rev.*, 2002, **102**, 3947-4007.
35. M. Alvaro, P. Ateizar, A. Corma, B. Ferrer, H. Garcia and M.T. Navarro, Photochemical Generation of Electrons and Holes in Germanium-Containing ITQ-17 Zeolite, *J. Phys. Chem. B*, 2004, **109**, 3696-3700.
36. I. Batonneau-Gener, A. Moissette, C. Bremard and G. Buntinx, Time resolved resonance Raman, transient diffuse reflectance and kinetic studies of species generated by UV laser photolysis of biphenyl occluded within dehydrated Y-faujasite zeolites, *J. Photochem. Photobiol. A-Chemistry*, 2008, **195**, 156-166.
37. R. Dabestani, M. Kidder and A.C. Buchanan, Pore size effect on the dynamics of excimer formation for chemically attached pyrene on various silica surfaces, *J. Phys. Chem. C*, 2008, **112**, 11468-11475.
38. J.M. Stokke and D.W. Mazyck, Photocatalytic degradation of methanol using silica-titania composite pellets: Effect of pore size on mass transfer and reaction kinetics, *Environmental Science and Technology*, 2008, **42**, 3808-3813.
39. J.P. Da Silva, E.V. Bastos, L.V.F. Ferreira and R.G. Weiss, Surface photochemistry of the herbicide napropamide. The role of the media and environmental factors in directing the fates of intermediates, *Photochem. Photobiol. Sci.*, 2008, **7**, 69-75.
40. L.F.V. Ferreira, J.P. Da Silva, I.F. Machadoa, T.J.F. Branco and J.C. Moreira, Surface photochemistry: Dibenzo-p-dioxin adsorbed onto silicalite, cellulose and silica, *J. Photochem. Photobiol. A: Chemistry*, 2007, **186**, 254-262.
41. L.F.V. Ferreira, J.P. Da Silva, I.F. Machadoa and T.J.F. Branco, Surface photochemistry: benzophenone as a probe for the study of silica and reversed-phase silica surfaces, *Photochem. Photobiol. Sci.*, 2006, **5**, 665-673.
42. M. Alvaro, P. Ateizar, A. Corma, B. Ferrer, H. Garcia and M.T. Navarro, Photochemical Generation of Electrons and Holes in Germanium-Containing ITQ-17 Zeolite, *J. Phys. Chem. B*, 2004, **109**, 3696-3700.
43. S. Takagi, D.A. Tryk and H. Inoue, Photochemical Energy Transfer of Cationic Porphyrin Complexes on Clay Surface, *J. Phys Chem. B*, 2002, **106**, 5455-5460.
44. S. Takagi, T. Shimada, M. Eguchi, T. Yui, H. Yoshida, D.A. Tryk and H. Inoue, High-Density Adsorption of Cationic Porphyrins on Clay Layer Surfaces without Aggregation: The Size-Matching Effect, *Langmuir*, 2002, **18**, 2265-2272.
45. M. Eguchi, S. Takagi, H. Tachibana and H. Inoue, The 'size matching rule' in di-, tri-, and tetra-cationic charged porphyrin/synthetic clay complexes: effect of the inter-charge distance and the number of charged sites, *Journal of Physics and Chemistry of Solids*, 2004, **65**, 403-407.
46. S. Takagi, M. Eguchi, D.A. Tryk and H. Inoue, Light-Harvesting Energy Transfer and Subsequent Electron Transfer of Cationic Porphyrin Complexes on Clay Surfaces, *Langmuir*, 2006, **22**, 1406-1408.
47. R.W. Kessler, G. Krabichler, S. Uhl, D. Oelkrug, W.P. Hagan, J. Hyslop and F. Wilkinson, Transient Decay Following Pulse Excitation of Diffuse- Scattering Samples, *Optica Acta*, 1983, **30**(8), 1099-1111.
48. H.K.A. Kan and T.P. Lin, Calculation of reflectance of a light diffuser with non-uniform absorption, *Journal of the optical society of america*, 1970, **60**, 1252-1256.
49. D. Oelkrug, W. Honnen, F. Wilkinson and C.J. Willsher, Modeling of Transient Production and Decay Following Laser Excitation of Opaque Materials, *Journal of the Chemical Society-Faraday Transactions II*, 1987, **83**, 2081-2095.
50. W.J. Albery, P.N. Bartlett, C.P. Wilde and J.R. Darwent, A General-Model for Dispersed Kinetics in Heterogeneous Systems, *J. Am. Chem. Soc.*, 1985, **107**(7), 1854-1858.
51. J.M. Turner, M.W. Karl and J.F. Kauffman, Spectroscopic signatures of protonated perylene in concentrated sulphuric acid, *J. Photochem. Photobiol. A:Chemistry*, 2004, **163**, 433-438.
52. P. Sotero and R. Acre, Surface and adsorbates effects on the photochemistry and photophysics of adsorbed perylene on unactivated silica gel and alumina, *J. Photochem. Photobiol. A:Chemistry*, 2004, **167**, 191-199.
53. K. Kawai, N. Yamamoto and H. Tsubomura, Simultaneous Formation of Perylene Cation and Anion by Flash Excitation of Perylene in Solutions, *Bulletin of the Chemical Society of Japan*, 1970 **43**, 2266-2268.
54. R.H. Clarke and R. Hochstrasser, Location and assignment of the Lowest triplet State of Perylene, *J. Mol. Spectroscopy*, 1969, **32**, 309-319.
55. Maciej Mazur and G. J. Blanchard, Oxidative Transformations of Surface-Bound Perylene, *Langmuir*, 2005, **21**, 1441-1447.