Supporting Information

Electroreductive Acylation of Aromatic Ketones with Acylimidazoles

Naoki Kise,* Syun Agui, Shinji Morimoto and Nasuo Ueda

Department of Biotechnology, Faculty of Engineering, Tottori University,
Koyama, Tottori 680-8552, Japan

Contents

A drawing of electrolysis cell	S2
Compound characterization data	S3-8

Electrolysis Cell

Compound Characterization Data

3-Phenyl-3-(trimethylsilyloxy)butan-2-one (2a'): Colorless paste. *R_f* 0.25 (hexane-ethyl acetate, 20:1). IR (neat) 1720, 1599, 1493, 949, 843, 756, 700 cm⁻¹. ¹H NMR (CDCl₃) δ 0.19 (s, 9 H), 1.71 (s, 3 H), 2.05 (s, 3 H), 7.19-7.45 (m, 5 H). ¹³C NMR (CDCl₃) δ 2.1 (q), 24.0 (q), 25.8 (q), 83.4 (s), 124.9 (d), 127.3 (d), 128.1 (d), 143.1 (s), 209.5 (s). Anal. Calcd for C₁₅H₂₆O₂Si: C, 66.05; H, 8.53. Found: C, 66.21; H, 8.48.

2-Hydroxy-4-methyl-2-phenylpentan-3-one (2d): Colorless paste. *R_f* 0.4 (hexane-ethyl acetate, 5:1). IR (neat) 3458, 1707, 1601, 1495, 928, 779, 741, 700 cm⁻¹. ¹H NMR (CDCl₃) δ 0.67 (d, 3 H, *J* = 6.8 Hz), 1.07 (d, 3 H, *J* = 6.8 Hz), 1.79 (s, 3 H), 2.83-2.98 (m, 1 H), 4.72 (s, 1 H), 7.26-7.46 (m, 5 H). ¹³C NMR (CDCl₃) δ 19.8 (q), 20.2 (q), 23.6 (q), 34.0 (d), 79.8 (s), 126.0 (d), 127.7 (d), 128.2 (d), 140.7 (s), 215.8 (s). Anal. Calcd for C₁₂H₁₆O₂: C, 74.87; H, 8.39. Found: C, 74.66; H, 8.43.

2-Hydroxy-2-phenyldecan-3-one (2e): Colorless paste. *R_f* 0.2 (hexane-ethyl acetate, 10:1). IR (neat) 3460, 1709, 1601, 1493, 758, 700 cm⁻¹. ¹H NMR (CDCl₃) δ 0.84 (t, 3 H, *J* = 6.8 Hz), 1.00-1.29 (m, 8 H), 1.34-1.58 (m, 2 H), 1.77 (s, 3 H), 2.23-2.48 (m, 2 H), 4.65 (s, 1 H), 7.25-7.45 (m, 5 H). ¹³C NMR (CDCl₃) δ 14.0 (q), 22.5 (t), 23.9 (t), 24.1 (q), 28.8 (t), 31.5 (t), 35.5 (t), 79.7 (s), 125.9 (d), 127.7 (d), 128.4 (d), 141.4 (s), 211.7 (s). Anal. Calcd for C₁₆H₂₄O₂: C, 77.38; H, 9.74. Found: C, 77.45; H, 9.78.

3-Hydroxy-4-methyl-3-phenylpentan-2-one (2h): Colorless paste. *R_f* 0.45 (hexane-ethyl acetate, 5:1). IR (neat) 3456, 1707, 1599, 1491, 756, 702 cm⁻¹. ¹H NMR (CDCl₃) δ 0.895 (d, 3 H, *J* = 6.8 Hz), 0.904 (d, 3 H, *J* = 6.8 Hz), 2.14 (s, 3 H), 2.74-2.89 (m, 1 H), 4.47 (s, 1 H), 7.24-7.41 (m, 3 H), 7.51-7.57 (m, 2 H). ¹³C NMR (CDCl₃) δ 16.4 (q), 16.5 (q), 23.5 (q), 32.5 (d), 85.4 (s), 126.5 (d), 127.4 (d), 128.6 (d), 140.1 (s), 209.4 (s). Anal. Calcd for C₁₂H₁₆O₂: C, 74.97; H, 8.39. Found: C, 75.08; H, 8.45.

3-Hydroxy-3-(4-methoxyphenyl)butan-2-one (2i): Colorless paste. *R_f* 0.35 (hexane-ethyl acetate, 2:1). IR (neat) 3458, 1713, 1611, 1583, 970, 916, 847, 783, 662 cm⁻¹. ¹H NMR (CDCl₃) δ 1.76 (s, 3 H), 2.08 (s, 3 H), 3.81 (s, 3 H), 4.52 (s, 1 H), 6.86-6.93 (m, 2 H), 7.31-7.38 (m, 2 H). ¹³C NMR (CDCl₃) δ 23.4 (q), 24.1 (q), 55.3 (q), 79.4 (s), 113.9 (d), 127.1 (d), 133.3 (s), 159.1 (s), 209.6 (s). Anal. Calcd for C₁₁H₁₄O₃: C, 68.02; H, 7.27. Found: C, 67.81; H, 7.36.

3-Hydroxy-3-(3-methoxyphenyl)butan-2-one (2j): Colorless paste. *R_f* 0.2 (hexane-ethyl acetate, 5:1). IR (neat) 3456, 1713, 1599, 1583, 1487, 795, 777, 702, 683 cm⁻¹. ¹H NMR (CDCl₃) δ 1.77 (s, 3 H), 2.10 (s, 3 H), 3.81 (s, 3 H), 4.52 (s, 1 H), 6.82-6.88 (m, 1 H), 6.96-7.04 (m, 2 H), 7.29 (t, 1 H, *J* = 7.8 Hz). ¹³C NMR (CDCl₃) δ 23.4 (q), 24.2 (q), 55.1 (q), 79.7 (s), 111.7 (d), 113.1 (d), 118.1 (d), 129.4 (d), 142.9 (s), 159.6 (s), 209.2 (s). Anal. Calcd for C₁₁H₁₄O₃: C, 68.02; H, 7.27. Found: C, 68.15; H, 7.39.

3-Hydroxy-3-(2-methoxyphenyl)butan-2-one (2k): Colorless paste. *R_f* 0.2 (hexane-ethyl acetate, 2:1). IR (neat) 3449, 1717, 1601, 1585, 1491, 914, 789, 756, 679 cm⁻¹. ¹H NMR (CDCl₃) δ 1.68 (s, 3 H), 2.08 (s, 3 H), 3.78 (s, 3 H), 4.34 (brs, 1 H), 6.89 (dd, 1 H, *J* = 1.1, 8.1 Hz), 7.01 (dt, 1 H, *J* = 1.1, 7.6 Hz), 7.32 (ddd, 1 H, *J* = 1.6, 7.6, 8.1 Hz), 7.48 (dd, 1 H, *J* = 1.6, 7.6 Hz). ¹³C NMR (CDCl₃) δ 23.3 (q), 23.6 (q), 55.1 (q), 78.3 (s), 111.0 (d), 120.7 (d), 126.6 (d), 129.4 (d), 130.3 (s), 156.2 (s), 210.1 (s). Anal. Calcd for C₁₁H₁₄O₃: C, 68.02; H, 7.27. Found: C, 68.09; H, 7.30.

3-(4-Fluorophenyl)-3-hydroxybutan-2-one (2l): Colorless paste. *R_f* 0.2 (hexane-ethyl acetate, 5:1). IR (neat) 3449, 1713, 1601, 1508, 916, 853, 837 cm⁻¹. ¹H NMR (CDCl₃) δ 1.77 (s, 3 H), 2.08 (s, 3 H), 4.52 (s, 1 H), 7.01-7.11 (m, 2 H), 7.37-7.46 (m, 2 H). ¹³C NMR (CDCl₃) δ 23.4 (q), 24.3 (q), 79.4 (s), 115.3 (d, *J*_{CCF} = 21.1 Hz), 127.7 (d, *J*_{CCCF} = 8.4 Hz), 137.2 (s, *J*_{CCCCF} = 3.3 Hz), 162.2 (s, *J*_{CF} = 245.5 Hz), 209.1 (s). Anal. Calcd for C₁₀H₁₁FO₂: C, 65.92; H, 6.09. Found: C, 66.07; H, 6.12.

3-(3,4-Dimethoxyphenyl)-3-hydroxybutan-2-one (2m): white solid. Mp 45-46 °C. *R_f* 0.25 (hexane-ethyl acetate, 2:1). IR (KBr) 3470, 1713, 1605, 1591, 1518, 970, 935, 889, 860, 816, 766 cm⁻¹. ¹H NMR (CDCl₃) δ 1.77 (s, 3 H), 2.09 (s, 3 H), 3.87 (s, 3 H), 3.88 (s, 3 H), 4.53 (brs, 1 H), 6.86 (d, 1 H, *J* = 8.6 Hz), 6.90 (d, 1 H, *J* = 1.9 Hz), 7.00 (dd, 1 H, *J* = 1.9, 8.6 Hz). ¹³C NMR (CDCl₃) δ 23.2 (q), 24.0 (q), 55.7 (q), 79.4 (s), 109.0 (d), 110.7 (d), 118.0 (d), 133.7 (s), 148.5 (s), 148.7 (s), 209.5 (s). Anal. Calcd for C₁₂H₁₆O₄: C, 64.27; H, 7.19. Found: C, 64.23; H, 7.14.

3-Hydroxy-3-(naphthalen-1-yl)butan-2-one (2n): Colorless paste. *R_f* 0.2 (hexane-ethyl acetate, 5:1). IR (neat) 3464, 1713, 1599, 1510, 995, 968, 912, 806, 781, 673 cm⁻¹. ¹H NMR (CDCl₃) δ 1.90 (s, 3 H), 1.92 (s, 3 H), 4.40 (s, 1 H), 7.40-7.53 (m, 3 H), 7.75 (dd, 1 H, *J* = 0.8, 7.3 Hz), 7.83-7.95 (m, 3 H). ¹³C NMR (CDCl₃) δ 24.1 (q), 26.2 (q), 80.5 (s), 124.67 (d), 124.70 (d), 125.0 (d), 125.5 (d), 126.4 (d), 128.7 (d),

129.6 (d), 131.1 (s), 134.2 (s), 135.7 (s), 212.0 (s). Anal. Calcd for C₁₄H₁₄O₂: C, 78.48; H, 6.59. Found: C, 78.64; H, 6.68.

3-Hydroxy-3-(naphthalen-2-yl)butan-2-one (2o): White solid. Mp 51-52 °C. *Rf* 0.2 (hexane-ethyl acetate, 5:1). IR (neat) 3454, 1713, 1599, 1506, 968, 953, 918, 899, 858, 822, 789, 748 cm⁻¹. ¹H NMR (CDCl₃) δ 1.90 (s, 3 H), 2.10 (s, 3 H), 4.64 (brs, 1 H), 7.44-7.55 (m, 3 H), 7.79-7.90 (m, 3 H), 7.95 (d, 1 H, *J* = 2.1 Hz). ¹³C NMR (CDCl₃) δ 23.5 (q), 24.2 (q), 80.0 (s), 123.6 (d), 124.8 (d), 126.16 (d), 126.19 (d), 127.3 (d), 128.0 (d), 128.3 (d), 132.6 (s), 132.9 (s), 138.7 (s), 209.4 (s). Anal. Calcd for C₁₄H₁₄O₂: C, 78.48; H, 6.59. Found: C, 78.52; H, 6.60.

1-(1-Hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)ethanone (2p): Colorless paste. *Rf* 0.3 (hexane-ethyl acetate, 5:1). IR (neat) 3462, 1709, 1491, 982, 789, 762, 733 cm⁻¹. ¹H NMR (CDCl₃) δ 1.84-2.06 (m, 4 H), 2.07 (s, 3 H), 2.73-2.98 (m, 2 H), 4.57 (s, 1 H), 6.88-6.93 (m, 1 H), 7.13-7.25 (m, 3 H). ¹³C NMR (CDCl₃) δ 18.8 (t), 24.6 (q), 29.5 (t), 34.5 (t), 78.5 (s), 126.5 (d), 127.5 (d), 128.0 (d), 129.4 (d), 135.1 (s), 137.4 (s), 210.5 (s). Anal. Calcd for C₁₂H₁₄O₂: C, 75.76; H, 7.42. Found: C, 75.70; H, 7.35.

Methyl 5-hydroxy-4-oxo-5-phenylhexanoate (2r): Colorless paste. *Rf* 0.45 (hexane-ethyl acetate, 2:1). IR (neat) 3474, 1738, 1717, 1601, 1495, 764, 702 cm⁻¹. ¹H NMR (CDCl₃) δ 1.80 (s, 3 H), 2.44-2.52 (m, 1 H), 2.55-2.68 (m, 2 H), 2.73-2.81 (m, 1 H), 3.63 (s, 3 H), 4.40 (s, 1 H), 7.28-7.33 (m, 1 H), 7.35-7.39 (m, 2 H), 7.34-7.37 (m, 1 H). ¹³C NMR (CDCl₃) δ 24.8 (q), 27.9 (t), 30.6 (t), 51.7 (q), 79.8 (s), 125.7 (d), 127.9 (d), 128.5 (d), 141.5 (s), 172.9 (s), 210.5 (s). Anal. Calcd for C₁₃H₁₆O₄: C, 66.09; H, 6.83. Found: C, 66.20; H, 6.88.

Methyl 6-hydroxy-5-oxo-6-phenylheptanoate (2s): Colorless paste. *Rf* 0.5 (hexane-ethyl acetate, 2:1). IR (neat) 3470, 1736, 1717, 1601, 1493, 762, 702 cm⁻¹. ¹H NMR (CDCl₃) δ 1.75-1.87 (m, 2 H), 1.77 (s, 3 H), 2.08-2.44 (m, 2 H), 2.37-2.45 (m, 1 H), 2.49-2.56 (m, 1 H), 3.60 (s, 3 H), 4.51 (s, 1 H), 7.29-7.33 (m, 2 H), 7.35-7.39 (m, 2 H), 7.40-7.43 (m, 2 H). ¹³C NMR (CDCl₃) δ 18.9 (t), 24.1 (q), 32.4 (t), 34.3 (t), 51.3 (q), 79.7 (s), 125.7 (d), 127.8 (d), 127.8 (d), 128.5 (d), 141.3 (s), 173.2 (s), 211.1 (s). Anal. Calcd for C₁₄H₁₈O₄: C, 67.18; H, 7.25. Found: C, 67.08; H, 7.31.

1-Hydroxy-1,1-diphenylbutan-2-one (3b): Colorless paste. *Rf* 0.3 (hexane-ethyl acetate, 10:1). IR (neat) 3449, 1709, 1599, 1491, 980, 766, 700 cm⁻¹.

^1H NMR (CDCl_3) δ 1.03 (t, 3 H, $J = 7.3$ Hz), 2.57 (q, 2 H, $J = 7.3$ Hz), 4.88 (s, 1 H), 7.32-7.40 (m, 10 H). ^{13}C NMR (CDCl_3) δ 8.3 (q), 31.8 (t), 85.4 (s), 127.9 (d), 128.0 (d), 128.3 (d), 141.6 (s), 211.9 (s). Anal. Calcd for $\text{C}_{17}\text{H}_{18}\text{O}_2$: C, 79.97; H, 6.71. Found: C, 80.10; H, 6.72.

1-Hydroxy-1,1-diphenylpentan-2-one (3c): Colorless paste. R_f 0.4 (hexane-ethyl acetate, 10:1). IR (neat) 3447, 1709, 1599, 1493, 767, 748, 700 cm^{-1} . ^1H NMR (CDCl_3) δ 0.80 (t, 3 H, $J = 7.5$ Hz), 1.47-1.55 (m, 2 H), 2.54 (t, 3 H, $J = 7.0$ Hz), 4.91 (s, 1 H), 7.32-7.40 (m, 10 H). ^{13}C NMR (CDCl_3) δ 13.4 (q), 17.7 (t), 40.1 (t), 85.4 (s), 127.96 (d), 127.99 (d), 128.3 (d), 141.5 (s), 211.0 (s). Anal. Calcd for $\text{C}_{17}\text{H}_{18}\text{O}_2$: C, 80.28; H, 7.13. Found: C, 80.16; H, 7.19.

1-Hydroxy-1,1-diphenylnonan-2-one (3e): Colorless paste. R_f 0.5 (hexane-ethyl acetate, 10:1). IR (neat) 3447, 1709, 1599, 1493, 764, 700 cm^{-1} . ^1H NMR (CDCl_3) δ 0.84 (t, 3 H, $J = 6.9$ Hz), 1.09-1.27 (m, 8 H), 1.41-1.49 (m, 2 H), 2.55 (t, 2 H, $J = 7.4$ Hz), 4.92 (s, 1 H), 7.32-7.40 (m, 10 H). ^{13}C NMR (CDCl_3) δ 14.0 (q), 22.5 (t), 24.4 (t), 28.8 (t), 28.9 (t), 31.5 (t), 38.4 (t), 85.5 (s), 128.1 (d), 128.11 (d), 128.4 (d), 141.5 (s), 211.3 (s). Anal. Calcd for $\text{C}_{21}\text{H}_{26}\text{O}_2$: C, 81.25; H, 8.44. Found: C, 81.36; H, 8.45.

Methyl 5-hydroxy-4-oxo-5,5-diphenylpentanoate (3f): Colorless paste. R_f 0.2 (hexane-ethyl acetate, 5:1). IR (neat) 3462, 1740, 1717, 1599, 1493, 768, 754, 702 cm^{-1} . ^1H NMR (CDCl_3) δ 2.57 (t, 2 H, $J = 6.9$ Hz), 2.93 (t, 2 H, $J = 6.9$ Hz), 3.67 (s, 2 H), 4.55 (s, 3 H), 7.32-7.41 (m, 10 H). ^{13}C NMR (CDCl_3) δ 28.1 (t), 33.2 (t), 51.9 (q), 85.4 (s), 127.9 (d), 128.2 (d), 128.4 (d), 141.7 (s), 172.9 (s), 209.9 (s). Anal. Calcd for $\text{C}_{18}\text{H}_{18}\text{O}_4$: C, 72.47; H, 6.08. Found: C, 72.34; H, 6.00.

Methyl 6-hydroxy-5-oxo-6,6-diphenylhexanoate (3g): Colorless paste. R_f 0.5 (hexane-ethyl acetate, 2:1). IR (neat) 3458, 1734, 1717, 1599, 1493, 766, 752, 700 cm^{-1} . ^1H NMR (CDCl_3) δ 1.80-1.87 (m, 2 H), 2.23 (t, 2 H, $J = 7.3$ Hz), 2.65 (t, 2 H, $J = 7.3$ Hz), 3.61 (s, 3 H), 4.73 (s, 1 H), 7.33-7.40 (m, 10 H). ^{13}C NMR (CDCl_3) δ 19.3 (t), 32.6 (t), 37.1 (t), 51.4 (q), 85.4 (s), 127.8 (d), 128.0 (d), 128.3 (d), 141.3 (s), 173.2 (s), 210.4 (s). Anal. Calcd for $\text{C}_{19}\text{H}_{20}\text{O}_4$: C, 73.06; H, 6.45. Found: C, 72.87; H, 6.40.

Methyl 2-hydroxy-2-(4-methoxyphenyl)propanoate (4c): Colorless paste. R_f 0.1 (hexane-ethyl acetate, 5:1). IR (neat) 3489, 1732, 1611, 1583, 1512, 978, 835, 806, 750 cm^{-1} . ^1H NMR (CDCl_3) δ 1.77 (s, 3 H), 3.73 (brs, 1 H), 3.77 (s, 3 H), 3.80 (s, 3 H),

6.86-6.90 (m, 2 H), 7.44-7.48 (m, 2 H). ^{13}C NMR (CDCl_3) δ 26.5 (q), 53.0 (q), 55.1 (q), 75.3 (s), 113.5 (d), 126.3 (d), 134.7 (s), 159.0 (s), 176.2 (s). Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_4$: C, 62.85; H, 6.71. Found: C, 62.77; H, 6.65.

Methyl 2-hydroxy-2-(3-methoxyphenyl)propanoate (4d): Colorless paste. *R_f* 0.2 (hexane-ethyl acetate, 5:1). IR (neat) 3466, 1732, 16.01, 1585, 1489, 978, 872, 837, 791, 735, 698 cm^{-1} . ^1H NMR (CDCl_3) δ 1.77 (s, 3 H), 3.72 (brs, 1 H), 3.78 (s, 3 H), 3.82 (s, 3 H), 6.82-6.86 (m, 1 H), 7.10-7.13 (m, 2 H), 7.25-7.29 (m, 1 H). ^{13}C NMR (CDCl_3) δ 26.6 (q), 53.2 (q), 55.2 (q), 75.7 (s), 111.1 (d), 113.0 (d), 117.5 (d), 129.3 (d), 144.3 (s), 159.5 (s), 175.9 (s). Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_4$: C, 62.95; H, 6.74. Found: C, 62.77; H, 6.65.

Methyl 2-hydroxy-2-(2-methoxyphenyl)propanoate (4e): White solid. Mp 91-92 $^{\circ}\text{C}$. *R_f* 0.1 (hexane-ethyl acetate, 5:1). IR (KBr) 3524, 1746, 1599, 1585, 974, 934, 800, 756 cm^{-1} . ^1H NMR (CDCl_3) δ 1.79 (s, 3 H), 3.71 (s, 3 H), 3.82 (s, 3 H), 6.87-6.92 (m, 1 H), 6.98-7.03 (m, 1 H), 7.29-7.34 (m, 1 H), 7.43-7.47 (m, 1 H). ^{13}C NMR (CDCl_3) δ 24.0 (q), 52.3 (q), 55.2 (q), 73.9 (s), 110.8 (d), 120.6 (d), 126.0 (d), 129.2 (d), 130.9 (s), 156.3 (s), 176.3 (s). Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_4$: C, 62.85; H, 6.71. Found: C, 62.90; H, 6.71.

Methyl 2-(4-fluorophenyl)-2-hydroxypropanoate (4f): Colorless paste. *R_f* 0.25 (hexane-ethyl acetate, 5:1). IR (neat) 3483, 1732, 1603, 1508, 978, 839, 820, 750 cm^{-1} . ^1H NMR (CDCl_3) δ 1.77 (s, 3 H), 3.78 (s, 3 H), 3.81 (brs, 1 H), 7.00-7.05 (m, 2 H), 7.50-7.55 (m, 2 H). ^{13}C NMR (CDCl_3) δ 26.8 (q), 53.3 (q), 75.3 (s), 115.0 (d, J_{CCF} = 21.9 Hz), 127.1 (d, J_{CCCF} = 8.6 Hz), 138.4 (s, J_{CCCCF} = 2.9 Hz), 162.3 (s, J_{CF} = 245.1 Hz), 175.9 (s). Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{FO}_3$: C, 60.60; H, 5.59. Found: C, 60.48; H, 5.65.

Methyl 2-(3,4-dimethoxyphenyl)-2-hydroxypropanoate (4g): Colorless paste. *R_f* 0.2 (hexane-ethyl acetate, 2:1). IR (neat) 3489, 1732, 1605, 1593, 1518, 814, 768, 743, 719 cm^{-1} . ^1H NMR (CDCl_3) δ 1.77 (s, 3 H), 3.74 (brs, 1 H), 3.78 (s, 3 H), 3.88 (s, 3 H), 3.90 (s, 3H), 6.84 (d, 1 H, J = 8.3 Hz), 7.06 (dd, 1 H, J = 2.3, 8.3 Hz), 7.10 (d, 1 H, J = 2.3 Hz). ^{13}C NMR (CDCl_3) δ 26.7 (q), 53.2 (q), 55.9 (q), 75.5 (s), 108.5 (d), 110.7 (d), 117.5 (d), 135.2 (s), 148.6 (s), 148.7 (s), 176.2 (s). Anal. Calcd for $\text{C}_{12}\text{H}_{16}\text{O}_5$: C, 59.99; H, 6.71. Found: C, 60.07; H, 6.68.

Methyl 1-hydroxy-1,2,3,4-tetrahydronaphthalene-1-carboxylate (4h): Colorless paste. *R_f* 0.25 (hexane-ethyl acetate, 5:1). IR (neat) 3503, 1732, 1493, 993, 949, 802, 775, 737, 669 cm⁻¹. ¹H NMR (CDCl₃) δ 1.93-2.29 (m, 4 H), 2.73-2.94 (m, 2 H), 3.76 (s, 3 H), 3.77 (s, 1 H), 7.10-7.30 (m, 4 H). ¹³C NMR (CDCl₃) δ 19.0 (t), 29.3 (t), 35.4 (t), 53.1 (q), 74.2 (s), 126.3 (d), 126.9 (d), 127.9 (d), 129.2 (d), 136.1 (s), 137.1 (s), 177.2 (s). Anal. Calcd for C₁₂H₁₄O₃: C, 69.88; H, 6.84. Found: C, 69.93; H, 6.81.

2-Hydroxy-2-phenylcyclobutanone (11): Colorless paste. *R_f* 0.5 (hexane-ethyl acetate, 5:1). IR (neat) 3412, 1786, 1601, 1495, 918, 849, 766, 700 cm⁻¹. ¹H NMR (CDCl₃) δ 2.34-2.42 (m, 1 H), 2.67-2.74 (m, 1 H), 2.93 (brs, 1 H), 2.96-3.02 (m, 2 H), 7.34-7.43 (m, 3 H), 7.45-7.49 (m, 1 H). ¹³C NMR (CDCl₃) δ 27.8 (t), 41.0 (t), 92.5 (s), 126.0 (d), 128.8 (d), 129.0 (d), 138.7 (s). Anal. Calcd for C₁₀H₁₀O₂: C, 74.06; H, 6.21. Found: C, 73.89; H, 6.06.

2-Hydroxy-2-phenylcycloheptanone (14): Colorless paste. *R_f* 0.35 (hexane-ethyl acetate, 10:1). IR (neat) 3476, 1705, 1491, 841, 762, 719, 696 cm⁻¹. ¹H NMR (CDCl₃) δ 1.36-1.57 (m, 3 H), 1.88-2.07 (m, 3 H), 2.25-2.37 (m, 2 H), 2.45-2.52 (m, 1 H), 2.79-2.87 (m, 1 H), 4.56 (s, 1 H), 7.27-7.32 (m, 1 H), 7.33-7.39 (m, 2 H), 7.43-7.47 (m, 2 H). ¹³C NMR (CDCl₃) δ 23.6 (t), 27.8 (t), 30.5 (t), 36.4 (t), 39.5 (t), 82.3 (s), 125.7 (d), 127.9 (d), 128.6 (d), 141.6 (s), 213.7 (s). Anal. Calcd for C₁₃H₁₆O₄: C, 76.44; H, 7.90. Found: C, 76.55; H, 8.01.