Supporting Information

Enantioselective 1,3-Dipolar Cycloaddition Reaction between Diazoacetates and α -Substituted Acroleins: Total Synthesis of Manzacidin A

Taichi Kano, Takuya Hashimoto, and Keiji Maruoka*

Department of Chemistry, Graduate School of Science, Kyoto University
Sakyo, Kyoto 606-8502, Japan

General Information. Infrared (IR) spectra were recorded on a Shimadzu IRPrestige-21 spectrometer. ^1H NMR spectra were measured on a JEOL JNM-FX400 (400 MHz) spectrometer. Data were reported as follows: chemical shifts in ppm from tetramethylsilane (in the case of CDCl_3) or CD_2HOD ($\delta = 3.30$, in the case of CD_3OD) as an internal standard, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, dd = double-doublet, m = multiplet, br = broad, app = apparent), coupling constants (Hz), and assignment. ^{13}C NMR spectra were measured on a JEOL JNM-FX400 (100 MHz) spectrometer with complete proton decoupling. Chemical shifts were reported in ppm from the residual solvent as an internal standard. High performance liquid chromatography (HPLC) was performed on Shimadzu 10A instruments at 280 nm using 4.6 mm x 25 cm Daicel Chiralpak and Chiralcel. High-resolution mass spectra (HRMS) were performed on Applied Biosystems Mariner 8295 API-TOF workstation. Optical rotations were measured on a JASCO DIP-1000 digital polarimeter. For thin layer chromatography (TLC) analysis throughout this work, Merck precoated TLC plates (silica gel 60 GF₂₅₄, 0.25 mm) were used. The products were purified by flash column chromatography on silica gel 60 (Merck 1.09386.9025, 230-400 mesh). 2-Substituted acroleins were purchased or prepared according to the procedures¹⁻³ and used after distillation or column chromatography on silica gel.

Preparation of (*S*)-BINOL/Ti(OPr^{*i*})₄ (1:1 molar ratio) Complex (2a).

To a stirred mixture of (*S*)-BINOL (0.10 mmol, 28.6 mg) in CH₂Cl₂ (1.0 mL) was added Ti(OPr^{*i*})₄ (0.10 mmol, 29.5 μL) at room temperature under argon. The mixture was then stirred for 2 h at room temperature to afford a orange solution of titanium (*S*)-BINOLate (2a).

Preparation of (*S*)-BINOL/Ti(OPr^{*i*})₄ (2:1 molar ratio) Complex (2b).

To a stirred mixture of (*S*)-BINOL (0.20 mmol, 57.2 mg) in CH₂Cl₂ (1.0 mL) was added Ti(OPr^{*i*})₄ (0.10 mmol, 29.5 μL) at room temperature under argon. The mixture was then stirred for 2 h at room temperature to afford a dark orange solution of titanium (*S*)-BINOLate (2b).

Preparation of Bis{((*S*)-binaphthoxy)(isopropoxy) titanium} Oxide (2c).

To a stirred mixture of Ag₂O (0.05 mmol, 11.6 mg) in CH₂Cl₂ (0.9 mL) was added 1.0 M CH₂Cl₂ solution of ClTi(OPr^{*i*})₃ (0.10 mmol, 100 μL) at room temperature under argon. After stirring for 5 h at room temperature, (*S*)-BINOL (0.10 mmol, 28.6 mg) was added to the mixture, which was then stirred for 2 h at room temperature to afford a dark orange colored solution of chiral bis-Ti() oxide (2c).

General Procedure.

To a catalyst solution prepared as described above were added the corresponding α-substituted acrolein (1.0 mmol) and alkyl diazoacetate (1.5 mmol) dropwise at –40 °C under argon. The reaction mixture was stirred at the same temperature until complete consumption of the aldehyde. The mixture was quenched with ice-cooled aqueous NaHCO₃ and extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to give the cycloadduct.

(*R*)-Δ²-5-Formyl-5-methyl-3-pyrazolinecarboxylic acid ethyl ester (Table 1, entry 4).

Prepared according to the general procedure with methacrolein (1.0 mmol, 82.8 μL) and

ethyl diazoacetate (1.5 mmol, 158 μ L) over the course of 2 h. The crude material was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 1:1) to give the title cycloadduct as a pale yellow oil [54% yield (99.4 mg)].

^1H NMR (400 MHz, CDCl_3) δ 9.54 (1H, s, **CHO**), 6.42 (1H, s, **NH**), 4.31 (2H, q, J = 7.2 Hz, **OCH₂**), 3.30 (1H, d, J = 17.6 Hz, **CHH**), 2.83 (1H, d, J = 17.6 Hz, **CHH**), 1.42 (3H, s, **CH₃**), 1.35 (3H, t, J = 7.2 Hz, **OCH₂CH₃**) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 198.1, 161.7, 142.4, 73.3, 61.5, 39.5, 19.8, 14.3 ppm; IR (neat) 3318, 2980, 2932, 1701, 1568, 1422, 1375, 1337, 1265, 1111, 818 cm^{-1} ; $[\alpha]_D^{27} = -283.9^\circ$ (c = 1.0, C_6H_6 ; 90% ee).

Full characterization including HRMS and HPLC analysis was carried out after reduction of the aldehyde to the corresponding alcohol. The absolute configuration was deduced from a total synthesis of manzacidin A.

(*R*)- Δ^2 -5-Hydroxymethyl-5-methyl-3-pyrazolinecarboxylic acid ethyl ester.

To a stirred solution of (*R*)- Δ^2 -5-formyl-5-methyl-3-pyrazolinecarboxylic acid ethyl ester (0.25 mmol, 46.5 mg) in ethyl acetate was added NaBH_4 (0.25 mmol, 10.5 mg) at -78°C , and the reaction mixture was warmed to 0°C within 30 min. The reaction solution was then quenched with aqueous NH_4Cl and extracted with ethyl acetate. The organic layer was dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 1:3) to give the title alcohol as a colorless oil [73% yield (34.0 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralpak AD-H, hexane/2-propanol = 10:1, flow rate = 0.5 mL/min, retention time; 25.3 min (minor) and 27.4 min (major)).

^1H NMR (400 MHz, CDCl_3) δ 6.18 (1H, s, **NH**), 4.29 (2H, q, J = 7.2 Hz, **OCH₂**), 3.51 (2H, s, **CH₂OH**), 2.98 (1H, d, J = 17.6 Hz, **CHH**), 2.67 (1H, d, J = 17.2 Hz, **CHH**), 1.34 (3H, t, J = 7.2 Hz, **OCH₂CH₃**), 1.27 (3H, s, **CH₃**) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 162.5, 142.5, 68.4, 67.0, 61.1, 39.7, 22.9, 14.3 ppm; IR (neat) 3327, 2980, 2934, 2874, 1697, 1558, 1422, 1377, 1339, 1277, 1117, 1051 cm^{-1} ; HRMS (ESI) exact mass calcd. for $\text{C}_8\text{H}_{14}\text{N}_2\text{O}_3$: m/z 209.0897 ($[\text{M} + \text{Na}]^+$), found: m/z 209.0897 ($[\text{M} + \text{Na}]^+$); $[\alpha]_D^{29} = -17.4^\circ$ (c = 1.0, C_6H_6 ; 90% ee).

Δ^2 -5-Formyl-5-methyl-3-pyrazolinecarboxylic acid *tert*-butyl ester (Table 1, entry 6).

Prepared according to the general procedure with methacrolein (1.0 mmol, 82.8 μ L) and *tert*-butyl diazoacetate (1.5 mmol, 213 mg) over the course of 1 h. The crude material was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 1:1) to give the title cycloadduct as a pale yellow oil [52 % yield (109.6 mg)].

^1H NMR (400 MHz, CDCl_3) δ 9.53 (1H, s, **CHO**), 6.47 (1H, s, **NH**), 3.25 (1H, d, J = 17.6 Hz, **CHH**), 2.78 (1H, d, J = 18.0 Hz, **CHH**), 1.54 (9H, s, *t*-Bu), 1.41 (3H, s, **CH₃**) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 198.4, 160.9, 143.7, 82.3, 73.1, 39.6, 28.1, 19.7 ppm; IR (neat) 3308, 2978, 2934, 1697, 1572, 1368, 1273, 1250, 1072, 845, 814 cm^{-1} ; $[\alpha]_{\text{D}}^{28} = -169^\circ$ (c = 1.0, C_6H_6 ; 91% ee).

Full characterization including HRMS and HPLC analysis was carried out after reduction of the aldehyde to the corresponding alcohol.

Δ^2 -5-Hydroxymethyl-5-methyl-3-pyrazolinecarboxylic acid *t*-butyl ester.

To the stirred solution of Δ^2 -5-formyl-5-methyl-3-pyrazolinecarboxylic acid *tert*-butyl ester (0.52 mmol, 109.6 mg) in ethyl acetate was added NaBH_4 (0.25 mmol, 21.0 mg) at 0 $^\circ\text{C}$ and the reaction mixture was stirred for 10 min at the same temperature. The reaction solution was then quenched with aqueous NH_4Cl and extracted with ethyl acetate. The organic layer was dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 1:2) to give the title alcohol as a white solid [44% yield (49.2 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel OD-H, hexane/2-propanol = 10:1, flow rate = 0.5 mL/min, retention time; 16.3 min (minor) and 19.0 min (major)).

^1H NMR (400 MHz, CDCl_3) δ 5.86 (1H, s, **NH**), 3.50 (2H, s, **CH₂OH**), 2.98 (1H, d, J = 17.6 Hz, **CHH**), 2.64 (1H, d, J = 17.6 Hz, **CHH**), 1.54 (9H, s, *t*-Bu), 1.25 (3H, s, **CH₃**) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 161.8, 144.0, 81.8, 68.1, 66.9, 39.8, 28.2, 22.9 ppm; IR (neat) 3325, 2976, 2932, 1694, 1562, 1368, 1281, 1250, 1123, 1051, 843, 812, 750 cm^{-1} ;

HRMS (ESI) exact mass calcd. for $C_{10}H_{18}N_2O_3$: m/z 237.1210 ($[M + Na]^+$), found: m/z 237.1211 ($[M + Na]^+$); $[\alpha]_D^{29} = -9.4^\circ$ ($c = 1.0$, C_6H_6 ; 91% ee).

Δ^2 -5-Formyl-5-ethyl-3-pyrazolinecarboxylic acid *tert*-butyl ester (Table 2, entry 3).

Prepared according to the general procedure with 2-ethylacrolein (1.0 mmol, 84.1 mg) and *tert*-butyl diazoacetate (1.5 mmol, 213 mg) over the course of 3 h. The crude material was purified by column chromatography on silica gel (eluting with hexane/diethyl ether = 1:1) to give the title cycloadduct as a pale yellow oil [63% yield (157.2 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel OJ-H, hexane/2-propanol = 10:1, flow rate = 0.5 mL/min, retention time; 17.7 min (major) and 19.4 min (minor)).

1H NMR (400 MHz, $CDCl_3$) δ 9.50 (1H, s, CHO), 6.48 (1H, s, NH), 3.22 (1H, d, $J = 18.0$ Hz, CHH), 2.83 (1H, d, $J = 18.0$ Hz, CHH), 1.83 (2H, m, CCH_2CH_3), 1.53 (9H, s, *t*-Bu), 0.93 (3H, t, $J = 7.2$ Hz, CCH_2CH_3) ppm; ^{13}C NMR (100 MHz, $CDCl_3$) δ 199.2, 161.0, 143.9, 82.3, 76.7, 37.0, 28.2, 26.6, 8.1 ppm; IR (neat) 3316, 2976, 2936, 1697, 1570, 1458, 1416, 1368, 1258, 1128, 843, 812, 750, 696 cm^{-1} ; HRMS (ESI) exact mass calcd. for $C_{11}H_{18}N_2O_3$: m/z 249.1210 ($[M + Na]^+$), found: m/z 249.1209 ($[M + Na]^+$); $[\alpha]_D^{25} = -151.6^\circ$ ($c = 1.0$, C_6H_6 ; 83% ee).

Δ^2 -5-Formyl-5-(2-benzyloxyethyl)-3-pyrazolinecarboxylic acid *tert*-butyl ester (Table 2, entry 5).

Prepared according to the general procedure with 2-(2-benzyloxyethyl)acrolein (1.0 mmol, 190 mg) and *tert*-butyl diazoacetate (1.5 mmol, 213 mg) over the course of 1 h. The crude material was purified by column chromatography on silica gel (eluting with hexane/diethyl ether = 1:1) to give the title cycloadduct as a pale yellow oil [81% yield (270.5 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralpak AD-H, hexane/2-propanol = 10:1, flow rate = 1.0 mL/min, retention time; 15.5 min (minor) and 17.2 min (major)).

1H NMR (400 MHz, $CDCl_3$) δ 9.55 (1H, s, CHO), 7.26-7.37 (5H, m, ArH), 6.85 (1H, s,

NH), 4.42 (2H, app dd, $J = 11.6, 16.8$ Hz, PhCH₂), 3.57 (2H, m, BnOCH₂), 3.18 (1H, d, $J = 18.0$ Hz, CHH), 2.82 (1H, d, $J = 18.0$ Hz, CHH), 2.19 (1H, m, BnOCH₂CHH), 2.10 (1H, m, BnOCH₂CHH), 1.52 (9H, s, *t*-Bu) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 199.1, 161.0, 143.4, 137.2, 128.4, 127.8, 127.7, 82.2, 75.3, 73.4, 65.8, 39.4, 35.1, 28.2 ppm; IR (neat) 3327, 2978, 2930, 2866, 1724, 1697, 1576, 1368, 1256, 1125, 845, 810, 746, 698 cm⁻¹; HRMS (ESI) exact mass calcd. for C₁₈H₂₄N₂O₄: m/z 355.1628 ([M + Na]⁺), found: m/z 355.1620 ([M + Na]⁺); [α]_D²⁹ = -87.0° ($c = 1.0$, C₆H₆; 80% ee).

Δ²-5-Formyl-5-(2-phenylethyl)-3-pyrazolinecarboxylic acid *tert*-butyl ester (Table 2, entry 6).

Prepared according to the general procedure with 2-(2-phenylethyl)acrolein (1.0 mmol, 160.2 mg) and *tert*-butyl diazoacetate (1.5 mmol, 213 mg) over the course of 4 h. The crude material was purified by column chromatography on silica gel (eluting with hexane/diethyl ether = 1:1) to give the title cycloadduct as a pale yellow oil [63% yield (189.9 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel OD-H, hexane/2-propanol = 20:1, flow rate = 1.0 mL/min, retention time; 29.8 min (minor) and 47.7 min (major)).

¹H NMR (400 MHz, CDCl₃) δ 9.50 (1H, s, CHO), 7.14-7.31 (5H, m, ArH), 6.53 (1H, s, NH), 3.26 (1H, d, $J = 18.0$ Hz, CHH), 2.88 (1H, d, $J = 18.0$ Hz, CHH), 2.62 (2H, m, PhCH₂), 2.12 (2H, m, PhCH₂CH₂), 1.53 (9H, s, *t*-Bu) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 198.8, 160.9, 143.8, 140.2, 128.6, 128.1, 126.4, 82.3, 76.1, 37.8, 35.7, 30.2, 28.2 ppm; IR (neat) 3318, 2976, 2930, 1703, 1572, 1447, 1364, 1260, 1126, 839, 748, 700 cm⁻¹; HRMS (ESI) exact mass calcd. for C₁₇H₂₂N₂O₃: m/z 325.1523 ([M + Na]⁺), found: m/z 325.1522 ([M + Na]⁺); [α]_D²⁷ = -85.8° ($c = 1.0$, C₆H₆; 82% ee).

Δ²-5-Formyl-5-isopropyl-3-pyrazolinecarboxylic acid *tert*-butyl ester (Table 2, entry 7).

Prepared according to the general procedure with 2-isopropylacrolein (1.0 mmol, 98.1 mg)

and *tert*-butyl diazoacetate (1.5 mmol, 213 mg) over the course of 3 h. The crude material was purified by column chromatography on silica gel (eluting with hexane/diethyl ether = 3:1 to 1:1) to give the title cycloadduct as a pale yellow oil [82% yield (209.1 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel OD-H, hexane/2-propanol = 10:1, flow rate = 0.5 mL/min, retention time; 14.4 min (minor) and 16.1 min (major)).

¹H NMR (400 MHz, CDCl₃) δ 9.55 (1H, s, **CHO**), 6.56 (1H, s, **NH**), 3.21 (1H, d, *J* = 18.4 Hz, **CHH**), 2.88 (1H, d, *J* = 18.0 Hz, **CHH**), 2.14 (1H, m, **CH(CH₃)₂**), 1.53 (9H, s, *t*-Bu), 0.98 (6H, app dd, *J* = 6.8, 11.2 Hz, **CH(CH₃)₂**) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 200.0, 161.1, 143.7, 82.2, 79.4, 34.6, 32.0, 28.2, 17.6, 17.1 ppm; IR (neat) 3302, 2972, 1724, 1699, 1568, 1368, 1267, 1258, 1138, 843 cm⁻¹; HRMS (ESI) exact mass calcd. for C₁₂H₂₀N₂O₃: *m/z* 263.1366 ([M + Na]⁺), found: *m/z* 263.1369 ([M + Na]⁺); [α]_D²⁷ = -213.2° (*c* = 1.0, C₆H₆; 92% ee).

Δ²-5-Formyl-5-cyclohexyl-3-pyrazolinecarboxylic acid *tert*-butyl ester (Table 2, entry 8).

Prepared according to the general procedure with 2-cyclohexylacrolein (1.0 mmol, 138 mg) and *tert*-butyl diazoacetate (1.5 mmol, 213 mg) over the course of 5 h. The crude material was purified by column chromatography on silica gel (eluting with hexane/diethyl ether = 2:1 to 3:2) to give the title cycloadduct as a pale yellow oil [77% yield (215.9 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel OD-H, hexane/2-propanol = 20:1, flow rate = 0.5 mL/min, retention time; 23.0 min (minor) and 31.3 min (major)).

¹H NMR (400 MHz, CDCl₃) δ 9.52 (1H, s, **CHO**), 6.55 (1H, s, **NH**), 3.19 (1H, d, *J* = 18.0 Hz, **CHH**), 2.90 (1H, d, *J* = 18.0 Hz, **CHH**), 1.69-1.83 (5H, m, Cy), 1.53 (9H, s, *t*-Bu), 1.00-1.26 (6H, m, Cy) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 200.3, 161.1, 143.8, 82.1, 79.3, 42.2, 34.8, 28.2, 27.9, 27.4, 26.3, 26.1, 26.0 ppm; IR (neat) 3310, 2928, 2857, 1713, 1570, 1447, 1364, 1260, 1125, 835, 743 cm⁻¹; HRMS (ESI) exact mass calcd. for C₁₅H₂₄N₂O₃: *m/z* 303.1679 ([M + Na]⁺), found: *m/z* 303.1679 ([M + Na]⁺); [α]_D²⁸ = -143° (*c* = 1.0, C₆H₆; 94% ee).

(3aR,6R)-6-Methoxy-3a-methyl-3a,4-dihydro-3H-pyrazolo[1,5-c]oxazole-2-carboxylic acid ethyl ester (6).

To a stirred solution of (*R*)- Δ^2 -5-hydroxymethyl-5-methyl-3-pyrazolinecarboxylic acid ethyl ester (0.18 mmol, 33.5 mg, 90% ee) in methyl orthoformate (1.0 mL) was added a catalytic amount of pyridinium *p*-toluenesulfonate. After stirring for 1 h at room temperature, the reaction mixture was treated with aqueous NaHCO₃ and the organic phase was extracted with ethyl acetate. The residue was concentrated under reduced pressure and purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 2:1) to give the title bicyclic compound **6** as a colorless oil [89% yield (36.4 mg)].

¹H NMR (400 MHz, CDCl₃) δ 5.95 (1H, s, CH₃OCH), 4.33 (2H, m, OCH₂CH₃), 3.94 (1H, d, *J* = 8.4 Hz, OCHH), 3.81 (1H, *J* = 8.4 Hz, OCHH), 3.44 (3H, s, OCH₃), 3.18 (1H, d, *J* = 18.0 Hz, N=CCHH), 2.90 (1H, d, *J* = 18.0 Hz, N=CCHH), 1.59 (3H, s, CCH₃), 1.36 (3H, t, 7.2 Hz, OCH₂CH₃) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 161.5, 148.2, 112.9, 76.5, 69.9, 61.8, 53.8, 45.5, 25.9, 14.3 ppm; IR (neat) 2974, 2931, 1711, 1585, 1450, 1290, 1242, 1157, 1092, 1009 cm⁻¹; HRMS (ESI) exact mass calcd. for C₁₀H₁₆N₂O₄: *m/z* 229.1183 ([M + H]⁺), found: *m/z* 229.1184 ([M + H]⁺); [α]_D²⁵ = -467.0° (*c* = 1.0, CHCl₃).

(4S,6R)-6-Hydroxymethyl-6-methyl-1,4,5,6-tetrahydro-pyrimidine-4-carboxylic acid (8).

To a solution of **6** (0.159 mmol, 36.4 mg) in isopropanol/H₂O (1.5 mL/0.5 mL) was added Raney nickel (50 mg, purchased from Aldrich). Hydrogen gas (balloon) was then charged to the reaction flask, and the mixture was stirred at 80 °C for 24 h. The reaction mixture was filtered, and the residue was washed with methanol. After concentration *in vacuo*, the residue was passed through column chromatography on Dowex 50W $\times$ 8 (100~200 mesh, H⁺ form, eluting with H₂O then 1N aqueous NH₃) to give the title compound **8** as a mixture of diastereomers (85:15), which was used for the next step without further purification.

Manzacidin A (1a).

Prepared according to the literature procedure.⁴ The crude product was purified by reverse phase HPLC (cosmosil 5C18-AR-II, eluting with H₂O/CH₃CN/TFA = 75:25:0.1) to give pure manzacidin A [50% yield (27.4 mg)].

¹H NMR (400 MHz, CD₃OD) δ 8.07 (1H, s, N=CH-NH), 7.03 (1H, d, J = 1.2 Hz, BrCCH), 6.92 (1H, d, J = 1.2 Hz, BrCCH), 4.47 (1H, dd, J = 5.2, 10.0 Hz, NCHCO₂H), 4.36 (1H, d, J = 12.0 Hz, OCHH), 4.24 (1H, d, J = 11.6 Hz, OCHH), 2.38 (1H, dd, J = 5.2, 14.0 Hz, CHHCHCO₂), 2.22 (1H, dd, J = 10.4, 14.0 Hz, CHHCHCO₂), 1.46 (3H, s, CH₃) ppm; ¹³C NMR (100 MHz, CD₃OD) δ 172.8, 160.4, 151.9, 125.2, 123.1, 118.4, 98.1, 68.7, 53.8, 31.2, 24.1 ppm (one ¹³C peak remains unidentified, which overlapped with the solvent peak)⁴; IR (neat) 3198, 2982, 1670, 1589, 1456, 1400, 1387, 1188, 1138, 1078 cm⁻¹; HRMS (ESI) exact mass calcd. for C₁₂H₁₄BrN₃O₄: m/z 344.0240 ([M + H]⁺) and 346.0221 ([M + H]⁺), found: m/z 344.0231 ([M + H]⁺) and 346.0223 ([M + H]⁺); [α]_D²⁹ = -23.0° (c = 0.50, CH₃OH).

Mechanistic Studies for the Enantioselective 1,3-Dipolar Cycloaddition between Diazoacetates and α -Substituted Acroleins Catalyzed by Ti-BINOL Complexes.

Since all Ti-BINOL complexes **2a-c** gave similar results in the cycloaddition between diazoacetates and methacrolein (Table 1 in the text), the following hypothetical transition state model is proposed as a common structure for **2a-c**, where one prochiral face of methacrolein is efficiently blocked by a binaphthyl moiety (Fig 1).

Fig 1

Although the present cycloaddition reaction was promoted by Ti-BINOL complexes, coordination of methacrolein or *t*-butyl diazoacetate to Ti-BINOL complex (*S*)-**2b** was not verified by ¹H-NMR analysis (Fig 2). Since coordination of a strongly coordinative nitron to the Ti-BINOL complex was observed in ¹H-NMR study, the coordination ability of *t*-butyl diazoacetate might be weaker than that of the nitron.

Based on these observations, we suppose that titanium Lewis acid catalysts activate aldehydes to facilitate the 1,3-cycloaddition reaction during the facile ligand exchange between methacrolein and *t*-butyl diazoacetate on the titanium center (Scheme 1).

Scheme 1

- (1) Takano, S.; Inomata, K.; Samizu, K.; Tomita, S.; Yanase, M.; Suzuki, M.; Iwabuchi, Y.; Sugihara, T.; Ogasawara, K. *Chem. Lett.* **1989**, 1283.
- (2) Basu, K.; Richards, J.; Paquette, L. A. *Synlett* **2004**, 2841.

- (3) Marvel, C. S.; Myers, R. L.; Saunders, J. H. *J. Am. Chem. Soc.* **1948**, 70, 1694.
- (4) Namba, K.; Shinada, T.; Teramoto, T.; Ohfuné, Y. *J. Am. Chem. Soc.* **2000**, 122, 10708.