Supporting Information

Enantioselective Addition of Terminal Alkynes to Isolated Isoquinoline Iminiums

Alexander M. Taylor, Stuart L. Schreiber*

*Howard Hughes Medical Institute,
Broad Institute of Harvard and MIT,
12 Oxford St., Cambridge, Mass. 02138*

stuart_schreiber@harvard.edu

Materials and methods. Except as otherwise noted, reactions were carried out under nitrogen with dry, freshly-purified solvents. Solvents were passed through a solvent column (activated alumina and Q5 reactant) prior to use. Triethylamine was distilled over calcium hydride under a Nitrogen atmosphere. NMR spectra were recorded at 600 MHz, 500 MHz, or 400 MHz using a Varian I-600, I-500, or M-400 instrument, respectively. Chemical shifts for proton NMR spectra are reported in parts per million downfield from tetramethylsilane and were referenced to residual protonated solvent (CHCl_3 : δ 7.26). Chemical shifts for carbon NMR spectra are reported in parts per million downfield from tetramethylsilane and referenced to protonated solvent (CHCl_3 : δ 77.0). Data are represented as follows: chemical shift (multiplicity, integration [br = broad, s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet], coupling constants in Hertz). HPLC separations were performed on an Agilent 1100 series with Chiralpak OD, OJ, ASH, and AD columns as noted. Isopropanol and hexanes were used as solvents, and flow rate was set at 1.0 mL/ min. UV absorbance was detected at 230 nm, except where noted. SFC/ MS chromatography was performed with a Berger analytic SFC using CO_2 and MeOH as the mobile phase and Berger Instruments 60A 6μ 4.6 x 150 mm silica, OD, and AD columns. Low-resolution mass spectra were obtained with JEOL AX-505H, SX-102A (CI/ EI), Micromass Platform II, and LCT (APCI/ ES/ LCMS). Only molecular ions, fragments from molecular ions, and other major peaks were reported. High-resolution mass spectra were obtained through the Harvard University mass spectrometry facility. Infrared spectra were obtained with a Thermo Nicolet Avatar 370 DTGS. Optical rotations were obtained with a Jasco DIP 370 digital polarimeter using a 2 mL cell with a 1 dm path length. Melting points were measured on a Mel-Temp apparatus and are uncorrected. Flash chromatography was performed with the indicated solvent on E. Merck silica gel 60 (40-63 mesh). Separations were monitored on E. Merck silica gel 60 F_{254} plates (0.25 mm) and visualized with a UV lamp and by staining with Dragendorff's reagent. Reagents were purchased from Aldrich, Acros, and Strem and used without further purification. Yeast growth was monitored by UV absorption at 660 nm with a Spectramax 384plus plate reader at 25 °C.

2-(2-bromo-5-methoxybenzyl)-3,4-dihydro-6,7-dimethoxyisoquinolinium bromide (3)

General procedure for preparation of isoquinolinium salts: An erlenmeyer flask was charged with a stirbar and 3,4-dihydro-6,7-dimethoxyisoquinoline hydrochloride (185 mg, 0.81 mmol), to which Na_2CO_3 (aq., sat., 10 mL) was added. After 10 of stirring, EtOAc (10 mL) was added, and the organic layer was separated. The aqueous layer was extracted 2x with EtOAc (10 mL), and the organic layers were combined, dried with magnesium sulfate, filtered, and concentrated *in vacuo* to give a clear oil. To this oil in a round bottom flask was added diethyl ether (8 mL) and 2-bromo-5-methoxybenzylbromide (250 mg, 0.89 mmol, 1.1 eq.) and a stirbar. The solution was heated to reflux overnight. After cooling, the precipitate was collected and washed 2x with diethyl ether to yield a yellow powder (quantitative); ^1H NMR (400 MHz, CDCl_3) δ 9.88 (s, 1H), 7.62 (d, 1H, $J = 2.8$ Hz), 7.56 (s, 1H), 7.44 (d, 1H, $J = 8.8$ Hz), 6.83 (dd, 1H, $J = 2.8, 8.8$ Hz), 6.79 (s, 1H), 5.50 (s, 2H), 3.97 (t, 2H, $J = 8.4$ Hz), 3.95 (s, 3H), 3.89 (s, 3H), 3.84 (s, 3H), 3.22 (t, 2H, $J = 8.4$ Hz); ^{13}C NMR (400 MHz, CDCl_3) δ 165.4, 159.6, 157.5, 148.8, 133.9, 132.3, 131.2, 118.4, 118.3, 117.0, 115.8, 114.9, 110.5, 62.7, 56.7, 56.5, 56.2, 47.6, 25.6; IR (CDCl_3 film, cm^{-1}): 3415 (m), 3011 (w), 2941 (w), 2837 (w), 2190 (w), 1647 (m), 1604 (m), 1565 (m), 1524 (m), 1480 (m), 1464 (m), 1403 (m), 1374 (m), 1341 (m), 1300 (m), 1280 (m), 1246 (m), 1173 (w), 1173 (m), 1142 (m), 1100(w), 1054 (m), 1019 (m), 1007 (m), 922 (m), 859 (w), 820 (w); LRMS (m/z) 390.0 [M-Br]; HRMS (TOF ES) calcd for $\text{C}_{19}\text{H}_{21}\text{Br}_2\text{NO}_3$ [M-Br] 390.0696, Found 390.0705 Error 2.3 ppm; mp = 195 – 196 °C.

3,4-dihydro-3,4-dimethoxy-2-methylisoquinolinium iodide (4)

The general procedure was used to add methyl iodide (400 μL , 6.4 mmol, 1.1 eq.) to 3,4-dihydro-6,7-dimethoxyisoquinoline hydrochloride (1.27 g, 5.6 mmol). The precipitate was collected as a yellow powder (1.5 g, 4.5 mmol, 80%); ^1H NMR (400 MHz, CDCl_3) δ 9.74 (s, 1H), 7.59 (s, 1H), 6.86 (s, 1H), 4.05 (t, 2H, $J = 8.4$ Hz), 3.98 (s, 3H), 3.91 (s, 3H), 3.89 (s, 3H), 3.33 (t, 2H, $J = 8$ Hz); ^{13}C NMR (400 MHz, CDCl_3) δ 164.6, 157.3, 148.7, 131.8, 117.0, 115.4, 110.8, 56.8, 56.7, 50.2, 47.8, 25.4; IR (CH_2Cl_2 film, cm^{-1}) 3022 (m), 3005 (m), 2960 (s), 2942 (s), 2835 (m), 1667 (s), 1568 (s), 1460 (s), 1450 (s), 1397 (s), 1377 (s), 1338 (s), 1300 (s), 1281 (s), 1261 (s), 1247 (s), 1210 (s), 1195 (s), 1188 (s), 1167 (s), 1129 (s), 1092 (m), 1053 (m), 1031 (m), 1009 (s), 987 (s), 939 (m), 894 (s), 872 (m); LRMS (m/z) 206.1 [M-I]; HRMS (TOF ES) calcd for $\text{C}_{12}\text{H}_{16}\text{INO}_2$ [M-I] 206.1181, Found 206.1190 Error 4.4 ppm; mp = 200 – 201 °C.

5

3,4-dihydro-6,7-dimethoxy-2-methyl-1-phenylisoquinolinium iodide (5)

The general procedure was used to add methyl iodide (79 μ L, 1.2 mmol, 1.1 eq.) to 3,4-dihydro-6,7-dimethoxy-1-phenylisoquinoline hydrochloride (300 mg, 1.1 mmol). The precipitate was isolated as a yellow powder (146 mg, 0.36 mmol, 32%); ^1H NMR (600 MHz, CDCl_3) δ 7.71 (m, 3H), 7.65 (m, 4H), 6.93 (s, 1H), 6.38 (s, 1H), 4.53 (t, 2H, $J = 7.8$ Hz), 4.03 (s, 3H), 3.75 (s, 3H), 3.59 (s, 3H), 3.53 (t, 2H, $J = 7.8$ Hz); ^{13}C NMR (400 MHz, CDCl_3) δ 137.1, 156.7, 148.0, 133.9, 131.9, 129.6, 129.1, 128.7, 119.6, 115.2, 111.0, 57.0, 55.9, 52.9, 47.0, 25.8; IR (CDCl_3 film, cm^{-1}): 3455 (w), 3017 (w), 2970 (w), 2940 (w), 2835 (w), 2189 (w), 1625 (m), 1603 (s), 1562 (s), 1522 (s), 1493 (m), 1464 (m), 1445 (m), 1386 (s), 1340 (s), 1293 (s), 1274 (s), 1222 (s), 1185 (m), 1113 (s), 1030 (m), 1009 (m), 993 (m), 951 (m), 920 (m), 869 (m), 826 (m); LRMS (m/z) 282.1 [M-I]; HRMS (TOF ES) calcd for $\text{C}_{25}\text{H}_{25}\text{Br}_2\text{NO}_3$ [M-I] 282.1494, Found 282.1504 Error 3.5 ppm; mp = 177 – 183 $^\circ\text{C}$.

2-(2-bromo-5-methoxy)-benzyl-7-methoxyisoquinolinium bromide (6)

The general procedure was used to add 2-bromo-5-methoxybenzylbromide (690 mg, 2.5 mmol, 1.1 eq.) to 7-methoxyisoquinoline (356 mg, 2.2 mmol). The precipitate was collected as an off-white powder (399 mg, 0.91 mmol, 41%); ^1H NMR (600 MHz, CDCl_3) δ 11.06 (s, 1H), 8.43 (d, 1H, $J = 6.6$ Hz), 7.07 (d, 1H, $J = 6.6$ Hz), 7.99 (d, 1H, $J = 3.6$ Hz), 7.94 (s, 1H), 7.72 (dd, 1H, $J = 2.4, 9$ Hz), 7.78 (d, 1H, $J = 8.4$ Hz), 6.88 (dd, 1H, $J = 3, 9$ Hz), 6.30 (s, 2H), 4.06 (s, 3H), 3.93 (s, 3H); ^{13}C NMR (400 MHz, CDCl_3) δ 161.0, 159.5, 148.0, 133.9, 133.1, 132.8, 132.1, 130.6, 129.5, 128.3, 125.4, 118.3, 118.1, 114.5, 137.4, 63.1, 56.3, 56.3; IR (CDCl_3 film, cm^{-1}) 3414 (m), 3012 (m), 2970 (m), 2941 (m), 2837 (m), 2194 (m), 1616 (9s), 1594 (m), 1545 (m), 1520 (m), 1486 (m), 1453 (m), 1408 (s), 1369 (m), 1313 (m), 1281 (m), 1248 (m), 1219 (m), 1173 (m), 1123 (m), 1056 (m), 1043 (m), 1020 (m), 995 (w), 922 (m), 854 (m), 822 (m); LRMS (m/z) 360.1 [M-Br]; HRMS (TOF ES) calcd for $\text{C}_{18}\text{H}_{17}\text{Br}_2\text{NO}_2$ [M-Br] 358.0443, Found 358.0432 Error 3.1 ppm; mp = 144 – 150 $^\circ\text{C}$.

Preparation of QUINAP:CuBr stock solution: To increase consistency and accuracy between experiments, stock solutions of R- and S-QUINAP with CuBr in methylene chloride were prepared. A round-bottom flask was charged with a weighed amount of CuBr. To this was added QUINAP (1.1 eq.) and a stirbar. The flask was sealed and purged with Nitrogen. Methylene chloride was added to make the

concentration of CuBr 1 mg/ mL. The mixture stirred until homogeneous. The final mixture was a clear yellow solution, which was transferred into a Sure/Stor flask. The stock solution was handled under Schlenk conditions.

2-(2-bromo-5-methoxybenzyl)-1,2,3,4-tetrahydro-6,7-dimethoxy-1-(2-phenylethynyl)isoquinoline (3a)

General procedure for the enantioselective alkyne addition to isoquinolinium salts A round-bottom flask was charged with iminium salt **3** (50 mg, 0.106 mmol) and a stirbar before being sealed and purged with Nitrogen. R-QUINAP: CuBr (718 μ L, 0.005 mmol, 5 mol%) was added from a stock solution, followed by methylene chloride (717 μ L) and phenylacetylene (117 μ L, 1.06 mmol, 10 eq.). The solution was cooled to -78 °C in a dry ice/ isopropanol bath before the addition of triethylamine (15 mL, 0.106 mmol, 1 eq.). The solution was placed in a cryocool bath, where it warmed to -55 °C and sat for 44 h. Upon completion, the solution was loaded directly onto a silica gel column for flash column chromatography (1:1:1% hexanes: diethyl ether: triethylamine) to yield **3a** (47 mg, 0.95 mmol, 91%) as an oil; ^1H NMR (600 MHz, CD_2Cl_2) δ 7.47 (m, 3H), 7.32 (t, 3H, J = 3.6 Hz), 7.20 (d, 1H, J = 3 Hz), 6.77 (s, 1H), 6.74 (dd, 1H, J = 3.6, 9 Hz), 4.79 (s, 1H), 4.00 (dd, 2H, J = 14.4, 21 Hz), 3.81 (s, 3H), 3.81 (s, 3H), 4.78 (s, 3H), 3.11 (td, 1H, J = 4.2, 11.4 Hz), 2.98 (m, 1H), 2.86 (dddd, 1H, J = 1.8, 6, 11.4 Hz), 2.72 (dt, 1H, J = 3.6, 16.2 Hz); ^{13}C NMR (400 MHz, CD_2Cl_2) δ 159.5, 148.8, 147.9, 139.2, 133.6, 132.0, 128.5, 127.5, 126.5, 123.5, 116.5, 115.3, 114.6, 112.0, 111.0, 88.1, 86.8, 59.3, 56.2, 56.1, 55.8, 54.6, 46.1, 29.0; IR (CH_2Cl_2 film, cm^{-1}) 3382 (m), 3059 (w), 2999 (m), 2916 (m), 2832 (m), 1611 (m), 1596 (m), 1572 (m), 1518 (s), 1489 (m), 1464 (s), 1442 (m), 1409 (m), 1378 (m), 1359 (m), 1337 (m), 1263 (s), 1225 (s), 1189 (m), 1161 (m), 1130 (m), 1097 (m), 1070 (w), 1053 (m), 1016 (m), 969 (w), 864 (m), 845 (m), 806 (m); Product ratio (from R-QUINAP synthesis) was determined by HPLC, Chiralpak AD column, 5% IPA in hexanes, 1.0 mL/ min, 230 nm, $t_{\text{R}}^{(\text{major})}$ = 14.96 min, area = 3.62×10^4 (97.9%), $t_{\text{R}}^{(\text{minor})}$ = 8.07 min, area = 7.70×10^2 (2.1%) (96% ee); LRMS (m/z) 492.1 [$\text{M}+1$]; HRMS (TOF ES) calcd for $\text{C}_{27}\text{H}_{26}\text{BrNO}_3$ [$\text{M}+1$] 492.1174, Found 492.1170 Error 0.8 ppm; Product of R-QUINAP: $[\alpha]_{\text{D}}^{25}$ = -53 ° (c = 0.7, CH_2Cl_2).

2-(2-bromo-5-methoxybenzyl)-1,2,3,4-tetrahydro-6,7-dimethoxy-1-(2-(trimethylsilyl)ethynyl)isoquinoline (3b)

The general procedure was used to add trimethylsilylacetylene (150 μ L, 1.06 mmol, 10 eq.) to iminium salt **3** (50 mg, 0.106 mmol). Following column chromatography (Si gel, 2:1:1% hexanes: diethyl ether: triethylamine), **3b** was isolated as a clear oil (50 mg, 0.102 mmol, 96%); ^1H NMR (600 MHz, CD_2Cl_2) δ 7.43 (d, 1H, $J = 8.4$ Hz), 7.14 (d, 1H, $J = 3$ Hz), 6.71 (dd, 1H, $J = 3.6, 9$ Hz), 6.68 (s, 1H), 6.56 (s, 1H), 4.53 (s, 1H), 3.89 (dd, 2H, $J = 12, 14.4$ Hz), 3.78 (s, 3H), 3.77 (s, 3H), 3.76 (s, 3H), 2.98 (m, 1H), 2.91 (m, 1H), 2.75 (m, 1H), 2.66 (dt, 1H, $J = 3.6, 16.2$ Hz), 0.18 (s, 9H); ^{13}C NMR (400 MHz, CD_2Cl_2) δ 159.4, 148.8, 147.9, 139.2, 133.6, 127.2, 126.5, 116.4, 115.3, 112.0, 111.0, 104.4, 90.9, 59.2, 56.2, 56.1, 54.8, 46.0, 29.0, 0.22; IR (CH_2Cl_2 film, cm^{-1}) 3352 (m), 2999 (w), 2954 (m), 2932 (m), 2835 (m), 2360 (m), 1674 (w), 1656 (m), 1650 (m), 1644 (m), 1634 (w), 1602 (s), 1574 (m), 1574 (m), 1514 (s), 1467 (s), 1439 (w), 1435 (w), 1377 (w), 1338 (m), 1277 (s), 1248 (m), 1162 (m), 1126 (m), 1100 (m), 1054 (m), 1018 (m), 844 (m); Product ratio (from R-QUINAP synthesis) was determined by HPLC, Chiralpak OD column, 5% IPA in hexanes, 1.0 mL/ min, 230 nm, $t_{\text{R}}^{(\text{major})} = 6.25$ min, area = 9.99×10^4 (99.6%), $t_{\text{R}}^{(\text{minor})} = 4.33$ min, area = 3.96×10^3 (0.4%) (99% ee); LRMS (m/z) 488.2 $[\text{M}+1]$; HRMS (TOF ES) calcd for $\text{C}_{24}\text{H}_{28}\text{BrNO}_3\text{Si}$ $[\text{M}+1]$ 488.1256, Found 488.1262 Error 1.2 ppm; Product of R-QUINAP: $[\alpha]_{\text{D}}^{25} = -18^\circ$ ($c = 0.2$, CH_2Cl_2).

2-(2-bromo-5-methoxybenzyl)-1-(2-ethoxyethynyl)-1,2,3,4-tetrahydro-6,7-dimethoxyisoquinoline (3c)

The general procedure was used for the addition of ethylethynyl ether (102 μ L, 1.06 mmol, 10 eq.) to iminium **3** (50 mg, 0.106 mmol). Following column chromatography (Si gel, 2:1:1% hexanes: diethyl ether: triethylamine), **3c** was isolated (46.5 mg, 0.101 mmol, 95%); ^1H NMR (400 MHz, CD_2Cl_2) δ 7.43 (d, 1H, $J = 8.4$ Hz), 7.15 (d, 1H, $J = 2.8$ Hz), 6.702 (dd, 1H, $J = 3.2, 8.8$ Hz), 6.67 (s, 1H), 6.57 (s, 1H), 4.57 (s, 1H), 4.10 (q, 1H, $J = 6.7$ Hz), 3.82 (d, 2H, $J = 3.2$ Hz), 3.77 (s, 3H), 3.77 (s, 3H), 3.75 (s, 3H), 2.97 (m, 2H), 2.72 (m, 1H), 2.65 (m, 1H), 1.37 (t, 3H, $J = 7.6$ Hz); ^{13}C NMR (400 MHz, CD_2Cl_2) δ 158.4, 148.4, 14738, 139.6, 133.5, 129.3, 126.1, 116.3, 115.2, 114.5, 111.8, 110.9, 95.3, 75.1, 59.2, 56.2, 56.1, 55.8, 53.5, 45.7, 36.3, 29.1, 14.6; IR (CH_2Cl_2 film, cm^{-1}) 3585 (w), 3381 (s), 2999 (m), 2934 (s), 2833 (s), 1654 (m), 1601 (m), 1572 (m), 1519 (s), 1464 (s), 1409 (w), 1377 (m), 1567 (m), 1338 (m), 1277 (m), 1263 (m), 1225 (m), 1187 (w), 1159 (m), 1134 (m), 1102 (m), 1053 (w), 1016 (m); Product ratio (from R-

QUINAP synthesis) was determined by HPLC, Chiralpak OD column, 5% IPA in hexanes, 1.0 mL/ min, 230 nm, $t_R^{(major)} = 11.1$ min, area = 1.06×10^4 (85.0%), $t_R^{(minor)} = 9.37$ min, area = 1.89×10^3 (15.0%) (70% ee); LRMS (m/z) 460.0 [M+1]; HRMS (TOF ES) calcd for $C_{23}H_{26}BrNO_4$ [M+1] 460.1123, Found 460.1125 Error 0.4 ppm; Product of S-QUINAP: $[\alpha]_D^{25} = +4^\circ$ (c = 0.5, CH_2Cl_2).

2-(2-bromo-5-methoxybenzyl)-1,2,3,4-tetrahydro-6,7-dimethoxy-1-(3-methoxyprop-1-ynyl)isoquinoline (3d)

The general procedure was used to add methylpropargyl ether (90 μ L, 1.06 mmol, 10 eq.) to isoquinolinium salt **3** (50 mg, 0.106 mmol). Following column chromatography (Si gel, 2:1:1% hexanes: diethyl ether: triethylamine), **3d** was isolated (39 mg, 0.85 mmol, 80%).; 1H NMR (500 MHz, CD_2Cl_2) δ 7.46 (d, 1H, $J = 9$ Hz), 7.16 (d, 1H, $J = 3$ Hz), 6.73 (dd, 1H, $J = 3, 8.5$ Hz), 6.70 (s, 1H), 6.61 (s, 1H), 4.64 (s, 1H), 4.12 (t, 2H, $J = 2$ Hz), 3.88 (s, 2H), 3.80 (s, 3H), 3.79 (s, 3H), 3.78 (s, 3H), 3.38 (s, 3H), 3.02 (m, 2H), 2.82 (m, 1H), 2.68 (m, 1H); ^{13}C NMR (500 MHz, CD_2Cl_2) δ 159.4, 148.7, 147.8, 139.1, 133.6, 127.4, 126.3, 116.4, 115.2, 114.6, 111.9, 110.7, 84.7, 82.7, 60.2, 58.2, 57.6, 56.1, 56.1, 55.7, 54.0, 45.9, 28.9; IR (CH_2Cl_2 film, cm^{-1}) 3424 (m), 2925 (s), 1728 (m), 1666 (w), 1612 (m), 1596 (m), 1572 (m), 1519 (s), 1464 (s), 1410 (m), 1377 (m), 1356 (m), 1262 (s), 1226 (m), 1187 (m), 1262 (s), 1226 (m), 1187 (m), 1133 (s), 1102 (s), 1016 (m), 945 (m), 863 (m), 807 (m); Product ratio (from S-QUINAP synthesis) was determined by HPLC, Chiralpak ASH column, 5% IPA in hexanes, 1.0 mL/ min, 230 nm, $t_R^{(major)} = 10.77$ min, area = 3.56×10^4 (99.6%), $t_R^{(minor)} = 16.73$ min, area = 1.35×10^2 (0.4%) (99% ee); LRMS (m/z) 460.0 [M+1]; HRMS (TOF ES) calcd for $C_{23}H_{26}BrNO_4$ [M+1] 460.1123, Found 460.1126 Error 0.7 ppm; Product of R-QUINAP: $[\alpha]_D^{25} = -30^\circ$ (c = 0.4, CH_2Cl_2).

1,2,3,4-tetrahydro-6,7-dimethoxy-1-(3-methoxyprop-1-ynyl)-2-methylisoquinoline (4d)

The general procedure was used to add methylpropargyl ether (127 μ L, 1.5 mmol, 10 eq.) to 3,4-dihydro-6,7-dimethoxy-2-methylisoquinolinium iodide (**4**, 50 mg, 0.15 mmol). Following column chromatography (Si gel, diethyl ether with 1% triethylamine), isoquinoline **4d** was isolated (29.3 mg, 0.106 mmol, 71%).; 1H NMR (500 MHz, CD_2Cl_2) δ 6.73 (s, 1H), 6.57 (s, 1H), 4.42 (s, 1H), 4.11 (t, 2H, $J = 1.5$ Hz), 3.78 (s, 3H), 3.77 (s, 3H), 3.32 (s, 3H), 2.88 (m, 3H), 2.72 (m, 3H), 2.48 (s, 3H); ^{13}C NMR (400 MHz, $CDCl_3$) δ

148.6, 147.8, 127.5, 126.1, 111.9, 110.7, 84.8, 82.2, 60.2, 87.5, 56.3, 56.2, 56.1, 48.6, 46.7, 28.7; IR (CH₂Cl₂, cm⁻¹) 3632 (w), 3403 (m), 2927 (s), 1641 (m), 1612 (m), 1503 (m), 1466 (m), 1409 (m), 1370 (m), 1257 (s), 1226 (s), 1098 (s), 944 (m), 805 (m); Product ratio (from R-QUINAP synthesis) was determined by HPLC, Chiralpak OJ column, 10% IPA in hexanes, 1.0 mL/ min, 230 nm, $t_R^{(major)} = 29.91$ min, area = 2.88×10^4 (98.7%), $t_R^{(minor)} = 17.99$ min, area = 3.8×10^2 (1.3%) (97% ee); LRMS (m/z) 276.2 [M+1]; HRMS (TOF ES) calcd for C₁₆H₂₁NO₃ [M+1] 276.1599, Found 276.1598 Error 0.4 ppm; $[\alpha]_D^{25} = +0.4^\circ$ (c = 0.245, CH₂Cl₂).

methyl 3-(1,2,3,4-tetrahydro-6,7-dimethoxy-2-methyl-1-phenylisoquinolin-1-yl)propionate (5e)

The general procedure was used to add methyl propionate (102 μ L, 1.22 mmol, 10 eq.) to 3,4-dihydro-6,7-dimethoxy-2-methyl-1-phenylisoquinolinium iodide (**5**, 50 mg, 0.122 mmol). Following column chromatography (Si gel, 1:1:1% hexanes: diethyl ether: triethylamine), **5e** (39.3 mg, 0.108 mmol, 88%) was isolated.; ¹H NMR (600 MHz, CD₂Cl₂) δ 7.58 (d, 1H, $J = 7.2$ Hz), 7.33 (t, 2H, $J = 7.2$ Hz), 7.28 (t, 1H, $J = 6.9$ Hz), 6.56 (s, 1H), 6.13 (s, 1H), 3.84 (s, 3H), 3.79 (s, 3H), 3.55 (s, 3H), 3.25 (m, 1H), 3.09 (td, 1H, $J = 3, 12$ Hz), 2.96 (q, 1H, $J = 6$ Hz), 2.71 (dd, 1H, $J = 1.2, 15.6$ Hz), 2.19 (s, 3H); ¹³C NMR (400 MHz, CD₂Cl₂) δ 154.3, 148.8, 147.9, 143.8, 130.9, 128.4, 128.3, 128.2, 126.8, 112.6, 111.3, 86.5, 82.5, 67.2, 56.1, 56.0, 52.9, 49.2, 40.3, 29.2; IR (CH₂Cl₂ film, cm⁻¹) 3407 (w), 3059 (m), 2999 (m), 2951 (s), 2851 (s), 2801 (m), 2706 (m), 2612 (w), 2298 (m), 2218 (s), 1745 (s), 1611 (m), 1523 (m), 1513 (s), 1488 (m), 1466 (m), 1448 (m), 1404 (m), 1369 (m), 1356 (m), 1331 (m), 1246 (s), 1219 (m), 1148 (m), 1124 (m), 1101 (m), 1081 (m), 1064 (m), 1028 (m), 1015 (m), 978 (m), 960 (w), 951 (m), 924 (w), 897 (m), 858 (m), 827 (m), 807 (w); Product ratio (from S-QUINAP synthesis) was determined by HPLC, Chiralpak ASH column, 2% IPA in hexanes, 1.0 mL/ min, 230 nm, $t_R = 9.68$ min, area = 3.2×10^7 (49%), $t_R = 11.14$ min, area = 3.3×10^7 (51%) (no ee); LRMS (m/z) 366.3 [M+1]; HRMS (TOF ES) calcd for C₂₂H₂₅NO₄ [M+1] 366.1705, Found 366.1702460.1125 Error 0.8 ppm.

2-(2-bromo-5-methoxybenzyl)-1,2-dihydro-7-methoxy-1-(2-(trimethylsilyl)ethynyl)isoquinoline (6b)

The general procedure was used to add trimethylsilyl acetylene (160 μ L, 1.139 mmol, 10 eq.) to 7-methoxy-2-(2-bromo-5-methoxy)-benzylisoquinolinium bromide **6** (50 mg, 0.114 mmol). The reaction was conducted at -20 °C. The crude solution was purified by column chromatography (Si gel, 2:1:1% hexanes:

ethyl acetate: triethylamine) to yield **6b** as a colorless oil that turned brown over time (34.8 mg, 0.076 mmol, 67%).; ¹H NMR (600 MHz, CD₂Cl₂) δ 7.46 (d, 1H, *J* = 8.4 Hz), 6.99 (d, 1H, *J* = 3 Hz), 6.88 (d, 1H, *J* = 7.8 Hz), 6.735 (m, 2H), 6.64 (d, 1H, *J* = 3 Hz), 6.07 (d, 1H, *J* = 7.2 Hz), 5.44 (d, 1H, *J* = 7.8 Hz), 5.08 (s, 1H), 4.50 (d, 1H, *J* = 15 Hz), 4.28 (d, 1H, *J* = 15 Hz), 3.74 (s, 3H), 3.72 (s, 3H), 0.15 (s, 9H); ¹³C NMR (400 MHz, CD₂Cl₂) δ 159.6, 158.2, 137.8, 134.9, 133.9, 129.1, 125.6, 124.8, 116.3, 114.9, 114.7, 113.9, 112.3, 103.1, 99.7, 90.5, 56.9, 55.8, 55.6, 52.6, 0.0; IR 3371 (m), 3061 (m), 3001 (m), 2957 (m), 2934 (m), 2900 (m), 2836 (m), 2158 (m), 1639 (m), 1626 (m), 1620 (m), 1613 (m), 1578 (m), 1573 (m), 1568 (m), 1475 (s), 1468 (m), 1464 (m), 1454 (m), 1443 (m), 1435 (m), 1428 (m), 1422 (m), 1414 (m), 1393 (m), 1370 (m), 1348 (m), 1291 (m), 1249 (s), 1214 (m), 1192 (m), 1154 (m), 1104 (m), 1054 (m), 1039 (m), 1019 (m), 990 (m), 934 (m), 844 (m), 815 (m); Product ratio (from S-QUINAP synthesis) was determined by HPLCC, Chiralpak OD column, 1% IPA in hexanes, 1.0 mL/ min, 230 nm, *t*_R^(major) = 8.41 min, area = 7.9 x 10³ (91.4%), *t*_R^(minor) = 7.57 min, area = 7.4 x 10² (8.6%) (83% ee); LRMS (*m/z*) 456.1 [M+1]; HRMS (TOF ES) calcd for C₂₃H₂₆BrNO₂Si [M+1] 456.0994, Found 456.0982 Error 2.6 ppm; Product of S-QUINAP [α]_D²⁵ = +13 ° (c = 0.44, CH₂Cl₂).

resin-bound 3,4-dihydro-6-methoxy-isoquinolin-7-ol (7)

A 15 mL plastic column was charged with functionalized 500 μm polystyrene resin (“macrobeads,” 300 mg, 0.435 mmol), capped with a septum and a 3-way stopcock. The beads were dried under high vacuum overnight before being swollen in dry methylene chloride (5 mL) for 1 h. The solvent was drained under positive Argon pressure before the addition of neat triflic anhydride (73 μL). The mixture stood 5 min before being drained under positive Argon pressure. The resin was suspended in 4% TfOH solution in methylene chloride (5.8 mL) and agitate slightly for 10 min. The mixture was then drained under positive Argon pressure, and the beads were washed 2x with methylene chloride (10 mL) before being taken up in a minimal amount of methylene chloride. Fresh 2,6-lutidine (405 μL, 3.48 mmol, 8.0 eq.) was added to the mixture, which caused a color change. 7-hydroxy-6-methoxy-3,4-dihydroisoquinoline (230 mg, 1.3 mmol, 3.0 eq.) was then added in DMF (2 mL), and the mixture was agitated 20 h at rt. The beads were drained again under positive Argon pressure before being washed with DMF (rinsed 3x 5 min, suspended 2x 20 min), THF (1x 15 min, 1x 120 min), methylene chloride (2x 5 min, 2x 120 min). The beads were then dried under house vacuum 30 min and high vacuum overnight.; MAS ¹H (600 MHz, CDCl₃): δ 8.2 (br), 6.8 (br), 6.6 (br), 3.7 (br), 2.7 (br).

resin-bound 2-(2-bromo)-benzyl-3,4-dihydro-7-hydroxy-6-methoxyisoquinolinium bromide (8)

A vial with a septum top was charged with resin **7** (148 mg, 0.214 mmol) before being sealed and purged with Argon. 2-bromo-benzylbromide (210 mg, 0.86 mmol, 4.0 eq.) was dissolved in diethyl ether (3 mL), which was added to the resin via syringe. The mixture was agitated at rt overnight. The beads were drained via canula under positive Argon pressure and washed 3x with methylene chloride (5 min) before being dried under house vacuum and high vacuum.; MAS ^1H (600 MHz, CDCl_3): δ 9.5 (br), 7.9 (br), 7.5 (br), 7.4 (br), 7.2 (br), 5.6 (br), 3.9 (br), 3.8 (br), 3.2 (br).

resin-bound 2-(2-bromo)-benzyl-1,2,3,4-tetrahydro-7-hydroxy-6-methoxy-1-(2-phenylethynyl)-isoquinoline (9a)

A 2 mL vial was charged with resin **8** (71 mg, 0.103 mmol) before the addition of S-QUINAP:CuBr stock solution (738 μL , 0.005 mmol, 5 mol%). Phenylacetylene (113 μL , 1.03 mmol, 10 eq.) was added, and the mixture was cooled to $-80\text{ }^\circ\text{C}$ before the addition of triethylamine (14 μL , 0.103 mmol, 1 eq.). The reaction proceeded at this temperature with occasional agitation 40 h before the solvent was removed via canula. The beads were washed 3x with methylene chloride (5 min). The resin was dried under house vacuum and high vacuum.; MAS ^1H (600 MHz, CDCl_3): δ 7.62 (br), 7.58 (br), 7.4 (br), 7.2 (br), 7.1 (br), 6.82 (br), 6.5 (br), 4.8 (br), 4.2 (br), 3.2 (br), 3.1 (br), 2.9 (br), 2.8 (br), 2.7 (br).

2-(2-bromo)-benzyl-1,2,3,4-tetrahydro-7-hydroxy-6-methoxy-1-(2-phenylethynyl)-isoquinoline (9b)

Following MAS ^1H NMR analysis, the compound was cleaved from the resin. An Eppendorf tube was charged with resin **9a** (27.4 mg, 0.027 mmol) followed by a cleavage cocktail solution (95/5 THF/HF-py v/v, 100 μL solution per 10 mg resin). The sample was vortexed at rt 1.5 h. The sample was treated with methoxytrimethylsilane (volume equivalent to that of the cleavage cocktail) and vortexed an addition 30 min. The sample was filtered through a pipette plugged with glass wool and celite. The resin and celite

were washed 2x with methylene chloride, 2x with acetonitrile, and 2x with methylene chloride. The filtrates were collected and concentrated *in vacuo* to yield a clear oil (10.3 mg, 0.023 mmol, 84%).; ¹H NMR (500 MHz, CDCl₃): δ 7.67 (dd, 1H, *J* = 9.5, 2 Hz), 7.62 (dd, 1H, *J* = 10, 1.5 Hz), 7.52 (m, 2H), 7.35 (m, 4H), 7.18 (td, 1H, *J* = 10, 2.5 Hz), 6.91 (s, 1H), 6.62 (s, 1H), 5.66 (s, 1H), 4.83 (s, 1H), 4.08 (d, 2H, *J* = 2.5 Hz), 3.86 (s, 3H), 3.20 (td, 1H, *J* = 13.5, 5 Hz), 3.05 (m, 1H), 2.88 (m, 1H), 2.76 (dt, 1H, *J* = 19, 4.5 Hz); ¹³C NMR (400 MHz, CDCl₃): δ 145.8, 143.7, 137.7, 132.7, 131.7, 130.6, 128.4, 128.1, 127.9, 127.9, 127.2, 125.3, 124.7, 123.1, 113.3, 110.6, 87.6, 86.4, 58.8, 55.8, 54.2, 45.8, 28.8; IR (CH₂Cl₂ film, cm⁻¹) 3678 (w), 3653 (w), 3414 (s), 2935 (m), 2836 (m), 1687 (s), 1656 (s), 1642 (m), 1598 (m), 1567 (m), 1548 (w), 1514 (m), 1488 (m), 1464 (m), 1441 (m), 1381 (m), 1342 (m), 1286 (s), 1228 (m), 1170 (m), 1127 (m), 1024 (m); Product ratio was determined to be 75% by HPLC, Chiralpak AD-RH column, 60% MeCN in H₂O, 1.0 mL/ min, 230 nm, *t*_R^(major) = 9.8 min, area = 1.17 x 10⁴ (88%), *t*_R^(minor) = 13.2 min, area = 1.6 x 10³ (12%); LRMS (*m/z*) 447.9 [M+1]; HRMS (TOF ES) calcd for C₂₅H₂₂BrNO₃ [M+1] 448.0912, Found 448.0917 Error 1.1 ppm; Product of R-QUINAP: [α]_D²⁵ = -65 ° (c = 0.2, CH₂Cl₂).

1,2,3,4-tetrahydro-6,7-dimethoxy-1-(2-(3,4-dimethoxyphenyl)ethynyl)-2-methylisoquinoline (10)

The general procedure was used to add 3,4-dimethoxy phenylacetylene (972 mg, 6 mmol, 10 eq.) to 3,4-dihydro-6,7-dimethoxy-2-methylisoquinolinium iodide (200 mg, 0.6 mmol). The crude solution was purified by column chromatography (Si gel, 5% methanol in methylene chloride, 1% triethylamine) to yield **12** as a white solid (205 mg, 0.56 mmol, 93%).; ¹H NMR (400 MHz, CDCl₃) δ 7.04 (dd, 1H, *J* = 1.6, 8.4 Hz), 6.91 (d, 1H, *J* = 1.6 Hz), 6.84 (s, 1H), 6.77 (d, 1H, *J* = 8.4 Hz), 6.59 (s, 1H), 4.59 (s, 1H), 3.86 (s, 3H), 3.84 (s, 3H), 3.05 (m, 1H), 2.93 (m, 2H), 2.70 (m, 1H), 2.61 (s, 3H); ¹³C NMR (400 MHz, CDCl₃) δ 149.2, 148.4, 148.0, 147.3, 127.2, 125.5, 124.9, 115.3, 114.4, 111.1, 110.8, 110.3, 85.9, 85.9, 56.6, 55.9, 55.8, 55.8, 55.8, 48.7, 43.7, 28.4; IR (CH₂Cl₂ film, cm⁻¹) 3273 (w), 2998 (m), 2936 (s), 2835 (s), 2835 (s), 2799 (s), 2214 (w), 1599 (s), 1578 (s), 1513 (s), 1464 (s), 1409 (s), 1369 (s), 1355 (s), 1321 (s), 1264 (s), 1243 (s), 1227 (s), 1204 (s), 1167 (s), 1138 (s), 1102 (m), 1062 (w), 1025 (s), 941 (w), 854 (m), 807 (m); Product ratio (from S-QUINAP synthesis) was determined by SFC, Chiralpak OD column, 4.5 min gradient: 5% methanol -> 50% methanol, 1.0 mL/ min, 230 nm, *t*_R^(major) = 3.92 min, area = 1.2 x 10⁵ (98%), *t*_R^(minor) = 4.81 min, area = 2.3 x 10³ (2%) (96% ee); LRMS (*m/z*) 368.2 [M+1]; HRMS (TOF ES) calcd for C₂₂H₂₉NO₄ [M+1] 368.1862, Found 368.1863 Error 0.3 ppm; Product of S-QUINAP: [α]_D²⁵ = -52 ° (c = 1.64, CH₂Cl₂); mp = 38 -42 °C.

homolaudanosine (11)

A flask was charged with propargyl amine **12** (133 mg, 0.362 mmol) and 5% Pd/C (154 mg, 0.072 mmol, 0.2 eq.) before being sealed and purged with Nitrogen. 1:1 ethanol: ethyl acetate (5 mL) was added, and the flask was evacuated and flushed with Hydrogen three times. The reaction stirred at rt 20 h before the catalyst was filtered off and the solution concentrated *in vacuo*. The crude product was purified by column chromatography (Si gel, 2% methanol in methylene chloride, 1% triethylamine) to yield a clear oil (102 mg, 0.27 mmol, 76%); Compare to reported compound: Meyers, A. I., et al. *Tetrahedron*. **1987**, *43*, 5095-5108.; ^1H NMR (600 MHz, CDCl_3) δ 6.78 (d, 1H, $J = 7.8$ Hz), 6.72 (dd, 2H, $J = 1.8$ Hz, 10.2 Hz), 6.56 (s, 1H), 6.53 (s, 1H), 3.84 (s, 3H), 3.84 (s, 3H), 3.83 (s, 3H), 3.82 (s, 3H), 3.42 (t, 1H, $J = 5.4$ Hz), 3.17 (m, 1H), 2.84 (m, 3H), 2.54 (m, 1H), 2.47 (s, 3H), 2.05 (m, 2H), 1.23 (t, 1H, $J = 7.2$ Hz); IR (CH_2Cl_2 film, cm^{-1}) 3415 (m), 2995 (s), 2935 (s), 2833 (s), 2787 (m), 2598 (w), 2278 (w), 2041 (w), 1843 (w), 1609 (s), 1590 (m), 1516 (w), 1464 (s), 1418 (m), 1372 (m), 1356 (m), 1338 (m), 1319 (m), 1260 (m), 1227 (s), 1155 (m), 1139 (m), 1104 (m), 1069 (m), 1000 (m), 931 (w), 859 (m), 806 (m); LRMS (m/z) 372.3 [$\text{M}+1$]; HRMS (TOF ES) calcd for $\text{C}_{22}\text{H}_{29}\text{NO}_4$ [$\text{M}+1$] 372.2155, Found 372.2167 Error 2.1 ppm; Product of S-QUINAP: $[\alpha]_D^{25} = +3^\circ$ ($c = 1.3$, CHCl_3).

2-(2-bromo)-benzyl-3,4-dihydro-6-methoxy-7-(*t*-butyl-dimethylsilyl ether)-isoquinolinium bromide(12)

A flask was charged with 3,4-dihydro-7-hydroxy-6-methoxyisoquinoline (2.5 g, 14 mmol), TBDMSCl (4.3 g, 28 mmol, 2 eq.), and imidazole (1.9 g, 28 mmol, 2 eq.) before being sealed and purged with Nitrogen. DMF (70 mL) was added, and the solution was stirred at rt overnight. The solution was partitioned between water and diethyl ether. The organic layer was separated, and the extraction was repeated twice. The organic layers were combined, dried with magnesium sulfate, and concentrated. The intermediate was purified by column chromatography (Si gel, 2.5:1 hexanes: diethyl ether $\rightarrow$ 1:1 hexanes: diethyl ether) to yield a clear oil (3.07, 75%). A flask was charged with this oil (3.05 g, 10.5 mmol) and distilled benzylbromide (3.5 g, 12 mmol, 1.3 eq.) before the addition of diethyl ether (30 mL). The reaction was heated to reflux overnight and stirred. Following cooling, the precipitate was collected, washed with diethyl ether, and dried under high vacuum to give an off-white solid (5.33 g, 94%). ^1H NMR (400 MHz, CDCl_3): δ 9.43 (s, 1H), 8.01 (dd, 1H, $J = 2, 7.6$ Hz), 7.61 (dd, 1H, $J = 1.2, 8$ Hz), 7.43 (td, 1H, $J = 1.2, 7.6$ Hz), 7.32

(td, 2H, $J = 2$, 7.8 Hz), 6.83 (s, 1H), 5.64 (s, 2H), 4.01 (t, 2H, $J = 8$ Hz), 3.91 (s, 3H), 3.27 (t, 2H, $J = 8$ Hz), 0.96 (s, 9H), 0.14 (s, 6H); ^{13}C NMR (400 MHz, CDCl_3): δ 164.7, 159.9, 144.9, 133.9, 133.4, 132.8, 131.6, 130.4, 128.7, 125.1, 125.1, 117.0, 111.3, 62.8, 56.2, 48.0, 25.7, 18.3, -0.07, -4.6; IR (CH_2Cl_2 film, cm^{-1}) 3415 (m), 3010 (m), 2953 (s), 2930 (s), 2894 (m), 2857 (s), 1646 (s), 1604 (s), 1559 (s), 1519 (s), 1491 (s), 1472 (s), 1454 (m), 1402 (s), 1336 (m), 1307 (s), 1282 (s), 1244 (s), 1148 (s), 1225 (m), 1205 (m), 1101 (m), 1029 (m), 1012 (m), 928 (m), 887 (m), 841 (m), 826 (m), 803 (m); LRMS (m/z) 460.1 [M-Br]; HRMS (TOF ES) calcd for $\text{C}_{23}\text{H}_{31}\text{Br}_2\text{NO}_2\text{Si}$ 460.1250 [M-Br], Found 460.1256 Error 1.3 ppm; mp = 184 – 185 °C.

2-(2-bromobenzyl)-1-ethynyl-1,2,3,4-tetrahydro-6-methoxyisoquinolin-7-ol (**13**)

The general procedure was used for the addition of trimethylsilylacetylene to iminium **13** (1.0 g, 1.8 mmol) to yield the corresponding propargyl amine (915 mg, 1.64 mmol, 89%) as an oil. A flask was charged with this oil (428 mg, 0.767 mmol) and methanol (20 mL), followed by potassium carbonate (320 mg, 2.3 mmol, 3 eq.). The reaction stirred at rt until disappearance of the starting material by TLC (24 h). The crude mixture was concentrated *in vacuo*, taken up in diethyl ether, dried with magnesium sulfate, filtered, and concentrated to give **14** as a white solid (281 mg, 0.75 mmol, 98%); ^1H NMR (400 MHz, CDCl_3): δ 7.563 (d, 2H, $J = 1.2$ Hz), 7.30 (td, 1H, $J = 1.2$, 7.6 Hz), 7.14 (td, 1H, $J = 1.2$, 7.6 Hz), 6.78 (s, 1H), 6.57 (s, 1H), 4.57 (s, 1H), 3.92 (s, 2H), 3.84 (s, 3H), 3.05 (m, 2H), 2.80 (m, 1H), 2.67 (m, 1H), 2.46 (d, 1H, $J = 2$ Hz); ^{13}C NMR (400 MHz, CDCl_3): δ 145.9, 143.8, 137.7, 132.8, 130.5, 128.5, 127.6, 127.2, 125.3, 124.8, 113.1, 110.7, 81.9, 74.3, 58.6, 55.9, 53.5, 45.4, 28.7; IR (CH_2Cl_2 film, cm^{-1}) 3212 (m), 3292 (s), 3058 (m), 2918 (m), 2835 (m), 2362 (w), 1622 (m), 1601 (m), 1566 (m), 1514 (s), 1464 (s), 1441 (s), 1378 (s), 1341 (m), 1288 (s), 1267 (s), 1232 (m), 1209 (m), 1166 (m), 1128 (m), 1113 (m), 1095 (m), 1044 (m), 1024 (m), 945 (m), 874 (m), 848 (m), 829 (m), 812 (m); LRMS (m/z) 372.1 [M+1]; HRMS (TOF ES) calcd for $\text{C}_{19}\text{H}_{18}\text{BrNO}_2$ [M+1] 372.0599, Found 372.0605 Error 1.6 ppm; Product of R-QUINAP: $[\alpha]_D^{25} = -57^\circ$ (c = 2.3, CH_2Cl_2), Product of S-QUINAP: $[\alpha]_D^{25} = +53^\circ$ (c = 1.2, CH_2Cl_2).

2-(2-bromobenzyl)-1,2,3,4-tetrahydro-6-methoxy-1-vinylisoquinolin-7-ol (**14**)

A flask was charged with tetrahydroisoquinoline **14** (23.6 mg, 0.063 mmol) and Lindlar's catalyst (27 mg, 0.013 mmol, 0.2 eq.), followed by 1:1 ethanol: ethyl acetate (2 mL). The flask was evacuated and flushed with Hydrogen three times before stirring at rt until the starting material disappeared, as visualized by TLC. The catalyst was filtered away, and the solution was concentrated *in vacuo*. The crude product was purified by chromatography (Si gel, 4:1 hexanes: ethyl acetate) to yield allyl amine **15** as an oil (9.4 mg, 0.025

mmol, 40%); ^1H NMR (400 MHz, CDCl_3): δ 7.60 (dd, 1H, $J = 1.2, 7.6$ Hz), 7.30 (t, 1H, $J = 7.6$ Hz), 7.12 (td, 1H, $J = 1.2, 7.6$ Hz), 6.71 (s, 1H), 6.58 (s, 1H), 5.91 (m, 1H), 5.31 (s, 1H), 5.28 (dd, 1H, $J = 1.6, 7.2$ Hz), 4.05 (d, 1H, $J = 8.4$ Hz), 4.039 (d, 1H, $J = 15.2$ Hz), 3.56 (s, 3H), 3.63 (d, 1H, $J = 15.2$ Hz), 3.04 (m, 1H), 2.85 (m, 1H), 2.74 (m, 1H), 2.58 (m, 1H); ^{13}C NMR (400 MHz, CDCl_3): δ 145.2, 143.5, 139.7, 138.9, 132.5, 130.3, 129.0, 128.1, 127.2, 126.2, 124.3, 118.2, 113.9, 110.5, 66.9, 57.9, 55.9, 46.9, 28.7; IR: (CH_2Cl_2 film, cm^{-1}) 3426 (s), 2958 (m), 2359 (w), 2339 (w), 2096 (m), 1644 (s), 1514 (m), 1464 (m), 1441 (m), 1377 (m), 1288 (m), 1261 (m), 1234 (m), 1127 (m), 1094 (m), 1023 (m), 945 (m), 846 (m); Product ratio (from S-QUINAP synthesis) was determined, Chiralpak OD column, 1% IPA in hexanes, 1.0 mL/min, 230 nm, $t_{\text{R}}^{(\text{major})} = 21.6$ min, area = 1.8×10^4 (99%), $t_{\text{R}}^{(\text{minor})} = 20.2$ min, area = 207 (1%) (98% ee); LRMS (m/z) 373.9 [M+1]; HRMS (TOF ES) calcd for $\text{C}_{19}\text{H}_{20}\text{BrNO}_2$ [M+1] 374.0755, Found 374.0755 Error 0.0 ppm; Product of S-QUINAP [α] $^{25}_{\text{D}} = +50^\circ$ ($c = 0.3$, CH_2Cl_2).

2-(2-bromo)-benzyl-3,4-dihydro-7-hydroxy-1-vinylisoquinoline (15)

Compare to reported compound: Taylor, S. J., Taylor, A. M., Schreiber, S. L. *Angew. Chem. Int. Ed.* **2004**, *43*, 1681.

2-(2-bromo)-benzyl-3,4-dihydro-6,7-dimethoxy-1-vinylisoquinoline (16)

Compound **17** was prepared using procedures similar to those used for the preparation of **15**, starting from 3,4-dihydro-6,7-dimethoxyisoquinoline hydrochloride.; ^1H NMR (600 MHz, CDCl_3): δ 7.60 (d, 1H, $J = 7.2$ Hz), 7.54 (d, 1H, $J = 8.4$ Hz), 7.30 (t, 1H, $J = 7.2$), 7.12 (t, 1H, $J = 7.2$), 6.62 (s, 1H), 6.61 (s, 1H), 5.93 (m, 1H), 5.33 (d, 1H, $J = 7.2$), 5.32 (d, 1H, $J = 16.8$), 4.08 (d, 1H, $J = 9$ Hz), 4.04 (d, 1H, $J = 15$ Hz), 3.86 (s, 3H), 3.83 (s, 3H), 3.65 (d, 1H, $J = 15$ Hz), 3.04 (m, 1H), 2.84 (m, 1H), 2.75 (m, 1H), 2.59 (m, 1H); ^{13}C NMR (400 MHz, CDCl_3): δ 147.6, 147.0, 139.9, 138.8, 132.5, 130.3, 128.1, 127.2, 126.8, 124.3, 118.2, 111.2, 111.1, 66.9, 57.9, 55.9, 55.8, 46.7, 28.5; IR (CH_2Cl_2 film, cm^{-1}) 3406 (m), 3073 (w), 2923 (m), 2782 (m), 1642 (w), 1612 (w), 1514 (m), 1464 (m), 1452 (m), 1442 (m), 1352 (m), 1298 (m), 1259 (m), 1227 (m), 1095 (m), 1024 (m), 806 (m); LRMS (m/z) 387.9 [M+1]; HRMS (TOF ES) calcd for $\text{C}_{20}\text{H}_{22}\text{BrNO}_2$ [M+1] 388.0912, Found 388.0905 Error 1.8 ppm; [α] $^{25}_{\text{D}} = +40^\circ$ ($c = 0.15$, CH_2Cl_2).

Yeast Growth Assays

Yeast extract media (YPD) was prepared by mixing yeast extract (10 g), peptone (20 g), and dextrose (20 g) in milliQ water (1.0 L) and stirring until homogeneous. YPD agar was prepared by adding BactoAgar

(20 g) to the YPD media, followed by sterilization in the autoclave. Aliquots (~30 mL) of this solution were poured into plastic petri dishes in a sterile hood and allowed to set. Yeast (BY4741) was cultured by swabbing a small amount of yeast from an existing colony onto the agar with a sterilized wire loop. The plates were incubated at 30 °C for ~48 h before a small amount was transferred by sterile wire loop to a covered but unsealed 50 mL plastic vial with YPD (~10 mL). This solution was agitated at 36 °C several hours before cell density was determined by analyzing the 600 nm absorption of an aliquot. An aliquot of this solution was diluted with YPD such that a 90 µL aliquot contained $\sim 2.5 \times 10^4$ cells. Inhibition experiments were performed in a 96-well plate. Each well was treated with yeast/ YPD (90 µL), followed by addition of small molecule in DMSO (10 µL) to make the final concentration of the compound 40 µM. Each compound was tested in triplicate. The plate was incubated at rt and at given time points was agitated on an orbital shaker to ensure even distribution of the yeast in the wells before UV absorption analysis.

Fig S1: Optical density of yeast as a function of time in the presence of tetrahydroisoquinoline derivatives