Enantioselective Cyanosilylation of α,α -Dialkoxy Ketones Catalyzed by Proline-Derived in Situ Prepared *N*-Oxide as Bifunctional Organocatalyst

Bo Qin,[†] Xiaohua Liu,[†] Jian Shi,[†] Ke Zheng,[†] Haitao Zhao[†] and Xiaoming Feng^{,†,‡}*

[†]Key Laboratory of Green Chemistry & Technology (Sichuan University), Ministry of Education,
College of Chemistry, Sichuan University, Chengdu 610064, China

[‡]State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Chengdu 610041,
China

xmfeng@scu.edu.cn

SUPPORTING INFORMATION

1. General.....	2
2. Typical procedure for catalyst preparation	2
3. Preparation of the acetal ketones	5
4. General procedure for the preparation of racemic cyanosilylation of acetal ketones.....	7
5. General procedure for the <i>N,N'</i> -dioxide-catalyzed asymmetric cyanosilylation of acetal ketones	8
6. General procedure for the conversion of cyanohydrin trimethylsilyl ether to its corresponding derivative.....	16
7. Character Data of Cyanohydrin 10.....	19
8. Copy of ¹ H NMR and ¹³ C NMR spectra for the catalytic mechanism	19

1. General

^1H NMR spectra were recorded on commercial instruments (400 or 300 MHz). Chemical shifts were reported in ppm from tetramethylsilane with the solvent resonance as the internal standard (CDCl_3 , $\delta = 7.26$). Spectra are reported as follows: chemical shift (δ ppm), multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet), coupling constants (Hz), integration, and assignment. ^{13}C NMR spectra were collected on commercial instruments (100 or 75 MHz) with complete proton decoupling. Chemical shifts are reported in ppm from the tetramethylsilane with the solvent resonance as internal standard (CDCl_3 , $\delta = 77.0$). The enantiomeric excesses were determined by HPLC analysis on chiral DAICEL CHIRALCEL OJ, OJ-H, OD, OD-H, OF or CHIRALPAK AD-H, AS-H column at 210 nm, 220 nm and 254 nm. Optical rotations were measured on a commercial polarimeter. All ketones, *L*-proline, (*S*)-1-*N*-Boc-piperidine-2-carboxylic acid, and trimethylsilyl cyanide (TMSCN) were commercially purchased and used without further purification. CH_2Cl_2 , and $\text{ClCH}_2\text{CH}_2\text{Cl}$ were distilled over CaH_2 before use.

2. Typical procedure for catalyst preparation

To a solution of *L*-*N*-Boc-proline (860 mg, 4 mmol) in CH_2Cl_2 (40 mL) was added Et_3N (0.62 mL, 4.4 mmol), isobutyl carbonochloridate (576 mg, 4.4 mmol) at $0\text{ }^\circ\text{C}$ under stirring. After 15 minutes, cyclopentanamine (374 mg, 4.4 mmol) was added. The reaction was allowed to warm to room

temperature and detected by TLC. The mixture were washed with 1M KHSO₄, saturated NaHCO₃ and brine, then dried over anhydrous MgSO₄ and concentrated. The residue in CH₂Cl₂ (4 mL) was added TFA (4 mL) and stirred until the reaction was finished. Then, the solvent was evaporated, and CH₂Cl₂ (10 mL) was added. The pH value was brought into the range of 8–10 by the addition of 1M NaOH. The aqueous phase was extracted with CH₂Cl₂ (5×20 mL). The combined organic phase was washed with brine, dried over anhydrous MgSO₄ and evaporated in vacuo. The residue was directly used for next step. To a solution of *L*-N-Boc-proline-2-carbonamide in CH₃CN (4 mL) was added K₂CO₃ (608 mg, 4.4 mmol) and 1, 3-dibromopropane (204 uL, 2 mmol) under stirring. It was kept at 80 °C, and monitored by TLC. Then, K₂CO₃ was removed by filtration. The residue was concentrated and purified by silica gel column chromatography (EtOAc) to give a white product **3a** (0.737 g, 91% yield).

To a solution of **3a** (405 mg, 1 mmol) in CH₂Cl₂ (20 mL) was added *m*-chlorobenzoperoxoic acid (*m*-CPBA) (383 mg, 2.2 mmol) at 0 °C under stirring. After the reaction was completed, the mixture was concentrated and purified by the basic aluminum oxide column chromatography (EtOAc / MeOH = 1:1) to give a white powder **2b** (379 mg, 87% yield).

catalyst **2b**: white powder; $[\alpha]_D^{25} = -34.1$ ($c = 0.41$, CH₂Cl₂); ¹H NMR (300 MHz, CDCl₃): $\delta = 10.66$ (d, $J = 2.9$ Hz, 2H), 4.11-4.13 (m, 2H), 3.45-3.54 (m, 6H), 3.14-3.33 (m, 4H), 2.56-2.62 (m, 2H), 2.33-2.37 (m, 6H), 1.84-1.85 (m, 6H), 1.53-1.62 (m, 8H) 1.37-1.39 (m, 4H); ¹³C NMR (75 MHz, CDCl₃): $\delta = 166.6, 77.4, 67.8, 64.8, 50.2, 33.0, 32.6, 27.4, 23.6, 23.5, 20.0, 19.7$; ESI-HRMS Calcd for (C₂₃H₄₀N₄O₄⁺) 436.3044, Found: 436.3109.

catalyst **3a**: white powder; m.p. 88-90 °C; $[\alpha]_D^{25} = -120.9$ ($c = 0.41$, CHCl₃); ¹H NMR (400 MHz, CDCl₃): $\delta = 7.26$ (s, 2H), 4.19 (d, $J = 5.6$ Hz, 2H), 3.17 (s, 2H), 2.97 (d, $J = 3.2$ Hz, 2H), 2.53 (s, 4H), 2.29 (d, $J = 5.6$ Hz, 2H), 1.64-2.18 (m, 22H), 1.37 (s, 4H); ¹³C NMR (100 MHz, CDCl₃): $\delta =$

174.0, 67.7, 54.0, 53.8, 50.1, 33.0 (d, $J = 5.3$ Hz), 30.4, 28.4, 24.0, 23.5 (d, $J = 3.0$ Hz); ESI-HRMS Calcd for ($C_{23}H_{40}N_4O_2+H^+$) 405.3224, Found: 405.3558.

catalyst **3b**: white powder; m.p. 84-86 °C; $[\alpha]_D^{25} = -143.2$ (c = 0.41, $CHCl_3$); 1H NMR (300 MHz, $CDCl_3$): $\delta = 7.26$ (m, 2H), 3.05-3.16 (m, 2H), 2.93-2.97 (m, 2H), 2.68-2.70 (m, 2H), 2.43-2.48 (m, 4H), 2.05-

2.23 (m, 4H), 1.56-1.80 (m, 8H), 0.75-0.78 (m, 4H), 0.43-0.45 (m, 4H); ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 176.2, 67.7, 54.0$ (d, $J = 5.7$ Hz), 30.5, 28.2, 24.1, 21.9, 6.5 (d, $J = 4.4$ Hz); ESI-HRMS Calcd. for ($C_{19}H_{32}N_4O_2+H^+$) 349.2598, Found: 349.4938.

catalyst **3c**: white powder; m.p. 110-112 °C; $[\alpha]_D^{25} = -102.4$ (c = 0.42, $CHCl_3$); 1H NMR (400 MHz, $CDCl_3$): $\delta = 7.21$ (d, $J = 8.8$ Hz, 2H), 3.67-3.74 (m, 2H), 3.14 (t, $J = 6.8$ Hz, 2H), 2.92-2.96 (m, 2H), 2.50 (t, $J = 7.2$ Hz, 4H), 2.23-2.29 (m, 2H), 2.08-2.18

(m, 4H), 1.57-1.84 (m, 16H), 1.31-1.40 (m, 4H), 1.07-1.21 (m, 6H); ^{13}C NMR (100 MHz, $CDCl_3$): $\delta = 173.6, 67.7, 54.1, 53.8, 47.0, 33.0$ (d, $J = 2.3$ Hz), 30.5, 28.4, 25.4, 24.6 (d, $J = 6.9$ Hz), 24.1; ESI-HRMS Calcd for ($C_{25}H_{44}N_4O_2^+$) 432.3464, Found: 432.3955.

catalyst **3d**: white powder; m.p. 72-73 °C; $[\alpha]_D^{25} = -131.4$ (c = 0.42, $CHCl_3$); 1H NMR (400 MHz, $CDCl_3$): $\delta = 7.19$ (s, 2H), 3.16 (s, 2H), 2.86 (d, $J = 5.6$ Hz, 2H), 2.53 (s, 2H), 2.28 (d, $J = 7.2$ Hz, 2H), 2.14

(m, 4H), 1.62-1.81 (m, 8H), 1.34 (s, 18H); ^{13}C NMR (100 MHz, $CDCl_3$): $\delta = 174.0, 68.5, 54.1, 53.8, 50.0, 30.4, 29.7, 28.4, 24.0$; ESI-HRMS Calcd for ($C_{21}H_{40}N_4O_2+H^+$) 381.3224, Found: 381.3796.

catalyst **3e**: white powder; m.p. 66-68 °C; $[\alpha]_D^{25} = -121.6$ (c = 0.41, $CHCl_3$); 1H NMR (400 MHz, $CDCl_3$): $\delta = 7.32$ (s, 2H), 3.15-3.27 (m, 6H), 2.99-3.03 (m, 2H), 2.47-2.56 (m, 4H), 2.27-2.33 (m, 2H), 2.11-

2.18 (m, 4H), 1.64-1.86 (m, 8H), 1.45-1.52 (m, 4H), 1.30-1.39 (m, 4H), 0.93 (t, $J = 7.2$ Hz, 6H); ^{13}C

NMR (100 MHz, CDCl₃): δ = 174.6, 67.8, 54.0, 53.9, 38.4, 21.7, 30.5, 28.5, 24.2, 20.0, 13.7; ESI-HRMS Calcd for (C₂₁H₄₀N₄O₂+H⁺) 381.3224, Found: 381.3652.

catalyst **3f**: white powder; m.p. 114-115 °C; $[\alpha]_D^{25} = -160.2$ (c = 0.5, CHCl₃); ¹H NMR (400 MHz, CDCl₃): δ = 7.63 (d, *J* = 8.4 Hz, 2H); 7.27-7.37 (m, 10H), 5.10-5.14 (m, 2H), 3.17 (t, *J* = 8.0 Hz,

2H), 3.05 (dd, *J* = 10.0, 4.4 Hz, 2H), 2.60 (t, *J* = 7.2 Hz, 4H), 2.31 (q, *J* = 9.2 Hz, 2H), 2.07-2.21 (m, 2H), 1.68-1.83 (m, 8H), 1.49 (d, *J* = 6.8 Hz, 6H), ¹³C NMR (100 MHz, CDCl₃): δ = 173.8, 143.4, 128.6, 127.1, 125.8, 67.7, 54.2, 54.0, 47.6, 30.5, 28.5, 24.2, 22.2; ESI-HRMS Calcd for (C₂₉H₄₀N₄O₂+H⁺) 477.3224, Found: 477.3580.

catalyst **3g**: white powder; m.p. 130-132 °C; $[\alpha]_D^{25} = -140.4$ (c = 0.50, CHCl₃); ¹H NMR (400 MHz, CDCl₃): δ = 6.53 (s, 2H), 4.18 (m, 2H), 3.04 (m, 2H), 2.61 (m, 2H), 2.40 (m, 2H), 2.16-2.20 (m,

4H), 1.94 (m, 8H), 1.67 (m, 10H), 1.24-1.36 (m, 12H); ¹³C NMR (100 MHz, CDCl₃): δ = 173.8, 68.1, 54.6, 51.3, 50.1, 32.8, 30.5, 24.7, 23.6, 23.5, 23.3; ESI-HRMS Calcd for (C₂₅H₄₄N₄O₂+H⁺): 433.3537, Found: 433.3543.

catalyst **9**: pale powder; $[\alpha]_D^{25} = -62.2$ (c = 0.47, CHCl₃); ¹H NMR (400 MHz, CDCl₃): δ = 10.81 (d, *J* = 7.2 Hz, 1H), 7.12 (d, *J* = 8.0 Hz, 1H), 4.17-4.20 (m, 2H), 3.54-3.67 (m, 1H), 3.47-3.54 (m, 1H), 3.27-

3.31 (m, 3H), 3.17-3.20 (m, 1H), 2.37-2.48 (m, 3H), 2.50-2.65 (m, 2H), 2.14-2.22 (m, 1H), 1.95-1.99 (m, 6H), 1.69-1.82 (m, 2H), 1.60-1.69 (m, 9H), 1.45-1.50 (m, 2H), 1.34-1.38 (m, 2H); ¹³C NMR (100 MHz, CDCl₃): δ = 173.9, 167.0, 75.8, 68.0, 65.3, 53.7, 53.2, 50.4, 50.4, 33.2, 33.2, 33.1, 33.0, 32.8, 30.6, 27.8, 24.1, 23.8, 23.7, 23.7, 23.6, 20.2; ESI-HRMS Calcd for (C₂₃H₄₀N₄O₃+H⁺): 421.3173, Found: 421.3179.

3. Preparation of the acetal ketones

a. Prepared from arylglyoxal monohydrate (**4**, **5**)

To a suspension of arylglyoxal monohydrate (1.34 g, 10.0 mmol) and PTS (*p*-toluenesulfonic acid, 0.5 mmol) in CH_2Cl_2 (40 mL) at room temperature was added dropwise HC(OMe)_3 (or HC(OEt)_3) (10.0 mmol). The mixture was stirred for 30 min. Triethylamine (0.20 mL) was added and the mixture was concentrated. The residue was purified by silica gel column chromatography (acetone:hexane/1:100) to give an α,α -diethoxymethyl aryl ketone.

b. Preparation of the α,α -dialkoxymethyl phenyl ketones (**6a-6k**)

To a 500 mL three-neck flask with a condenser was added 1,4-dioxane (125 mL), SeO_2 (27.5 g, 0.25 mol) and water (5 mL). The mixture was heated at 50-55 °C and stirred until the solid was dissolved, then acetophenone (30 g, 0.25 mol) was added and the reaction mixture was refluxed for 4 h. The solid was removed by filtration, the filtrate was concentrated. The residue was purified by silica gel column chromatography (acetone:hexane/1:10) to give arylglyoxal monohydrate (23 g, yield 69%).

To a suspension of arylglyoxal monohydrate (1.34 g, 10.00 mmol) and PTS (1.0 mmol) in toluene (30 mL) was added benzyl alcohol (10.80 g, 100 mmol). The mixture was refluxed and dehydrated by water separator. After stirring for 6 h, the reaction was finished and the mixture was concentrated. The

residue was purified by silica gel column chromatography (acetone:hexane/1:50) to give acetal ketones **6a-6k**.

c. Prepared from α,α -dipropoxyacetone (**7b**, **6l-6p**)

To a solution of 1,3-dihydroxypropan-2-one (9.0 g, 0.1 mmol) and ⁱPrOH (31 mL, 0.4 mol) in high-pressure reactor was added a suspension of H⁺-cation resin (1.72 g, 10 mol%) in Toluene (20 mL). The mixture was heated at 110 °C for 6 h and then cooled with ice-water bath. Then, the solid was removed by filtration and the filtrate was concentrated. The residue was purified by silica gel column chromatography (acetone:hexane/1:100) to give α,α -dipropoxyacetone **7b**.

To a mixture of α,α -dipropoxyacetone **7b** (348 mg, 2.0 mmol) and aryl aldehyde (2.5 mmol) was slowly added piperidine (2.0 mmol). The mixture was heated at 100 °C until α,α -dipropoxyacetone was consumed completely and monitored with TLC. The mixture was cooled to room temperature and the solid was removed by filtration and the filtrate was concentrated. The residue was purified by silica gel column chromatography (acetone:hexane/1:100) to give an alkenyl ketone **6l-6p**.

4. General procedure for the preparation of racemic cyanosilylation of acetal ketones

To a mixture of α,α -dialkoxyl ketone **6,7** (0.20 mmol) and *N*-oxide **11** (5 mol%) in CH_2Cl_2 at room temperature was slowly added TMSCN (1.5 eq.). The mixture was stirred for 1 h and then purified by silica gel column chromatography (ether:hexane = 1:100-1:50) to give a racemic cyanohydrin trimethylsilyl ethers in quantitative yield.

5. General procedure for the *N,N'*-dioxide-catalyzed asymmetric cyanosilylation of acetal ketones

The mixture of chiral diamide **3a** (8.0 mg, 10 mmol%) and *m*-CPBA (7.0 mg, 20 mol%) in $\text{ClCH}_2\text{CH}_2\text{Cl}$ (0.4 mL) were stirred at $-20\text{ }^\circ\text{C}$ for 10 min. Then, *t*-BuOMe (0.2 mL) was added to the mixture. Subsequently, TMSCN (56 μL , 0.4 mmol) and acetal ketone **6** (0.2 mmol) added at $-45\text{ }^\circ\text{C}$. After being stirred for 10-36 h, the reaction mixture was directly purified by column chromatography on silica gel eluted (ether:hexane = 1:50-1:5) to give the chiral cyanohydrin trimethylsilyl ethers.

Products:

3,3-bis(benzyloxy)-2-phenyl-2-(trimethylsilyloxy)propanenitrile **(+)-8a**

was obtained as a colorless oil in 99% isolated yield and 92% ee [determined by converting this compound to its corresponding β -aminoalcohol derivative

8aY], $[\alpha]_{\text{D}}^{25} = +10.2$ ($c = 0.92$, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.56\text{--}7.60$ (m, 2H), 7.35-7.40 (m, 8H), 7.19-7.22 (m, 3H), 6.87-6.90 m, 2H), 4.76 (d, $J = 15.7$ Hz, 2H), 4.58 (s, 1H), 4.52 (d, $J = 12.3$ Hz, 1H), 4.33 (d, $J = 12.3$ Hz, 1H), 0.16 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 137.5$, 137.3, 136.7, 128.9, 128.4, 128.2, 128.1, 127.9, 127.8, 127.7, 127.7, 126.7, 119.1, 104.0, 78.7, 71.8, 70.7, 0.9; ESI-HRMS Calcd for $(\text{C}_{26}\text{H}_{29}\text{NO}_3\text{Si}+\text{Na}^+)$ 454.1809, Found: 454.1795.

3,3-bis(benzyloxy)-2-(4-chlorophenyl)-2-(trimethylsilyloxy) propane-

nitrile (+)-8b was obtained as a colorless oil in 83% isolated yield and

88% ee [determined with chiral HPLC, Daicel Chiralpak AD-H,

isopropanol:hexane/1:99, 1.0 mL/min, $\lambda = 220$ nm, tr (minor) = 6.4 min, tr (major) = 9.6 min]; $[\alpha]_D^{25} = +3.3$ (c = 0.66, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.47$ -7.50 (m, 2H), 7.32-7.37 (m, 8H), 7.22-7.26 (m, 2H), 6.92-6.93 (m, 2H), 4.74 (d, $J = 2.0$ Hz, 2H), 4.55 (s, 1H), 4.56 (d, $J = 12.2$ Hz, 1H), 4.36 (d, $J = 12.0$ Hz, 1H), 0.16 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 137.1$, 136.5, 136.0, 135.0, 128.7, 128.4, 128.3, 128.2, 127.9, 127.8, 118.8, 103.7, 78.1, 71.8, 71.1, 0.9; ESI-HRMS Calcd for $(\text{C}_{26}\text{H}_{28}\text{NO}_3\text{Si}+\text{NH}_4^+)$ 483.1865, Found: 483.1869.

3,3-bis(benzyloxy)-2-(3-chlorophenyl)-2-(trimethylsilyloxy)propane

nitrile (+)-8c was obtained as a colorless oil in 99% isolated yield and 93%

ee [determined by converting this compound to its corresponding β -

aminoalcohol derivative **8cY**]; $[\alpha]_D^{25} = +10.1$ (c = 0.84, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.47$ -7.53 (m, 2H), 7.31-7.38 (m, 7H), 7.23-7.26 (m, 3H), 6.91-6.93 (m, 2H), 4.71-4.80 (m, 2H), 4.57 (d, $J = 12.5$ Hz, 1H), 4.55 (s, 1H), 4.37 (d, $J = 12.2$ Hz, 1H), 0.18 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 139.5$, 137.1, 136.4, 134.2, 129.4, 129.1, 128.3, 127.9, 127.8, 127.8, 126.9, 125.1, 118.6, 103.7, 78.1, 71.9, 71.0, 0.9; ESI-HRMS Calcd for $(\text{C}_{26}\text{H}_{28}\text{NO}_3\text{Si}+\text{NH}_4^+)$ 483.1865, Found: 483.1856.

3,3-bis(benzyloxy)-2-(4-fluorophenyl)-2-(trimethylsilyloxy)propane-

nitrile(+)-8d was obtained as a colorless oil in 99% isolated yield and 89%

ee [determined with chiral HPLC, Daicel Chiralpak ADH,

isopropanol:hexane/1:99, 1.0 mL/min, $\lambda = 210$ nm, tr (minor) = 6.6 min, tr (major) = 9.2 min]; $[\alpha]_D^{25} = +12.2$ (c = 0.61, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.53$ -7.58 (m, 2H), 7.36-7.38 (m, 5H),

7.23-7.25 (m, 3H), 6.95-7.10 (m, 2H), 6.94-6.95 (m, 2H), 4.76 (m, 2H), 4.58 (s, 1H), 4.58 (d, $J = 12.2$ Hz, 1H), 4.39 (d, $J = 12.2$ Hz, 1H), 0.17 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 164.6, 161.4, 137.1, 136.5, 133.2, 128.7, 128.6, 128.4, 128.2, 127.9, 127.8, 119.0, 115.2, 114.9, 103.8, 78.1, 71.8, 71.0, 0.9$; ESI-HRMS Calcd for $(\text{C}_{26}\text{H}_{28}\text{FNO}_3\text{Si} + \text{Na}^+)$: 472.1715, Found: 472.1720.

3,3-bis(benzyloxy)-2-(2-fluorophenyl)-2-(trimethylsilyloxy)propane-

nitrile(+)-8e was obtained as a colorless oil in 92% isolated yield and 87% ee [determined with chiral HPLC, Daicel Chiralcel ODH,

isopropanol:hexane/1:99, 0.5 mL/min, $\lambda = 210$ nm, tr (minor) = 12.2 min, tr (major) = 12.8 min]; $[\alpha]_D^{25} = +25.8$ ($c = 0.42$, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.72$ -7.73 (m, 1H), 7.04-7.39 (m, 10H), 7.00-7.04 (m, 1H), 6.86-6.88 (m, 2H), 4.99 (d, $J = 1.0$ Hz, 1H), 4.79 (dd, $J = 22.1, 12.1$ Hz, 2H), 4.60 (d, $J = 12.2$ Hz, 1H), 4.38 (d, $J = 12.2$ Hz, 1H), 0.13 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 161.5, 158.2, 137.4, 136.6, 131.0$ (d, $J = 8.5\text{Hz}$), 129.4, 128.4, 128.2, 128.0, 127.7, 127.7, 124.1 (d, $J = 3.1$ Hz), 118.0, 116.5 (d, $J = 22.1$ Hz), 101.8 (d, $J = 3.9$ Hz), 77.8, 72.2, 70.3, 0.8; ESI-HRMS Calcd for $(\text{C}_{26}\text{H}_{28}\text{FNO}_3\text{Si} + \text{Na}^+)$ 472.1715, Found: 472.2135.

3,3-bis(benzyloxy)-2-(3-nitrophenyl)-2-(trimethylsilyloxy)propane-

nitrile (+)-8f was obtained as a colorless oil in 99% isolated yield and 85% ee [determined with chiral HPLC, Daicel Chiralpak ADH,

isopropanol:hexane/1:99, 1.0 mL/min, $\lambda = 220$ nm, tr (minor) = 14.0 min, tr (major) = 16.7 min]; $[\alpha]_D^{25}$

= +23.7 (c = 0.22, CH₂Cl₂); ¹H NMR (300 MHz, CDCl₃): δ = 8.37-8.38 (m, 1H), 8.21-8.25 (m, 1H), 7.92-7.96 (m, 1H), 7.52-7.58 (m, 1H), 7.21-7.38 (m, 8H), 6.93-6.97 (m, 2H), 4.76 (s, 2H), 4.65 (s, 1H), 4.63 (d, *J* = 11.7 Hz, 1H), 4.45(d, *J* = 12.0 Hz, 1H), 0.21 (s, 9H); ¹³C NMR (75 MHz, CDCl₃): δ = 148.0, 139.4, 136.7, 136.2, 133.1, 129.0, 128.8, 128.5, 128.4, 128.1, 127.9, 127.9, 121.9, 118.4, 103.3, 71.9, 71.6, 15.2, 0.9; ESI-HRMS Calcd for (C₂₆H₂₈N₂O₅Si+NH₄⁺): 494.2106, Found: 494.2098.

3,3-bis(benzyloxy)-2-*p*-tolyl-2-(trimethylsilyloxy)propanenitrile(+)-8g

was obtained as a colorless oil in 73% isolated yield and 90% ee [determined with chiral HPLC, Daicel Chiralpak AD-H, isopropanol:hexane/1:99, 1.0 mL/min, λ = 210 nm, tr (minor) = 7.8 min, tr (major) = 11.3 min]; [α]_D²⁵ = +3.86 (c = 0.44, CH₂Cl₂); ¹H NMR (300 MHz, CDCl₃): δ = 7.44-7.47 (m, 2H), 7.34-7.37 (m, 5H), 7.17-7.22 (m, 5H), 6.90-6.93 (m, 2H), 4.73-4.74 (m, 2H), 4.57 (s, 1H), 4.52 (d, *J* = 12.3 Hz, 1H), 4.34 (d, *J* = 12.3 Hz, 1H), 2.39 (s, 3H), 0.15 (s, 9H); ¹³C NMR (75 MHz, CDCl₃): δ = 138.9, 137.4, 136.8, 134.5, 128.8, 128.4, 128.2, 127.9, 127.8, 127.6, 126.6, 119.2, 104.0, 78.6, 71.6, 70.8, 21.2, 1.0; ESI-HRMS Calcd for (C₂₇H₃₁NO₃Si+NH₄⁺) 463.2411, Found: 463.2415.

3,3-bis(benzyloxy)-2-(4-methoxyphenyl)-2-(trimethylsilyloxy)propanenitrile(+)-8h

was obtained as a colorless oil in 99% isolated yield and 90% ee [determined with chiral HPLC, Daicel Chiralpak ADH, isopropanol:hexane/1:99, 1.0 mL/min, λ = 220 nm, tr (minor) = 11.9min, tr (major) = 20.0 min]; [α]_D²⁵

= +0.90 (c = 0.97, CH₂Cl₂); ¹H NMR (300 MHz, CDCl₃): δ = 7.47-7.50 (m, 2H), 7.34-7.38 (m, 5H), 7.21-7.26 (m, 3H), 6.88-6.95 (m, 4H), 4.74 (m, 2H), 4.56 (s, 1H), 4.53 (d, *J* = 12.5 Hz, 1H), 4.35 (d, *J* = 12.3 Hz, 1H), 3.84 (s, 3H), 0.15 (s, 9H); ¹³C NMR (75 MHz, CDCl₃): δ = 160.0, 137.4, 136.8, 129.5, 128.4, 128.2, 128.0, 127.9, 127.8, 127.7, 119.3, 113.4, 104.0, 78.3, 71.6, 70.9, 55.3, 1.0; ESI-HRMS Calcd for (C₂₇H₃₁NO₄Si+NH₄⁺) 479.2361, Found: 479.2357.

3,3-bis(benzyloxy)-2-(3,4-dimethoxyphenyl)-2-(trimethylsilyloxy)

propanenitrile (-)-8i was obtained as a colorless oil in 94% isolated yield and 88% ee [determined with chiral HPLC, Daicel Chiralpak

ADH, isopropanol:hexane/10:90, 1.0 mL/min, λ = 220 nm, tr (minor) = 4.8 min, tr (major) = 6.4 min]; [α]_D²⁵ = -34.2 (c = 0.32, CH₂Cl₂); ¹H NMR (300 MHz, CDCl₃): δ = 7.36-7.38 (m, 5H), 7.15-7.24 (m, 4H), 6.94-6.99 (m, 3H), 6.85 (d, *J* = 8.5 Hz, 1H), 4.77 (m, 2H), 4.58 (s, 1H), 4.57 (d, *J* = 11.9 Hz, 1H), 4.35 (d, *J* = 12.1 Hz, 1H), 3.80 (s, 3H), 3.77 (s, 3H), 0.16 (s, 9H); ¹³C NMR (75 MHz, CDCl₃): δ = 149.4, 148.4, 137.3, 136.7, 129.7, 128.3, 128.1, 127.8, 127.7, 119.2, 110.3, 109.8, 104.0, 78.3, 71.6, 71.0, 55.8, 55.7, 0.9; ESI-HRMS Calcd for (C₂₈H₃₃NO₅Si+NH₄⁺): 509.2466, Found: 509.2467.

3,3-bis(benzyloxy)-2-(naphthalen-2-yl)-2-(trimethylsilyloxy)propane-

nitrile(-)-8j was obtained as a light yellow powder in 85% isolated yield and 93% ee [determined by converting this compound to its corresponding

β-aminoalcohol derivative **8jY**]; [α]_D²⁵ = -6.90 (c = 0.32, CH₂Cl₂); ¹H NMR (300 MHz, CDCl₃): δ =

8.09 (d, $J = 1.5$ Hz, 1H), 7.84-7.90 (m, 3H), 7.62-7.66 (m, 1H), 7.54-7.58 (m, 2H), 7.04-7.39 (m, 8H), 6.79-6.82 (m, 2H), 4.80 (m, 2H), 4.68 (s, 1H), 4.53 (d, $J = 12.5$ Hz, 1H), 4.33 (d, $J = 12.2$ Hz, 1H), 0.18 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 137.3, 136.5, 134.9, 133.4, 132.7, 128.5, 128.4, 128.1, 127.9, 127.8, 127.7, 127.6, 126.7, 126.4, 126.4, 123.9, 119.1, 104.0, 78.9, 71.8, 70.8, 1.0$; ESI-HRMS Calcd for $(\text{C}_{30}\text{H}_{31}\text{NO}_3\text{Si}+\text{NH}_4^+)$: 499.2411, Found: 499.2407.

3,3-bis(benzyloxy)-2-(furan-2-yl)-2-(trimethylsilyloxy)propanenitrile (+)-8k was obtained as a white powder in 99% isolated yield and 90% ee (determined by converting this compound to its corresponding β -aminoalcohol

derivative **8kY**); $[\alpha]_{\text{D}}^{25} = +18.2$ ($c = 1.08, \text{CH}_2\text{Cl}_2$); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.33$ -7.39 (m, 6H), 7.23-7.27 (m, 3H), 6.97-7.00 (m, 2H), 6.68 (d, $J = 3.3$ Hz, 1H), 6.42-6.49 (m, 1H), 4.89(s, 1H), 4.86 (d, $J = 12.0$ Hz, 1H), 4.78 (d, $J = 12.0$ Hz, 1H), 4.60(d, $J = 12.3$ Hz, 1H), 4.41(d, $J = 12.3$ Hz, 1H), 0.09 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 149.1, 143.0, 137.4, 136.7, 128.4, 128.2, 127.9, 127.8, 127.8, 117.6, 110.8, 110.6, 101.9, 73.5, 72.0, 70.1, 29.7, 0.3$; ESI-HRMS Calcd for $(\text{C}_{24}\text{H}_{27}\text{NO}_4\text{Si}+\text{NH}_4^+)$: 439.2048, Found: 439.2055.

(E)-2-(diisopropoxymethyl)-4-phenyl-2-(trimethylsilyloxy)but-3-

enenitrile (-)-8l was obtained as a white powder in 92% isolated yield and 92% ee [determined with chiral HPLC, Daicel Chiralcel OF,

isopropanol:hexane/1:99, 1.0 mL/min, $\lambda = 210$ nm, tr (major) = 4.6 min, tr (minor) = 8.8 min], $[\alpha]_{\text{D}}^{25} = -38.2$ ($c = 0.35, \text{CH}_2\text{Cl}_2$); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.42$ -7.44 (m, 2H), 7.26-7.33 (m, 3H), 6.92 (d, $J = 16.2$ Hz, 1H), 6.23 (d, $J = 16.1$ Hz, 1H), 4.60 (s, 1H), 3.92-4.02 (m, 2H), 1.18-1.26 (m, 12H), 0.25 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 137.5, 137.3, 136.7, 128.9, 128.4, 128.2, 128.1, 127.9, 127.8, 127.7, 127.7, 126.7, 119.1, 104.0, 78.7, 71.8, 70.7, 0.9$; ESI-HRMS Calcd for $(\text{C}_{20}\text{H}_{31}\text{NO}_3\text{Si}+\text{H}^+)$ 362.2146, Found: 362.2136.

(E)-4-(4-chlorophenyl)-2-(diisopropoxymethyl)-2-

(trimethylsilyloxy) but-3-enenitrile (-)-8m was obtained as a

white powder in 88% isolated yield and 87% ee [determined with

chiral HPLC, Daicel Chiralcel OF, isopropanol:hexane/1:99, 1.0 mL/min, $\lambda = 210$ nm, tr (major) = 4.7 min, tr (minor) = 8.6 min], $[\alpha]_D^{25} = -94.2$ ($c = 0.36$, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.26$ -7.37 (m, 4H), 6.86 (d, $J = 16.1$ Hz, 1H), 6.21 (d, $J = 16.1$ Hz, 1H), 4.60 (s, 1H), 3.91-4.01 (m, 2H), 1.17-1.25 (m, 12H), 0.23 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 134.2$, 134.0, 132.5, 128.8, 128.2, 125.3, 118.9, 101.9, 71.8, 70.7, 23.0, 22.9, 22.5, 21.8, 1.33; ESI-HRMS Calcd for $(\text{C}_{20}\text{H}_{30}\text{ClNO}_3\text{Si}+\text{Na}^+)$ 418.1576, Found: 418.1582.

(E)-2-(diisopropoxymethyl)-4-(4-nitrophenyl)-2-(trimethylsilyloxy)

but-3-enenitrile(-)-8n was obtained as a white powder in 99%

isolated yield and 93% ee [determined with chiral HPLC, Daicel

Chiralcel OF, isopropanol:hexane/5:95, 1.0 mL/min, $\lambda = 300$ nm, tr (major) = 11.7 min, tr (minor) = 14.8 min], $[\alpha]_D^{25} = -64.7$ ($c = 0.46$, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 8.21$ (d, $J = 8.8$ Hz, 2H), 7.56 (d, $J = 8.8$ Hz, 2H), 6.95 (d, $J = 16.2$ Hz, 1H), 6.41 (d, $J = 16.4$ Hz, 1H), 4.64 (s, 1H), 3.92-4.01 (m, 2H), 1.18-1.32 (m, 12H), 0.25 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 147.5$, 141.8, 131.5, 129.4, 127.6, 124.0, 118.6, 101.9, 76.8, 72.1, 70.9, 23.0, 22.9, 22.5, 21.7, 1.3; ESI-HRMS Calcd for $(\text{C}_{20}\text{H}_{30}\text{N}_2\text{O}_5\text{Si}+\text{Na}^+)$: 429.1816, Found: 429.1822.

(E)-2-(diisopropoxymethyl)-4-(4-methoxyphenyl)-2-

(trimethylsilyl-oxy)but-3-enenitrile (-)-8o was obtained as a white

powder in 85% isolated yield and 90% ee [determined with chiral

HPLC, Daicel Chiralcel OF, isopropanol:hexane/5:95, 1.0 mL/min, $\lambda = 254$ nm, tr (major) = 4.5 min, tr (minor) = 8.3 min], $[\alpha]_D^{25} = -44.7$ (c = 0.55, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 7.35$ -7.38 (m, 2H), 6.83-6.90 (m, 3H), 6.08 (d, $J = 16.1$ Hz, 1H), 4.59 (s, 1H), 3.90-4.01 (m, 2H), 3.77 (s, 3H), 1.17-1.27 (m, 12H), 0.23 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 158.8, 133.3, 128.3, 122.2, 119.1, 114.0, 102.0, 77.4, 71.6, 70.7, 55.3, 23.0, 22.9, 22.4, 21.8, 1.4$; ESI-HRMS Calcd for $(\text{C}_{21}\text{H}_{33}\text{NO}_4\text{Si}+\text{Na}^+)$: 414.2071, Found: 414.2086.

(E)-2-(diisopropoxymethyl)-4-(pyridin-3-yl)-2-(trimethylsilyloxy)but-3-

enenitrile (-)-8p was obtained as a light yellow powder in 95% isolated

yield and 85% ee [determined with chiral HPLC, Daicel Chiralcel OF,

isopropanol:hexane/8:92, 1.0 mL/min, $\lambda = 254$ nm, tr (major) = 12.1 min, tr (minor) = 18.0 min], $[\alpha]_D^{25} = -20.4$ (c = 0.76, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 8.64$ (d, $J = 2.0$ Hz, 1H), 8.50-8.52 (m, 1H), 7.72-7.76 (m, 1H), 7.26-7.30 (m, 1H), 6.88 (d, $J = 16.2$ Hz, 1H), 6.33 (d, $J = 16.2$ Hz, 1H), 4.61 (s, 1H), 3.92-3.99 (m, 2H), 1.17-1.25 (m, 12H), 0.24 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 149.4, 148.7, 133.5, 131.2, 130.2, 127.0, 123.5, 118.7, 101.9, 77.4, 71.8, 70.8, 23.0, 22.9, 22.5, 21.7, 1.3$; ESI-HRMS Calcd for $(\text{C}_{19}\text{H}_{30}\text{N}_2\text{O}_3\text{Si}+\text{H}^+)$: 363.2098, Found: 363.2113.

3,3-diisopropoxy-2-methyl-2-(trimethylsilyloxy)propanenitrile(+)-8q

was

obtained as a colorless oil in 90% isolated yield and 92% ee [determined by converting this compound to its corresponding β -aminoalcohol derivative **7bY**],

$[\alpha]_D^{25} = +12.5$ ($c = 0.83$, CH_2Cl_2); ^1H NMR (300 MHz, CDCl_3): $\delta = 4.47$ (s, 1H), 3.89-3.96 (m, 2H), 1.53 (s, 3H), 1.22 (d, $J = 6.1$ Hz, 9H), 1.17(d, $J = 6.2$ Hz, 3H), 0.25 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3): $\delta = 121.0, 101.1, 72.9, 71.4, 70.4, 23.1, 22.9, 22.4, 22.1, 21.7, 1.28$; ESI-HRMS Calcd for $(\text{C}_{13}\text{H}_{27}\text{NO}_3\text{Si}+\text{Na}^+)$: 296.1652, Found: 296.1663.

6. General procedure for the conversion of cyanohydrin trimethylsilyl ether to its corresponding derivative

To a solution of cyanohydrin trimethylsilyl ether (0.20 mmol) in dry ether (30.0 mL) at room temperature was added LiAlH_4 (0.60 mmol). The mixture was stirred for 1.0 h, and cooled with ice-water bath. Water (0.40 mL) was slowly added and the resulting mixture was filtered. The filtrate was dried over anhydrous K_2CO_3 , and then filtered. To this filtrate was added triethyl amine (1.2 mmol) and benzoyl chloride (0.24 mmol). After being stirred at room temperature for 0.5 h, the mixture was filtered and concentrated. The residue was purified by silica gel column chromatography (acetone: hexane/1:20 to 1:5) to give the β -aminoalcohol derivative.

N-(3,3-bis(benzyloxy)-2-hydroxy-2-phenylpropyl)benzamide **8aY**

was

obtained as a white powder in 90% isolated yield; mp: 129-130 $^\circ\text{C}$; ^1H NMR (300 MHz, CDCl_3) $\delta = 7.60$ -7.63 (m, 2H), 7.46-7.50 (m, 2H), 7.24-7.44 (m,

12H), 7.17-7.23 (m, 4H), 6.38 (br, 1H), 4.77 (s, 1H), 4.65 (d, $J = 11.4, 3.6$ Hz, 2H), 4.46 (dd, $J = 28.6, 11.6$ Hz, 2H), 4.20 (dd, $J = 14.1, 4.1$ Hz, 1H), 4.02 (dd, $J = 14.1, 4.1$ Hz, 1H), 3.58 (br, 1H); ^{13}C NMR (75 MHz, CDCl_3) $\delta = 168.4, 140.6, 137.2, 137.1, 134.3, 131.3, 130.1, 128.6, 128.5, 128.4, 128.1, 127.9, 127.6, 126.8, 126.4, 106.1, 78.1, 72.3, 71.7, 45.4$; ESI-HRMS Calcd for $(\text{C}_{30}\text{H}_{29}\text{NO}_4 + \text{H}^+)$: 466.2019, Found: 466.2105. HPLC conditions: Daicel Chiralcel OJ-H, isopropanol:hexane/15:85, 1.0 mL/min, $\lambda = 254$ nm, tr (minor) = 19.0 min, tr (major) = 20.9 min.

***N*-(3,3-bis(benzyloxy)-2-(3-chlorophenyl)-2-hydroxypropyl)benzamide 8cY** was obtained as a white powder in 82% isolated yield; ^1H NMR (300 MHz, CDCl_3) $\delta = 7.66$ (s, 1H), 7.49-7.52 (m, 4H), 7.20-7.34 (m, 14H), 6.52 (br, 1H), 4.68-4.74 (m, 2H), 4.71 (s, 1H), 4.57 (d, $J = 11.2$ Hz, 1H), 4.45 (d, $J = 11.3$ Hz, 1H), 4.12 (dd, $J = 14.2, 7.2$ Hz, 1H), 3.97 (dd, $J = 14.2, 4.4$ Hz, 1H), 3.89 (br, 1H); ^{13}C NMR (75 MHz, CDCl_3) $\delta = 168.7, 143.0, 137.1, 136.9, 134.1, 134.0, 131.5, 129.3, 128.6, 128.5, 128.4, 128.1, 128.0, 127.7, 127.0, 126.9, 124.7, 105.8, 78.2, 72.4, 71.9, 45.5$; ESI-HRMS Calcd for $(\text{C}_{30}\text{H}_{28}\text{ClNO}_4 + \text{Na}^+)$: 524.1599, Found: 524.1590. HPLC conditions: Daicel Chiralcel OJ-H, isopropanol:hexane/8:92, 1.0 mL/min, $\lambda = 254$ nm, tr (minor) = 14.0 min, tr (major) = 16.0 min.

***N*-(3,3-bis(benzyloxy)-2-(naphthalen-2-yl)propyl)benzamide 8jY** was obtained as a white powder in 82% isolated yield; ^1H NMR (300 MHz, CDCl_3) $\delta = 8.13$ (s, 1H), 7.82-7.85 (m,

3H), 7.70-7.74 (m, 1H), 7.47-7.50 (m, 4H), 7.37-7.42 (m, 1H), 7.15-7.30 (m, 11H), 6.58 (br, 1H), 4.80 (d, $J = 11.3$ Hz, 1H), 4.40(d, $J = 11.3$ Hz, 1H), 4.70 (d, $J = 11.4$ Hz, 1H), 4.40 (d, $J = 11.3$ Hz, 1H), 4.66 (d, $J = 11.4$ Hz, 1H), 4.40 (d, $J = 11.3$ Hz, 1H), 4.29 (dd, $J = 14.2, 7.1$ Hz, 1H), 4.14 (dd, $J = 14.2, 4.3$ Hz, 1H), 3.80 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) $\delta = 168.5, 138.1, 137.2, 137.1, 133.1, 132.8, 131.3, 128.5, 128.4, 128.4, 128.1, 127.9, 127.5, 126.8, 126.1, 126.0, 125.8, 124.4, 106.2, 78.3, 72.3, 71.9, 45.4$; ESI-HRMS Calcd for $(\text{C}_{34}\text{H}_{31}\text{NO}_4 + \text{Na}^+)$: 540.2145, Found: 540.2140; HPLC conditions: Daicel Chiralcel OD-H, isopropanol:hexane/10:90, 1.0 mL/min, $\lambda = 254$ nm, tr (major) = 24.5 min, tr (minor) = 29.5 min.

***N*-(3,3-bis(benzyloxy)-2-(furan-2-yl)-2-hydroxypropyl)benzamide 8kY** was

obtained as a white powder in 85% isolated yield; ^1H NMR (300 MHz, CDCl_3) $\delta = 7.53$ -7.55 (m, 2H), 7.41-7.42 (m, 2H), 7.26-7.35 (m, 10H), 7.18-7.20 (m, 2H), 6.73 (br, 1H), 6.48 (d, $J = 3.3$ Hz, 1H), 6.4-6.38 (m, 1H), 4.98 (s, 1H), 4.58-4.74 (m, 3H), 4.45 (d, $J = 11.6$ Hz, 1H), 4.18 (dd, $J = 14.0, 6.8$ Hz, 1H), 4.02 (dd, $J = 14.1, 4.6$ Hz, 1H), 3.66 (br, 1H); ^{13}C NMR (75 MHz, CDCl_3) $\delta = 168.3, 153.8, 142.2, 137.2, 137.1, 134.1, 131.3, 128.6, 128.4, 128.1, 128.0, 127.9, 126.9, 110.6, 108.1, 104.5, 77.4, 72.2, 71.5, 43.6$; ESI-HRMS Calcd for $(\text{C}_{28}\text{H}_{27}\text{NO}_5 + \text{Na}^+)$: 480.1781, Found: 480.1790; HPLC conditions: Daicel Chiralcel OD-H, isopropanol:hexane/3:97, 1.0 mL/min, $\lambda = 254$ nm, tr (major) = 40.2 min, tr (minor) = 49.0 min.

***N*-(2-hydroxy-3,3-diisopropoxy-2-methylpropyl)benzamide 8qY** was obtained as a white powder in 83% isolated yield; ^1H NMR (300 MHz, CDCl_3) δ = 7.76-7.80 (m, 2H), 7.38-7.48 (m, 3H), 6.97 (s, 1H), 4.47 (s, 1H), 3.89-3.95 (m, 2H), 3.63 (dd, J = 13.7, 6.4 Hz, 1H), 3.53 (dd, J = 13.6, 4.9 Hz, 1H), 2.70 (br, 1H), 1.23 (s, 3H), 1.17-1.20 (m, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ = 167.7, 134.7, 131.3, 128.4, 126.9, 130.6, 73.4, 71.2, 70.9, 45.6, 23.5, 23.4, 22.3, 22.2, 21.3 ; ESI-HRMS Calcd. for $(\text{C}_{17}\text{H}_{27}\text{NO}_4 + \text{Na}^+)$ 332.1832, Found: 332.1837; HPLC conditions: Daicel Chiralcel AD, isopropanol:hexane/10:90, 1.0 mL/min, λ = 254 nm, tr (minor) = 6.4 min, tr (major) = 7.3 min.

7. Character Data of Cyanohydrin 10

3,3-bis(benzyloxy)-2-hydroxy-2-phenylpropanenitrile 10 was obtained as a colorless oil by silica gel column chromatography (ether:hexane/1:6); ^1H NMR (300 MHz, CDCl_3) δ = 7.63-7.64 (m, 2H), 7.41-7.44 (m, 3H), 7.25-7.37 (m, 8H), 7.11-7.13 (m, 2H), 4.71 (d, J = 11.8Hz, 1H), 4.66 (s, 1H), 4.49 (d, J = 11.8Hz, 1H), 4.45 (d, J = 11.6Hz, 1H), 4.35 (d, J = 11.9Hz, 1H), 3.77 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ = 136.4, 136.1, 135.1, 129.4, 128.6, 128.5, 128.4, 128.3, 128.1, 127.9, 127.9, 126.4, 119.1, 103.7, 763.4, 72.0, 71.2; ESI-HRMS Calcd for $(\text{C}_{23}\text{H}_{21}\text{NO}_3 + \text{Na}^+)$ 382.1414, Found: 382.1430.

8. Copy of ^1H NMR and ^{13}C NMR spectra for the catalytic mechanism

- Copy of ^1H NMR spectra of *N,N'*-dioxide **2b**: δ_{NH} = 10.68 ppm (d, J = 2.9 Hz)
- Copy of ^1H NMR spectra of *N,N'*-dioxide **2b** and TMSCN (ratio: 1/1): δ_{NH} = 4.5 ppm (br);
- Copy of ^1H NMR spectra of the mixture of *N,N'*-dioxide **2b**, TMSCN and acetal ketone **6a** (ratio: 1/1/1): δ_{NH} = 7.89 ppm (br) and 8.23 ppm (br).
- Copy of ^1H NMR spectra of acetal ketone **6a**;
- Copy of ^1H NMR spectra of product **8a**;

f. Copy of ^1H NMR and ^{13}C NMR spectra of the cyanohydrin **10**.

a.Copy of ^1H NMR spectra of N,N' -dioxide 2b

b. Copy of ^1H NMR spectra of *N,N'*-dioxide 2b and TMS-CN (ratio: 1/1)

c. Copy of ^1H NMR spectra of the mixture of N,N' -dioxide 2b, TMSCN and acetal ketone 6a (ratio: 1/1/1)

d. Copy of ^1H NMR spectra of acetal ketone 6a

e. Copy of ^1H NMR spectra of product 8a

Current Data Parameters
NAME fkm-2006-286
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20061026
Time 11:52
INSTRUM av300
PROBHD 5 mm QNP 1H/13
PULPROG zg30
TD 32768
SOLVENT CDCl3
NS 16
DS 0
SWH 5995.204 Hz
FIDRES 0.182959 Hz
AQ 2.7329011 sec
RG 256
DM 83.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 3.00 usec
PL1 -2.00 dB
SFO1 300.1320882 MHz

F2 - Processing parameters
SI 32768
SF 300.1300124 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 9.00 cm
F1P 10.000 ppm
F1 3001.30 Hz
F2P -0.500 ppm
F2 -150.06 Hz
PPMCM 0.52500 ppm/cm
HZCM 157.56825 Hz/cm

QB-PhOH

136.382
136.069
135.111
129.419
128.633
128.523
128.445
128.267
128.117
127.941
127.907
126.365
119.056
103.674

77.422
77.000
76.575
76.346
72.026
71.150

15.174

Current Data Parameters
NAME fxm-2006-286
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20061026
Time 11.56
INSTRUM av300
PROBHD 5 mm QNP 1H/13
PULPROG zgpg
TO 65536
SOLVENT CDCl3
NS 205
DS 4
SWH 22675.736 Hz
FIDRES 0.346004 Hz
AQ 1.445186 sec
RG 8192
DM 22.050 usec
DE 6.00 usec
TE 300.0 K
D1 2.00000000 sec
d11 0.03000000 sec
d12 0.0002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 5.50 usec
PL1 -6.00 dB
SF01 75.4775998 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 80.00 usec
PL2 -2.00 dB
PL12 17.70 dB
PL13 17.71 dB
SF02 300.1312005 MHz

F2 - Processing parameters
SI 65536
SF 75.4677563 MHz
MDM EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

10 NMR plot parameters
CX 20.00 cm
CY 4.00 cm
F1P 200.500 ppm
F1 15131.29 Hz
F2P -0.500 ppm
F2 -37.73 Hz
PPMCM 10.05000 ppm/cm
HZCM 758.45099 Hz/cm