Supplementary Material

Enantioselective Epoxidation of α,β -Enones Promoted by α,α -Diphenyl-L-Prolinol as Bifunctional Organocatalyst

Alessandra Lattanzi

*Dipartimento di Chimica, Università di Salerno, Via S. Allende, I-84081 Baronissi, Salerno, Italy
lattanzi@unisa.it*

General Information

All the reactions were carried out under air using freshly distilled solvents. Reactions were monitored by thin layer chromatography (TLC) on Merck silica gel plates (0.25 mm) and visualized by UV light or by 10% H₂SO₄/ethanol spray test. Flash chromatography was performed on Merck silica gel (60, particle size: 0.040-0.063 mm). ¹H NMR and ¹³C NMR spectra were recorded in CDCl₃ solutions on a Bruker DRX 400 spectrometer (400 MHz) at room temperature. The following abbreviations were used to explain the multiplicities: s=singlet, d=doublet, dq=doublet of quartets, t=triplet, dt= doublet of triplets, q=quartet, m=multiplet. Optical rotations were performed on a Jasco Dip-1000 using the Na lamp. Absolute configurations were determined by optical rotations or by chiral HPLC (Waters 486, UV detector) using Daicel Chiralcel OD and Daicel Chiralpak AD columns. The diastereoisomeric ratio of the epoxides (*trans/cis* > 98/<2) was determined by ¹H-NMR analysis of the crude reaction mixtures. The enantioselectivities were determined by chiral HPLC analysis. Catalyst **3** was prepared according to the literature procedure.¹ Catalysts **1**, **2**, N-benzyl L-prolinol, TBHP, CHP were commercially obtained from Aldrich and used without further purification. α,β -Enones which were not commercially available were prepared via aldol condensation using standard conditions.²

General Procedure for the Enantioselective Epoxidation of α,β -Enones. To a solution of catalyst **2** (11.5 mg, 0.045 mmol) and *trans*-chalcone (**4a**) (31 mg, 0.150 mmol) in distilled hexane (0.8 mL) (hexane of HPLC grade furnished comparable results) was added TBHP (5-6 M decane solution, 33 μ L, 0.18 mmol) at room temperature (20°C) and stirring was maintained for the indicated time. The crude reaction mixture was then purified by flash chromatography on silica gel (petroleum ether/ diethyl ether 99/1) to provide the epoxy ketone **5a**.

Epoxides in Table 2 are known compounds, their analytical data are identical with those reported in the literature.³ The configuration of the predominant enantiomer was determined by comparison with the HPLC retention times or optical rotations reported in the literature.^{3,4-6}

Data for epoxides in Table 2:

***trans*-(2*R*,3*S*)-Epoxy-1,3-diphenylpropan-1-one^{3d}:** ¹H NMR (400 MHz) δ 4.08 (d, *J* 1.9 Hz, 1H), 4.29 (d, *J* 1.9 Hz, 1H), 7.35-7.63 (m, 8H), 8.00-8.03 (m, 2H).

The absolute configuration was determined by chiral HPLC using Chiralcel OD column and compared with the literature data.^{3d}

***trans*-(2*R*,3*S*)-Epoxy-3-phenyl-1-(4-bromophenyl)-propan-1-one^{3a}:** ¹H NMR (400 MHz) δ 4.05 (d, *J* 1.9 Hz, 1H), 4.22 (d, *J* 1.9 Hz, 1H), 7.33-7.41 (m, 5H), 7.60-7.65 (m, 2H), 7.86-7.90 (m, 2H).

The absolute configuration was determined by comparison with the optical rotation reported in the literature $[\alpha]_D^{15} = -162.1$ (*c* 0.46, CHCl₃) [lit.^{3a} for (2*S*,3*R*)-epoxy-3-phenyl-1-(4-bromophenyl)-propan-1-one: $[\alpha]_D = +194$ (*c* 1.5, CHCl₃) for 88% ee].

***trans*-(2*R*,3*S*)-Epoxy-3-phenyl-1-(3-methylphenyl)-propan-1-one^{3b}:** ¹H NMR (400 MHz) δ 2.41 (s, 3H), 4.07 (d, *J* 1.9 Hz, 1H), 4.29 (d, *J* 1.9 Hz, 1H), 7.34-7.82 (m, 9H).

The absolute configuration was determined by chiral HPLC using Chiralcel OD column and compared with the literature data.^{3b}

***trans*-(2*R*,3*S*)-Epoxy-3-phenyl-1-(2-naphthyl)-propan-1-one^{3c}:** ¹H NMR (400 MHz) δ 4.15 (d, *J* 2.0 Hz, 1H), 4.44 (d, *J* 2.0 Hz, 1H), 7.33-7.65 (m, 6H), 7.81-7.96 (m, 4H), 8.01-8.12 (m, 1H), 8.56 (s, 1H).

The absolute configuration was determined by comparison with the optical rotation reported in the literature $[\alpha]_D^{23} = -90.1$ (*c* 1.0, CH₂Cl₂) [lit.^{3c} for (2*R*,3*S*)-epoxy-3-phenyl-1-(2-naphthyl)-propan-1-one: $[\alpha]_D^{25} = -100$ (*c* 1, CH₂Cl₂) for 73% ee].

***trans*-(2*R*,3*S*)-Epoxy-3-(4-methoxyphenyl)-1-phenyl-1-propanon-1-one^{3a}:** ¹H NMR (400 MHz) δ 3.80 (s, 3H), 4.02 (d, *J* 2.0 Hz, 1H), 4.28 (d, *J* 2.0 Hz, 1H), 6.86-6.96 (m, 2H), 7.23-7.34 (m, 2H), 7.40-7.63 (m, 3H), 7.96-8.04 (m, 2H).

The absolute configuration was determined by comparison with the optical rotation reported in the literature $[\alpha]_D^{20} = -128.2$ (*c* 0.61, CHCl₃) [lit.^{3a} for (2*S*,3*R*)-epoxy-3-(4-methoxyphenyl)-1-phenyl-1-propanon-1-one: $[\alpha]_D = +131$ (*c* 0.70, CHCl₃) for 82% ee].

***trans*-(2*R*,3*S*)-Epoxy-3-(4-chlorophenyl)-1-phenyl-1-propanon-1-one^{3c}:** ¹H NMR (400 MHz) δ 4.06 (d, *J* 1.8 Hz, 1H), 4.24 (d, *J* 1.8 Hz, 1H), 7.25-7.68 (m, 7H), 7.97-8.05 (m, 2H).

The absolute configuration was determined by chiral HPLC using Chiralcel OD column and compared with the literature data.⁶

***trans*-(3*R*,4*S*)-Epoxy-4-phenylbutan-2-one**^{3d}: ¹H NMR (400 MHz) δ 2.18 (s, 3H), 3.49 (d, *J* 2.0 Hz, 1H), 4.00 (d, *J* 2.0 Hz, 1H), 7.19-7.46 (m, 5H).

The absolute configuration was determined by comparison with the optical rotation reported in the literature $[\alpha]_D^{18} = -75.9$ (*c* 1.0, CHCl₃) [lit.^{3d} for (3*S*,4*R*)-epoxy-4-phenylbutan-2-one: $[\alpha]_D^{25} = +96.5$ (*c* 1.0, CHCl₃) for 94% ee].

***trans*-(2*R*,3*S*)-Epoxy-1-phenylbutan-1-one**^{3e}: ¹H NMR (400 MHz) δ 1.52 (d, *J* 5.2 Hz, 3H), 3.23 (dq, *J* 2.0, 5.2 Hz, 1H), 3.98 (d, *J* 2.0 Hz, 1H), 7.48-7.65 (m, 3H), 7.99-8.03 (m, 2H).

The absolute configuration was determined by comparison with the optical rotation reported in the literature $[\alpha]_D^{25} = -10.0$ (*c* 0.6, CHCl₃) [lit.^{3e} for (2*S*,3*R*)-epoxy-1-phenylbutan-1-one: $[\alpha]_D^{25} = +1.5$ (*c* 0.56, CHCl₃) for 10% ee].

***trans*-(3*R*,4*S*)-Epoxy-6-phenylhexan-2-one**^{3d}: ¹H NMR (400 MHz) δ 1.95 (dt, *J* 5.6, 7.6 Hz, 2H), 2.02 (s, 3H), 2.77-2.82 (m, 2H), 3.08 (dt, *J* 2.0, 5.6 Hz, 1H), 3.17 (d, *J* 2.0 Hz, 1H), 7.16-7.34 (m, 5H).

The absolute configuration was determined by comparison with the optical rotation reported in the literature $[\alpha]_D^{19} = +15.5$ (*c* 1.23, CHCl₃) [lit.^{3d} for (3*S*,4*R*)-epoxy-6-phenylhexan-2-one: $[\alpha]_D^{26} = -21.9$ (*c* 1.24, CHCl₃) for 88% ee].

(2*S*,3*R*)-2-Methyl-2,3-epoxy-1,4-naphthoquinone⁷: ¹H NMR (400 MHz) δ 1.74 (s, 3H), 3.86 (s, 1H), 7.15-7.80 (m, 2H), 7.92-7.98 (m, 1H), 8.00-8.06 (m, 1H).

The absolute configuration was determined by chiral HPLC using Chiralpak AD column and compared with the literature data.^{3b}

Table 1 shows HPLC conditions and retention times for each epoxy ketone:

Table 1. HPLC Conditions and Retentions Times for the Determination of Enantiomeric Excess of the Epoxy Ketones^a

epoxy ketone		chiral column	eluent (hexane/2-propanol)	flow rate (mL/min)	retention time (min)		ref.
R ¹	R ²				(2 <i>S</i> ,3 <i>R</i>)	(2 <i>R</i> ,3 <i>S</i>)	
Ph	Ph	Chiralcel OD	98/2	1	17.3	18.5	3d
<i>p</i> -Br-C ₆ H ₄	Ph	Chiralcel OD	95/5	0.8	22.6	20.2	note b
<i>m</i> -Me-C ₆ H ₄	Ph	Chiralcel OD	30/1	0.8	17.8	18.8	3b
β-Naphthyl	Ph	Chiralpak AD	95/5	1	19.4	22.2	4
Ph	<i>p</i> -MeO-C ₆ H ₄	Chiralcel OD	95/5	0.8	21.7	23.1	5
Ph	<i>p</i> -Cl-C ₆ H ₄	Chiralcel OD	55/1	0.5	54.5	57.0	6
Me	Ph	Chiralpak AD	98/2	1	13.7 (3 <i>S</i> ,4 <i>R</i>)	10.9 (3 <i>R</i> ,4 <i>S</i>)	3d
Ph	Me	Chiralcel OD	94/6	0.8	10.7	11.6	5
Me	Ph(CH ₂) ₂	Chiralpak AD	98/2	1	12.4 (3 <i>S</i> ,4 <i>R</i>)	10.0 (3 <i>R</i> ,4 <i>S</i>)	3f
Chiralpak AD	20/1	1	10.6	11.7	3b

^a Detection at 254 nm. ^b This work. The corresponding enantiomers of the racemic and enantiomerically enriched epoxide were cleanly separated under conditions used for the analysis. See below for the HPLC trace.

CHANNEL A

STORED TO BIN # 3

DATA SAVED TO BIN # 3

CH= "A" PS= 1.

FILE	1.	METHOD	0.	RUN	3	INDEX	3	BIN	3
PEAK#	AREA%	RT	AREA	BC					
1	0.155	16.65	571	02					
2	0.245	17.67	901	03					
3	51.277	21.39	188917	02					
4	48.323	23.5	178035	03					
TOTAL	100.		368424						

Figure 1. HPLC (Chiralcel OD) trace of racemic *trans*-2,3-epoxy-3-phenyl-1-(4-bromophenyl)-propan-1-one

CH= "A" RS= 1

FILE 1.	METHOD 0.	RUN 1	INDEX 1	BIN 1
PEAK#	AREA%	RT	AREA	BO
1	0.024	3.81	79.01	
2	0.044	4.58	146.02	
3	0.056	4.91	186.03	
4	0.037	7.3	122.01	
5	87.118	20.29	289803.01	
6	12.721	22.69	42518.01	
TOTAL	100.		382654	

Figure 2. HPLC (Chiralcel OD) trace of enantiomerically enriched *trans*-(2R,3S)-epoxy-3-phenyl-1-(4-bromophenyl)-propan-1-one

NLE investigation (Figure 2 of the paper) is reported in tabulated form:

Table 2. Linear effect in the Epoxidation of **4a** Using Catalyst **2** of Different Degrees of Optical Purity^a

entry	ee 2 %	t(h)	yield 5a % ^b	ee 5a % ^c
1	100	48	53	76
2	82	48	51	66
3	58	48	47	46
4	34	48	51	28
5	18	48	49	14

^a The reaction was carried out in hexane at room temperature with 1.2 equiv of TBHP in the presence of 30 mol% of **2** at C=0.2 M of **4a**. ^b Yield of isolated product. ^c The absolute configuration of the predominant enantiomer was 2*R*,3*S*.

References

1. Bailey, D.; O'Hagan, D.; Tavash, M. *Tetrahedron: Asymmetry* **1997**, 8, 149.
2. (a) Kohler, E. P.; Chadwell, H. M. *Org. Synth. Col. Vol. I*, **1932**, 78. (b) Riahi, A.; Thorey, C.; Hénin, F.; Muzart, J. *Synth. Commun.* **1998**, 28, 4339.
3. (a) Lygo, B.; Wainwright, P. G. *Tetrahedron* **1999**, 55, 6289. (b) Arai, S.; Tsuge, H.; Oku, M.; Miura, M.; Shioiri, T. *Tetrahedron* **2002**, 58, 1623. (c) Kumaraswamy, G.; Sastry, M. N. V.; Nivedita, J.; Ravi Kumar K.; Vairamani, M. *Tetrahedron: Asymmetry* **2003**, 14, 3797. (d) Bougauchi, M.; Watanabe, S.; Arai, T.; Sasai, H.; Shibasaki, M. *J. Am. Chem. Soc.* **1997**, 119, 2329. (e) Watanabe, S.; Arai, T.; Sasai, H.; Bougauchi, M.; Shibasaki, M. *J. Org. Chem.* **1998**, 63, 8089. (f) Nemoto, T.; Ohshima, T.; Yamaguchi, K.; Shibasaki, M. *J. Am. Chem. Soc.* **2001**, 123, 2725.
4. Lasterra-Sánchez, M. E.; Felfer, U.; Mayon, P.; Roberts, S. M.; Thorton, S. R.; Todd, C. J. *J. Chem. Soc., Perkin Trans. 1* **1996**, 343.
5. Adam, W.; Bheema Rao, P.; Degen, H.-G.; Saha-Möller, C. R. *J. Am. Chem. Soc.* **2000**, 122, 5654.
6. Ooi, T.; Ohara, D.; Tamura, M.; Maruoka, K. *J. Am. Chem. Soc.* **2004**, 126, 6844.
7. (a) Snatzke, G.; Wynberg, H.; Feringa, B.; Marsman, B. G.; Greydanus, B.; Pluim, H.; *J. Org. Chem.* **1980**, 45, 4094.