Supporting Information for:

Enantioselective Rh-Catalyzed Arylation of *N*-Tosylarylimines with Arylboronic Acids

Hai-Feng Duan, Yi-Xia Jia, Li-Xin Wang, Qi-Lin Zhou*

State Key Laboratory and Institute of Elemento-organic Chemistry, Nankai University

Tianjin 300071, China

CONTENTS

(A) General Procedure for Rhodium-Catalyzed Arylation of *N*-Tosylarylimines with Arylboronic Acids

(B) Analytical Data of Products

(C) Deprotection of 4-Methylbenzenesulfonyl Group

(D) Crystal Structure and Structure Refinement for [Rh(COD)((*S*)-1a)₂]BF₄

General:

All reactions and manipulations were performed using standard Schlenk techniques. Melting points were measured on a RY-I apparatus and uncorrected. NMR spectra were recorded with a Varian or Bruker spectrometer at 400 MHz (^1H NMR), 100 MHz (^{13}C NMR) in CDCl_3 . Chemical shifts were reported in ppm down field from internal Me_4Si . Optical rotations were determined using a Perkin Elmer 341 MC polarimeter. Elemental analyses were performed on the Yanaco CDRDER MT-3 instrument. Mass spectra were recorded on the VG-7070E spectrometer. HPLC analyses were performed using a Hewlett Packard Model HP 1100 Series or Waters 2996 instruments. $\text{MeOCH}_2\text{CH}_2\text{OMe}$, $\text{ClCH}_2\text{CH}_2\text{Cl}$, THF, dioxane, 2-PrOH and toluene were simply distilled before use. $[\text{RhCl}(\text{C}_2\text{H}_4)_2]_2$,¹ $\text{Rh}(\text{acac})(\text{C}_2\text{H}_4)_2$,¹ $\text{Rh}(\text{COD})(\text{MeCN})_2[\text{BF}_4]$,² and *N*-Tosylarylimine³ were prepared according to the reported procedures. Ligand (*S*)-**1** can be obtained from J&K Chemical Co.⁴ Phenylboronic acid and 4-fluorophenylboronic acid were purchased from Shijiazhuang TianYue Fine Chemicals Co., Ltd. All other arylboronic acids were purchased from Aldrich and used as received.

(A) General Procedure for Rhodium-Catalyzed Arylation of *N*-Tosylarylimines with Arylboronic Acids

A Schlenk tube was charged with $\text{Rh}(\text{acac})(\text{CH}_2\text{CH}_3)_2$ (1.5 mg, 0.0057 mmol, 3 mol % Rh) and **1** ((*S*)-ShiP) (4.5 mg, 0.0114 mmol, 2 equiv to Rh) in nitrogen atmosphere. Toluene (1 mL) was then added and the mixture was stirred at 35 °C for 10 min. Water (1 mL), $\text{PhB}(\text{OH})_2$ (0.38 mmol), KF (44 mg, 0.72 mmol) and 4-chlorobenzaldehyde *N*-tosylimine **7a** (0.19 mmol) were added sequentially and the mixture was stirred for 20 h at 35 °C. The mixture was extracted with CH_2Cl_2 , dried over MgSO_4 . After concentration the residue was chromatographed on silica gel column with petroleum ether/ethyl acetate (8:1) to give the product **9aa** as a white solid (85 % yield).

(B) Analytical Data of Products

(*R*)-*N*-[(4-Chlorophenyl)phenylmethyl]-4-methylbenzenesulfonamide (9aa).

85% yield, white solid, m.p. 120-122 °C. $[\alpha]_D^{20} = +4.8$ (*c* 0.97, CHCl₃), 93% *ee* [HPLC conditions: Chiracel OD-H column, hexane/2-propanol = 93:7, flow = 1.0 mL/min, wavelength = 230 nm, t_R = 18.8 min for (*S*)-enantiomer and t_R = 26.7 min for (*R*)-enantiomer]. ¹H NMR δ 7.54 (d, *J* = 8.0 Hz, 2H), 7.22–7.11 (m, 7H), 7.02–7.06 (m, 4H), 5.53 (d, *J* = 6.4 Hz, 1H), 5.41 (br s, 1H), 2.38 (s, 3H).

(*R*)-*N*-[(4-Fluorophenyl)phenylmethyl]-4-methylbenzenesulfonamide (9ba).

80% yield, white solid, m.p. 117-119 °C. $[\alpha]_D^{20} = -4.1$ (c 1.02, CHCl_3), 93% *ee* [HPLC conditions: Chiralcel OD-H column, hexane/2-propanol = 80:20, flow = 0.8 mL/min, wavelength = 230 nm, t_R = 10.0 min for (*S*)-enantiomer, t_R = 12.0 min for (*R*)-enantiomer]. ^1H NMR δ 7.55 (d, J = 8.4 Hz, 2H), 7.22–7.20 (m, 3H), 7.15 (d, J = 8.0 Hz, 2H), 7.10–7.04 (m, 4H), 6.91–6.86 (m, 2H), 5.55 (d, J = 6.8 Hz, 1H), 5.16 (d, J = 6.8 Hz, 1H), 2.38 (s, 3H). ^{13}C NMR δ 143.2, 140.3, 137.3, 136.4, 129.3, 129.1, 129.0, 128.6, 127.6, 127.2, 127.1, 115.4, 115.1, 60.7, 21.4. MS (EI) m/z 355 (M^+). Anal. Calcd for $\text{C}_{20}\text{H}_{18}\text{FNO}_2\text{S}$: C, 67.58; H, 5.10; N, 3.94; Found: C, 67.80; H, 5.19; N, 3.84.

(*R*)-*N*-[(4-Bromophenyl)phenylmethyl]-4-methylbenzenesulfonamide (9ca).

82% yield, white solid, m.p. 121-123 °C. $[\alpha]_D^{20} = +4.1$ (*c* 0.81, CHCl₃), 93% *ee* [HPLC conditions: Chiralcel OD-H column, hexane/2-propanol = 80:20, flow = 0.5 mL/min, wavelength = 230 nm, t_R = 32.0 min for (*S*)-enantiomer, t_R = 44.8 min for (*R*)-enantiomer. ¹H NMR δ 7.55 (d, *J* = 8.4 Hz, 2H), 7.33 (d, *J* = 8.0 Hz, 2H), 7.22–7.20 (m, 3H), 7.16 (d, *J* = 8.4 Hz, 2H), 7.04–7.02 (m, 2H), 7.00 (d, *J* = 8.0 Hz, 2H), 5.51 (d, *J* = 6.8 Hz, 1H), 5.03 (d, *J* = 6.8 Hz, 1H), 2.39 (s, 3H). ¹³C NMR δ 143.4, 140.0, 139.5, 137.2, 131.5, 129.4, 129.1, 128.7, 127.8, 127.2, 127.1, 121.5, 60.8, 21.5. MS (EI) *m/z* 415 (*M*⁺). Anal. Calcd for C₂₀H₁₈BrNO₂S: C, 57.70; H, 4.36; N, 3.36; Found: C, 58.13; H, 4.10; N, 3.11.

(*R*)-*N*-[(4-Methylphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9da).⁵

77% yield, white solid, m.p. 114-116 °C. $[\alpha]_D^{20} = +11.0$ (c 0.7, CHCl_3), 94% *ee* [HPLC conditions: Chiralcel OD-H column, hexane/2-propanol = 95:5, flow = 1.0 mL/min, wavelength = 230 nm, t_R = 17.5 min for (*S*)-enantiomer, t_R = 25.9 min for (*R*)-enantiomer]. ^1H NMR δ 7.56 (d, J = 8.3 Hz, 2H), 7.20–7.08 (m, 7H), 7.02–6.95 (m, 4H), 5.52 (d, J = 6.8 Hz, 1H), 5.02 (d, J = 6.8 Hz, 1H), 2.37 (s, 3H), 2.27 (s, 3H).

(*R*)-*N*-[(4-Methoxyphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9ea).

65% yield, white solid, m.p. 145-147 °C. $[\alpha]_D^{20} = +13.6$ (*c* 0.58, CHCl₃), 93% *ee* [HPLC conditions: Chiralcel OD-H column, hexane/2-propanol = 80:20, flow = 0.5 mL/min, wavelength = 230 nm, t_R = 20.0 min for (*S*)-enantiomer, t_R = 31.7 min for (*R*)-enantiomer]. ¹H NMR δ 7.56 (d, *J* = 8.8 Hz, 2H), 7.20–7.18 (m, 3H), 7.14 (d, *J* = 8.8 Hz, 2H), 7.11–7.08 (m, 2H), 6.99 (d, *J* = 8.8 Hz, 2H), 6.73 (d, *J* = 8.8 Hz, 2H), 5.52 (d, *J* = 6.8 Hz, 1H), 5.03 (br s, 1H), 3.74 (s, 3H), 2.37 (s, 3H).

(*R*)-*N*-[(3-Methoxyphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9fa).

72% yield, white solid, m.p. 153-155 °C. $[\alpha]_D^{20} = +2.9$ (*c* 1.34, CHCl₃), 96% *ee* [HPLC conditions: Chiralcel OD column, hexane/2-propanol = 95;5, flow = 0.8 mL/min, wavelength = 230 nm, $t_R = 29.7$ min for (*R*)-enantiomer, $t_R = 34.4$ min for (*S*)-enantiomer. ¹H NMR δ 7.57 (d, *J* = 8.8 Hz, 2H), 7.21–7.18 (m, 3H), 7.14–7.09 (m, 5H), 6.73–6.71 (m, 1H), 6.67 (d, *J* = 8.0 Hz, 1H), 6.60 (s, 1H), 5.53 (d, *J* = 6.8 Hz, 1H), 5.06 (br s, 1H), 3.68 (s, 3H), 2.37 (s, 3H). ¹³C NMR δ 159.6, 143.1, 142.0, 140.4, 137.4, 129.5, 129.3, 128.5, 127.5, 127.3, 127.2, 119.7, 113.1, 112.9, 61.3, 55.1, 21.4. MS (EI) *m/z* 367 (*M*⁺). Anal. Calcd for C₂₁H₂₁NO₃S: C, 68.64; H, 5.76; N, 3.81; Found: C, 68.45; H, 5.93; N, 3.72.

(*R*)-*N*-[(2-Chlorophenyl)phenylmethyl]-4-methylbenzenesulfonamide (9ga).

79% yield, white solid, m.p. 169-171 °C. $[\alpha]_D^{20} = -16.5$ (c 0.49, CHCl_3), 95% *ee* [HPLC conditions: Chiralpak AD-H column, hexane/2-propanol = 90:10, flow = 0.7 mL/min, wavelength = 230 nm, t_R = 36.4 min for (*S*)-enantiomer, t_R = 41.8 min for (*R*)-enantiomer]. ^1H NMR δ 7.62 (d, J = 8.8 Hz, 2H), 7.35–7.32 (m, 1H), 7.23–7.21 (m, 4H), 7.16–7.14 (m, 4H), 7.06–7.04 (m, 2H), 5.90 (d, J = 6.8 Hz, 1H), 5.22 (d, J = 6.8 Hz, 1H), 2.37 (s, 3H).

(*R*)-*N*-[(2-Bromophenyl)phenylmethyl]-4-methylbenzenesulfonamide (9ha).

76% yield, white solid, m.p. 172-174 °C. $[\alpha]_D^{20} = -23.3$ (c 0.54, CHCl_3), 93% *ee* [HPLC conditions: Chiralpak AD-H column, hexane/2-propanol = 90:10, flow = 0.7 mL/min, wavelength = 230 nm, t_R = 38.8 min for (*S*)-enantiomer, t_R = 42.8 min for (*R*)-enantiomer]. ^1H NMR δ 7.63 (d, J = 7.6 Hz, 2H), 7.43–7.34 (m, 2H), 7.23–7.15 (m, 6H), 7.09–7.04 (m, 3H), 5.91 (d, J = 7.2 Hz, 1H), 5.23 (br s, 1H), 2.37 (s, 3H). ^{13}C NMR δ 143.3, 139.3, 139.1, 137.0, 133.1, 129.5, 129.4, 129.0, 128.6, 127.8, 127.5, 127.4, 127.2, 123.1, 60.5, 21.4; MS (EI) m/z 415(M^+). Anal. Calcd for $\text{C}_{20}\text{H}_{18}\text{BrNO}_2\text{S}$: C, 57.70; H, 4.36; N, 3.36; Found: C, 57.93; H, 4.17; N, 3.11.

(*R*)-*N*-[(2-Methylphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9ia).⁶

73% yield, white solid, m.p. 136-138 °C. $[\alpha]_{\text{D}}^{20} = -10.0$ (c 0.98, CHCl_3), 95% *ee* [HPLC conditions: Chiralcel OD-H column, hexane/2-propanol = 80:20, flow = 0.5 mL/min, wavelength = 230 nm, $t_{\text{R}} = 12.0$ min for (*R*)-enantiomer, $t_{\text{R}} = 15.1$ min for (*S*)-enantiomer]. ^1H NMR δ 7.54 (d, $J = 9.2$ Hz, 2H), 7.19–7.17 (m, 3H), 7.12–7.09 (m, 4H), 7.06–7.03 (m, 4H), 5.79 (d, $J = 7.2$ Hz, 1H), 5.23 (br s, 1H), 2.35 (s, 3H), 2.15 (s, 3H).

(*R*)-*N*-[(2-Methoxyphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9ja).

67% yield, white solid, m.p. 127-128 °C. $[\alpha]_{\text{D}}^{20} = +23.9$ (c 0.51, CHCl_3), 92% *ee* [HPLC conditions: Chiralcel OD-H column, hexane/2-propanol = 90:0, flow = 0.8 mL/min, wavelength = 230 nm, $t_{\text{R}} = 14.8$ min for (*R*)-enantiomer, $t_{\text{R}} = 18.3$ min for (*S*)-enantiomer. ^1H NMR δ 7.52 (d, $J = 8.8$ Hz, 2H), 7.22–7.12 (m, 6H), 7.04 (d, $J = 8.8$ Hz, 2H), 6.98 (d, $J = 6.4$ Hz, 1H), 6.76 (t, $J = 7.6$ Hz, 1H), 6.66 (d, $J = 7.6$ Hz, 1H), 6.64 (d, $J = 9.2$ Hz, 1H), 5.67 (d, $J = 9.2$ Hz, 1H), 3.57 (s, 3H), 2.31 (s, 3H).

(*R*)-*N*-[(1-Naphthyl)phenylmethyl]-4-methylbenzenesulfonamide (9ka).

72% yield, white solid, m.p. 176-177 °C. $[\alpha]_D^{20} = +3.9$ (*c* 0.50, CHCl₃), 96% *ee* [HPLC conditions: Chiralcel OD-H column, hexane/2-propanol = 80:20, flow = 0.5 mL/min, wavelength = 230 nm, t_R = 25.2 min for (*R*)-enantiomer, 32.2 min for (*S*)-enantiomer]. ¹H NMR δ 7.85–7.78 (m, 3H), 7.73 (d, *J* = 7.6 Hz, 1H), 7.51–7.36 (m, 4H), 7.30–7.23 (m, 2H), 7.20–7.18 (m, 2H), 7.15–7.12 (m, 2H), 7.07 (d, *J* = 8.0 Hz, 2H), 6.31 (d, *J* = 6.8 Hz, 1H), 5.14 (d, *J* = 6.8 Hz, 1H), 2.35 (s, 3H).

(*R*)-*N*-[(2-Furyl)phenylmethyl]-4-methylbenzenesulfonamide (9la).

84% yield, brown solid, m.p. 134-136 °C. $[\alpha]_D^{20} = +4.6$ (c 0.50, CHCl_3), 85% *ee* [HPLC conditions: Chiralcel OD-H column, hexane/2-propanol = 95:5, flow = 0.8 mL/min, wavelength = 230 nm, t_R = 25.5 min for (*S*)-enantiomer, t_R = 27.8 min for (*R*)-enantiomer]. ^1H NMR δ 7.58 (d, J = 8.4 Hz, 2H), 7.25–7.22 (m, 5H), 7.18–7.14 (m, 5H), 5.61 (d, J = 7.2 Hz, 1H), 5.12 (d, J = 7.2 Hz, 1H), 2.37 (s, 3H).

(S)-N-[(4-Fluorophenyl)phenylmethyl]-4-methylbenzenesulfonamide (9mb).

The enantiomer of **9ba**, 56% yield, 94% ee, $[\alpha]_{\text{D}}^{20} = +4.5$ (*c* 0.97, CHCl₃).

(S)-N-[(4-Methylphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9mc).

The enantiomer of **9da**, 75% yield, 92% ee, $[\alpha]_{\text{D}}^{20} = -9.5$ (*c* 0.68, CHCl₃).

(S)-N-[(4-Methoxyphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9md).

The enantiomer of **9ea**, 70% yield, 95% ee, $[\alpha]_{\text{D}}^{20} = -14.8$ (*c* 0.75, CHCl₃).

(S)-N-[(3-Methoxyphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9me).

The enantiomer of **9fa**, 68% yield, 91% ee, $[\alpha]_{\text{D}}^{20} = -2.4$ (*c* 1.10, CHCl₃).

(S)-N-[(2-Methylphenyl)phenylmethyl]-4-methylbenzenesulfonamide (9mf).

The enantiomer of **9ia**, 65% yield, 93% ee, $[\alpha]_{\text{D}}^{20} = +9.6$ (*c* 0.85, CHCl₃) for 93% ee.

(C) Deprotection of 4-Methylbenzenesulfonyl Group ⁵

A 0.1 M solution of SmI₂ in THF (27 mL, 2.7 mmol) and HMPA (1.3 mL, 7.43 mmol) were added to sulfonamide **9da** (87 mg, 0.25 mmol, 94% ee) at room temperature. After refluxing for 6 h, the mixture was cooled and quenched with brine and extracted with ether. The combined organic layer was washed with saturated aq NaHCO₃ and brine, dried over Na₂SO₄, and finally purified by chromatography on silica gel with petroleum ether/ethyl acetate/Et₃N (2:1:0.05) to give (*R*)-(4-methylphenyl)phenylmethylamine **10** as pale yellow oil (40 mg, 82 % yield). $[\alpha]_{\text{D}}^{20} = +1.3$ (*c* 2.0, CHCl₃). ¹H NMR (CDCl₃): 7.38–7.35 (m, 2H), 7.32–7.26 (m, 3H, 7.24–7.22 (m, 2H), 7.12 (d, *J* = 8.1Hz, 2H), 5.17 (s, 1H), 2.31 (s, 3), 1.99 (br s, 2H). Protection of **10** with p-toluenesulfonyl chloride according to reported procedure ⁷ to give (*R*)-N-[(4-methylphenyl)phenylmethyl]-4-methylbenzenesulfonamide in 88% yield with 94% ee.

Deprotection of Ts group of sulfonamide **9ka** was carried out by the same procedure as **9da**, giving (*R*)-(1-naphthyl)phenylmethylamine **11** as a white solid (74% yield). m.p. 65-68 °C. $[\alpha]_{\text{D}}^{20} = +40.9$ (*c* 1.0, CHCl₃) (lit.⁸ $[\alpha]_{\text{D}}^{20} = +42.2$ (*c* 1.0, CHCl₃) for (*R*) isomer 96% ee). ¹H NMR (CDCl₃): 8.07–8.04 (m, 1H), 7.86–7.84 (m, 1H), 7.77 (d, d, *J* = 8.1Hz, 1H), 7.65

(dd, $J = 8.1\text{ Hz}$, 1H), 7.50–7.37 (m, 4H), 7.32–7.21 (m, 4H), 5.95 (s, 1H), 1.90 (s, 1H). The ee value of **11** was determined by using the same method as **10**. (*R*)-*N*-[(1-naphthyl)-phenylmethyl]-4-methylbenzenesulfonamide was obtained in 81% yield with 96% ee.

(D) Crystal Structure and Structure Refinement for $[\text{Rh}(\text{COD})((S)\text{-1})_2]\text{BF}_4$

Crystal Structure of $[\text{Rh}(\text{COD})((S)\text{-1})_2]\text{BF}_4$ (Anion was omitted for clarity)

Crystal Data and Structure Refinement for $[\text{Rh}(\text{COD})((S)\text{-1})_2]\text{BF}_4$

Empirical formula	$\text{C}_{55.50} \text{H}_{53} \text{B} \text{Cl}_3 \text{F}_4 \text{O}_6 \text{P}_2 \text{Rh}$
Formula weight	1173.99
Temperature	294(2) K
Wavelength	0.71073 Å
Crystal system, space group	Monoclinic, $P2(1)$
Unit cell dimensions	$a = 15.362(2) \text{ Å}$ $\alpha = 90^\circ$ $b = 10.6888(14) \text{ Å}$ $\beta = 92.150(2)^\circ$ $c = 16.338(2) \text{ Å}$ $\gamma = 90^\circ$
Volume	$2680.7(6) \text{ Å}^3$
Z, Calculated density	2, 1.454 Mg/m^3
Absorption coefficient	0.591 mm^{-1}
$F(000)$	1202
Crystal size	$0.32 \times 0.24 \times 0.20 \text{ mm}$
Theta range for data collection	1.79 to 26.37° .
Limiting indices	$-15 \leq h \leq 19$, $-13 \leq k \leq 13$, $-20 \leq l \leq 11$
Reflections collected / unique	15197 / 10282 [$R(\text{int}) = 0.0357$]
Completeness to $\theta = 26.37$	99.4 %

Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	1.000000 and 0.771606
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	10282 / 65 / 667
Goodness-of-fit on F ²	1.028
Final R indices [I>2σ(I)]	R1 = 0.0494, wR2 = 0.1137
R indices (all data)	R1 = 0.0880, wR2 = 0.1320
Absolute structure parameter	0.03(3)
Largest diff. peak and hole	0.713 and -0.434 e. Å ⁻³

References

- (1) Cramer, R. *J. Am. Chem. Soc.* **1964**, 86, 217.
- (2) Oi, S.; Moro, M.; Fukuhara, H.; Kawanishi, T.; Inoue, Y. *Tetrahedron* **2003**, 59, 4351.
- (3) (a) Vishwakarama, L. C.; Stringer, O. D.; Davis, F. A. *Organic Syntheses* **1987**, 66, 203.
(b) Brain, E. L.; Prasad, S. R.; Thomas, C. W. *Synlett* **1994**, 493. (c) Chemla, F.; Hebbe, V.; Normant, J.-F. *Synthesis* **2000**, 75.
- (4) <http://www.jkchemical.com/products/ChiralSpiroPhosphorusLigands.asp>.
- (5) Kuriyama, M.; Soeta, T.; Hao, X. Y.; Chen, O.; Tomioka, K. *J. Am. Chem. Soc.* **2004**, 126, 8128.
- (6) Tokunaga, N.; Otomaru, Y.; Okamoto, K.; Ueyama, K.; Shintani, R.; Hayashi, T. *J. Am. Chem. Soc.* **2004**, 126, 13584.
- (7) Hayashi, T.; Ishigedani, M. *J. Am. Chem. Soc.* **2000**, 122, 976.
- (8) Hayashi, T.; Kawai, M.; Tokunaga, N. *Angew. Chem., Int. Ed.* **2004**, 43, 6125.