Enantioselective Synthesis of P-Stereogenic Benzophospholanes via Palladium-Catalyzed Intramolecular Cyclization

Tim J. Brunker,^a Brian J. Anderson,^a Natalia F. Blank,^a David S. Glueck,^{a} and Arnold L.
Rheingold^b*

^a6128 Burke Laboratory, Department of Chemistry, Dartmouth College, Hanover, New Hampshire,
03755 ^b Department of Chemistry, University of California, San Diego, La Jolla, California, 92093

Supporting Information

1. Experimental Section
2. X-ray Crystallographic Details
3. NMR Spectra

General Experimental Details. Unless otherwise noted, all reactions and manipulations were performed in dry glassware under a nitrogen atmosphere at 20 °C in a dry box or using standard Schlenk techniques. Petroleum ether (bp 38-53 °C), CH₂Cl₂, ether, THF, and toluene were dried over alumina columns similar to those described by Grubbs.¹ NMR spectra were recorded using Varian 300 or 500 MHz spectrometers. ¹H or ¹³C NMR chemical shifts are reported vs Me₄Si and were determined by reference to the residual ¹H or ¹³C solvent peaks. ³¹P NMR chemical shifts are reported vs H₃PO₄ (85%) used as an external reference. Coupling constants are reported in Hz, as absolute values unless noted otherwise. Unless indicated, peaks in NMR spectra are singlets.

Instead of the expected quartet resonances² in the ³¹P NMR spectra of the phosphine-boranes, poorly resolved multiplets were often observed; in some cases the signal was an apparent doublet. Elemental analyses were provided by Schwarzkopf Microanalytical Laboratory or Quantitative Technologies Inc. Mass spectra were recorded at the University of Illinois, Urbana-Champaign (<http://www.scs.uiuc.edu/~msweb>).

The following compounds were prepared by literature methods: *o*-IC₆H₄CH₂CH₂I,³ PH₂Ph(BH₃),⁴ PH₂Men(BH₃),⁵ Pd(diphos*)(stilbene) and Pd(Et-FerroTANE)(Ph)(I) catalyst precursors,⁶ Pd(dba)₂,⁷ and Pd((S)-Me₂NCH(Me)C₆H₄)(μ-Cl))₂.⁸

***o*-IC₆H₄CH₂CH₂PH(Ph)(BH₃) (**8a**)**

PH₂Ph(BH₃) (1.09 g, 8.55 mmol) was dissolved in THF (10 mL) and added slowly to a stirred suspension of NaH (205 mg, 8.55 mmol) in THF (20 mL) at 0 °C. H₂ evolution was observed immediately and by the end of the addition a purple-grey solution formed. The solution was stirred at 0 °C for a further 45 min and then a solution of 2-(2'-iodophenyl)ethyl iodide (3.06 g, 8.55 mmol) in THF (30 mL) was added dropwise over 10 min. The cooling bath was then removed and the reaction mixture was stirred for 18 h at room temperature. The pale green mixture was quenched by addition of H₂O (50 mL), transferred to a separating funnel and Et₂O (60 mL) was added. The organic layer was then separated and the aqueous layer was washed with additional Et₂O (60 mL). The organic extracts were washed with brine (2 x 50 mL) and dried over MgSO₄. The solution was then filtered and all solvent removed *in vacuo* to give a pale orange oil. Phosphine-borane **8a** (R_f = 0.26, 10% ethyl acetate/hexane) was separated from a small amount of iodide starting material by dry-column flash chromatography on silica. A small amount of unreacted 2-(2'-iodophenyl)ethyl iodide was eluted with hexane. Elution with 10% ethyl acetate/hexane yielded, after removal of solvent *in vacuo*, 2.33 g (6.58 mmol, 77%) of the product as a colorless cloudy oil, shown to be essentially pure by NMR spectroscopy.

Anal calcd for $C_{14}H_{17}BIP$: C, 47.50; H, 4.84. Found: C, 47.58; H, 4.99. HRMS calcd. for $C_{14}H_{16}BIP$ (M-H)⁺: m/z 353.0127. Found: m/z 353.0143. $^{31}P\{^1H\}$ NMR ($CDCl_3$): δ -3.3 (br d, J = 55). 1H NMR ($CDCl_3$): δ 7.82-7.71 (m, 3H, Ar), 7.58-7.46 (m, 3H, Ar), 7.31-7.20 (m, 2H, Ar), 6.94-6.88 (m, 1H, Ar), 5.57 (d of m, $^1J_{PH}$ = 340, 1H, PH), 3.05-2.85 (m, 2H, CH_2), 2.37-2.16 (m, 2H, CH_2), 1.5-0.3 (v. br q, 3H, BH_3). $^{13}C\{^1H\}$ NMR ($CDCl_3$): δ 143.1 (d, J = 13, Ar q), 139.9 (Ar), 133.1 (d, J = 9, Ar), 132.0 (d, J = 3, Ar), 129.6 (Ar), 129.3 (d, J = 10, Ar), 128.9 (Ar), 128.7 (Ar), 125.2 (d, J = 56, Ar q), 100.0 (Ar-I), 35.4 (d, J = 2, CH_2), 24.2 (d, J = 34, CH_2).

***o*-IC₆H₄CH₂CH₂PH(Cy)(BH₃) (8b)**

Cyclohexylphosphine (0.368 g, 3.17 mmol) was dissolved in THF (10 mL) and cooled to 0 °C. BH_3-SMe_2 (1.58 mL, 3.17 mmol, 2.0 M solution in THF) was then added by syringe and the mixture was stirred at 0 °C for 2 h. The solution was then cooled to -78 °C and *s*-BuLi (2.26 mL, 3.17 mmol, 1.4 M solution in cyclohexane) was added by syringe. After 40 min at -78 °C the solution was warmed to room temperature and stirred for an additional 20 min. The solution was then re-cooled to -78 °C and a solution of 2-(2'-iodophenyl)ethyl iodide (1.13 g, 3.17 mmol) in THF (10 mL) was added dropwise. The reaction mixture was allowed to warm up slowly and stir overnight. The reaction mixture was then quenched by the addition of H_2O (30 mL) and Et_2O (25 mL). The layers were separated, and the aqueous layer was extracted with Et_2O (2 x 25 mL). The combined organic extracts were washed with brine (30 mL) and dried over $MgSO_4$. The solution was filtered and all volatiles removed *in vacuo* to give a colorless oil which was purified by gradient column chromatography on silica. Elution with hexane first gave unreacted 2-(2'-iodophenyl)ethyl iodide (73 mg). Elution with 10% ethyl acetate/hexanes (product R_f = 0.46) then yielded 780 mg (2.17 mmol, 68% yield based on starting phosphine) of pale yellow viscous oil shown to be pure product by NMR spectroscopy.

HRMS m/z calcd. for $C_{14}H_{22}BIP$ (M-H)⁺: m/z 359.0597. Found: m/z 359.0611. $^{31}P\{^1H\}$ NMR (C_6D_6): δ 5.2 (br d, J = 57). 1H NMR (C_6D_6): δ 7.60 (dd, J = 8, 1, 1H, Ar), 6.92-6.84 (m,

2H, Ar), 6.49 (m, 1H, Ar), 4.00 (d of m, $^1J_{\text{PH}} = 351$, PH), 2.94-2.68 (m, 2H, CH₂), 1.75-1.52 (m, 2H), 1.49-1.39 (m, 5H), 1.37-1.26 (m, 1H), 1.20-1.02 (m, 2H), 1.00-0.85 (m, 3H), 1.8-0.8 (br q, 3H, BH₃). $^{13}\text{C}\{^1\text{H}\}$ NMR (C₆D₆): δ 143.9 (d, J = 12, Ar q), 139.8 (Ar), 129.8 (Ar), 128.8 (Ar), 128.5 (Ar), 100.2 (Ar q), 35.9 (CH₂), 30.6 (d, J = 35, CH), 28.5 (d, J = 2, CH₂), 27.8 (CH₂), 26.5 (d, J = 2, CH₂), 26.4 (d, J = 3, CH₂), 25.8 (d, J = 1, CH₂), 19.3 (d, J = 31, CH₂).

***o*-IC₆H₄CH₂CH₂PH(Men)(BH₃) (**8c**)**

A degassed solution of PH₂(Men)(BH₃) (0.455 g, 2.44 mmol) in dry THF (20 mL) was cooled to -78°C at which temperature *sec*-BuLi (2 mL, 2.8 mmol, 1.4 M solution in cyclohexanes, 1.2 equiv) was added via syringe. The reaction mixture was stirred at this temperature for 40 min and then was warmed up and stirred for 30 min at room temperature to ensure the completion of the reaction. The mixture was cooled again to -78°C and 2-(2'-iodophenyl)ethyl iodide (874 mg, 2.44 mmol) was added dropwise as a solution in THF (10 mL). The mixture was stirred overnight while warming up to room temperature and quenched with water. The mixture was extracted with ether (3x25 mL). The combined organic layers were washed with water (3x50 mL) and a saturated solution of NH₄Cl (50 mL) and dried over MgSO₄. The product was purified by column chromatography. Small amounts of unreacted 2-(2'-iodophenyl)ethyl iodide and PH₂(Men)(BH₃) as well as PH₂(Men) (deprotected during the reaction) were eluted with hexane. Elution with 2% ethyl acetate/hexanes yielded, after removal of solvent *in vacuo*, 438 mg (43%) of the product as a colorless oil, shown to be an essentially pure 1:1 mixture of diastereomers by NMR spectroscopy.

Anal. calcd for C₁₈H₃₁BIP: C, 51.95; H, 7.51. Found: C, 52.37; H, 7.39. HRMS calcd for C₁₈H₂₈BIP (M-3H)⁺: *m/z* 413.1067. Found: *m/z* 413.1062. Note: we observed similar mass spectroscopic behavior (loss of 3H) with other menthylphosphine-boranes.⁵ $^{31}\text{P}\{^1\text{H}\}$ NMR (toluene-*d*₈): δ -4.3 (br m), -6.3 (br m). ^1H NMR (toluene-*d*₈): δ 7.55 (1H, CH, Ar), 7.53 (1H, CH, Ar), 6.89 (d, J = 7.5, 1H, CH, Ar), 6.87 (1H, CH, Ar), 6.86 (1H, CH, Ar), 6.78-6.76 (m, 1H, CH, Ar), 6.52-6.44 (m, 2H, Ar), 4.43 (dm, J = 350, 1H, P-H), 4.36 (br dm, J = 355, 1H, P-H), 3.00-2.90

(m, 1H), 2.81-2.60 (m, 2H), 1.98-1.76 (m, 3H), 1.76-1.42 (m, 11 H), 1.32-1.12 (m, 7H), 0.94-0.82 (m, 10 H), 0.79 (d, J = 7, 3H, Me), 0.77 (d, J = 7, 3H, Me, overlapping with the previous signal), 0.62 (d, J = 7, 3H, Me), 0.54 (d, J = 7, 3H, Me). $^{13}\text{C}\{^1\text{H}\}$ NMR (toluene- d_8): δ 143.8 (d, J = 10, CH), 143.7 (d, J = 10, CH, overlapping with previous signal), 139.9 (CH), 137.4 (quat), 129.9 (CH), 129.6 (CH), 129.1 (CH), 128.7 (d, J = 4, CH), 128.4 (d, J = 5, CH), 128.2 (CH), 100.2 (Ar-I), 100.1 (Ar-I), 43.3 (d, J = 3, CH), 43.1 (CH), 36.8 (d, J = 23, CH_2), 36.6 (d, J = 22, CH_2), 34.7 (CH_2), 34.5 (d, J = 11, CH_2), 34.3 (d, J = 31, CH), 33.5 (d, J = 13, CH), 32.6 (d, J = 32, CH), 31.9 (CH_2), 29.4 (CH_2), 28.4 (d, J = 10, CH), 27.9 (d, J = 5, CH), 24.7 (d, J = 10, CH_2), 24.6 (d, J = 11, CH_2), 23.0 (CH_2), 22.4 (Me), 22.3 (Me), 21.3 (Me), 21.2 (Me), 19.8 (d, J = 4, CH_2), 19.6 (CH_2), 15.1 (Me), 14.3 (Me).

***o*-IC₆H₄CH₂CH₂PH(Ph) (**9a**)**

(2-(2'-iodophenyl)ethyl)phenylphosphine-borane (**8a**, 0.22 g, 0.63 mmol) and DABCO (84 mg, 0.75 mmol) were dissolved in benzene (4 mL) under N_2 and stirred at 50 °C for 1 h. The solvent was then removed *in vacuo* and the residue was dissolved in petroleum ether (5 mL) and filtered. The filtrate was pumped down to yield a sticky oil which was then redissolved in petroleum ether (2 mL) and passed through a thick plug of silica eluting with petroleum ether (15 mL). The solvent was removed in vacuo to yield (2-(2'-iodophenyl)ethyl)phenylphosphine **9a** (175 mg, 0.51 mmol, 81% yield) as a colorless mobile oil shown to be essentially pure by NMR spectroscopy. Oxidation of this air-sensitive phosphine was observed by both elemental analysis and mass spectroscopy. Satisfactory elemental analysis of a derivative, the phosphonium salt **9a**• HBF_4 is reported below.

Anal. Calcd for $\text{C}_{14}\text{H}_{14}\text{IP}$: C, 49.44; H, 4.15. Anal. Calcd for $\text{C}_{14}\text{H}_{14}\text{IOP}$: C, 47.21; H, 3.96. Found: C, 47.51; H, 3.87. HRMS calcd for $\text{C}_{14}\text{H}_{15}\text{IOP} (\text{MOH})^+$: m/z 356.9905. Found: m/z 356.9922. $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ -50.5. ^1H NMR (C_6D_6): δ 7.61 (dd, J = 8, 1, 1H, Ar), 7.39-7.35 (m, 2H, Ar), 7.09-7.05 (m, 3H, Ar), 6.86 (td, J = 8, 1, 1H, Ar), 6.73 (dd, J = 8, 2, 1H, Ar), 6.45 (td, J = 8, 2, 1H, Ar), 4.13 (ddd, $^1J_{\text{PH}} = 206$, $^3J_{\text{HH}} = 8, 6$, 1H, PH), 2.78-2.64 (m, 2H, CH_2), 1.94-1.78

(m, 2H, CH₂). ¹³C{¹H} NMR (C₆D₆): δ 145.1 (d, J = 8, Ar q), 139.7 (Ar), 135.6 (d, J = 12, Ar q), 134.1 (d, J = 16, Ar), 129.5 (Ar), 128.6 (d, J = 6, Ar), 128.4 (Ar), 128.3 (Ar), 128.0 (Ar, obscured by C₆D₆ peaks), 100.4 (Ar q), 40.1 (d, J = 8, CH₂), 24.2 (d, J = 15, CH₂).

(2-(2'-iodophenyl)ethyl)phenylphosphonium tetrafluoroborate (9a•HBF₄)

(2-(2'-iodophenyl)ethyl)phenylphosphine (**9a**, 80 mg, 0.24 mmol) was dissolved in Et₂O (1 mL) and added dropwise to a solution of HBF₄•OMe₂ (35 mg, 0.26 mmol) in Et₂O (1 mL). A white precipitate immediately formed. The mixture was stirred for an additional 5 min, and then the precipitate was allowed to settle and the solution was decanted. The solid was washed with additional Et₂O (2 mL) and then dried in vacuo to give analytically pure product as a white microcrystalline solid. Yield: 80 mg (0.19 mmol, 80%).

Anal. calcd for C₁₄H₁₅PIBF₄: C, 39.29; H, 3.53. Found: C, 39.50; H, 3.59. HRMS calcd for C₁₄H₁₅PI: m/z 340.9956. Found: m/z 340.9961. ³¹P{¹H} NMR (CD₃CN): δ -18.9 (broad). ³¹P NMR (CD₃CN): δ -18.9 (br t, ¹J_{PH} = 517). ¹H NMR (CD₃CN): δ 7.92-7.79 (m, 4H, Ar), 7.70-7.64 (m, 2H, Ar), 7.40-7.30 (m, 2H, Ar), 7.03-6.97 (m, 1H, Ar), 7.01 (broad d of m, ¹J_{PH} = 517, 2H, PH₂), 3.14-3.05 (m, 2H), 2.92-2.80 (br m, 2H). ¹³C{¹H} NMR (CD₃CN): δ 142.4 (d, J = 16, Ar), 141.2 (Ar), 136.2 (br, Ar), 135.5 (d, J = 11, Ar), 131.3 (Ar), 131.2 (Ar), 131.1 (Ar), 130.6 (Ar), 130.4 (Ar), 100.7 (C-I), 35.1 (CH₂), 18.7 (d, J = 45, CH₂).

***o*-IC₆H₄CH₂CH₂PH(Cy) (**9b**)**

(2-(2'-iodophenyl)ethyl)cyclohexylphosphine-borane (**8b**, 320 mg, 0.89 mmol) and diethylamine (0.92 mL, 8.89 mmol) were placed in an ampule under N₂ and heated to 50 °C for 4 h. The reaction mixture was then cooled to room temperature and excess diethylamine was removed *in vacuo*. The residue was dissolved in petroleum ether and filtered through a thick plug of silica in a glovebox.

The silica was washed with ca. 30 mL of petroleum ether and the filtrate was then pumped to dryness *in vacuo* to give 200 mg (0.58 mmol, 65% yield) of the product as a colorless mobile oil shown to be pure by NMR spectroscopy. For elemental analysis, the air-sensitive oil was converted to its white crystalline HBF₄ salt **9b•HBF₄** (see below).

³¹P{¹H} NMR (C₆D₆): δ -48.4. ¹H NMR (C₆D₆): δ 7.65 (dd, J = 7, 1, 1H, Ar), 6.93-6.87 (m, 2H, Ar), 6.50-6.47 (m, 1H, Ar), 3.07 (d of m, ¹J_{PH} = 194, PH), 2.83-2.76 (m, 2H), 1.87-1.79 (m, 2H), 1.77-1.73 (m, 2H), 1.64-1.54 (m, 4H), 1.20-1.08 (m, 5H). ¹³C{¹H} NMR (C₆D₆): δ 145.6 (d, J = 9, Ar q), 139.8 (Ar), 129.5 (Ar), 128.5 (Ar), 128.0 (Ar, obscured by solvent), 100.5 (Ar q), 40.3 (d, J = 12, CH₂), 32.8 (d, J = 7, CH₂), 32.1 (d, J = 17, CH₂), 31.4 (d, J = 8, CH), 27.3 (d, J = 4, CH₂), 27.2 (d, J = 9, CH₂), 26.5 (d, J = 1, CH₂), 19.7 (d, J = 15, CH₂).

(2-(2'-iodophenyl)ethyl)cyclohexylphosphonium tetrafluoroborate (9b•HBF₄)

(2-(2'-iodophenyl)ethyl)cyclohexylphosphine (**9b**, 80 mg, 0.23 mmol) was dissolved in Et₂O (1 mL) and added dropwise to a solution of HBF₄•OMe₂ (35 mg, 0.26 mmol) in Et₂O (1 mL). A white fluffy precipitate immediately formed. The mixture was stirred for 30 min, and then the solvent was removed *in vacuo*. The white residue was washed with Et₂O (2 x 2 mL), dried *in vacuo* and recrystallized by layering a solution in MeCN (2 mL) with Et₂O (10 mL) and storing at -30 °C for one day. The product was isolated as colorless needles and dried. Yield: 77 mg (0.19 mmol, 77%).

Anal. calcd for C₁₄H₂₁PIBF₄: C, 38.75; H, 4.88. Found: C, 38.67; H, 4.94. ³¹P{¹H} NMR (CD₃CN): δ -11.3 (broad). ¹H NMR (CD₃CN): δ 7.91 (dd, J = 8, 1, 1H, Ar), 7.43-7.37 (m, 2H, Ar), 7.06-7.03 (m, 1H, Ar), 5.94 (broad d, J = 454, 2H, PH), 3.16-3.11 (m, 2H), 2.61-2.55 (m, 2H), 2.55-2.48 (m, 1H), 2.03-1.96 (m, 2H), 1.87-1.80 (m, 2H), 1.74-1.69 (m, 1H), 1.54-1.45 (m, 2H), 1.42-1.33 (m, 2H), 1.32-1.25 (m, 1H). ¹³C{¹H} NMR (CD₃CN): δ 142.4 (d, J = 14, Ar q), 141.0 (Ar), 130.8 (Ar), 130.4 (Ar), 130.3 (Ar), 100.4 (Ar-I), 35.3 (d, J = 4), 28.2 (d, J = 3), 27.4 (d, J = 44), 26.5 (d, J = 15), 25.5 (d, J = 2), 14.3 (d, J = 44).

***o*-IC₆H₄CH₂CH₂PH(Men) (9c)**

A degassed solution of *o*-IC₆H₄CH₂CH₂PH(Men)(BH₃) (**8c**, 438.1 mg, 1.05 mmol) in toluene (10 mL) was heated to 60 °C, at which temperature a solution of DABCO (130 mg, 1.5 mmol, 1.1 equiv) in THF (5 mL) was added via syringe. The reaction mixture was stirred overnight at 60 °C. The solvent was removed in vacuo and the reaction flask was brought into the glove box. The product was purified by passing it through a short pad of silica (in a pipette, 5 mm x 30 mm) and washing the pad further with 40 mL of hexanes. Drying in vacuo resulted in 348 mg (82%) of the desired product, isolated as a colorless oil. The diastereomeric ratio of the product was 1.2:1. Spectra are reported for the mixture of diastereomers.

Anal. Calcd for C₁₈H₂₈IP: C, 53.74; H, 7.02. Found: C, 52.48; H, 6.59. HRMS calcd for C₁₈H₂₉IP (M+H)⁺: m/z 403.1052. Found: m/z 403.1039. ³¹P{¹H} NMR (THF): δ -73.6 (minor), -80.2 (major). ¹H NMR (C₆D₆): δ 7.63 (d, J = 8, 2H, Ar), 6.94-6.86 (m, 4H, Ar), 6.51-6.45 (m, 2H, Ar), 3.31 (dm, J = 198, 1H, P-H), 3.04 (dm, J = 160, 1H, P-H), 2.88-2.73 (m, 5H, overlapped with previous signal), 2.42-2.32 (m, 1H, CH), 2.14-2.04 (m, 1H, CH), 1.99-1.68 (m, 6H), 1.68-1.52 (m, 7H), 1.49-1.39 (m, 1H, CH), 1.30-1.13 (m, 4H), 1.13-1.02 (m, 2H), 1.02-0.94 (m, 1H), 0.90 (d, J = 7, 3H, Me), 0.88 (d, J = 7, 3H, Me), 0.84 (d, J = 6.5, 3H, Me), 0.83 (d, J = 6.5, 3H, Me, overlapping with previous signal), 0.78 (d, J = 7, 3H, Me), 0.74 (d, J = 7, 3H, Me). ¹³C{¹H} NMR (C₆D₆): δ 145.5 (d, J = 8, quat, Ar), 145.4 (d, J = 10, quat, Ar), 139.7 (Ar), 129.5 (Ar), 129.4 (Ar), 128.5 (Ar), 128.4 (Ar), 128.2 (Ar), 128.0 (Ar), 127.9 (Ar), 100.5 (quat, Ar), 100.4 (quat, Ar), 47.3 (d, J = 12, CH), 46.3 (d, J = 10, CH), 43.4 (d, J = 1, CH₂), 40.9 (d, J = 8, CH₂), 40.6 (d, J = 9, CH₂), 40.3 (d, J = 14, CH₂), 35.3 (CH₂), 35.2 (CH₂), 35.3 (d, J = 12, CH, overlapped with previous signal), 34.1 (d, J = 5, CH), 33.8 (d, J = 11, CH), 33.6 (d, J = 7, CH), 28.8 (d, J = 14, CH), 28.7 (d, J = 11, CH), 25.3 (d, J = 8, CH₂), 25.0 (d, J = 8, CH₂), 22.8 (Me), 22.7 (Me), 21.9 (Me), 21.6 (Me), 20.3 (d, J = 15, CH₂), 18.9 (d, J = 15, CH₂), 15.5 (Me), 15.3 (Me).

Alternative Method for Phosphine-Borane Deprotection (Conversion of **8a-c** into **9a-c**)

The general procedure is described for R = Ph (**8a**); the table also summarizes results for **8b-c**.⁹ Under N₂, piperazinomethyl polystyrene (Matrix Innovation (www.matrix-innovation.com) catalog number 3-221-1330, 1% DVB, 100-200 mesh, 2.7 mmol/g loading, 1.0 g, 2.7 mmol) was suspended in 15 mL of dry toluene. A solution of **8a** (0.48 g, 1.4 mmol) in 10 mL of dry toluene was added via cannula. The mixture was stirred at 50 °C for 12 h; the resin particles swelled. After the borane was removed, as monitored by ³¹P NMR spectroscopy, the suspension was filtered via cannula and the toluene was removed *in vacuo* to give the desired phosphine as a colorless oil (390 mg, 84%).

R	mmol 8	mmol resin	T (°C)	Solvent	Time (h)	Yield (%)
Ph (8a)	1.4	2.7	50	toluene	12	84
Cy (8b)	0.53	1.1	60	toluene	48	80
Men (8c)	0.84	1.7	60	toluene	48	80

Additional Information for Scheme 2: Pd-Catalyzed Intramolecular Phosphination

Figure S1. Structures of the diphosphine ligands in the catalyst precursors Pd(diphos*)(*trans*-stilbene).¹⁰

Catalyst Screening in Cyclization of 9a (General Method for 9a-c) A solution of **9a** (40 mg, 0.12 mmol) in toluene (0.5 mL) was added to a solution of the catalyst precursor Pd(diphos*)(*trans*-stilbene), 5 mol %) in a few drops of toluene. Typically on addition of the phosphine the solution darkened somewhat initially to orange and then the color faded back to yellow after a few seconds. NaOSiMe₃ (118 μL , 1.0 M solution in THF, 1.18×10^{-4} mol) was then added causing the solution to darken somewhat, and the mixture was transferred to an NMR tube. The reaction mixture was then monitored periodically by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy. A white precipitate (NaI) formed during the course of catalysis. When the reaction was complete, the mixture was pumped to dryness *in vacuo*. The residue was then slurried in petroleum ether (1 mL) and the catalyst byproduct removed by chromatography on a small silica column eluting with petroleum ether (15-20 mL). The clear solution was then pumped to dryness in a tared vial, to calculate the yield, to give a colorless oily product, which generally solidified on standing. The

product was confirmed to be benzophospholane **10a** by comparison of NMR spectroscopic data to the literature data (see below). Note: the reactions in Table 1 have not been optimized, so the reported times do not necessarily correspond to relative rates. Unlike the screening reactions, done in NMR tubes as described above, larger-scale reactions, such as entry 2, were carried out in flasks with stirring, which may affect the reaction rate.

ee determination for 10a and 10b: The general method is described for **10a**. Phosphine **10a** (typically 20 mg) as a solution in C₆D₆ (0.5 mL) was added to Pd((S)-Me₂NCH(Me)C₆H₄)(μ-Cl))₂ (0.6 equiv), and the mixture was shaken until all the Pd complex dissolved. This solution displayed well-resolved resonances at δ 45.4 (**a**) and 45.3 (**b**) in the ³¹P{¹H} NMR spectrum and at δ 8.30 (t, J = 8, **b**) and 7.95 (t, J = 8, **a**) in the ¹H NMR spectrum (where (**a**) and (**b**) represent the resonances belonging to each diastereomer) from which the ee was determined. The ee values in Table 1 in the text are those obtained from analysis of the ¹H NMR spectra, in which the signals were better baseline resolved; the ³¹P NMR spectra gave consistent results.

Note: catalysts with (*R,R*)-Me-DuPhos and (*R,R*)-Me-BPE ligands gave preferentially **10a** which yielded diastereomer **b** of the Pd(**10a**)(*C~N)(Cl) complex, while catalysts with (*R,R*)-*i*-Pr-DuPhos and (*S,S*)-Me-5-Fc favored the other enantiomer of **10a**, which gave diastereomer **a** of the Pd complex.

Table S2. Selected NMR Data (C₆D₆) Used to Determine the ee of Phosphines **10a** and **11a**

Phosphine	³¹ P NMR	¹ H NMR
10a	45.4 (a) 45.3 (b)	7.95 (t, J = 8, a) 8.30 (t, J = 8, b)
10b	55.5 (a) 54.8 (b)	8.46 (t, J = 7, a) 8.58 (t, J = 7, b)

Catalyst Screening in Cyclization of 9b. The general procedure described above for **9a** was followed. No color change from pale yellow was observed on addition of the phosphine. For all

catalysts tested, a white precipitate of NaI was immediately observed on addition of base and the reaction was shown to be complete by $^{31}\text{P}\{^1\text{H}\}$ NMR after 20 min; workup gave **10b** as a colorless oil. Its ee was determined as described above by analysis of ^1H and ^{31}P NMR spectra after complexation to a chiral Pd fragment (Table S2); in all cases racemic (within experimental error) phosphine **10b** was formed.

Characterization Data for the Benzophospholanes

Phosphine **10a** is a known compound (Quin, L. D.; Rao, N. S. *J. Org. Chem.* **1983**, *48*, 3754-3759), which was isolated as a colorless oily solid. $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ -3.0 (lit. δ -3.9). The ^1H NMR spectrum was not reported by Quin and Rao. ^1H NMR (C_6D_6): δ 7.53 (m, 1H, Ar), 7.22-6.94 (overlapping m, 8H, Ar), 2.97-2.85 (m, 1H), 2.74-2.62 (m, 1H), 2.09-1.92 (m, 1H), 1.84-1.76 (m, 1H). Our ^{13}C NMR data is consistent with that of Quin and Rao, although they reported only 6 of the expected 10 Ar peaks. $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ 150.1 (d, J = 2, Ar), 141.2 (d, J = 8, Ar), 140.8 (d, J = 24, Ar), 132.2 (Ar), 132.1 (Ar), 132.0 (Ar), 129.6 (Ar), 128.9 (d, J = 6, Ar), 127.4 (d, J = 8, Ar), 125.5 (d, J = 2, Ar), 34.9 (d, J = 6, CH_2), 28.3 (d, J = 10, CH_2).

Colorless oil, $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ 4.3. ^1H NMR (C_6D_6): δ 7.56-7.54 (m, 1H, Ar), 7.12-7.03 (m, 3H, Ar), 3.12-3.05 (m, 1H), 2.86-2.78 (m, 1H), 1.90-1.80 (m, 1H), 1.75-1.57 (m, 5H), 1.53-1.51 (m, 1H), 1.41-1.34 (m, 1H), 1.25-1.12 (m, 2H), 1.11-1.03 (m, 2H), 0.99-0.94 (m, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ 148.9 (Ar q), 142.1 (d, J = 13, Ar q), 131.3 (d, J = 24, Ar), 128.6 (Ar), 126.4 (d, J = 7, Ar), 124.8 (d, J = 2, Ar), 36.4 (d, J = 16), 35.0 (d, J = 6), 30.3 (d, J = 15), 29.6 (d, J = 13), 27.1 (d, J = 11), 26.9 (d, J = 10), 26.6 (d, J = 1), 21.7 (d, J = 11).

1-menthylbenzophospholane (**10c**)

Method I (Pd((*R,R*)-Me-DuPhos)(*trans*-stilbene) as a catalyst precursor): To a stirring slurry of Pd((*R,R*)-Me-DuPhos)(*trans*-stilbene) (7 mg, 0.01 mmol, 5 mol %) in toluene (0.5 mL) was added *o*-IC₆H₄CH₂CH₂PH(Men) (**9c**, 92.4 mg, 0.23 mmol) as a solution in toluene (1 mL) followed by NaOSiMe₃ (230 μ L, 0.23 mmol, 1 M solution in THF). The reaction mixture changed color from slightly yellow to dark-orange and precipitate was observed after 15 min. The reaction was monitored by ³¹P NMR spectroscopy. After the reaction was done (ca. 1 h) the product phosphine was isolated by column chromatography on a pipette-size column (silica, petroleum ether as an eluent) as a colorless oil (30 mg, 48%). The compound was formed as a 1:1 ratio of diastereomers. The spectra are reported for the mixture of diastereomers, unless otherwise stated.

Anal. Calcd. For C₁₈H₂₇P: C, 78.79; H, 9.92. Found: C, 78.66; H, 10.45. HRMS (EI) calcd for C₁₈H₂₇P: *m/z* 274.1850. Found: *m/z* 274.1851. ³¹P{¹H} NMR (toluene-*d*₈): δ 1.9, 0.6. ¹H NMR (C₆D₆): δ 7.58-7.53 (m, 1H), 7.45-7.40 (m, 1H), 7.12-7.03 (m, 6H), 3.22-3.13 (m, 1H), 3.13-3.03 (m, 1H), 2.93-2.79 (m, 3H), 2.79-2.69 (m, 1H), 2.06-1.94 (m, 1H), 1.90-1.73 (m, 3H), 1.71-1.42 (m, 7H), 1.42-1.26 (m, 2H), 1.21-1.09 (m, 1H), 1.09-1.00 (m, 2H), 1.00 (d, *J* = 6.5, 3H, Me, overlapped with previous signal), 0.97 (d, *J* = 7.5, 3H, Me), 0.94 (d, *J* = 6.5, 3H, Me), 0.94-0.88 (m, 2H, overlapped with the previous signal), 0.86 (d, *J* = 7, 3H, Me), 0.80-0.69 (m, 4H), 0.69 (d, *J* = 6.5, 3H, Me, overlapped with previous signal), 0.66 (d, *J* = 7, 3H, Me). ¹³C{¹H} NMR (C₆D₆): δ 149.6 (quat, Ar), 148.7 (d, *J* = 4, quat, Ar), 142.3 (d, *J* = 12, quat, Ar), 140.8 (d, *J* = 15, quat, Ar), 132.0 (d, *J* = 23, Ar), 130.1 (d, *J* = 23, Ar), 128.3 (Ar), 127.9 (Ar), 126.7 (d, *J* = 7, Ar), 126.2 (d, *J* = 7, Ar), 124.7 (d, *J* = 2, Ar), 124.6 (d, *J* = 2, Ar), 47.0 (d, *J* = 15, CH), 46.1 (d, *J* = 10, CH), 40.5 (d, *J* = 24, CH), 38.3 (d, *J* = 22, CH), 37.8 (CH₂), 36.7 (d, *J* = 4, CH₂), 35.4 (d, *J* = 5, CH₂), 35.3 (d, *J* = 5, CH₂), 35.3 (CH₂, overlapped with previous signal), 35.0 (CH₂), 33.7 (d, *J* = 2, CH), 33.6 (d, *J* = 4, CH), 28.7 (d, *J* = 17, CH), 28.3 (d, *J* = 20, CH), 25.7 (d, *J* = 7, CH₂), 25.4 (d, *J* = 8, CH₂), 23.2

(d, $J = 10$, CH_2), 22.7 (Me), 22.6 (Me), 21.8 (Me), 21.7 (Me), 18.4 (d, $J = 13$, CH_2), 15.6 (Me), 15.5 (d, $J = 2$, Me).

Method II ($\text{Pd}(\text{OAc})_2$ as a catalyst precursor, larger scale): To a stirring slurry of $\text{Pd}(\text{OAc})_2$ (6 mg, 0.03 mmol, 9 mol %) in toluene (0.5 mL) was added *o*- $\text{IC}_6\text{H}_4\text{CH}_2\text{CH}_2\text{PH}(\text{Men})$ (132 mg, 0.329 mmol) as a solution in toluene (1 mL) followed by NaOSiMe_3 (330 μL , 0.330 mmol, 1 M solution in THF). The reaction mixture changed color from slightly yellow to orange and precipitate was observed after 15 min. The reaction was monitored by ^{31}P NMR spectroscopy. After the reaction was done (ca. 20 h) the product phosphine was isolated by column chromatography on a 0.5 inch x 4 inch column (silica, petroleum ether as an eluent) as a colorless oil (44 mg, 49%). The compound was formed as a 1.2:1 ratio of diastereomers.

1-phenylbenzophospholane-borane (**11a**)

Phosphine-borane **11a** was prepared either by treatment of **10a** with $\text{BH}_3(\text{SMe}_2)$, as described in detail below for **11b**, or by direct cyclization of **8a** (see below).

Anal calc for $\text{C}_{14}\text{H}_{16}\text{BP}$: C, 74.38; H, 7.13. Found: C, 73.95; H, 7.05. HRMS m/z calcd. for $\text{C}_{14}\text{H}_{15}\text{BP}$ (M-H^+): m/z 225.1004. Found: m/z 225.1025. $^{31}\text{P}\{^1\text{H}\}$ NMR (CDCl_3): δ 38.0 (br m). ^1H NMR (CDCl_3): δ 7.63 (br t, $J = 8$, 1H, Ar), 7.57-7.47 (m, 4H, Ar), 7.45-7.35 (m, 4H, Ar), 3.40-3.32 (m, 2H), 2.56-2.33 (m, 2H), 1.62-0.60 (br q, 3H, BH_3). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3): δ 148.7 (d, $J = 15$, Ar q), 131.9 (d, $J = 8$, Ar), 131.9 (d, $J = 2$, Ar), 131.5 (d, $J = 45$, Ar q), 131.5 (d, $J = 2$, Ar), 131.1 (d, $J = 7$, Ar q), 130.5 (d, $J = 13$, Ar), 129.0 (d, $J = 10$, Ar), 128.2 (d, $J = 9$, Ar), 125.9 (d, $J = 8$, Ar), 31.9 (d, $J = 4$, CH_2), 26.2 (d, $J = 38$, CH_2).

1-cyclohexylbenzophospholane-borane (**11b**)

A solution of 1-cyclohexylbenzophospholane (**10b**, 30 mg, 1.37×10^{-4} mol) in THF (0.5 mL) was placed in an NMR tube, which was sealed with a rubber septum. 0.5 mL of $\text{BH}_3(\text{SMe}_2)$ (1×10^{-3} mol, 2.0 M solution in THF) was then added by syringe and the reaction was observed to go to completion by NMR spectroscopy. The solution was then poured into H_2O (10 mL) cautiously and extracted with Et_2O (3 x 10 mL). The combined Et_2O extracts were dried over MgSO_4 , filtered and pumped to dryness *in vacuo* to yield 46 mg of the crude product as a white sticky oil. The oil was dissolved in petroleum ether and passed through a silica plug column eluting with 20 mL of petroleum ether to remove impurities. The product was then removed from the silica by eluting with CH_2Cl_2 and the solution was dried *in vacuo* to yield 25 mg (78% yield) of a colorless oil.

Anal. Calcd for $\text{C}_{14}\text{H}_{22}\text{BP}$: C, 72.44; H, 9.55. Found: C, 72.08; H, 9.09. HRMS calcd for $\text{C}_{14}\text{H}_{22}\text{BPNa}$ ($\text{M}+\text{Na}$) $^+$: m/z 255.1450. Found: m/z 255.1438. $^{31}\text{P}\{^1\text{H}\}$ NMR (CDCl_3): δ 46.2 (br m). ^1H NMR (CDCl_3): δ 7.62-7.59 (m, 1H, Ar), 7.45-7.42 (m, 1H, Ar), 7.36-7.32 (m, 2H, Ar), 3.34-3.22 (m, 2H), 2.32-2.26 (m, 1H), 2.22-2.14 (m, 1H), 1.87 (m, 1H), 1.82-1.73 (br m, 4H), 1.69 (br m, 1H), 1.37-1.28 (br m, 1H), 1.25-1.13 (m, 4H), 0.98-0.42 (br q, $J = 93$, 3H, BH_3). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3): δ 148.3 (d, $J = 13$, Ar q), 131.4 (d, $J = 2$, Ar), 130.9 (d, $J = 54$, Ar q), 130.6 (d, $J = 12$, Ar), 127.5 (d, $J = 9$, Ar), 125.7 (d, $J = 7$, Ar), 35.4 (d, $J = 29$, CH), 32.4 (d, $J = 4$, CH_2), 26.9 (CH_2), 26.63 (d, $J = 21$, CH_2), 26.6 (d, $J = 2$, CH_2), 26.5 (d, $J = 1$, CH_2), 25.9 (d, $J = 1$, CH_2), 20.3 (d, $J = 37$, CH_2).

Phenylbenzophospholane sulfide (**12**)

Phosphine-borane **11a** (0.65 g, 2.9 mmol) was dissolved in diethyl ether (5 mL), diethylamine (3 mL, 29 mmol) was added, and the mixture was stirred overnight. ^{31}P NMR monitoring showed that deprotection was complete. Solid sulfur (140 mg, 4.35 mmol) was added to the phosphine solution, which turned darker orange, with some orange solid present. After stirring for 4 h, the ^{31}P NMR spectrum of the mixture showed that reaction was complete. The solvent was removed on a rotary

evaporator. The residue was dissolved in CH_2Cl_2 and the solution was filtered through Celite, then concentrated under vacuum and treated with ether to precipitate insoluble impurities. The solution was decanted and the remaining solid was washed with ether. The combined solutions were concentrated under vacuum to give 670 mg of crude product as an orange oil. Dry-column flash chromatography with CH_2Cl_2 gave 600 mg of white solid (85% yield).

This reaction was also successfully scaled up, with a simpler workup: **11a** (3.1 g, 13.7 mmol) was dissolved in diethyl ether (10 mL), diethylamine (14.2 mL, 137 mmol) was added, and the mixture was stirred for 5 h. Sulfur (673 mg, 20.6 mmol) was degassed in 10 mL of dry ether and transferred via cannula to the phosphine solution. The yellow solution was stirred for 12 h after which it was orange. It was filtered through Celite, and the plug was washed with dichloromethane. Water (100 mL) and dichloromethane (100 mL) were added and the organic layer was separated, washed with 100 mL of brine, and dried. The solvent was removed to yield 4.2 g of an orange oil. Column chromatography was used to isolate the product ($R_f = 0.28$, in 10% ethyl acetate/petroleum ether) as a yellow-orange oil. The oil was dissolved in dichloromethane and layered with ether and left at 0 °C overnight to give white crystals suitable for X-ray crystallography; yield was comparable to the smaller-scale reaction.

Anal. Calcd for $\text{C}_{14}\text{H}_{13}\text{PS}$: C, 68.83; H, 5.36. Found: C, 68.66; H, 5.11. HRMS m/z calcd. for $\text{C}_{14}\text{H}_{13}\text{PS}$: m/z 244.0476. Found: m/z 244.0484. $^{31}\text{P}\{^1\text{H}\}$ NMR (CDCl_3): δ 58.7. ^1H NMR (CDCl_3): δ 7.70-7.65 (m, 3H, Ar), 7.54-7.47 (m, 2H, Ar), 7.44-7.39 (m, 4H, Ar), 3.49-3.41 (m, 1H), 3.31-3.25 (m, 1H), 2.79-2.72 (m, 1H), 2.67-2.58 (m, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3): δ 147.0 (d, $J = 28$, Ar q), 134.6 (d, $J = 87$, Ar q), 133.7 (d, $J = 76$, Ar q), 132.5 (d, $J = 3$, Ar), 131.8 (d, $J = 3$, Ar), 130.9 (d, $J = 11$, Ar), 129.3 (d, $J = 11$, Ar), 128.8 (d, $J = 12$, Ar), 128.7 (d, $J = 11$, Ar), 126.2 (d, $J = 11$, Ar), 35.0 (d, $J = 57$), 30.2 (d, $J = 1$).

Pd-catalyzed Asymmetric Cyclization of Phosphine-Borane 8a (Table 2) A solution of **8a** (50 mg, 1.41×10^{-4} mol) in THF (0.5 mL) was added to a solution of the catalyst precursor (Pd(diphos*)(*trans*-stilbene), 5 mol %) in a few drops of THF. On addition of the phosphine-borane the solution turned from dark yellow to a pale orange color. Proton sponge (32 mg, $1.48 \times$

10^{-4} mol) in THF (1 mL) was then added to the solution; no further color change was observed. The mixture was transferred to an NMR tube, then monitored periodically by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy. Crystals of protonated proton sponge formed after about 30 min at room temperature. After several days a few gas bubbles were also observed in the solution. When the reaction was complete the mixture was filtered in air to remove the base•HI adduct and the residue was washed with additional THF (5 mL); the orange solution was then pumped to dryness. Column chromatography of the residue on silica eluting with ethyl acetate (10%)/hexanes yielded pure phosphine-borane **11a** ($R_f = 0.31$) as a colorless oily solid. Yields are reported in Table 2.

Effect of Added $\text{BH}_3(\text{SMe}_2)$ on $\text{Pd}((R,R)\text{-Me-DuPhos})$ -catalyzed Cyclization of **8a** A solution of **8a** (50 mg, 1.41×10^{-4} mol) in THF (0.5 mL) was charged to a NMR tube and $\text{BH}_3(\text{SMe}_2)$ (2.0 M in THF, 70 μL , 1.40×10^{-4} mol) was added via syringe. A solution of the catalyst precursor $\text{Pd}((R,R)\text{-Me-DuPhos})(\text{trans-stilbene})$ (4 mg, 0.007 mmol, 5 mol %) in a few drops of THF was added, turning the solution yellow. Then proton sponge (32 mg, 1.48×10^{-4} mol) dissolved in 0.5 mL of THF was added. The reaction mixture was then monitored periodically by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy. After 11 d, the solution remained yellow with no visible salt formation. Only a trace (ca. 5%, by ^{31}P NMR integration) of cyclized product was observed; the major P-containing compound was starting material **8a**. Note: in a control experiment, treatment of proton sponge (30 mg, 1.40×10^{-4} mol) with $\text{BH}_3(\text{SMe}_2)$ (70 μL of a 2.0 M solution in THF, 1.40×10^{-4} mol) in 1 mL of THF- d_8 led to minimal reaction (7% of a new proton-sponge containing species, by ^1H NMR integration) after 11 d.

Ee determination for **11a:** Under N_2 , phosphine-borane **11a** was treated with NEt_2H (10 equiv) or DABCO (1.2 equiv) as a solution in toluene (1 mL). The flask was sealed and heated at 50 $^\circ\text{C}$ (NEt_2H) or 80 $^\circ\text{C}$ (DABCO) for 2 h. All volatiles were then removed *in vacuo*. The residue was dissolved in petroleum ether (1 mL) in the glovebox and passed through a column of silica, eluting with petroleum ether (15 mL). The solvent was then removed in vacuo to yield phosphine **10a**, whose ee was determined by NMR spectroscopy according to the procedure described above.

Formation of Racemic 11a by Pd-catalyzed Cyclization of 8a. Under N₂, phosphine-borane **8a** (3.0 g, 8.5 mmol) was dissolved in 10 mL of THF. NaOSiMe₃ (8.9 mL of 1.0 M THF solution, 8.9 mmol), and the catalyst precursor Pd(dba)₂ (244 mg, 0.425 mmol, 5 mol %) were added and the solution was stirred for 24 h. The mixture was filtered through Celite. The volatile materials were removed under vacuum to give a brown oil. ³¹P NMR spectroscopy (toluene) showed the desired product, plus impurities at δ 26.2 (major impurity) and 17.2. The oil was loaded on a silica column with minimal CH₂Cl₂, and the product (1.1 g, 58% yield) was isolated as a colorless oil by dry column flash chromatography eluting with 10% ethyl acetate/hexane.

Synthesis of 13 by Deprotonation/methylation of Phosphine-Borane 11a

A solution of **11a** (110 mg, 0.49 mmol) in 20 mL of ether was cooled to -78 °C. *Sec*-butyl lithium (1.0 mL of 1.4 M solution in cyclohexane, 1.4 mmol, 3 equiv) was added dropwise and the solution turned yellow. The mixture was stirred for 1 h at -78 °C before MeI (0.3 mL, 4.9 mmol, 10 equiv) was added via a syringe. The solution was stirred at -78 °C for 1 h; the yellow color faded. The colorless solution was then warmed to room temperature. Water (30 mL) and ether (30 mL) were added, and the layers were separated. The aqueous layer was extracted with 2 x 15 mL of ether and combined with the first organic fraction. The ether phase was washed with 30 mL of brine, dried with magnesium sulfate and concentrated to give 100 mg of a colorless oil that contained two diastereomers in a 7:1 ratio (75% de) according to integration of the ¹H and ³¹P NMR spectra. Column chromatography eluting with 5% ethyl acetate/hexanes gave a colorless oil which was then recrystallized from petroleum ether at -10 °C to give diastereomerically pure white needles of the major diastereomer (75 mg, 64% yield).

Anal. Calcd for C₁₅H₁₈BP: C, 75.04; H, 7.56. Found: C, 75.12; H, 7.66. HRMS m/z calcd. for C₁₅H₁₇BP (M-H)⁺: 239.1161. Found: m/z 239.1155. ³¹P{¹H} NMR (CDCl₃): δ 45.6 (br apparent d, J = 60). ¹H NMR (CDCl₃): δ 7.58-7.52 (m, 3H, Ar), 7.50-7.47 (m, 2H, Ar), 7.43-7.35 (m, 4H,

Ar), 3.51-3.44 (m, $J_{\text{PH}} = 11$, $J_{\text{HH}} = 16$, 8, 1H), 3.01-2.95 (m, $J_{\text{PH}} = 7$, $J_{\text{HH}} = 16$, 6.3, 1H), 2.76-2.70 (m, $J_{\text{PH}} = 10.5$, $J_{\text{HH}} = 8$, 6.3, 7.2, 1H), 1.37 (dd, $J_{\text{PH}} = 17$, $J_{\text{HH}} = 7$, 3H), 1.4-0.4 (br m, 3H, BH_3). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3): δ 147.8 (d, $J = 15$, Ar q), 132.2 (d, $J = 10$, Ar), 131.8 (d, $J = 2$, Ar), 131.5 (d, $J = 2$, Ar), 131.0 (d, $J = 40$, Ar q), 130.7 (d, $J = 10$, Ar), 130.4 (Ar q), 129.0 (d, $J = 10$, Ar), 128.2 (d, $J = 9$, Ar), 125.9 (d, $J = 8$, Ar), 40.6 (CH_2), 34.9 (d, $J = 37$, CHMe), 15.3 (d, $J = 5$, Me).

The minor diastereomer was not obtained in pure form, but it could be characterized by NMR spectroscopy of the reaction mixture. $^{31}\text{P}\{^1\text{H}\}$ NMR (CDCl_3): δ 34.1-33.6 (br m).¹¹ Selected ^1H NMR (CDCl_3) (the aryl resonances of the two diastereomers overlapped): δ 3.43-3.34 (m, $J_{\text{PH}} = 11$, $J_{\text{HH}} = 16$, 8, 1H), 2.95-2.90 (m, $J_{\text{PH}} = 7$, $J_{\text{HH}} = 16$, 10, 1H), 2.67-2.61 (m, $J_{\text{PH}} = 10.5$, $J_{\text{HH}} = 10$, 8, 7.2, 1H), 1.37 (dd, $J_{\text{PH}} = 15$, $J_{\text{HH}} = 7.2$, 3H).

Stereochemistry of Deprotonation/Methylation The trans stereochemistry of the P-Ph and C-Me groups in **13** was determined by X-ray crystallography on diastereomerically pure **13**. These results were consistent with NMR studies. The ^1H NMR assignments in Figure S2 were made using DEPT, C-H correlation, H-H correlation, and 1D-NOE Experiments.

Figure S2. Selected ^1H NMR Data (CDCl_3) for the major diastereomer of **13**, and Important NOEs Observed

Assignment of the trans stereochemistry of the Me and P-Ph groups was consistent with these observations:

1. The Me group showed a strong NOE to H_A , but only a weak one to H_B , which suggests H_A and Me are cis.

2. H_C showed a strong NOE to H_B, but a weak one to H_A, suggesting that H_B and H_C are cis.
3. H_B, but not H_A, showed an NOE to H_D, suggesting a cis relationship between these hydrogens.

Table S2. Summary of 1D-NOESY Data for the major diastereomer of **13** (CDCl₃)^a

Signal	NOE (in order of intensity)
1.37 (Me)	2.73 (H _C) > 2.98 (H _A) > 7.58-7.52 (H _D)
2.73 (H _C)	1.37 (Me) > 7.58-7.52 (H _D) > 3.48 (H _B) > 2.98 (H _A)
2.98 (H _A)	3.48 (H _B) > 1.37 (Me) > 7.43-7.35 (H _E) > 2.73 (H _C)
3.48 (H _B)	2.98 (H _A) > 7.43-7.35 (H _E) > 2.73 (H _C) > 7.58-7.52 (H _D) > 1.37 (Me)

^a For clarity, not all of these NOEs are shown in Figure S2.

Figure S3. ORTEP diagram of the benzophospholane sulfide **12**.

Table 1. Crystal data and structure refinement for Ph-benzophospholane sulfide, glu227 (**12**).

Identification code	glu227	
Empirical formula	C ₁₄ H ₁₃ P S	
Formula weight	244.27	
Temperature	208(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2(1)	
Unit cell dimensions	a = 8.3368(16) Å	$\alpha = 90^\circ$
	b = 9.1981(17) Å	$\beta = 115.081(3)^\circ$
	c = 8.8279(17) Å	$\gamma = 90^\circ$
Volume	613.1(2) Å ³	
Z	2	
Density (calculated)	1.323 g/cm ³	
Absorption coefficient	0.362 mm ⁻¹	
F(000)	256	
Crystal size	0.41 x 0.37 x 0.21 mm ³	
Theta range for data collection	2.55 to 28.33°	
Index ranges	-11 ≤ h ≤ 10, -11 ≤ k ≤ 11, -11 ≤ l ≤ 11	
Reflections collected	4124	
Independent reflections	2289 [R(int) = 0.0266]	
Completeness to theta = 25.00°	98.7 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.943 and 0.887	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	2289 / 1 / 145	
Goodness-of-fit on F ²	1.163	
Final R indices [I > 2σ(I)]	R ₁ = 0.0339, wR ₂ = 0.0845	
R indices (all data)	R ₁ = 0.0344, wR ₂ = 0.0850	
Absolute structure parameter	0.01(7)	
Largest diff. peak and hole	0.578 and -0.183 e Å ⁻³	

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for Ph-benzophospholane sulfide, glu227. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
S(1)	8197(1)	9610(1)	2354(1)	37(1)
P(1)	6645(1)	7914(1)	1637(1)	26(1)
C(1)	7726(3)	6200(2)	1764(3)	27(1)
C(2)	8605(3)	5312(3)	3147(3)	32(1)
C(3)	9338(3)	4013(2)	2941(3)	37(1)
C(4)	9222(3)	3638(3)	1384(3)	40(1)
C(5)	8390(3)	4538(3)	7(3)	37(1)
C(6)	7608(2)	5823(2)	189(3)	30(1)
C(7)	6577(3)	6875(3)	-1187(3)	35(1)
C(8)	5363(3)	7742(2)	-608(2)	31(1)
C(9)	5196(2)	7719(2)	2682(2)	27(1)
C(10)	5397(3)	8630(3)	3994(3)	34(1)
C(11)	4249(3)	8495(3)	4759(3)	41(1)
C(12)	2926(3)	7466(3)	4209(3)	42(1)
C(13)	2736(3)	6555(3)	2915(3)	38(1)
C(14)	3862(3)	6670(2)	2142(3)	32(1)

Table 3. Bond lengths [Å] and angles [deg] for Ph-benzophospholane sulfide, glu227.

S(1)-P(1)	1.9531(7)	C(6)-C(7)	1.505(3)
P(1)-C(1)	1.796(2)	C(7)-C(8)	1.536(3)
P(1)-C(9)	1.813(2)	C(9)-C(10)	1.381(3)
P(1)-C(8)	1.816(2)	C(9)-C(14)	1.395(3)
C(1)-C(2)	1.392(3)	C(10)-C(11)	1.391(3)
C(1)-C(6)	1.395(3)	C(11)-C(12)	1.376(4)
C(2)-C(3)	1.388(3)	C(12)-C(13)	1.370(3)
C(3)-C(4)	1.380(4)	C(13)-C(14)	1.379(3)
C(4)-C(5)	1.389(3)		
C(5)-C(6)	1.392(3)		
C(1)-P(1)-C(9)	107.84(9)	C(10)-C(9)-C(14)	120.09(19)
C(1)-P(1)-C(8)	92.25(10)	C(10)-C(9)-P(1)	119.79(16)
C(9)-P(1)-C(8)	109.64(9)	C(14)-C(9)-P(1)	120.11(15)
C(1)-P(1)-S(1)	115.76(7)	C(9)-C(10)-C(11)	119.3(2)
C(9)-P(1)-S(1)	114.02(7)	C(12)-C(11)-C(10)	120.3(2)
C(8)-P(1)-S(1)	115.21(8)	C(13)-C(12)-C(11)	120.4(2)
C(2)-C(1)-C(6)	121.1(2)	C(12)-C(13)-C(14)	120.2(2)
C(2)-C(1)-P(1)	129.35(18)	C(13)-C(14)-C(9)	119.7(2)
C(6)-C(1)-P(1)	109.52(15)		
C(3)-C(2)-C(1)	119.2(2)		
C(4)-C(3)-C(2)	119.8(2)		
C(3)-C(4)-C(5)	121.4(2)		
C(4)-C(5)-C(6)	119.3(2)		
C(5)-C(6)-C(1)	119.18(19)		
C(5)-C(6)-C(7)	126.0(2)		
C(1)-C(6)-C(7)	114.80(18)		
C(6)-C(7)-C(8)	107.24(17)		
C(7)-C(8)-P(1)	104.04(13)		

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for Ph-benzophospholane sulfide, glu227. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
S(1)	38(1)	33(1)	39(1)	-3(1)	15(1)	-10(1)
P(1)	27(1)	26(1)	25(1)	0(1)	11(1)	-1(1)
C(1)	23(1)	27(1)	33(1)	0(1)	13(1)	0(1)
C(2)	27(1)	38(1)	31(1)	3(1)	11(1)	-2(1)
C(3)	26(1)	34(1)	47(1)	9(1)	11(1)	2(1)
C(4)	29(1)	32(1)	59(2)	-7(1)	19(1)	1(1)
C(5)	31(1)	43(1)	39(1)	-13(1)	16(1)	-6(1)
C(6)	27(1)	32(1)	34(1)	-6(1)	15(1)	-6(1)
C(7)	36(1)	45(1)	27(1)	-1(1)	16(1)	0(1)
C(8)	33(1)	34(1)	25(1)	2(1)	10(1)	-1(1)
C(9)	25(1)	28(1)	27(1)	2(1)	10(1)	3(1)
C(10)	35(1)	37(1)	29(1)	-1(1)	13(1)	2(1)
C(11)	49(1)	50(1)	29(1)	-2(1)	20(1)	3(1)
C(12)	42(1)	55(1)	39(1)	8(1)	26(1)	6(1)
C(13)	34(1)	35(1)	48(1)	6(1)	19(1)	-2(1)
C(14)	37(1)	28(1)	34(1)	0(1)	18(1)	3(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for Ph-benzophospholane sulfide, glu227.

	x	y	z	U(eq)
H(2A)	8699	5584	4196	39
H(3A)	9905	3398	3849	45
H(4A)	9710	2764	1254	48
H(5A)	8356	4284	-1026	45
H(7A)	5878	6354	-2211	42
H(7B)	7377	7525	-1396	42
H(8A)	4265	7224	-864	38
H(8B)	5092	8690	-1138	38
H(10A)	6290	9326	4363	41
H(11A)	4376	9102	5645	49
H(12A)	2156	7388	4719	51
H(13A)	1845	5858	2558	46
H(14A)	3732	6051	1265	38

Figure S4. ORTEP diagram of methylated benzophospholane-borane **13**.

Table 1. Crystal data and structure refinement for methylated Ph-benzophospholane-borane, glu244 (**13**).

Identification code	glu244	
Empirical formula	C ₁₅ H ₁₈ B P	
Formula weight	240.07	
Temperature	208(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	P2(1)2(1)2(1)	
Unit cell dimensions	a = 9.4071(12) Å	$\alpha = 90^\circ$
	b = 11.3293(16) Å	$\beta = 90^\circ$
	c = 12.9898(16) Å	$\gamma = 90^\circ$
Volume	1384.4(3) Å ³	
Z	4	
Density (calculated)	1.152 g/cm ³	
Absorption coefficient	0.173 mm ⁻¹	
F(000)	512	
Crystal size	0.30 x 0.12 x 0.08 mm ³	
Theta range for data collection	2.67 to 28.22°	
Index ranges	-1 ≤ h ≤ 12, -13 ≤ k ≤ 12, -7 ≤ l ≤ 15	
Reflections collected	3698	
Independent reflections	2448 [R(int) = 0.0173]	
Completeness to theta = 25.00°	98.4 %	
Absorption correction	multi-scan	
Max. and min. transmission	0.9863 and 0.9498	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	2448 / 0 / 166	
Goodness-of-fit on F ²	1.030	
Final R indices [I > 2σ(I)]	R1 = 0.0424, wR2 = 0.1092	
R indices (all data)	R1 = 0.0477, wR2 = 0.1129	
Absolute structure parameter	-0.19(14)	
Largest diff. peak and hole	0.243 and -0.219 e Å ⁻³	

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for methylated Ph-benzophospholane-borane, glu244. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
P(1)	7535(1)	2304(1)	1918(1)	36(1)
C(1)	7851(2)	3446(3)	1007(2)	35(1)
C(2)	8562(2)	4522(3)	1154(2)	42(1)
C(3)	8769(2)	5259(3)	319(3)	51(1)
C(4)	8270(3)	4943(3)	-636(3)	52(1)
C(7)	6610(2)	1949(3)	0(2)	44(1)
C(8)	7173(2)	1211(3)	913(2)	39(1)
C(10)	5477(3)	1927(3)	3383(2)	53(1)
C(11)	4160(3)	2058(4)	3847(3)	66(1)
C(12)	3204(3)	2851(4)	3457(3)	59(1)
C(13)	3540(3)	3519(4)	2618(2)	54(1)
C(14)	4858(3)	3391(3)	2137(2)	44(1)
C(15)	8514(3)	522(3)	625(2)	54(1)
B(1)	9009(3)	1989(5)	2897(3)	57(1)
C(9)	5825(2)	2578(3)	2524(2)	38(1)
C(5)	7564(3)	3885(3)	-794(2)	46(1)
C(6)	7348(2)	3135(3)	37(2)	38(1)

Table 3. Bond lengths [Å] and angles [deg] for methylated Ph-benzophospholane-borane, glu244.

P(1)-C(1)	1.779(3)	C(8)-P(1)-B(1)	118.84(18)
P(1)-C(9)	1.817(2)	C(6)-C(1)-C(2)	120.3(3)
P(1)-C(8)	1.831(3)	C(6)-C(1)-P(1)	111.1(2)
P(1)-B(1)	1.916(3)	C(2)-C(1)-P(1)	128.5(2)
C(1)-C(6)	1.391(3)	C(3)-C(2)-C(1)	119.0(3)
C(1)-C(2)	1.403(4)	C(4)-C(3)-C(2)	120.2(3)
C(2)-C(3)	1.382(4)	C(3)-C(4)-C(5)	121.5(3)
C(3)-C(4)	1.374(5)	C(6)-C(7)-C(8)	107.4(2)
C(4)-C(5)	1.385(4)	C(15)-C(8)-C(7)	111.8(2)
C(7)-C(6)	1.513(4)	C(15)-C(8)-P(1)	111.47(19)
C(7)-C(8)	1.545(4)	C(7)-C(8)-P(1)	104.2(2)
C(8)-C(15)	1.530(4)	C(9)-C(10)-C(11)	120.6(3)
C(10)-C(9)	1.377(4)	C(12)-C(11)-C(10)	119.7(3)
C(10)-C(11)	1.386(4)	C(13)-C(12)-C(11)	120.5(3)
C(11)-C(12)	1.368(6)	C(12)-C(13)-C(14)	120.4(3)
C(12)-C(13)	1.364(5)	C(9)-C(14)-C(13)	119.2(3)
C(13)-C(14)	1.396(4)	C(10)-C(9)-C(14)	119.5(2)
C(14)-C(9)	1.389(4)	C(10)-C(9)-P(1)	118.1(2)
C(5)-C(6)	1.389(4)	C(14)-C(9)-P(1)	122.4(2)
		C(4)-C(5)-C(6)	119.0(3)
C(1)-P(1)-C(9)	108.18(13)	C(5)-C(6)-C(1)	119.9(3)
C(1)-P(1)-C(8)	92.82(14)	C(5)-C(6)-C(7)	125.8(2)
C(9)-P(1)-C(8)	105.10(11)	C(1)-C(6)-C(7)	114.2(3)
C(1)-P(1)-B(1)	117.14(16)		
C(9)-P(1)-B(1)	112.57(15)		

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for methylated Ph-benzophospholane-borane, glu244. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
P(1)	32(1)	44(1)	32(1)	0(1)	0(1)	1(1)
C(1)	29(1)	40(2)	37(2)	-4(2)	2(1)	0(1)
C(2)	36(1)	47(2)	45(2)	-7(2)	1(1)	-3(1)
C(3)	42(1)	44(2)	67(2)	2(2)	8(1)	-4(1)
C(4)	44(2)	54(3)	57(2)	17(2)	8(1)	4(1)
C(7)	41(1)	52(2)	39(2)	-3(2)	-3(1)	-8(1)
C(8)	40(1)	38(2)	40(2)	-1(2)	4(1)	-4(1)
C(10)	54(2)	56(3)	50(2)	11(2)	10(1)	3(1)
C(11)	67(2)	66(3)	65(3)	7(2)	27(2)	-7(2)
C(12)	44(2)	67(3)	67(2)	-16(2)	18(1)	-3(1)
C(13)	41(1)	67(3)	52(2)	-18(2)	-2(1)	8(1)
C(14)	43(1)	57(2)	31(2)	-5(2)	0(1)	3(1)
C(15)	52(2)	54(3)	54(2)	-5(2)	12(1)	4(1)
B(1)	45(2)	76(3)	49(2)	2(2)	-14(1)	11(2)
C(9)	34(1)	42(2)	36(2)	-5(1)	0(1)	-3(1)
C(5)	42(1)	59(2)	38(2)	9(1)	1(1)	3(1)
C(6)	31(1)	43(2)	39(2)	0(1)	0(1)	0(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for methylated Ph-benzophospholane-borane, glu244.

	x	y	z	U(eq)
H(2A)	8893	4739	1810	51
H(3A)	9252	5978	405	61
H(4A)	8411	5456	-1196	62
H(7A)	6817	1550	-653	52
H(7B)	5579	2052	57	52
H(8A)	6426	658	1150	47
H(10A)	6138	1389	3656	64
H(11A)	3924	1603	4428	79
H(12A)	2309	2936	3770	71
H(13A)	2881	4069	2363	64
H(14A)	5088	3849	1557	52
H(15A)	8287	-49	93	80
H(15B)	8874	115	1227	80
H(15C)	9231	1064	370	80
H(5A)	7236	3678	-1453	55
H(1C)	9050(30)	2750(40)	3320(30)	71(12)
H(1B)	8640(30)	1100(40)	3350(30)	78(12)
H(1A)	9970(50)	1900(50)	2560(40)	106(13)

9b•HBF₄

9c

9c

9c

10a

10a

11b

12

References

1. Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, *15*, 1518-1520.
2. Brunel, J. M.; Faure, B.; Maffei, M. *Coord. Chem. Rev.* **1998**, *178-180*, 665-698.
3. Ripa, L.; Hallberg, A. *J. Org. Chem.* **1996**, *61*, 7147-7155.
4. Bourumeau, K.; Gaumont, A.-C.; Denis, J.-M. *J. Organomet. Chem.* **1997**, *529*, 205-213.
5. Blank, N. F.; McBroom, K. C.; Glueck, D. S.; Kassel, W. S.; Rheingold, A. L. *Organometallics* **2006**, *25*, 1742-1748.
6. Bruner, T. J.; Blank, N. F.; Moncarz, J. R.; Scriban, C.; Anderson, B. J.; Glueck, D. S.; Zakharov, L. N.; Golen, J. A.; Sommer, R. D.; Incarvito, C. D.; Rheingold, A. L. *Organometallics* **2005**, *24*, 2730-2746.
7. Rettig, M. F.; Maitlis, P. M. *Inorg. Synth.* **1990**, *28*, 110-113.
8. Roberts, N. K.; Wild, S. B. *J. Am. Chem. Soc.* **1979**, *101*, 6254-6260.
9. Sayalero, S.; Pericas, M. A. *Synlett* **2006**, 2585-2588.
10. The ligands were commercially available, except for the ferrocenyl-phospholane 5-Fc (see ref. 16 of the text for its synthesis, and Boulton, L. T.; Lennon, I. C.; McCague, R. *Org. Biomol. Chem.* **2003**, *1*, 1094-1096 for the name.)
11. The large ^{31}P NMR chemical shift difference between the diastereomers of **13** is consistent with literature precedent. See: Quin, L. D., *The Heterocyclic Chemistry of Phosphorus: Systems Based on the Phosphorus-Carbon Bond*. Wiley-Interscience: New York, 1981, Chapter 5.