Enantioselective Total Synthesis of Erogorgiaene: Applications of Asymmetric Cu-Catalyzed Conjugate Additions of Alkylzincs to Acyclic Enones

Richard R. Cesati III, Judith de Armas and Amir H. Hoveyda*

*Department of Chemistry, Merkert Chemistry Center, Boston College
Chestnut Hill, Massachusetts 02467*

Supporting Information

General. Infrared (IR) spectra were recorded on a Perkin-Elmer 781 spectrophotometer, $\nu_{\max}$ in cm^{-1} . Bands are characterized as broad (br), strong (s), medium (m), and weak (w). ^1H NMR spectra were recorded on a Varian Unity INOVA 400 (400 MHz) spectrometer. Chemical shifts are reported in ppm from tetramethylsilane with the residual solvent resonance resulting from incomplete deuteration as the internal standard (CDCl_3 ; δ 7.26 ppm). Data are reported as follows: chemical shift, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), and coupling constants. ^{13}C NMR spectra were recorded on a Varian Unity INOVA 400 (100 MHz) with complete proton decoupling. Chemical shifts are reported in ppm from tetramethylsilane with the solvent as the internal reference (CDCl_3 ; δ 77.0 ppm). Enantiomer and diastereomer ratios were determined by chiral GLC analysis (Alltech Associates Chiraldex GTA column (30 m x 0.25 mm), Chiraldex BPH column (30 m x 0.25 mm) or Supelco Betadex 120 column (30 m x 0.25 mm)) or chiral HPLC analysis (Chiral Technologies Chiralcel OD column (0.46 cm x 25 cm) or Chiralcel OJ column (0.46 cm x 25 cm)) in comparison with authentic racemic materials. Microanalyses were performed by Robertson Microlit Laboratories (Madison, NJ). High-resolution mass spectrometry was performed by the University of Illinois Mass Spectrometry Laboratories (Urbana, IL) or at Boston College Mass Spectrometry Facility on a Micromass LCT ESI-MS (positive mode). Optical rotation values are recorded on a Rudolf Research Analytical Autopol IV polarimeter.

Unless otherwise stated, all reactions were conducted in oven- (135 °C) and flame-dried glassware under an inert atmosphere of dry nitrogen. Indicated temperatures refer to those of the reaction bath, while ambient laboratory temperature is noted as 22 °C. Solvents are purified under a positive pressure of dry argon by a modified Advanced ChemTech purification system – toluene and benzene are purified through Cu and alumina columns, dichloromethane, THF and Et_2O are purified through two alumina columns – with the following exceptions noted below. Dimethylformamide was distilled from CaH_2 immediately prior to use. Methanol was distilled from Mg turnings activated with iodine.

The following is a description of reagents, which required prior preparation or purification.¹ 3-Bromo-4-iodotoluene (**4**) was obtained from Matrix Scientific and purified by passage through a column of silica gel and neutral alumina using pentane as eluent. Copper (I) triflate (benzene complex, 2:1) was prepared according to a published procedure.² Peptidic Schiff-base ligands were prepared according to published procedures.³ 2,6-Di-*tert*-butyl-4-methylphenol (BHT) and PPh₃ were recrystallized from hexanes. Butyllithium (Strem; solution in hexanes) was titrated using *N*-pivaloyl-*o*-toluidine according to the method of Suffert.⁴ 2,2,6,6-Tetramethylpiperidine (TMP), *i*-Pr₂NH, pyridine, Et₃N, chlorotrimethylsilane (TMSCl), and methyl vinyl ketone were distilled from CaH₂ immediately prior to use. Methyllithium (Aldrich; solution in Et₂O) was titrated with endpoint indication by *N*-phenylnaphthylamine. Anhydrous LiCl was obtained through recrystallization from H₂O and drying at 120 °C under reduced pressure (0.1 mm Hg). Tributylstannane was distilled under reduced pressure to remove higher boiling impurities. Anhydrous NH₃ was condensed into a reaction mixture following passage through a column of NaOH pellets. Dess-Martin periodinane was prepared through a standard procedure.⁵ Oxygen was passed through P₂O₅ and Drierite as it was introduced into a reaction mixture or an ozone generator. Imidazole was recrystallized from benzene. All Grignard reagents were freshly prepared by using flame-dried Mg turnings purchased from Strem and titrated immediately prior to use with endpoint indication by *N*-phenylnaphthylamine. Cuprous iodide was recrystallized from a saturated aqueous solution of sodium iodide. Sodium hydride (Strem) was washed with anhydrous hexanes and stored under a nitrogen atmosphere. Biscyclopentadienylzirconium dichloride (Strem) was recrystallized from anhydrous toluene. All other reagents were used as obtained from Aldrich, Acros, Fisher or Strem Chemicals.

(E)-4-(2-Bromo-4-methylphenyl)but-3-en-2-one (5): Activated 4Å molecular sieves (5.50 g) were massed into a 50 mL round bottom flask and suspended in DMF (14.0 mL). To this suspension were added NaHCO₃ (2.89 g, 34.4 mmol) and *n*-Bu₄NCl (3.83 g, 13.8 mmol). After stirring 15 min at 22 °C, the reaction mixture was treated with 3-bromo-4-iodotoluene (2.00 mL, 13.8 mmol) and methyl vinyl ketone (2.29 mL, 27.5 mmol). Following an additional 15 min stirring at 22 °C, 155 mg Pd(OAc)₂ (0.688 mmol; 5 mol %) was added in one portion and the reaction mixture immediately heated to 60 °C and maintained for 4 h. The resulting brown mixture was cooled to 22 °C and diluted with H₂O and Et₂O (20 mL each) and transferred to a separatory funnel, the layers separated and the aqueous layer washed with Et₂O (3 x 25 mL). The combined organic layers were

¹ A general text covering the techniques described herein is available: Armarego, W. L. F.; Perrin, D. D. *Purification of Laboratory Chemicals*, 4th ed.; Butterworth-Heinemann: Oxford, U. K., 1998.

² (a) Salomon, R. G.; Kochi, J. K. *J. Am. Chem. Soc.* **1973**, *95*, 1889. (b) Salomon, R. G.; Kochi, J. K. *J. Am. Chem. Soc.* **1973**, *95*, 3300.

³ (a) Mizutani, H.; Degrado, S. J.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2002**, *124*, 779. (b) Luchaco-Cullis, C. A.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2002**, *124*, 8192.

⁴ Suffert, J. J. *Org. Chem.* **1989**, *54*, 509.

⁵ (a) Dess, D. B.; Martin, J. C. *J. Am. Chem. Soc.* **1991**, *113*, 7277. (b) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155.

dried over MgSO_4 , filtered and concentrated in vacuo to afford a yellow oil. Purification by chromatography on silica (4 : 1 hexanes/ Et_2O ; R_f = 0.3 in 9 : 1 hexanes/ EtOAc) afforded a white crystalline solid (2.94 g, 12.3 mmol; 89.3% yield). Mp 32.0–33.0 °C. IR (NaCl): 3050 (w), 3031 (w), 2993 (w), 2911 (w), 1671 (s), 1608 (s), 1485 (s), 1358 (s), 1312 (m), 1284 (m), 1256 (w), 1217 (m), 1175 (m), 1041 (m), 974 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.87 (1H, d, J = 16.3 Hz), 7.52 (1H, d, J = 8.1 Hz), 7.46 (1H, s), 7.14 (1H, d, J = 8.1 Hz), 6.59 (1H, d, J = 16.3 Hz), 2.41 (3H, s), 2.35 (3H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 198.2, 142.2, 141.7, 133.8, 131.3, 128.8, 128.7, 127.3, 125.5, 27.0, 20.9. HRMS Calcd. for $\text{C}_{11}\text{H}_{11}\text{BrO}$: 237.9993. Found: 237.9993. Anal. Calcd. for $\text{C}_{11}\text{H}_{11}\text{BrO}$: C, 55.25; H, 4.64. Found: C, 55.18; H, 4.63.

(S)-4-(2-Bromo-4-methylphenyl)pentan-2-one (6): A 50 mL round bottom flask was charged with $(\text{CuOTf})_2 \cdot \text{PhH}$ (21.0 mg, 0.0420 mmol; 1 mol %) and **2** (68.0 mg, 0.100 mmol; 2.4 mol %) under an atmosphere of nitrogen in a glove box. The flask was sealed with a septum and Teflon tape prior to removal from the glove box. The solid mixture was then dissolved in toluene (16.0 mL), stirred 5 min at 22 °C and treated with Me_2Zn (12.5 mmol; 6.27 mL of a 2 M solution in toluene). Following the addition, the reaction vessel was cooled to –78 °C in a dry ice/acetone bath. To the reaction mixture was then transferred a solution of **5** (1.00 g, 4.18 mmol) in toluene (5.0 mL) via cannula, with an additional 0.60 mL toluene used to quantitate the transfer. The reaction mixture was then warmed to –15 °C and maintained for 48 h. The yellow solution was then cooled to –78 °C and treated with a saturated solution of NH_4Cl (5 mL). The cooling bath was then removed and the excess Me_2Zn consumed as the solution slowly warmed to 22 °C. The now orange reaction mixture was diluted with H_2O and Et_2O (25 mL each) and transferred to a separatory funnel. The layers were separated and the aqueous layer was washed with Et_2O (3 x 25 mL). The combined organic layers were dried over MgSO_4 , filtered and concentrated in vacuo to afford a yellow oil. Purification by chromatography on silica (4 : 1 hexanes/ Et_2O ; R_f = 0.3 in 9 : 1 hexanes/ EtOAc) afforded a colorless oil (1.00 g, 3.92 mmol; 93.7% yield). IR (NaCl): 2968 (s), 2924 (s), 2873 (m), 1715 (s), 1608 (m), 1495 (s), 1363 (m), 1168 (m), 1042 (m), 859 (m), 822 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.38 (1H, s), 7.12–7.07 (2H, m), 3.78–3.69 (1H, m), 2.78 (1H, dd, J = 16.1, 5.5 Hz), 2.57 (1H, dd, J = 16.0, 8.7 Hz), 2.29 (3H, s), 2.12 (3H, s), 1.24 (3H, d, J = 7.0 Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 207.3, 141.5, 137.7, 133.4, 128.5, 126.8, 123.9, 50.9, 33.8, 29.9, 20.5(2). HRMS Calcd. for $\text{C}_{12}\text{H}_{15}\text{BrO}$: 257.0363. Found: 257.0364. Anal. Calcd. for $\text{C}_{12}\text{H}_{15}\text{BrO}$: C, 56.49; H, 5.93. Found: C, 56.74; H, 6.01. $[\alpha]_D^{20}$ +8.2° (c 0.61, CHCl_3) for a >98% ee sample.

Note added in proof: The adduct was determined to be *S* stereochemistry through comparison of the optical rotation obtained following reductive removal of bromine (Bu_3SnH , AIBN, C_6H_6 , 80 °C, 2 h) to natural (+)-curcumone. Found: $[\alpha]_D^{20}$ +43.2° (c 0.16, CHCl_3) for a 97% ee sample. Lit.⁶ +40.2° (c 2.35, CHCl_3) for a 98% ee sample.

⁶ Asaoka, M.; Shima, K.; Fujii, N.; Takei, H. *Tetrahedron* **1988**, *44*, 4757-4766.

4-[4-Methyl-2-(trimethylsilanylethynyl)phenyl]pentan-2-one

(7): Pd(PPh₃)₄ (404 mg, 0.349 mmol; 5 mol %) and BHT (15.4 mg, 0.070 mmol; 1 mol %) were massed into a 50 mL round bottom flask under a nitrogen atmosphere in a glove box. The flask was sealed with a septum and teflon tape, removed from the glove box and the contents dissolved in toluene (15.0 mL). A solution of **6** (1.78 g, 6.99 mmol) in toluene (10.0 mL) was transferred to the yellow reaction mixture via cannula, with an additional 5.0 mL toluene used to quantitate the transfer. 1-Tributylstannyl-2-trimethylsilylacetylene (3.25 g, 8.39 mmol) as a solution in toluene (5.0 mL) was added and the mixture heated to 80 °C and maintained 12 h. After cooling to 22 °C, all volatiles were removed in vacuo to afford a brown oil. Purification by chromatography on silica (4 : 1 hexanes/Et₂O; *R_f* = 0.3 in 4 : 1 hexanes/Et₂O) afforded a colorless oil (1.83 g, 6.72 mmol; 96.3% yield). *Note: Typically, two columns are required to obtain material free of tin-containing byproducts.* IR (NaCl): 2961 (m), 2149 (m), 1718 (s), 1249 (m), 854 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.28 (1H, m), 7.09 (2H, m), 3.79 (1H, dqd, *J* = 9.1, 7.1, 5.3 Hz), 2.83 (1H, dd, *J* = 15.7, 5.5 Hz), 2.59 (1H, dd, *J* = 15.7, 9.0 Hz), 2.28 (3H, s), 2.12 (3H, s), 1.27 (3H, d, *J* = 7.0 Hz), 0.25 (9H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 207.7, 145.0, 135.5, 133.4, 129.8, 125.6, 121.7, 103.7, 98.4, 51.4, 32.9, 29.8, 20.6, 20.2, -0.07(3). HRMS Calcd. for C₁₇H₂₄OSiNa (M+Na⁺): 295.1494. Found: 295.1495. [α]_D²⁰ +12.5° (*c* 0.35, CHCl₃).

4-Methyl-1-[1-methyl-3-trimethylsilyloxybut-3-enyl]-2-(trimethylsilanylethynyl)benzene (8)

(8): BuLi (2.98 mmol; 2.05 mL of a 1.45 M solution in hexanes) was added dropwise over 5 min to a solution of TMP (503 μL, 2.98 mmol) in THF (3.0 mL) at 0 °C. After stirring an additional 5 min at 0 °C, the yellow LiTMP solution was cooled to -78 °C. To this solution was transferred a solution of **7** (6.5 x 10² mg, 2.39 mmol) and TMSCl (403 μL, 3.17 mmol) in THF (12.0 mL) dropwise over 30 min. An additional 1.0 mL of THF was used to quantitate the transfer. Following the addition, the solution was allowed to stir for 10 min at -78 °C at which time TLC analysis indicated complete consumption of the starting material (product *R_f* = 0.8 in 9 : 1 hexanes/Et₂O). The solution was diluted with wet Et₂O (2 mL) and warmed to 22 °C. All volatiles were removed in vacuo to afford a white residue from which the product can be distilled directly under reduced pressure. The colorless oil thus obtained (8.00 x 10² mg, 2.32 mmol; 97.3%) was shown to be a 9 : 1 mixture of enol silyl ethers by ¹H NMR. IR (NaCl): 2960 (s), 2149 (s), 1653 (m), 1621 (m), 1251 (s), 1012 (s), 940 (m), 844 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.27 (1H, m), 7.08 (2H, m), 4.06 (1H, d, *J* = 0.5 Hz), 4.04 (1H, d, *J* = 0.7 Hz), 3.53 (1H, dqd, *J* = 8.6, 7.0, 6.8 Hz), 2.46 (1H, dd, *J* = 14.1, 6.4 Hz), 2.27 (3H, s), 2.25 (1H, dd, *J* = 14.1, 8.6 Hz), 1.22 (3H, d, *J* = 7.0 Hz), 0.25 (9H, s), 0.17 (9H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 157.9, 146.3, 134.9, 133.3, 129.6, 125.7, 121.9, 104.3, 97.5, 91.0, 43.3, 34.2, 20.8, 20.6, 0.07(3), 0.03(3). HRMS Calcd. for C₂₀H₃₂OSi₂Na (M+Na⁺): 367.1889. Found: 367.1883. [α]_D²⁰ -30.8° (*c* 0.45, CHCl₃).

Trifluoromethanesulfonic acid 1-{2-[4-methyl-2-(trimethylsilyl)ethynyl]phenyl}propyl}-vinyl ester:

MeLi (3.27 mmol; 2.11 mL of a 1.55 M solution in Et₂O) in a 50 mL round bottom flask was diluted with THF (2.0 mL) and immediately cooled to -78 °C. To this white suspension was transferred a solution of **8** (9.40×10^2 mg, 2.73 mmol) in THF (9.0 mL) dropwise over 10 min. An additional 0.5 mL THF was used to quantitate the transfer. Following the addition, the solution was warmed to 4 °C in a cold room and maintained for 3 h. The homogeneous solution thus obtained was cooled to -78 °C and treated with a solution of Tf₂NPh (1.17 g, 3.27 mmol) in THF (9.0 mL) in a dropwise fashion over 10 min. The resulting mixture was returned to the cold room (4 °C) and warmed slowly from -78 °C over 14 h as the dry ice/acetone bath evaporated. The resulting solution was diluted with wet Et₂O (5 mL) and warmed to 22 °C. All volatiles were removed in vacuo, and the residue purified by rapid chromatography on silica (19 : 1 hexanes/Et₂O; *R_f* = 0.7 in 9 : 1 hexanes/Et₂O) to afford a colorless oil (762 mg, 1.88 mmol; 69.1%) IR (NaCl): 2963 (s), 2150 (s), 1669 (m), 1418 (s), 1250 (m), 1213 (m), 1141 (m), 953 (w), 910 (w), 845 (w). ¹H NMR (CDCl₃, 400 MHz): δ 7.30 (1H, m), 7.10 (1H, dd, *J* = 8.0, 1.6 Hz), 7.07 (1H, d, *J* = 7.9 Hz), 5.07 (1H, d, *J* = 3.7 Hz), 4.88 (1H, ddd, *J* = 3.7, 0.9, 0.9 Hz), 3.59 (1H, ddq, *J* = 7.3, 7.3, 7.1 Hz), 2.80 (1H, ddd, *J* = 15.7, 7.5, 0.7 Hz), 2.58 (1H, ddd, *J* = 15.7, 7.4, 0.7 Hz), 2.29 (3H, s), 1.32 (3H, d, *J* = 7.1 Hz), 0.26 (9H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 155.4, 143.7, 135.8, 133.6, 129.9, 125.5, 122.0, 118.5 (q, *J* = 320 Hz), 105.6, 103.4, 98.5, 41.0, 34.3, 20.6 (2), -0.15(3). HRMS Calcd. for C₁₈H₂₃F₃O₃SSi: 404.1089. Found: 404.1082. [α]_D²⁰ -9.2° (c 0.49, CHCl₃).

Trimethyl[5-methyl-2-(1-methylbut-3-enyl)phenylethynyl]silane:

Pd(PPh₃)₄ (88.9 mg, 0.0769 mmol; 5 mol %), BHT (16.9 mg, 0.0769 mmol; 5 mol %) and LiCl (196 mg, 4.61 mmol) were massed into a 50 mL round bottom flask under a nitrogen atmosphere in a glove box. The flask was sealed with a septum and teflon tape, removed from the glove box and the contents dissolved in THF (15.0 mL). A solution of enol triflate (622 mg, 1.54 mmol) in THF (10.0 mL) was transferred to the yellow reaction mixture via cannula, with an additional 5.0 mL THF used to quantitate the transfer. Bu₃SnH (496 μL, 1.85 mmol) was added in one portion and the reaction flask immediately warmed to 60 °C and maintained for 30 min at which time TLC analysis indicated complete consumption of starting material (product *R_f* = 0.4 in hexanes). The solution was cooled to 22 °C and all volatiles removed in vacuo. Purification by chromatography on silica (pentane) afforded a colorless oil (325 mg, 1.27 mmol; 82.3%). IR (NaCl): 2961 (s), 2925 (w), 2150 (s), 1250 (s), 853 (s). ¹H NMR (CDCl₃, 400 MHz): δ 7.28 (1H, m), 7.10 (2H, m), 5.76 (1H, dddd, *J* = 17.0, 9.9, 7.2, 6.5 Hz), 5.01 (1H, ddt, *J* = 17.0, 2.2, 1.5 Hz), 4.97 (1H, ddt, *J* = 10.1, 2.2, 1.1 Hz), 3.36 (1H, dqd, *J* = 7.7, 7.0, 6.8 Hz), 2.44 (1H, dddd, *J* = 12.8, 7.8, 6.4, 1.5, 1.3 Hz), 2.28 (3H, s), 2.24 (1H, m), 1.24 (3H, d, *J* = 7.0 Hz), 0.25 (9H, s). ¹³C NMR (CDCl₃, 100 MHz): δ 146.2, 137.3, 135.0, 133.1, 129.6, 125.5, 122.0, 115.8, 104.2, 97.8, 41.8, 36.3,

20.6, 20.0, 0.00(3). HRMS Calcd. for $C_{17}H_{24}Si$: 256.1647. Found: 256.1647. $[\alpha]_D^{20}$ -60.0° (c 0.42, $CHCl_3$).

2-Ethynyl-4-methyl-1-(1-methylbut-3-enyl)benzene (9):

A solution of protected enyne (311 mg, 1.21 mmol) in 4 : 1 THF/MeOH (6.0 mL) at 22 °C was treated with K_2CO_3 (502 mg, 3.64 mmol) in one portion. Sufficient amount of water was added to make the solution homogeneous. Reaction progress was monitored by GC/MS, which indicated complete conversion within 8 h. The mixture was diluted with Et_2O and H_2O (5 mL each) and the layers separated. The aqueous layer was washed with Et_2O (3 x 10 mL) and the combined organic layers dried over $MgSO_4$, filtered and concentrated in vacuo. The crude oil thus obtained was purified by filtration through a plug of silica (pentane elution) to afford pure enyne as a colorless oil (214 mg, 1.16 mmol; 95.8%). IR (NaCl): 3295 (s), 2962 (m), 2925 (m), 2105 (w), 1640 (w), 1494 (m), 1454 (m), 993 (m), 914 (m), 820 (m). 1H NMR ($CDCl_3$, 400 MHz): δ 7.30 (1H, m), 7.13 (2H, m), 5.75 (1H, dddd, $J = 17.0, 10.0, 7.1, 6.6$ Hz), 5.00 (1H, dddd, $J = 17.0, 2.0, 1.5, 1.3$ Hz), 4.96 (1H, dddd, $J = 10.1, 1.3, 1.1, 0.9$ Hz), 3.40 (1H, ddq, 7.3, 7.0, 7.0 Hz), 3.23 (1H, s), 2.43 (1H, dddd, $J = 14.1, 6.6, 1.5, 1.3$ Hz), 2.30 (3H, s), 2.28 (1H, dddd, $J = 13.9, 7.3, 1.3, 1.1$ Hz), 1.24 (3H, d, $J = 7.0$ Hz). ^{13}C NMR ($CDCl_3$, 100 MHz): δ 146.4, 137.1, 135.1, 133.5, 129.9, 125.5, 121.0, 115.9, 82.6, 80.5, 41.7, 36.2, 20.7, 20.6. HRMS Calcd. for $C_{14}H_{16}$: 184.1252. Found: 184.1245. $[\alpha]_D^{20}$ -42.4° (c 0.58, $CHCl_3$).

1,6-Dimethyl-4-vinyl-1,2-dihydronaphthalene (10):

A solution of enyne **9** (184 mg, 0.998 mmol) in CH_2Cl_2 (20.0 mL) at 22 °C was treated with Ru complex **37** (31.2 mg, 0.0498 mmol; 5 mol %) in one portion. Complete consumption of the starting enyne required 1 h at 22 °C. *Note: the minor component of the reduced enol mixture was not consumed in the reaction.* The volatiles were then removed in vacuo, and the residue directly purified by chromatography on silica ($R_f = 0.5$ in pentane) to afford a colorless oil (154 mg, 0.835 mmol; 83.6%). This material was used immediately in the cross-metathesis reaction. IR (NaCl): 3031 (w), 2958 (m), 2865 (m), 2829 (w), 1451 (m), 812 (m). 1H NMR ($CDCl_3$, 400 MHz): δ 7.17 (1H, s), 7.11 (1H, d, $J = 7.7$ Hz), 7.03 (1H, d, $J = 7.7$ Hz), 6.65 (1H, dddd, $J = 17.4, 10.8, 1.1, 1.1, 1.1$ Hz), 6.11 (1H, dd, $J = 4.8, 4.8$ Hz), 5.53 (1H, dd, $J = 17.4, 1.8$ Hz), 5.20 (1H, dd, $J = 10.8, 1.8$ Hz), 2.87 (1H, ddq, $J = 7.1, 7.1, 7.0$ Hz), 2.45 (1H, ddd, $J = 16.8, 5.6, 4.9$ Hz), 2.33 (3H, s), 2.11 (1H, ddd, $J = 16.7, 7.3, 5.4$ Hz), 1.24 (3H, d, $J = 7.0$ Hz). ^{13}C NMR ($CDCl_3$, 100 MHz): δ 138.3, 136.1, 135.7, 135.5, 133.2, 127.8, 126.0, 124.9, 124.8, 115.0, 31.7, 31.3, 21.2, 19.8. $[\alpha]_D^{20}$ -45.7° (c 0.15, $CHCl_3$).

⁷ Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2000**, *122*, 8168.

4-(4,7-Dimethyl-3,4-dihydronaphthalen-1-yl)but-3-en-2-one (11): In a nitrogen-filled glove box, a CH_2Cl_2 (3.7 mL) solution of **10** (137 mg, 0.742 mmol) and methyl vinyl ketone (123 μL , 1.48 mmol) was treated with Ru complex **3** (46.5 mg, 0.0743 mmol) in one portion. The reaction mixture was loosely capped, to allow ethylene to escape, and stirred 24 h. The now brown solution was then removed from the glove box, and in vacuo. Purification by chromatography on silica ($R_f = 0.2$ in 9 : 1 hexanes/EtOAc) afforded a colorless oil (125 mg, 0.551 mmol; 74.3%). This material was used directly in the conjugate addition reaction. IR (NaCl): 2959 (m), 2923 (m), 2864 (m), 1668 (s), 1598 (m), 1360 (m), 1253 (m), 979 (w), 814 (w). ^1H NMR (CDCl_3 , 400 MHz): δ 7.48 (1H, dd, $J = 16.1, 1.1$ Hz), 7.17 (1H, s), 7.14 (1H, d, $J = 7.7$ Hz), 7.08 (1H, d, $J = 7.7$ Hz), 6.56 (1H, d, $J = 16.1$ Hz), 6.44 (1H, dd, $J = 4.9, 4.9$ Hz), 2.88 (1H, qdd, $J = 7.1, 6.8, 6.6$ Hz), 2.51 (1H, ddd, $J = 17.4, 6.2, 5.1$ Hz), 2.37 (3H, s), 2.35 (3H, s), 2.19 (1H, ddd, $J = 17.3, 7.5, 5.1$ Hz), 1.23 (3H, d, $J = 7.0$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 198.7, 142.3, 138.4, 135.9, 134.3, 132.2, 131.9, 128.6, 128.3, 126.2, 124.9, 31.8, 31.6, 27.6, 21.2, 19.7. HRMS Calcd. for $\text{C}_{16}\text{H}_{18}\text{O}$: 226.1358. Found: 226.1361. $[\alpha]_D^{20} -23.9^\circ$ (c 0.09, CHCl_3).

4-(4,7-Dimethyl-3,4-dihydronaphthalen-1-yl)pentan-2-one (13): A 13 x 100 mm test tube was charged with $(\text{CuOTf})_2 \cdot \text{PhH}$ (2.0 mg, 3.9×10^{-3} mmol; 5 mol %) and **12** (4.3 mg, 9.4×10^{-3} mmol; 12 mol %) under an atmosphere of nitrogen in a glove box. The contents of the test tube are dissolved in toluene (200 μL) and the resulting solution was allowed to stir 5 min at 22 $^\circ\text{C}$. To this solution was then added **11** (78.2 μmol ; 204 μL of 0.383 M solution in toluene). The test tube was sealed with a septum and Teflon tape prior to removal from the glove box. The mixture obtained was cooled to -30°C and treated with Me_2Zn (0.235 mmol; 117 μL of a 2 M solution in toluene). The resulting yellow mixture was warmed to 4 $^\circ\text{C}$ and allowed to remain at this temperature for 24 h. The solution was treated with a saturated solution of NH_4Cl (1 mL) and diluted with Et_2O (2 mL). The layers were separated and the organic layer removed with a Pasteur pipette and filtered through a plug of silica. This washing/drying process was repeated (4 x 2 mL Et_2O) and the resulting eluent concentrated in vacuo. The crude oil thus obtained was further purified by chromatography on silica ($R_f = 0.4$ in 9 : 1 hexanes/EtOAc) to afford a colorless oil (9.5 mg, 3.9×10^{-2} mmol; 50%). The diastereomer ratio was determined by GLC analysis of the corresponding tetrahydronaphthalene. IR (NaCl): 2960 (m), 2922 (m), 2873 (w), 2829 (m), 1715 (s), 1491 (m), 1453 (m), 1358 (m), 1161 (m), 816 (w). ^1H NMR (CDCl_3 , 400 MHz): δ 7.15 (1H, s), 7.08 (1H, d, $J = 7.5$ Hz), 7.00 (1H, d, $J = 7.5$ Hz), 5.78 (1H, dd, $J = 4.6, 4.2$ Hz), 3.45-3.37 (1H, m), 2.81 (1H, qdd, $J = 7.0, 6.8, 6.6$ Hz), 2.72 (1H, dd, $J = 16.8, 4.6$ Hz), 2.45 (1H, dd, $J = 16.7, 9.2$ Hz), 2.41 (1H, dddd, $J = 16.5, 6.5, 3.8, 1.6$ Hz), 2.33 (3H, s), 2.14 (3H, s), 2.08 (1H, ddd, $J = 16.6, 5.9, 5.8$ Hz), 1.17 (3H, d, $J = 6.8$ Hz), 1.15 (3H, d, $J = 7.1$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 208.3, 139.8, 139.1, 135.7, 133.2, 127.6, 126.5, 123.2, 121.0,

50.5, 31.9, 30.9, 30.5, 29.2, 21.4, 20.1, 19.8. HRMS Calcd. for $C_{17}H_{22}O$: 242.1671. Found: 242.1677. $[\alpha]_D^{20}$ -58.0° (c 0.16, $CHCl_3$).

4-(4,7-Dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)pentan-2-one

(14): To a solution of **13** (77.1 mg, 0.318 mmol) in MeOH (6.4 mL) at 0 °C was added $NaBH_4$ (24.1 mg, 0.636 mmol) in two equal portions. TLC indicated complete conversion to product (R_f = 0.1 in 9 : 1 hexanes/EtOAc) after 15 min. Excess reductant was then consumed by the addition of H_2O (1 mL) and MeOH was removed in vacuo. The resulting solution was washed with Et_2O (3 x 25 mL) and the combined organic layers dried over $MgSO_4$. After concentration in vacuo, the pale yellow residue was filtered through a pad of silica (4 : 1 hexanes/EtOAc elution) to afford a colorless oil (77.7 mg). This material was used in the subsequent step without further purification.

The unpurified material obtained above was transferred as a solution in THF (0.5 mL) to a two-neck round bottom flask fitted with a cold finger. The solution was cooled to $-78^\circ C$ and anhydrous NH_3 (5 mL) condensed into the mixture. Lithium dispersion (30 wt % in mineral oil; 0.2% Na content) was added portion-wise until a blue color persisted (typically 10 equiv were required). The reaction mixture was maintained at $-78^\circ C$ for 15 min and was subsequently treated with a large excess of solid NH_4Cl . The solution immediately went from blue to colorless upon the addition. The resulting suspension was warmed to $22^\circ C$ and diluted with H_2O (5 mL) and Et_2O (25 mL) with transfer to a separatory funnel. The layers separated and the aqueous layer washed with Et_2O (2 x 25 mL). The combined organic layers were dried over $MgSO_4$ and concentrated in vacuo. The unpurified yellow oil was passed through a plug of silica (4 : 1 hexanes/EtOAc elution) to remove the mineral oil. The resulting colorless oil (75.1 mg) was used in the subsequent step without further purification.

The crude oil obtained above was dissolved in CH_2Cl_2 (3.0 mL) at $22^\circ C$ and treated with Dess-Martin periodinane (155 mg, 0.366 mmol) in one portion. Following complete dissolution, the reaction mixture was treated with H_2O (7 μL). Within 1 min, the solution turned cloudy and a white precipitate formed. Following an additional 30 min at $22^\circ C$, TLC indicated complete conversion to product (R_f = 0.4 in 9 : 1 hexanes/EtOAc). The resulting suspension was diluted with Et_2O (10 mL) and concentrated in vacuo to a white solid residue. Purification by chromatography on silica (19 : 1 hexanes/EtOAc) afforded the desired stereoisomer (**14**) as a colorless oil (42.7 mg, 0.175 mmol; 57.3%) the undesired stereoisomer as a colorless oil (6.1 mg, 0.0250 mmol; 8.2%) and recovered starting material (7.3 mg, 0.030 mmol; 9.9%). The following characterization data are given for the major diastereomer obtained above. IR (NaCl): 2958 (m), 2928 (m), 2871 (w), 1715 (s), 1457 (m), 1374 (m), 1355 (m). 1H NMR ($CDCl_3$, 400 MHz): δ 7.13 (1H, d, J = 7.9 Hz), 7.07 (1H, br s), 6.96 (1H, dd, J = 7.8, 1.7 Hz), 2.75 (3H, m), 2.53 (1H, ABX, J_{AB} = 16.2 Hz, J_{AX} = 5.6 Hz), 2.41 (1H, ABX, J_{AB} = 16.2 Hz, J_{BX} = 8.3 Hz), 2.31 (3H, s), 2.16 (3H, s), 1.96 (1H, dddd, J = 12.8, 6.6, 5.0, 3.5 Hz), 1.80 (1H, dddd, J = 13.3, 6.6, 6.6, 3.5 Hz), 1.58 (1H, dddd, J = 13.2, 10.7, 8.3, 3.6 Hz), 1.33

(1H, dddd, $J = 12.8, 10.6, 9.2, 3.5$ Hz), 1.26 (3H, d, $J = 7.0$ Hz), 0.70 (3H, d, $J = 6.8$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 208.8, 140.2, 138.8, 134.8, 128.4, 126.7, 126.5, 49.2, 42.0, 33.2, 32.5, 31.0, 30.5, 21.0(2), 21.1, 14.9. HRMS Calcd. for $\text{C}_{17}\text{H}_{24}\text{ONa}$ ($\text{M}+\text{Na}^+$): 267.1725. Found: 267.1725. $[\alpha]_D^{20} +27.1^\circ$ (c 0.68, CHCl_3).

4-(4,7-Dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)pentan-2-one: The following characterization data are given for the minor diastereomer obtained above. IR (NaCl): 3005 (w), 2959 (m), 2926 (m), 2873 (m), 1714 (s), 1499 (m), 1455 (m), 1355 (m), 1156 (m), 816 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.08 (1H, s), 7.04 (1H, d, $J = 7.7$ Hz), 6.96-6.94 (1H, m), 2.91-2.74 (3H, m), 2.28 (3H, s), 2.20 (1H, s), 2.18 (1H, d, $J = 3.5$ Hz), 2.04 (3H, s), 1.86-1.73 (2H, m), 1.69-1.52 (2H, m), 1.25 (3H, d, $J = 7.1$ Hz), 1.05 (3H, d, $J = 6.8$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 209.2, 140.6, 138.4, 135.4, 129.1, 128.0, 126.8, 46.1, 42.9, 32.6, 32.1, 30.6, 29.1, 23.6, 21.4, 19.5, 18.7. HRMS Calcd. for $\text{C}_{17}\text{H}_{24}\text{ONa}$ ($\text{M}+\text{Na}^+$): 267.1725. Found: 267.1724. $[\alpha]_D^{20} -15.6^\circ$ (c 0.08, CHCl_3).

{1-[2-(4,7-Dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)propyl]vinyl}trimethylsilane: To a solution of $i\text{-Pr}_2\text{NH}$ (29 μL , 0.21 mmol) in THF (0.8 mL) at -78°C , was added BuLi (0.208 mmol; 149 μL of a 1.40 M solution in hexanes) dropwise over 5 min. The resulting solution was then warmed to -30°C , allowed to stir for 15 min, and then recooled to -78°C . To this solution of LDA was then transferred a solution of **14** (40.6 mg, 0.166 mmol) in THF (0.7 mL) and TMSCl (28 μL , 0.22 mmol) dropwise via cannula over 15 min. An additional 0.1 mL THF was used to quantitate the transfer. Following the addition, the reaction mixture was stirred for 10 min at -78°C , at which time TLC analysis indicated complete consumption of the starting ketone. The solution was diluted with wet Et_2O (2 mL) and allowed to warm to 22°C . All volatiles were removed in vacuo, to afford a white solid from which the enol silyl ether can be directly obtained by distillation under reduced pressure ($\text{bp} \approx 160^\circ\text{C}$ at 0.6 mm Hg). The colorless oil thus obtained (49.8 mg, 0.157 mmol; 94.8%) was shown to be a single regioisomer by ^1H NMR. IR (NaCl): 2958 (s), 2928 (s), 2871 (m), 1614 (m), 1303 (m), 1252 (m), 1015 (m), 847 (m), 810 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.13 (1H, d, $J = 7.5$ Hz), 7.03 (1H, s), 6.94 (1H, d, $J = 7.7$ Hz), 4.09 (2H, s), 2.91-2.87 (1H, m), 2.76-2.70 (1H, m), 2.50-2.43 (1H, m), 2.30 (3H, s), 2.12 (1H, dd, $J = 13.9, 7.1$ Hz), 2.02 (1H, dd, $J = 13.9, 7.5$ Hz), 1.96-1.89 (1H, m), 1.86-1.79 (1H, m), 1.58-1.49 (1H, m), 1.32 (1H, dddd, $J = 12.8, 11.4, 9.7, 3.3$ Hz), 1.26 (3H, d, $J = 6.8$ Hz), 0.65 (3H, d, $J = 7.0$ Hz), 0.24 (9H, s). ^{13}C NMR (CDCl_3 , 100 MHz): δ 158.4, 140.4, 139.8, 134.7, 128.2, 126.4, 126.0, 90.9, 42.2, 40.8, 34.8, 32.8, 31.6, 21.9, 21.4, 21.1, 14.3, 0.15(3). HRMS Calcd. for $\text{C}_{20}\text{H}_{32}\text{ONa}$ ($\text{M}+\text{Na}^+$): 339.2113. Found: 339.2118. $[\alpha]_D^{20} +19.5^\circ$ (c 0.12, CHCl_3).

3-(4,7-Dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)butyric acid: A suspension of the enol silyl ether (49.8 mg, 0.157 mmol) and NaHCO_3 (132 mg, 1.57 mmol) in CH_2Cl_2 (1.6 mL) was cooled to -78°C and treated with MeOH (1.6 mL) in a dropwise fashion over 5 min. *Note: the solution of the enol silyl ether must be cold prior to the addition of MeOH in order to avoid protodesilylation.* A vigorous stream of ozone was then bubbled through the reaction mixture for 30 sec, followed by immediate purging of the solution with

a rapid stream of dry nitrogen; nitrogen purge was maintained for 5 min. The resulting solution was treated with Me_2S (114 μL , 1.57 mmol) and allowed to warm slowly to 22°C , as the dry ice/acetone bath evaporated. The solution was diluted with CH_2Cl_2 (2 mL) and acidified ($\text{pH} < 2$) by the slow addition of a 10% aqueous solution of HCl. After dilution with H_2O (5 mL), the layers were separated, and the aqueous layer washed with CH_2Cl_2 (5 x 10 mL). The combined organic layers were dried over MgSO_4 , filtered and concentrated in vacuo to afford a colorless oil. Purification by chromatography on silica (7 : 3 hexanes/ Et_2O containing 0.1% AcOH; $R_f = 0.2$ in 15 : 1 hexanes/ EtOAc containing 0.1% AcOH) afforded a colorless oil (32.0 mg, 0.130 mmol; 82.7%) which solidifies in the freezer to fine white needles.⁸ IR (NaCl): 2959 (br), 2928 (m), 2972 (w), 1706 (s), 1498 (m), 1454 (m), 1411 (m), 1378 (m), 1306 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.13 (1H, d, $J = 7.9$ Hz), 7.06 (1H, s), 6.96 (1H, d, $J = 7.9$ Hz), 2.91-2.86 (1H, m), 2.78-2.65 (2H, m), 2.48 (1H, dd, $J = 15.2, 6.2$ Hz), 2.34 (1H, dd, $J = 15.2, 8.6$ Hz), 2.30 (3H, s), 1.96 (1H, dddd, $J = 12.8, 6.4, 4.9, 3.5$ Hz), 1.84 (1H, dddd, $J = 13.2, 6.8, 6.4, 3.4$ Hz), 1.59 (1H, dddd, $J = 13.2, 10.7, 8.4, 3.4$ Hz), 1.34 (1H, dddd, $J = 12.8, 10.8, 9.3, 3.5$ Hz), 1.26 (3H, d, $J = 6.8$ Hz), 0.78 (3H, d, $J = 6.8$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 178.3, 140.2, 138.5, 134.9, 128.4, 126.7, 126.5, 41.7, 39.6, 34.3, 32.5, 31.0, 22.0, 21.7, 21.1, 14.8. HRMS Calcd. for $\text{C}_{16}\text{H}_{22}\text{O}_2$: 246.1620. Found: 246.1623. $[\alpha]_D^{20} +43.2^\circ$ (c 0.16, CHCl_3).

3-(4,7-Dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)butan-1-ol: To a suspension of LiAlH_4 (21.2 mg, 0.558 mmol) in Et_2O (1.0 mL) at 22°C was transferred a solution of acid (0.112 mmol; 1.00 mL of a 0.112 M solution in Et_2O) dropwise via cannula over 5 min. An additional amount of Et_2O (2 x 0.25 mL) was used to quantitate the transfer. The resulting suspension was allowed to stir for 1 h at 22°C to ensure complete the reduction. Excess LiAlH_4 was consumed by the careful addition of H_2O (25 μL). The resulting white

suspension was successively treated with 15% aqueous NaOH (25 μL) and H_2O (75 μL) and allowed to stir for 15 min to a fine white slurry. The resulting mixture was filtered through a pad of Celite and concentrated in vacuo. The resulting oil was further purified by chromatography on silica (4 : 1 hexanes/ EtOAc ; $R_f = 0.1$ in 9 : 1 hexanes/ EtOAc) to afford a colorless oil (25.5 mg, 0.110 mmol; 98.0%). IR (NaCl): 3326 (br), 2956 (m), 2927 (m), 2871 (m), 1497 (w), 1453 (w), 1376 (w),

⁸ The enol silyl ether can also be converted to the carboxylic acid utilizing a two-step protocol: i) 1.5 equiv *m*-CPBA, 3.0 equiv NaHCO_3 , CH_2Cl_2 , 0°C ; K_2CO_3 / MeOH ii) 1.2 equiv NaIO_4 , $\text{MeOH}/\text{H}_2\text{O}$, 22°C .

1060 (w), 811 (w). ^1H NMR (CDCl_3 , 400 MHz): δ 7.14 (1H, d, $J = 7.7$ Hz), 7.05 (1H, s), 6.95 (1H, d, $J = 7.7$ Hz), 3.83-3.70 (2H, m), 2.88-2.83 (1H, m), 2.77-2.71 (1H, m), 2.36-2.29 (1H, m), 2.30 (3H, s), 1.94 (1H, dddd, $J = 12.7, 6.2, 4.8, 3.5$ Hz), 1.83 (1H, dddd, $J = 13.0, 6.6, 6.0, 3.3$ Hz), 1.72 (1H, dddd, $J = 13.6, 7.3, 7.1, 5.7$ Hz), 1.62-1.52 (2H, m), 1.33 (1H, dddd, $J = 13.0, 11.5, 9.9, 3.3$ Hz), 1.27 (3H, d, $J = 6.8$ Hz), 1.22 (1H, t, $J = 5.3$ Hz), 0.68 (3H, d, $J = 7.0$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 140.4, 139.4, 134.8, 128.2, 126.5, 126.2, 61.6, 42.0, 38.1, 33.5, 32.8, 31.6, 21.9, 21.8, 21.1, 14.4. HRMS Calcd. for $\text{C}_{16}\text{H}_{24}\text{O}$: 232.1827. Found: 232.1828. $[\alpha]_D^{20} +53.6^\circ$ (c 0.12, CHCl_3).

3-(3-Iodo-1-methylpropyl)-1,6-dimethyl-1,2,3,4-tetrahydronaphthalene:

To a solution of alcohol (23.0 mg, 99.0 μmol) in benzene (2.0 mL) was successively added PPh_3 (33.8 mg, 0.129 mmol), imidazole (12.1 mg, 0.178 mmol) and iodine (30.1 mg, 0.119 mmol) at 22 $^\circ\text{C}$. The resulting solution was stirred for 10 min at which time TLC indicated complete conversion to product ($R_f = 0.8$ in 9 : 1 hexanes/EtOAc). The suspension obtained was diluted with Et_2O (2 mL) and transferred to a separatory funnel with exhaustive washing with Et_2O (5 x 5 mL). The organic solution was successively washed with H_2O , 10% aqueous citric acid, saturated aqueous NaHCO_3 and saturated aqueous NaCl (10 mL each). After drying over MgSO_4 , filtration and concentration in vacuo, final purification was performed by chromatography on silica (50 : 1 hexanes/EtOAc) to afford a white solid (29.2 mg, 0.0853 mmol; 86.2%). *Note: this material should be stored protected from light.* IR (NaCl): 3004 (w), 2957 (m), 2926 (m), 2869 (m), 1497 (m), 1455 (m), 1376 (m), 1244 (m), 1181 (m), 811 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.14 (1H, d, $J = 7.9$ Hz), 7.05 (1H, s), 6.96 (1H, d, $J = 7.9$ Hz), 3.27 (2H, ABqdd, $J_{AB} = 19.1$ Hz, $J_{dd} = 8.1, 6.7$ Hz), 2.09-2.85 (1H, m), 2.78-2.69 (1H, m), 2.31 (3H, s), 2.30-2.26 (1H, m), 2.00-1.76 (4H, m), 1.55 (1H, dddd, $J = 13.4, 11.4, 9.0, 3.5$ Hz), 1.33 (1H, dddd, $J = 13.0, 11.4, 9.9, 3.5$ Hz), 1.27 (3H, d, $J = 7.0$ Hz), 0.67 (3H, d, $J = 6.8$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 140.6, 139.1, 135.1, 128.3, 126.8, 126.6, 41.1, 39.0, 38.6, 32.9, 31.7, 22.0(2), 21.4, 14.1, 5.9. HRMS Calcd. for $\text{C}_{16}\text{H}_{23}\text{I}$: 342.0845. Found: 342.0847. $[\alpha]_D^{20} +14.8^\circ$ (c 0.16, CHCl_3).

(+)-Erogorgiaene (1):

A 10 x 75 mm test tube was charged with CuI (1.9 mg, 9.8 μmol ; 20 mol %) under a nitrogen atmosphere in a glove box. The test tube was sealed with a septum and Teflon tape prior to removal from the glove box. To this vessel was then transferred a solution of iodide (48.8 μmol ; 140 μL of a 0.349 M solution in THF) via cannula. An additional amount of THF (2 x 50 μL) was used to quantitate the transfer. The resulting suspension was then cooled to -78°C and treated with isobutenylmagnesium bromide (73.2 μmol ; 84 μL of a 0.87 M solution in THF). The reaction mixture was transferred to a cold room, warmed to 4 $^\circ\text{C}$ and maintained at that temperature for 8 h. The resulting gray reaction mixture was diluted with Et_2O (2 mL) treated with a saturated aqueous solution of NH_4Cl (2 mL). The mixture was transferred to a separatory funnel with exhaustive washing with Et_2O (5 x 2 mL). The layers were separated and the aqueous layer washed with Et_2O (5 x 5 mL). The combined organic layers were dried over MgSO_4 , filtered and concentrated in vacuo. Initial purification was performed by chromatography on silica (hexanes; $R_f = 0.6$ in 49 : 1 hexanes/EtOAc). Final purification was performed by argention⁹ chromatography (19 : 1 hexanes/ Et_2O) to afford pure erogorgiaene as a colorless oil (7.6 mg, 28.1 μmol ; 57.5%). IR (NaCl): 2960 (m), 2923 (m), 2853 (m), 1495 (m), 1451 (m), 1376 (m), 810 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.14 (1H, d, $J = 7.9$ Hz), 7.03 (1H, s), 6.94 (1H, d, $J = 7.9$ Hz), 5.17 (1H, br t, $J = 7.1$

⁹ Mander, L., N.; Williams, C. M. *Tetrahedron*, 2001, 57, 425-477.

Hz), 2.90-2.86 (1H, m), 2.77-2.68 (1H, m), 2.30 (3H, s), 2.18-2.00 (3H, m), 1.92 (1H, dddd, $J = 12.6, 5.9, 5.9, 3.3$ Hz), 1.81 (1H, dddd, $J = 13.0, 6.6, 6.2, 3.2$ Hz), 1.72 (3H, br s), 1.64 (3H, br s), 1.58-1.51 (1H, m), 1.49-1.40 (1H, m), 1.38-1.30 (2H, m), 1.27 (3H, d, $J = 7.0$ Hz), 0.64 (3H, d, $J = 6.8$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 140.4, 139.9, 134.7, 131.3, 128.1, 126.4, 126.0, 124.9, 41.5, 37.0, 35.2, 32.8, 31.8, 26.3, 25.7, 21.9, 21.5, 21.1, 17.7, 14.5. HRMS Calcd. for $\text{C}_{20}\text{H}_{30}$: 270.2348. Found: 270.2352. $[\alpha]_D^{20} +40.6^\circ$ (c 0.14, CHCl_3). Lit: $[\alpha]_D^{20} +24.4^\circ$ (c 3.2, CHCl_3).

3-(2-Bromo-4-methylphenyl)butyric acid: To a solution of $i\text{-Pr}_2\text{NH}$ (0.635 mL, 4.53 mmol) in THF (20.1 mL) at -78°C was added BuLi (4.53 mmol; 2.59 mL of a 1.75 M solution in hexanes) dropwise over 10 min. The resulting solution was then warmed to -30°C , maintained 30 min, then recooled to -78°C . To the resulting solution of LDA was transferred a solution of **6** (0.770 g, 3.02 mmol) in THF (5.0 mL) dropwise via cannula. Upon completion of the addition, TMSCl (574 μL , 4.53 mmol) was added to the reaction mixture, which was subsequently warmed to -30°C and maintained for 2 h. The resulting solution was treated with wet Et_2O (10 mL) and warmed to 22°C . All volatiles were removed in vacuo to afford a pale yellow oil (1.10 g). The unpurified enol silyl ether, determined to be of single regiochemistry by ^1H NMR, was used without further purification in the subsequent step.

The crude enol silyl ether was dissolved in CH_2Cl_2 (33.6 mL) and cooled to -78°C . Pyridine (0.408 mL, 5.04 mmol) and MeOH (22.4 mL, 105 mmol) were successively added to this pale yellow reaction mixture. A vigorous stream of ozone was then introduced through the mixture, using a large gauge needle, until the persistence of a blue color. A stream of oxygen was utilized to purge the excess ozone from the mixture. The now colorless solution was treated with Me_2S (1.09 mL, 13.4 mmol) and allowed to warm to 22°C over 12 h as the dry ice/acetone bath evaporated. A short path distillation apparatus was attached to the flask and all volatile material was removed by distillation under reduced pressure (0.1 mm). The remaining residue was dissolved in EtOAc and transferred to a separatory funnel. The EtOAc layer was successively washed with 10% aqueous HCl (3 x 30 mL) and 1 M NaOH (3 x 30 mL). The NaOH washes were pooled and washed with pentane (3 x 30 mL). The NaOH layer was then acidified ($\text{pH} < 2$) with concentrated HCl and exhaustively washed with EtOAc. The combined EtOAc washes were dried over MgSO_4 , filtered and concentrated in vacuo to afford a colorless oil (0.747 g, 2.91 mmol; 96.2% yield). IR (NaCl): 3029 (br), 2969 (w), 2924 (w), 1709 (s), 1490 (w), 1414 (w), 1298 (w). ^1H NMR (CDCl_3 , 400 MHz): δ 7.39 (1H, m), 7.13-7.08 (2H, m), 3.77-3.68 (1H, m), 2.74 (1H, dd, $J = 15.6, 5.9$ Hz), 2.52 (1H, dd, $J = 15.7, 8.9$ Hz), 2.29 (3H, s), 1.30 (3H, d, $J = 7.0$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 177.2, 140.9, 137.9, 133.5, 128.6, 126.6, 124.0, 41.0, 34.4, 20.5(2). HRMS Calcd. for $\text{C}_{11}\text{H}_{13}\text{BrO}_2$: 256.0097. Found: 256.0099. Anal. Calcd. for $\text{C}_{11}\text{H}_{13}\text{BrO}_2$: C, 51.38; H, 5.10. Found: C, 51.40; H, 5.20.

3-(2-Bromo-4-methylphenyl)butan-1-ol: A 25 mL round bottom flask was charged with LiAlH_4 (0.158 g, 4.16 mmol) under a nitrogen atmosphere in a glove box. The flask was sealed with a septum and Teflon tape prior to removal from the glove box. The LiAlH_4 was suspended in Et_2O (4.7 mL) and allowed to cool to -50°C . To this suspension was then transferred a solution of acid (823 mg, 3.20 mmol) in THF (7.1 mL) dropwise via cannula over 30 min. The mixture was warmed to 4°C and maintained at this temperature for 4 h. The resulting grey suspension was cooled to -30°C and successively treated with H_2O (0.158 mL), 15% aqueous NaOH (0.158 mL) and H_2O (0.474 mL). After the mixture was allowed to stir for 15 min, the resulting white suspension was filtered through a pad of Celite (exhaustive washing with EtOAc). The filtrate was dried over MgSO_4 and concentrated in vacuo to afford a pale yellow oil. Purification by radial chromatography on silica (CH_2Cl_2) afforded a colorless oil (0.700 g, 2.88 mmol; 90.0% yield). IR (NaCl): 3321 (br), 2961 (w), 2926 (m), 2871 (m), 1608 (m), 1489 (w), 1455 (m), 1048 (m), 819 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.37 (1H, s), 7.14-7.08 (2H, m), 3.62-3.55 (2H, m), 3.39 (1H, ddq, $J = 7.3, 7.1, 7.0$ Hz), 2.29 (3H, s), 1.88-1.83 (2H, m), 1.28 (1H, dd, $J = 5.9, 5.5$ Hz), 1.24 (3H, d, $J = 7.0$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 142.4, 137.5, 133.2, 128.6, 127.0, 124.3, 61.0, 40.4, 34.2, 21.4, 20.5. HRMS Calcd. for $\text{C}_{11}\text{H}_{15}\text{BrO}$: 242.0307. Found: 242.0306. Anal. Calcd. for $\text{C}_{11}\text{H}_{15}\text{BrO}$: C, 54.34; H, 6.22. Found: C, 54.43; H, 6.39.

3-(4-Methyl-2-vinylphenyl)butan-1-ol: A 100 mL round bottom flask was charged with $\text{PdCl}_2(\text{PPh}_3)_2$ (289 mg, 0.411 mmol; 10 mol %) under a nitrogen atmosphere in a glove box. The flask was sealed with a septum and Teflon tape prior to removal from the glove box. To this mixture was transferred a solution of bromide (1.00 g, 4.11 mmol) in DMF (7.0 mL) dropwise via cannula, followed by the addition of Et_3N (1.72 mL, 12.3 mmol). The flask was placed in a stainless steel Parr reactor and charged with an atmosphere of ethylene (700 psi). The reactor was heated to 90°C and maintained for 12 h. After cooling to 22°C , the reactor was vented and the flask removed. The resulting orange suspension was diluted with Et_2O and H_2O (25 mL each) with transfer to a separatory funnel. The layers were separated and the aqueous layer washed with Et_2O (3 x 25 mL). The combined organic layers were washed with 10% aqueous HCl (3 x 25 mL), dried over MgSO_4 and concentrated in vacuo to afford a red/orange residue. Purification by chromatography on silica (4 : 1 hexanes/ Et_2O) afforded a colorless oil (0.730 g, 3.84 mmol; 93.3%). IR (NaCl): 3333 (br), 2961 (m), 2926 (m), 2872 (m), 1492 (m), 1455 (m), 1043 (m), 991 (m), 910 (m), 818 (m). ^1H NMR (CDCl_3 , 400 MHz): δ 7.26 (1H, m), 7.14-7.07 (3H, m), 5.58 (1H, dd, $J = 17.3, 1.6$ Hz), 5.29 (1H, dd, $J = 11.0, 1.6$ Hz), 3.63-3.47 (2H, m), 3.26 (1H, ddq, $J = 7.1, 7.1, 7.0$ Hz), 2.33 (3H, s), 1.85 (2H, dd, $J = 13.6, 6.7$ Hz), 1.25 (3H, d, $J = 7.0$ Hz), 1.17 (1H, t, $J = 5.5$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz): δ 141.0, 136.5, 135.4, 135.2, 128.9, 127.0, 125.5, 116.0, 61.1, 40.7, 30.4, 21.9, 21.0. HRMS Calcd. for $\text{C}_{13}\text{H}_{18}\text{ONa}$ ($\text{M}+\text{Na}^+$): 213.1255. Found: 213.1253. $[\alpha]_D^{20} +39.0^\circ$ (c 0.12, CHCl_3).

Toluene-4-sulfonic acid 3-(4-methyl-2-vinylphenyl)butyl ester:

A solution of alcohol (0.700 g, 3.68 mmol) in pyridine (1.79 mL, 22.1 mmol) was cooled to 0 °C and treated in one portion with TsCl (842 mg, 4.41 mmol). After stirring for 0.5 h at 0 °C, the reaction mixture was warmed to 22 °C and maintained for 2.5 h. The white suspension thus obtained was partitioned between H₂O (25 mL) and Et₂O (50 mL) and transferred to a separatory funnel. The layers were separated and the aqueous layer washed with Et₂O (2 x 25 mL). The combined organic layers were successively washed with 10% aqueous HCl, saturated aqueous NaHCO₃ and H₂O (3 x 25 mL each), dried over MgSO₄ and concentrated in vacuo. Purification by chromatography on silica (hexanes → 9 : 1 hexanes/EtOAc gradient; *R_f* = 0.2 in 9 : 1 hexanes/EtOAc) afforded a colorless oil (1.15 g, 3.34 mmol; 90.7%). IR (NaCl): 2964 (m), 2930 (m), 1361 (s), 1177 (s), 1097 (m), 942 (m), 892 (m), 816 (m). ¹H NMR (CDCl₃, 400 MHz): δ 7.72 (2H, AB, *J*_{AB} = 8.4 Hz), 7.30 (2H, AB, *J*_{AB} = 8.1 Hz), 7.21 (1H, s), 7.03 (1H, dd, *J* = 8.1, 1.3 Hz), 6.99 (1H, d, *J* = 7.7 Hz), 6.94 (1H, dd, *J* = 17.2, 11.0 Hz), 5.54 (1H, dd, *J* = 17.2, 1.5 Hz), 5.26 (1H, dd, *J* = 11.0, 1.6 Hz), 3.94 (2H, Abqdd, *J*_{AB} = 9.9 Hz, *J*_{dd} = 6.4, 6.4 Hz), 3.16 (1H, ddq, *J* = 7.1, 7.1, 7.0 Hz), 2.44 (3H, s), 2.31 (3H, s), 1.99-1.87 (2H, m), 1.16 (3H, d, *J* = 7.0 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ 144.5, 139.6, 136.5, 135.7, 134.7, 133.1, 129.7, 128.9, 127.8, 127.1, 125.2, 116.2, 68.9, 36.4, 30.2, 21.6(2), 20.9. HRMS Calcd. for C₂₀H₂₄O₃SNa (M+Na⁺): 367.1344. Found: 367.1342.

4,7-Dimethyl-1-hydroxymethyl-1,2,3,4-tetrahydronaphthalene:

A 100 mL round bottom flask fitted with a Liebig condenser was charged with tosylate (1.17 g, 3.40 mmol), Cp₂ZrCl₂ (49.6 mg, 0.170 mmol; 5 mol %) and 33.5 mL of THF at 22 °C. (2-Cyclohexyl)ethylmagnesium bromide (17.0 mmol; 11.8 mL of a 1.44 M solution in THF) was then added in one portion, the resulting solution heated to 55 °C and was maintained at this temperature for 14 h. The resulting red reaction mixture was then cooled to 0 °C and O₂ was bubbled through the solution for 5 min. The resulting colorless suspension was then diluted with H₂O and Et₂O (25 mL each) and transferred to a separatory funnel. The layers were separated and the aqueous layer washed with Et₂O (3 x 25 mL). The combined organic layers were dried over MgSO₄, filtered and concentrated in vacuo to afford a yellow oil. Purification by chromatography on silica (*R_f* = 0.4 in 19 : 1 DCM/Et₂O) afforded a colorless oil (395 mg, 2.07 mmol; 61.0% yield). ¹H NMR indicated the syn/anti diastereomer ratio to be 87 : 13. IR (NaCl): 3323 (br), 2925 (m), 2869 (m), 1501 (w), 1453 (w), 1039 (m), 814 (m). ¹H NMR (CDCl₃, 400 MHz): δ 7.16 (1H, d, *J* = 7.9 Hz), 7.04 (1H, s), 7.00 (1H, d, *J* = 7.9 Hz), 3.83 (2H, dd, *J* = 6.4, 5.7 Hz), 2.92 (1H, dddd, *J* = 6.0, 6.0, 6.0, 6.0 Hz), 2.85 (1H, qdd, *J* = 7.0, 6.8, 6.6 Hz), 2.31 (3H, s), 2.02-1.96 (1H, m), 1.89-1.79 (2H, m), 1.62-1.55 (1H, m), 1.40 (1H, dd, *J* = 6.0, 5.9 Hz), 1.29 (3H, d, *J* = 7.0 Hz). ¹³C NMR (CDCl₃, 100 MHz): δ

140.1, 136.2, 135.1, 129.1, 128.0, 127.3, 67.1, 40.7, 32.5, 28.3, 23.1, 22.4, 21.0. HRMS Calcd. for $C_{13}H_{18}O$: 190.1356. Found: 190.1358.

4,7-Dimethyl-1,2,3,4-tetrahydronaphthalene-1-carbaldehyde: To a solution of alcohol (0.280 g, 1.47 mmol) in CH_2Cl_2 (15 mL) at 22 °C, was added Dess-Martin periodinane (749 mg, 1.77 mmol) in one portion. Upon complete dissolution of the reaction mixture, H_2O (27 μ L) was then added. Within 1 min, the solution became cloudy and a white precipitate formed. Following an additional 30 min at 22 °C, TLC indicated complete conversion to product (R_f =

0.4 in 9 : 1 hexanes/EtOAc). The resulting suspension was diluted with Et_2O (10 mL) and concentrated in vacuo to a white solid residue. Purification by chromatography on silica (9 : 1 hexanes/EtOAc) afforded a colorless oil (265 mg, 1.40 mmol; 95.6%). 1H NMR indicated the syn/anti diastereomer ratio to be 87 : 13. IR (NaCl): 2957 (m), 2929 (m), 2861 (m), 1723 (s), 1503 (m), 1455 (w), 818 (w). 1H NMR ($CDCl_3$, 400 MHz): δ 9.71 (1H, d, J = 2.4 Hz), 7.21 (1H, d, J = 7.9 Hz), 7.07 (1H, br d, J = 7.9 Hz), 6.94 (1H, br s), 3.53 (1H, br dd, J = 5.7, 4.4 Hz), 2.88 (1H, qdd, J = 7.0, 6.8, 6.4 Hz), 2.32 (3H, s), 2.29-2.21 (1H, m), 1.94-1.84 (2H, m), 1.56-1.48 (1H, m), 1.29 (3H, d, J = 7.0 Hz). ^{13}C NMR ($CDCl_3$, 100 MHz): δ 202.3, 139.9, 135.7, 130.3, 130.0, 128.4, 128.3, 52.2, 32.0, 28.9, 22.0, 20.9, 20.7. HRMS Calcd. for $C_{13}H_{16}O$: 188.1201. Found: 188.1196. $[\alpha]_D^{20}$ -22.0° (c 0.10, $CHCl_3$).

4-(4,7-Dimethyl-1,2,3,4-tetrahydronaphthalen-1-yl)but-3-en-2-one: To a suspension of NaH (52.1 mg, 2.17 mmol) in THF (10 mL) at 22 °C was added dimethyl-2-oxopropylphosphonate (327 μ L, 2.37 mmol) dropwise over 5 min. After hydrogen gas evolution ceased (~1 h), to the resulting white suspension was transferred a solution of aldehyde (371 mg, 1.97 mmol) in THF (10 mL) dropwise via cannula over 10 min. Following 12 h of reaction time, the resulting colorless solution was treated with saturated aqueous NH_4Cl (25 mL) and

transferred to a separatory funnel. The layers separated and the aqueous layer washed with Et_2O (3 x 25 mL). The combined organic layers were dried over $MgSO_4$ and concentrated in vacuo to afford a pale yellow oil. Purification by chromatography on silica (19 : 1 hexanes/EtOAc; R_f = 0.3 in 9 : 1 hexanes/EtOAc) afforded a colorless oil (312 mg, 1.37 mmol; 69.4%). 1H NMR analysis indicated the product to be a 1 : 1 mixture of diastereomers. IR (NaCl): 2957 (w), 2929 (m), 2863 (m), 1674 (s), 1622 (s), 1360 (m), 1254 (m), 979 (m), 822 (m). 1H NMR ($CDCl_3$, 400 MHz): δ 7.15 (2H, d, J = 7.9 Hz), 7.02 (2H, dd, J = 7.9, 1.3 Hz), 6.85 (1H, dd, J = 11.5, 8.2 Hz), 6.83 (2H, s), 6.81 (1H, dd, J = 11.5, 8.2 Hz), 6.07 (1H, dd, J = 11.4, 1.1 Hz), 6.03 (1H, dd, J = 11.2, 1.1 Hz), 3.63-3.55 (2H, m), 2.93-2.88 (2H, m), 2.28-2.26 (12H, m), 2.12-2.04 (1H, m), 2.03-1.84 (4H, m), 1.70 (1H, dddd, J = 13.0, 8.9, 6.7, 2.4 Hz), 1.64-1.75 (1H, m), 1.50 (1H, dddd, J = 13.1, 9.0, 6.6, 2.6 Hz), 1.29 (3H, d, J = 7.1 Hz), 1.28 (3H, d, J = 7.0 Hz). ^{13}C NMR ($CDCl_3$, 100 MHz): δ 198.9(2), 151.9(2), 139.0(2), 135.5(2), 1353(2), 131.6, 131.4, 129.9, 129.8, 128.4, 128.3, 127.7(2), 42.5(2), 32.1, 32.0, 28.6, 28.5,

269(2), 26.4, 26.3, 22.9, 22.8, 20.9(2). HRMS Calcd. for $C_{16}H_{20}ONa$ ($M+Na^+$): 251.1412. Found: 251.1403.

¹H NMR Data and GLC Traces (Table 1)

4-(2-Bromo-4-methylphenyl)pentan-2-one (6): ¹H NMR (CDCl₃, 400 MHz): δ 7.38 (1H, s), 7.12–7.07 (2H, m), 3.78–3.69 (1H, m), 2.78 (1H, dd, *J* = 16.1, 5.5 Hz), 2.57 (1H, dd, *J* = 16.0, 8.7 Hz), 2.29 (3H, s), 2.12 (3H, s), 1.24 (3H, d, *J* = 7.0 Hz).

The optical purity of the conjugate addition adduct was established by chiral GLC analysis (Chiraldex-GTA, 115 °C, 15 psi); chromatograms are illustrated below.

from ACA: >98% ee

authentic racemic

4-Phenylpentan-2-one: ¹H NMR (CDCl₃, 400 MHz): δ 7.31 (5H, m), 3.30 (1H, dqd, *J* = 7.3, 7.0, 6.8 Hz), 2.76 (1H, dd, *J* = 16.3, 6.4 Hz), 2.66 (1H, dd, *J* = 16.3, 7.9 Hz), 2.07 (3H, s), 1.26 (3H, d, *J* = 7.0 Hz).

The optical purity of the conjugate addition adduct was established by chiral GLC analysis (β-dex, 110 °C, 15 psi); chromatograms are illustrated below.

from ACA: 86% ee

authentic racemic

4-(4-Trifluoromethylphenyl)pentan-2-one: ^1H NMR (CDCl_3 , 400 MHz): δ 7.55 (2H, AB, $J_{AB} = 8.1$ Hz), 7.33 (2H, AB, $J_{AB} = 8.1$ Hz), 3.38 (1H, dqd, $J = 7.1, 7.0, 6.8$ Hz), 2.73 (2H, ABX, $J_{AB} = 16.8$ Hz, $J_{AX} = 7.5$ Hz, $J_{BX} = 6.8$ Hz), 2.09 (3H, s), 1.27 (3H, d, $J = 7.0$ Hz).

The optical purity of the conjugate addition adduct was established by chiral GLC analysis (β -dex, 100 $^\circ\text{C}$, 15 psi); chromatograms are illustrated below.

from ACA: 89% ee

authentic racemic

4-(2-Methoxyphenyl)pentan-2-one: ^1H NMR (CDCl_3 , 400 MHz): δ 7.19 (1H, ddd, $J = 8.1, 7.3, 1.7$ Hz), 7.16 (1H, dd, $J = 7.5, 1.8$ Hz), 6.92 (1H, ddd, $J = 7.5, 7.5, 1.1$ Hz), 6.86 (1H, d, $J = 8.1, 0.9$ Hz), 3.83 (3H, s), 3.68 (1H, dqd, $J = 8.6, 7.0, 5.5$ Hz), 2.77 (1H, dd, $J = 15.7, 5.5$ Hz), 2.60 (1H, dd, $J = 15.7, 8.6$ Hz), 2.10 (3H, s), 1.24 (3H, d, $J = 7.0$ Hz).

The optical purity of the conjugate addition adduct was established by chiral HPLC analysis (Chiralcel OJ, 99.0:1.0 hexanes/isopropanol, 1.0 mL/min, 254 nm); chromatograms are illustrated below.

from ACA: 94% ee

authentic racemic

4-(2-Phenoxyphenyl)pentan-2-one: ^1H NMR (CDCl_3 , 400 MHz): δ 7.34-7.27 (3H, m), 7.18-7.05 (3H, m), 6.97-6.95 (2H, m), 6.87 (1H, dd, $J = 8.1, 1.5$ Hz), 3.67 (1H, dqd, $J = 8.1, 6.8, 6.6$ Hz).

Hz), 2.82 (1H, dd, $J = 15.9, 6.0$ Hz), 2.65 (1H, dd, $J = 16.1, 8.2$ Hz), 2.06 (3H, s), 1.26 (3H, d, $J = 7.0$ Hz).

The optical purity of the conjugate addition adduct was established by chiral HPLC analysis (Chiralcel OD, 99.5:0.5 hexanes/isopropanol, 1.0 mL/min, 254 nm); chromatograms are illustrated below.

from ACA: 98% ee

authentic racemic

4-(4-Fluorophenyl)pentan-2-one: ^1H NMR (CDCl_3 , 400 MHz): δ 7.23-7.15 (2H, m), 7.07 (1H, ddd, $J = 7.5, 7.5, 1.3$ Hz), 7.00 (1H, dd, $J = 10.8, 8.2, 1.3$ Hz), 3.58 (1H, dqd, $J = 7.9, 6.8, 6.8$ Hz), 2.82 (1H, dd, $J = 16.5, 6.3$ Hz), 2.71 (1H, dd, $J = 16.7, 8.2$ Hz), 2.11 (3H, s), 1.28 (3H, d, $J = 7.0$ Hz).

The optical purity of the conjugate addition adduct was established by chiral GLC analysis (Chiraldex-GTA, 80 °C, 15 psi); chromatograms are illustrated below.

from ACA: 97% ee

authentic racemic

4-o-Tolylpentan-2-one: ^1H NMR (CDCl_3 , 400 MHz): δ 7.18-7.07 (4H, m), 3.56 (1H, dqd, $J = 8.2, 6.8, 6.0$ Hz), 2.76 (1H, dd, $J = 16.5, 6.0$ Hz), 2.66 (1H, dd, $J = 16.5, 8.2$ Hz), 2.37 (3H, s), 2.09 (3H, s), 1.22 (3H, d, $J = 6.8$ Hz).

The optical purity of the conjugate addition adduct was established by chiral GLC analysis (Chiraldex-GTA, 90 °C, 15 psi); chromatograms are illustrated below.

Peak #	RT (min)	Type	Width (min)	Area	Area %	Name
1	47.104	1	1.155	517.974	95.300	
2	54.348	1	0.938	25.275	4.700	

from ACA: 91% ee

Peak #	RT (min)	Type	Width (min)	Area	Area %	Name
1	47.104	1	1.155	1185.206	49.975	
2	54.348	1	1.559	1194.465	50.025	

authentic racemic

¹H NMR Spectrum of Synthetic Erogorgiaene

¹³C NMR Spectrum of Synthetic Erogorgiaene