Supporting Information

Ene-Like Addition of an Oxoammonium Cation to Alkenes: Highly Selective Route to Allylic Alkoxyamines

Priya P. Pradhan, James M. Bobbitt,* and William F. Bailey*

Department of Chemistry, The University of Connecticut, Storrs, Connecticut 06269

Table of Contents

Experimental Section	S2 – S6
¹ H and ¹³ C NMR Spectra	S6 – S22

Experimental Section

General. Compound numbers and Tables refer to those contained in the main body of the paper. HPLC grade acetonitrile was used as received; reactions were conducted in capped Erlenmeyer flasks. The majority of alkenes used in this study were commercially available; literature procedures, incorporating some minor modifications, were followed for the preparation of butylidenecyclohexane,¹ 2-methyl-2-decene,² and (2,2-dimethylpropylidene)cyclopentane;³ 2,2,6,6-tetramethylpiperidine-1-oxoammonium tetrafluoroborate (**1**) was prepared following the procedure of Bobbitt and Merbough.⁴ Gas-liquid chromatography (GC) was performed on a Hewlett-Packard 5890 gas chromatograph equipped with a HP-5, 25 m X 0.2 mm fused-silica glass capillary column and a flame-ionization detector.

General Procedure for Ene-Like Addition of **1** to Tetrasubstituted Alkenes.

A solution of 1.00 g (3.33 mmol) of 4-acetamido-2,2,6,6-tetramethylpiperidine-1-oxoammonium tetrafluoroborate (**1**) and 3.33 mmol of an alkene in 15 mL of acetonitrile was stirred at room temperature for a period of time (Table 1) until starch-KI paper indicated complete consumption of the oxoammonium salt. The reaction mixture was concentrated at reduced pressure, the residual semisolid was stirred in 15 mL of dichloromethane for 1 h and filtered. The solid was discarded and the filtrate was washed sequentially with three 10-mL portions of saturated aqueous Na₂CO₃, two 10-mL

---

(1) Anselmi, C.; Centini, M.; Fedeli, P.; Paoli, M. L.; Segal, A.; Scesa, C.; Pelosi, P. *J. Agric. Food Chem.* **2000**, *48*, 1285.

(2) Hercouet, A.; Corre, M. *Phosphorus Sulfur Relat. Elem.* **1987**, *29*, 111.

(3) Armesto, D.; Gallego, M. G.; Horspool, W. M.; Agarrabeitia, A. R. *Tetrahedron* **1995**, *51*, 9223.

(4) Bobbitt, J. M.; Merbough, N. *Organic Syntheses*; Wiley: New York, 2005; Vol. 82, p 80.

portions of water, one 10-mL portion of brine, then dried (Mg_2SO_4) and concentrated under reduced pressure. The residue was purified by recrystallization or by silica gel chromatography as appropriate. The structures and purity of the products were established on the basis of the following data; yields may be found in Table 1.

2-Methyl-3-[4-acetamido-(2,2,6,6-tetramethylpiperidin-1-yl)oxy]-1-pentene

(Table 1, entry 1). Recrystallization from hexane gave a white solid, mp 108-109 °C, in 93 % yield: ^1H NMR δ 5.18 (br s, 1H), 4.85 (br s, 1H), 4.82 (br s, 1H), 4.14 – 4.06 (m, 1H), 3.91 (dd, $J = 3.8$ Hz, $J = 10.3$ Hz, 1H), 1.93 (s, 3H), 1.87 – 1.72 (m, 3H), 1.66 (s, 3H), 1.54 – 1.46 (m, 1H), 1.32 – 1.17 [overlapping patterns, i.e. 1.32 – 1.22 (m, 2H), 1.20 (s, 3H), 1.19 (s, 3H), 1.17 (s, 3H)], 1.08 (s, 3H), 0.72 (t, $J = 7.5$, 3H); ^{13}C NMR δ 169.4, 144.6, 113.8, 90.9, 60.2, 60.0, 46.7, 46.6, 41.3, 34.3, 34.2, 24.2, 23.8, 21.1, 21.1, 17.7, 10.0. Anal. Calcd for $\text{C}_{17}\text{H}_{32}\text{N}_2\text{O}_2$: C, 68.88; H, 10.88; N, 9.45. Found: C, 68.54; H, 10.50; N, 9.45.

1-{1-[4-Acetamido(2,2,6,6-tetramethylpiperidin-1-yl)oxy]butyl}cyclohexene

(Table 1, entry 2). Recrystallization from hexane gave a white solid, mp 114-116 °C, in 60 % yield: ^1H NMR δ 5.55 (s, 1H), 5.20 (br s, 1H), 4.13 – 4.09 (m, 1H), 3.87 (dd, $J = 3.9$ Hz, $J = 10.9$ Hz, 1H), 2.10 – 2.02 (m, 3H), 1.95 (s, 3H), 1.81 – 1.69 (m, 6H), 1.57 – 1.44 (m, 3H), 1.35 – 1.25 (m, 2H), 1.22 – 1.07 [overlapping pattern, i.e. 1.22 (s, 3H), 1.21 (s, 3H), 1.17 (s, 3H), 1.07 (s, 3H), 1.22 – 1.07 (m, 2H)], 0.89 (t, $J = 7.3$ Hz, 3H); ^{13}C NMR δ 169.4, 137.5, 125.9, 90.1, 60.5, 59.6, 46.7, 46.5, 41.4, 34.4, 34.0, 33.5, 25.4, 23.8, 23.5, 22.8, 22.6, 21.2, 21.0, 19.2, 14.3. HRMS-ES (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{21}\text{H}_{38}\text{N}_2\text{O}_2$, 351.3011; found, 351.2983.

1-{1-[4-Acetamido(2,2,6,6-tetramethylpiperidin-1-yl)oxy]-2,2-dimethylpropyl}cyclopentene (Table 1, entry 3). Recrystallization from hexane gave a white solid, mp 154-155 °C, in 80 % yield: ¹H NMR δ 5.70 (s, 1H), 5.27 (br s, 1H), 4.26 (s, 1H), 4.14 – 4.12 (m, 1H), 2.44 – 2.42 (m, 2H), 2.30 – 2.28 (m, 2H), 1.96 (s, 3H), 1.92 – 1.88 (m, 1H), 1.77 – 1.72 (m, 3H), 1.39 – 1.24 [overlapping pattern, i.e. 1.39 – 1.24 (m, 2H), 1.34 (s, 3H), 1.29 (s, 3H)], 1.19 (s, 3H), 1.03 (s, 9H), 1.02 (s, 3H); ¹³C NMR δ 169.5, 143.2, 134.7, 87.9, 61.4, 59.6, 47.4, 46.7, 41.3, 35.8, 35.4, 34.9, 33.3, 31.6, 27.7, 24.1, 23.8, 21.7, 21.4. Anal. Calcd for C₂₁H₃₈N₂O₂: C, 71.95; H, 10.93; N, 7.99. Found: C, 71.58; H, 10.54; N, 8.07.

2-Methylene-1-[4-acetamido(2,2,6,6-tetramethylpiperidin-1-yl)oxy]-cyclohexane (Table 1, entry 4). Recrystallization from hexane gave a white solid, mp 109-111 °C, in 95 % yield: ¹H NMR δ 5.17 (br s, 1H), 4.89 (s, 1H), 4.73 (s, 1H), 4.16 – 4.05 (m, 2H), 2.36 – 2.31 (m, 1H), 2.05 – 1.99 (m, 1H), 1.94 – 1.89 (m, 4H), 1.82 – 1.72 (m, 4H), 1.53 – 1.42 (m, 3H), 1.29 (dt, J = 4.0 Hz, J = 12.0, 2H), 1.23 (s, 3H), 1.23 (s, 3H), 1.19 (s, 3H), 1.11 (s, 3H); ¹³C NMR δ 169.4, 150.1, 107.6, 85.1, 60.4, 60.0, 46.6, 41.3, 34.8, 34.5, 34.4, 34.3, 28.4, 23.8, 23.6, 21.2. Anal. Calcd for C₁₈H₃₂N₂O₂: C, 70.09; H, 10.46; N, 9.08. Found: C, 70.18; H, 10.06; N, 9.02.

2-Methyl-3-[4-acetamido-(2,2,6,6-tetramethylpiperidin-1-yl)oxy]-1-octene (Table 1, entry 5). Recrystallization from hexane gave a white solid, mp 99-100 °C, in 86 % yield: ¹H NMR δ 5.33 (br s, 1H), 4.82 (br s, 1H), 4.80 (br s, 1H), 4.09 – 4.08 (m, 1H), 3.95 (dd, J = 3.6 Hz, J = 10.6 Hz, 1H), 1.92 (s, 3H), 1.78-1.72 (m, 3H), 1.65 (s, 3H), 1.50 – 1.48 (m, 1H), 1.32 – 1.19 [overlapping patterns, i.e. 1.32 – 1.23 (m, 12H), 1.19 (s, 3H), 1.18 (s, 3H)], 1.15 (s, 3H), 1.06 (s, 3H), 0.86 (t, J = 6.9 Hz, 3H); ¹³C NMR δ 169.5,

145.0, 113.8, 89.7, 60.3, 59.8, 46.7, 46.5, 41.3, 34.3, 34.1, 32.0, 31.4, 29.7, 29.5, 25.7, 23.7, 22.8, 21.1, 21.0, 17.5, 14.3. Anal. Calcd for $C_{22}H_{42}N_2O_2$: C, 72.08; H, 11.55; N, 7.64. Found: C, 72.00; H, 11.18; N, 7.65.

(E)-3-Ethyl-4-[4-acetamido(2,2,6,6-tetramethylpiperidin-1-yl)oxy]-2-pentene

(Table 1, entry 6). Isolated as a white solid, mp 118-119 °C, in 75 % yield: ^1H NMR δ 5.41 – 5.36 (m, 1H), 5.30 (br s, 1H), 4.16 – 4.11 (m, 2H), 2.15 – 2.00 (m, 2H), 1.95 (s, 3H), 1.81 – 1.73 (m, 2H), 1.61 (d, J = 4.0 Hz, 3H). 1.34 – 1.26 (m, 5H), 1.20 – 1.18 (m, 9H), 1.05 – 1.01 (m, 6H); ^{13}C NMR δ 169.5, 144.0, 120.8, 85.7, 66.0, 60.0, 46.6, 46.4, 41.4, 34.4, 31.1, 23.7, 21.2, 21.1, 20.5, 15.5, 14.2, 13.0. Anal. Calcd for $C_{18}H_{34}N_2O_2$: C, 69.63; H, 11.04; N, 9.02. Found: C, 69.15; H, 10.66; N, 9.27.

3,7-Dimethyl-6-[4-acetamido-(2,2,6,6-tetramethylpiperidinyloxy)]-7-octenoic acid

(Table 1, entry 7). Recrystallization from 95% ethanol gave a white solid, mp 82-83 °C, in 94 % yield: ^1H NMR δ 5.4 (br s, 1H), 4.90 – 4.88 (m, 2H), 4.17 – 4.15 (m, 1H), 4.06 – 4.03 (m, 1H), 2.42 – 2.32 (m, 1H), 2.22 – 2.09 (m, 1H), 2.02 – 1.92 [overlapping patterns, i.e. 2.02-1.92 (m, 1H), 1.99 (s, 3H)], 1.92 – 1.79 (m, 3H), 1.71 (s, 3H), 1.60 – 1.54 (m, 1H), 1.47 – 1.34 (m, 3H), 1.29 – 1.06 [overlapping patterns, i.e. 1.26 (s, 6H), 1.22 (s, 3H), 1.29 – 1.22 (m, 1H), 1.14 (s, 3H), 1.09 – 1.06 (m, 1H)], 1.01 (d, J = 6.5 Hz, 3H); ^{13}C NMR δ 178.2, 178.1, 169.7, 144.4, 114.4, 89.7, 41.9, 41.4, 41.3, 33.9, 32.7, 32.5, 30.4, 30.3, 28.1, 23.7, 21.1, 20.2, 19.8, 17.6, 17.4. HRMS-ES (m/z): $[M + H]^+$ calcd for $C_{21}H_{38}N_2O_4$, 383.2910; found, 383.2905.

2-Methyl-3-[4-acetamido-(2,2,6,6-tetramethylpiperidinyloxy)]-1,7-octadiene

(Table 1, entry 8). Flash chromatography on silica gel (hexane : EtOAc, 70:30) gave a white solid: R_f = 0.5 (hexane : EtOAc, 1:9), mp 98-100 °C: ^1H NMR δ 5.83 – 5.77 (m,

1H), 5.18 (br s, 1H), 5.02 – 4.94 (m, 2H), 4.86 – 4.85 (m, 2H), 4.13 – 4.11 (m, 1H), 4.01 – 4.00 (m, 1H), 2.05 – 1.99 (m, 2H), 1.95 (s, 3H), 1.82 – 1.74 (m, 3H), 1.69 (s, 3H), 1.55 – 1.53 (m, 1H), 1.35 – 1.19 (m, 13H), 1.09 (s, 3H); ¹³C NMR δ 169.5, 144.9, 139.1, 114.7, 114.0, 89.5, 60.4, 59.9, 46.7, 46.6, 41.4, 34.4, 34.2, 33.9, 31.0, 25.1, 23.8, 21.1, 17.6. Anal. Calcd for C₂₀H₃₆N₂O₂: C, 71.38; H, 10.78; N, 8.32. Found: C, 71.44; H, 10.53; N, 8.50.

Table 1, entry 1

Table 1, entry 1

Table 1, entry 2

Table 1, entry 2

Table 1, entry 3

Table 1, entry 3

Table 1, entry 4

Table 1, entry 4

Table 1, entry 5

Table 1, entry 6

Table 1, entry 6

Table 1, entry 7

Table 1, entry 7

Table 1, entry 8