This Appendix is for the paper “Extending the Mann-Whitney-Wilcoxon Rank Sum Test to Longitudinal Data Analysis with Covariates,” by Chen, R., Chen, T., Lu, N., Zhang, H., Wu, P., Feng, C. and Tu, X.M.

Appendix

We provide a sketch of the proof for Theorem 2 from which Theorem 1 follows as a special case.

Let $\hat{\eta}$ be the solution to the (31). Also, let $\mathbf{W}_n$ denote the normalized $\frac{1}{n}\mathbf{W}_n$. By a stochastic Taylor series expansion of the estimating equations in (25) and solving for $\hat{\eta} - \eta$, we obtain

$$\sqrt{n}(\hat{\eta} - \eta) = - \left(\frac{\partial}{\partial \eta} \mathbf{W}_n \right)^{-\top} \frac{\sqrt{n}}{n} \sum_{l=1}^K \sum_{i=1}^{n_l} \mathbf{W}_{n,li}(\eta_l) + \mathbf{o}_p(1),$$

where $\mathbf{o}_p(\cdot)$ denotes the stochastic $\mathbf{o}(\cdot)$ (see Kowalski and Tu, 2007, Chap. 1). We have

$$\begin{aligned} \frac{\partial}{\partial \eta} \mathbf{W}_n &= \frac{1}{n} \sum_{l=1}^K \sum_{i=1}^{n_l} \frac{\partial}{\partial \eta} \mathbf{W}_{n,li}(\eta_l) = \sum_{l=1}^K \frac{n_l}{n} \left(\frac{1}{n_l} \sum_{i=1}^{n_l} \frac{\partial}{\partial \eta} \mathbf{W}_{n,li}(\eta_l) \right) \\ &\rightarrow_p \sum_{l=1}^K \rho_l^2 E \left(\frac{\partial}{\partial \eta} \mathbf{W}_{n,li}(\eta_l) \right) = H. \end{aligned}$$

Thus, it follows that

$$\sqrt{n}(\hat{\eta} - \eta) = -H^{-1} \frac{\sqrt{n}}{n} \sum_{l=1}^K \sum_{i=1}^{n_l} \mathbf{W}_{n,li}(\eta_l) + \mathbf{o}_p(1).$$

By applying a Taylor expansion of multivariate U-statistic (Kowalski and Tu, 2007), we have:

$$\begin{aligned} \sqrt{n}\mathbf{U}_n &= - \left(\frac{\partial}{\partial \theta} \mathbf{U}_n \right)^{\top} \sqrt{n}(\hat{\theta} - \theta) - \left(\frac{\partial}{\partial \eta} \mathbf{U}_n \right)^{\top} \sqrt{n}(\hat{\eta} - \eta) - \\ &\quad - \left(\frac{\partial}{\partial \varphi} \mathbf{U}_n \right)^{\top} \sqrt{n}(\hat{\varphi} - \varphi) + \mathbf{o}_p(1). \end{aligned}$$

Since

$$\frac{\partial}{\partial \theta} \mathbf{U}_n = \sum_{k=2}^K \frac{1}{n_1 n_k} \sum_{\mathbf{i} \in C^{1k}} \frac{\partial}{\partial \theta} \mathbf{U}_{kn,\mathbf{i}}(\theta) = \sum_{k=2}^K \frac{1}{n_1 n_k} \sum_{\mathbf{i} \in C^{1k}} \frac{\partial}{\partial \theta} (D_{k\mathbf{i}} V_{k\mathbf{i}}^{-1} \Delta_{k\mathbf{i}} S_{k\mathbf{i}}),$$

it follows from the properties of U-statistics that

$$\begin{aligned} \frac{1}{n_1 n_k} \sum_{\mathbf{i} \in C^{1k}} \frac{\partial}{\partial \theta} \mathbf{U}_{kn,\mathbf{i}}(\theta) &\rightarrow_p E \left(\frac{\partial}{\partial \theta} \mathbf{U}_{kn,\mathbf{i}}(\theta) \right) = E \left(\frac{\partial}{\partial \theta} (D_{k\mathbf{i}} V_{k\mathbf{i}}^{-1} \Delta_{k\mathbf{i}} S_{k\mathbf{i}}) \right) \\ &= E \left(\left(\frac{\partial}{\partial \theta} S_{k\mathbf{i}} \right) \Delta_{k\mathbf{i}} V_{k\mathbf{i}}^{-1} D_{k\mathbf{i}}^{\top} \right) = -E \left(D_{k\mathbf{i}} \Delta_{k\mathbf{i}} V_{k\mathbf{i}}^{-1} D_{k\mathbf{i}}^{\top} \right). \end{aligned}$$

Since a linear combination of consistent estimates is again consistent, it follows that

$$\frac{\partial}{\partial \theta} \mathbf{U}_n \rightarrow_p \sum_{k=2}^K -E \left(D_{ki} \mathbf{\Delta}_{ki} V_{ki}^{-1} D_{ki}^\top \right) = -B.$$

Similarly,

$$\frac{\partial}{\partial \eta} \mathbf{U}_n \rightarrow_p E \left(\frac{\partial}{\partial \eta} \mathbf{U}_{n,i} \right) = \sum_{k=2}^K E \left(\left(\frac{\partial}{\partial \eta} (\mathbf{\Delta}_{ki} S_{ki}) \right) V_{ki}^{-1} D_{ki}^\top \right) = C.$$

Also, it can be shown that $\frac{\partial}{\partial \varphi} \mathbf{U}_n = \mathbf{o}_p(1)$. It follows from this and the $\sqrt{n}$ -consistency of $\hat{\varphi}$ that

$$\left(\frac{\partial}{\partial \varphi} \mathbf{U}_n \right)^\top \sqrt{n} (\hat{\varphi} - \varphi) = \mathbf{o}_p(1).$$

Now consider the U-statistics-like quantity $\sqrt{n} \mathbf{U}_n = \sqrt{n} \sum_{k=2}^K \frac{1}{n_1 n_k} \sum_{\mathbf{i} \in C^{1k}} \mathbf{U}_{kn,i}(\theta)$. It follows from the properties of the projection of U-statistics-like quantities that

$$\begin{aligned} \frac{\sqrt{n}}{n_1 n_k} \sum_{\mathbf{i} \in C^{1k}} \mathbf{U}_{kn,i}(\theta) &= \sqrt{n} \left(\frac{1}{n_1} \sum_{i=1}^{n_1} E(\mathbf{U}_{kn,i} | y_{1i}, \mathbf{x}_{1i}) + \frac{1}{n_k} \sum_{i=1}^{n_k} E(\mathbf{U}_{kn,i} | y_{ki}, \mathbf{x}_{ki}) \right) + \mathbf{o}_p(1) \\ &= \frac{\sqrt{n}}{\sqrt{n_1}} \frac{\sqrt{n_1}}{n_1} \sum_{i=1}^{n_1} E(\mathbf{U}_{kn,i} | y_{1i}, \mathbf{x}_{1i}) + \frac{\sqrt{n}}{\sqrt{n_k}} \frac{\sqrt{n_k}}{n_k} \sum_{i=1}^{n_k} E(\mathbf{U}_{kn,i} | y_{ki}, \mathbf{x}_{ki}) + \mathbf{o}_p(1). \end{aligned}$$

Thus, it follows that

$$\begin{aligned} \sqrt{n} \mathbf{U}_n &= \sum_{k=2}^K \frac{\sqrt{n}}{n_1 n_k} \sum_{\mathbf{i} \in C^{1k}} \mathbf{U}_{kn,i}(\theta) \\ &= \frac{\sqrt{n}}{\sqrt{n_1}} \frac{\sqrt{n_1}}{n_1} \sum_{k=2}^K E(\mathbf{U}_{kn,i} | y_{1i}, \mathbf{x}_{1i}) + \sum_{l=2}^K \frac{\sqrt{n}}{\sqrt{n_l}} \frac{\sqrt{n_l}}{n_l} E(\mathbf{U}_{kn,i} | y_{li}, \mathbf{x}_{li}) + \mathbf{o}_p(1) \\ &= \sum_{l=1}^K \frac{\sqrt{n}}{\sqrt{n_l}} \frac{\sqrt{n_l}}{n_l} \sum_{i=1}^{n_l} \mathbf{v}_{li} + \mathbf{o}_p(1). \end{aligned}$$

Thus, we have:

$$\begin{aligned} \sqrt{n} (\hat{\theta} - \theta) &= B^{-1} [\sqrt{n} \mathbf{U}_n + C \sqrt{n} (\hat{\eta} - \eta)] + \mathbf{o}_p(1) \\ &= B^{-1} \left[\sum_{l=1}^K \frac{\sqrt{n}}{\sqrt{n_l}} \frac{\sqrt{n_l}}{n_l} \sum_{i=1}^{n_l} \mathbf{v}_{li} - CH^{-1} \sum_{l=1}^K \frac{\sqrt{n}}{\sqrt{n_l}} \frac{\sqrt{n_l}}{n_l} \sum_{i=1}^{n_l} \mathbf{W}_{n,li}(\eta_l) \right] + \mathbf{o}_p(1) \\ &= B^{-1} \left[\sum_{l=1}^K \frac{\sqrt{n}}{\sqrt{n_l}} \frac{\sqrt{n_l}}{n_l} \sum_{i=1}^{n_l} (\mathbf{v}_{li} - CH^{-1} \mathbf{W}_{n,li}(\eta_l)) \right] + \mathbf{o}_p(1) \end{aligned} \tag{1}$$

By applying the central limit theorem to the above, we obtain the asymptotic normality of $\hat{\theta}$ with the asymptotic variance given in (32).