Supplementary material for Pupillo E. Extrapyrarnidal and cognitive signs in amyotrophic lateral sclerosis: A population based cross-sectional study. Amyotroph Lateral Scler Frontotemporal Degener. 2015; doi: 10.3109/21678421.2015.1040028.

Supplementary Table I. Selected baseline characteristics in the present sample and in an incidence study from the same origin populations.

A. LOMBARDIA Variable	Present sample (n = 135)		Incidence study* (n = 517)	
	n (%)	(95% CI)	n (%)	(95% CI)
Gender				
M	75 (60)	(51.7–68.3)	293 (56.7)	(52.4–61)
F	50 (40)	(31.7–48.3)	224 (43.3)	(39–47.6)
Onset				
Bulbar/generalized	49 (39.2)	(31.0–47.4)	182 (35.2)	(31.1–39.3)
Spinal	76 (60.8)	(52.6–69.0)	335 (64.8)	(60.7–68.9)
	Mean	Std (95% CI)	Mean	Std (95% CI)
Age at diagnosis (years)	66.1	11.1 (64.2–68.0)	63.6	NA NA
Disease duration at diagnosis (months)	11.9	9.5 (10.3–13.5)	10.6	9.4 (9.8–11.4)

95% CI: 95% confidence interval; Std: standard deviation.

M: male; F: female.

Reference: (18).

Supplementary Table II. Selected baseline characteristics in the present sample and in an incidence study from the same origin populations.

B. EMILIA ROMAGNA Variable	Present sample (n = 21)		Incidence study* (n = 347)	
	n (%)	(95% CI)	n (%)	(95%1 CI)
Gender				
M	16 (76.2)	(58.0–94.5)	187 (53.9)	(48.7–59.1)
F	5 (23.8)	(5.6–42.0)	160 (46.1)	(40.9–51.3)
Onset				
Bulbar/generalized	10 (47.6)	(26.2–69.0)	122 (35.2)	(30.2–40.2)
Spinal	11 (52.4)	(31.0–73.8)	225 (64.8)	(59.8–69.8)
	Mean	Std (95% CI)	Mean	Std (95% CI)
Age at diagnosis (years)	64.8	15.7 (57.7–71.9)	67.9	11.2 (66.7–69.1)
Disease duration at diagnosis (months)	11.4	11.6 (6.1–16.7)	12.4	11.9 (11.1–13.7)

95% CI: 95% confidence interval; Std: standard deviation.

M: male; F: female.

Reference: (19).