Supporting Information for:

A Facile Approach to 2-Acetamido-2-deoxy- β -D-Glucopyranosides via a Furanosyl Oxazoline

Ye Cai, Chang-Chun Ling and David R. Bundle*

Alberta Ingenuity Center for Carbohydrate Science, Department of Chemistry, University of Alberta,
Edmonton, AB T6G 2G2 Canada

Email Address: Dave.Bundle@ualberta.ca

General methods

Optical rotations were measured at 22 ± 2 °C. Analytical TLC was performed on Silica Gel 60-F₂₅₄ (Merck, Darmstadt) with detection by quenching of fluorescence and/or by charring with 5% sulfuric acid in water or with a ceric ammonium molybdate dip. All commercial reagents were used as supplied. *p*-TsOH was dehydrated by azeotropic distillation with benzene,¹ and the anhydrous acid was recrystallized from benzene and dried under vacuum at 50 °C. Column chromatography was performed on Silica Gel 60 (Silicycle, Ontario). ¹H NMR spectra were recorded at 300, 500, or 600 MHz. The first order proton chemical shifts δ_{H} are referenced to either residual CHCl₃ (δ_{H} 7.24, CDCl₃) or residual CD₂HOD (δ_{H} 3.30, CD₃OD), or internal acetone (δ_{H} 2.225, D₂O). *J*-values are given in Hz. The assignment of resonances for all compounds was made by two-dimensional homonuclear and heteronuclear chemical shift correlation experiments. Molecular sieves are stored in an oven at 150° C and flame-dried under vacuum before use. Organic solutions from extractions were dried with anhydrous Na₂SO₄ prior to concentration under vacuum at < 40 °C (bath). Microanalyses and electrospray mass spectroscopy were performed by the analytical services of this Department.

2-Methyl-(1,2-dideoxy-5,6-O-isopropylidene- α -D-glucofurano)-[2,1-*d*]-2-oxazoline (6)

The oxazoline **6** was prepared according to a literature procedure² but on a 30g scale. The ¹H NMR is identical to what was reported.²

1 Perrin, D. D.; Armarego, W. F. L. *Purification of Laboratory Chemicals*, Pergamon Press, 3rd Edition, Page 291

2 Furneaux, R. H.; Gainsford, G. J.; Lynch, G. P.; Yorke, S. C.; *Tetrahedron*, **1993**, 42, 9605-9612.

Methyl 2-acetamido-2-deoxy- β -D-glucopyranoside (8)

Oxazoline **6** (20.0 g) was dissolved in anhydrous methanol (500 mL), *p*-TsOH (0.2 eq.) was added, and the mixture was stirred at room temperature for 5 hours. The reaction was monitored by TLC and quenched with Et₃N to adjust the pH to 7.5. The solvent was removed under reduced pressure and the residue was washed several times with CH₂Cl₂ to afford a white solid (~19.0 g, yield 98%). $[\alpha]_{\text{D}}^{25} -39^{\circ}$ (*c* 1.1, CH₃OH); lit.³: $[\alpha]_{\text{D}}^{25} -42^{\circ}$ (*c* 1, H₂O). ¹H NMR (500 MHz, D₂O): δ 4.44 (d, 1H, *J* = 8.5 Hz, H-1), 3.93 (dd, 1H, *J* = 12.2, 2.0 Hz, H-6a), 3.76 (dd, 1H, *J* = 12.2, 5.6 Hz, H-6b), 3.69 (dd, 1H, *J* = 10.2, 8.5 Hz, H-2), 3.54 (dd, 1H, *J* = 10.2, 8.5 Hz, H-3), 3.50 (s, 3H, OMe), 3.46 (ddd, 1H, *J* = 10.0, 5.6, 2.2 Hz, H-5), 3.43 (dd, 1H, *J* = 10.0, 8.2 Hz, H-4), 2.03 (s, 3H, Ac); ¹³C NMR (125 MHz, D₂O): δ 175.56, 102.81, 76.77, 74.84, 70.83, 61.63, 57.93, 56.33, 23.03; HRMS *m/z* calcd. for C₉H₁₇NO₆Na (M+Na⁺) 258.0948, found 258.0948.

If the reaction was quenched with Dowex-2 (OH⁻) before it went to completion, compounds **7** and **9** could be isolated.

Methyl 2-acetamido-2-deoxy-5,6-O-isopropylidene- β -D-glucofuranoside (7)

$[\alpha]_{\text{D}}^{25} -36^{\circ}$ (*c* 0.9, CH₃OH) lit.² $[\alpha]_{\text{D}}^{25} -37^{\circ}$ (*c* 1.2, CH₃OH); ¹H NMR (500 MHz, CDCl₃): δ 5.51 (d, 1H, *J* = 6.1 Hz, NH), 4.84 (s, 1H, H-1), 4.39 (ddd, 1H, *J* = 7.9, 5.7, 5.7 Hz, H-5), 4.32 (d, 1H, *J* = 7.0 Hz, H-2), 4.21 (d, 1H, *J* = 4.9 Hz, H-3), 4.16 (dd, 1H, *J* = 8.5, 6.3 Hz, H-6a), 4.14 (dd, 1H, *J* = 7.6, 4.8 Hz, H-4), 4.02 (dd, 1H, *J* = 8.6, 5.4 Hz, H-6b), 3.39 (s, 3H, OMe), 1.98 (s, 3H, Ac), 1.44, 1.37 (s, 2 $\times$ 3H, isopropyl); ¹³C NMR (125 MHz, CDCl₃): δ 169.89, 109.34, 107.67, 83.30, 77.26, 77.00, 76.74, 75.17, 74.04, 66.99, 61.92, 55.58, 26.80, 25.22, 23.16; HRMS *m/z* calcd. for C₁₂H₂₁NO₆Na (M+Na⁺) 298.1261, found 298.1258.

Methyl 2-acetamido-2-deoxy- β -D-glucofuranoside (9)

$[\alpha]_{\text{D}}^{25} -54^{\circ}$ (*c* 1, CH₃OH), lit.³ $[\alpha]_{\text{D}}^{25} -56^{\circ}$ (*c* 1, CH₃OH); ¹H NMR (500 MHz, CD₃OD): δ 4.77 (d, 1H, *J* = 1.0 Hz, H-1), 4.17 (dd, 1H, *J* = 1.7, 4.8 Hz, H-3), 4.13 (s, 1H, *J* = 1.0 Hz, H-2), 4.05

(dd, 1H, $J = 4.8, 8.8$ Hz, H-4), 3.94 (ddd, 1H, $J = 3.0, 5.9, 8.8$ Hz, H-5), 3.82 (dd, 1H, $J = 3.0, 11.5$ Hz, H-6a), 3.64 (dd, 1H, $J = 5.9, 11.5$ Hz, H-6b), 3.34 (s, 3H, CH₃), 1.96 (s, 3H, Ac); ¹³C NMR (125 MHz, CD₃OD): δ 173.16, 109.32, 82.50, 76.13, 71.84, 65.22, 64.07, 55.57, 22.44; HRMS m/z calcd. for C₉H₁₇NO₆Na (M+Na⁺) 258.0948, found 258.0947.

General procedure for *b*-glycosylations with oxazoline 6

To a mixture of oxazoline **6** (1.0 mmol) in anhydrous alcohol (2.0 mL) or in a mixture of anhydrous alcohol and CH₂Cl₂, was added molecular sieves (4 Å, 300 mg), and anhydrous *p*-TsOH or CSA (0.5-1 eq.) was added. The mixture was stirred at room temperature overnight. To quench the reaction when the TLC indicated that reaction was finished, Dowex 2 (OH⁻) resin was added until the pH reached 7.0. After removing the solid by filtration, the reaction mixture was evaporated under reduced pressure. The residue was either washed several times with CH₂Cl₂ or purified by flash chromatography (CH₂Cl₂: CH₃OH 10:1) to afford a white solid.

Allyl 2-acetamido-2-deoxy-*b*-D-glucopyranoside

The title compound (931 mg) was prepared according to the general procedure and purified by washing with dichloromethane; yield 71%. If purified by chromatography, the yield was 82%. $[\alpha]_D^{25} -25^\circ$ (c 1.0, CH₃OH); lit.⁴: $[\alpha]_D^{25} -23^\circ$ (c 1.0, CH₃OH); ¹H NMR (500 MHz, D₂O) δ 5.90 (m, 1H, allyl), 5.30 (m, 1H, allyl), 5.26 (m, 1H, allyl), 4.56 (d, 1H, $J = 8.5$ Hz, H-1), 4.33 (m, 1H, allyl), 4.16 (m, 1H, allyl), 3.95 (dd, 1H, $J = 12.4, 1.2$ Hz, H-6a), 3.74 (high order m, 1H, H-6b), 3.70 (dd, 1H, $J = 10.3, 8.5$ Hz, H-2), 3.53 (high order m, 1H, H-3), 3.41-3.47 (m, 2H, H-4 + H-5), 2.02 (s, 3H, Ac); ¹³C NMR (125 MHz, D₂O): δ 175.49, 134.24, 119.02, 100.95, 76.76, 74.75, 71.30, 70.81, 61.63, 56.41, 32.01; HRMS m/z calcd. for C₁₁H₁₉NO₆Na (M+Na⁺) 284.1105, found 284.1105.

Benzyl 2-acetamido-2-deoxy-*b*-D-glucopyranoside

The title compound (955 mg) was prepared according to the general procedure, and purified by washing with dichloromethane, yield 60%. If purified by chromatography, the yield was 73%. $[\alpha]_D^{25} -29^\circ$ (c 0.9, CH₃OH); lit.⁵: $[\alpha]_D^{25} -50^\circ$ (c 0.3, H₂O); lit.⁶: $[\alpha]_D^{25} -29^\circ$ (c 0.93, H₂O); ¹H

(4) Gan, Z.; Cao, S.; Wu, Q.; Roy, R.J. *Carbohydr. Chem.* **1999**, *18*, 755-774.

(5) Vikha, I. V.; Sakharovsky, V. G.; Bystrov, V. F.; Khorlin, A. Ya. *Carbohydr. Res.* **1972**, *25*, 143-152.

NMR (600 MHz, D₂O) δ 7.43-7.50 (m, 3H, Bn), 7.40 (m, 2H, Bn), 4.90 (d, 1H, J = 12.2, Bn), 4.69 (d, 1H, J = 12.2 Hz, Bn), 4.55 (d, 1H, J = 8.5 Hz, H-1), 3.97 (dd, 1H, J = 12.4, <1.0 Hz, H-6a), 3.79 (high order m, 1H, H-6b), 3.72 (dd, 1H, J = 9.0, 9.0 Hz, H-2), 3.45-3.52 (m, 3H, H-3 + H-4 + H-5), 1.95 (s, 3H, Ac); ¹³C NMR (125 MHz, D₂O): δ 175.37, 137.66, 129.61, 129.47, 129.33, 100.80, 76.80, 74.62, 72.32, 70.80, 61.68, 56.44, 23.01; HRMS m/z calcd. for C₁₅H₂₁NO₆Na (M+Na⁺) 334.1261, found 334.1257.

Octyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (206 mg) was prepared according to the general procedure and purified by chromatography, yield 69%. $[\alpha]_D^{25}$ -22.8° (c 1.0, CH₃OH); lit.⁷: $[\alpha]_D^{25}$ -19.4° (c 2.8, pyridine); ¹H NMR (500 MHz, CD₃OD): δ 4.38 (d, 1H, J = 8.4 Hz, H-1), 3.84-3.90 (m, 2H, H-6a + OCH_aCH_b), 3.67 (dd, 1H, J = 11.9, 5.7 Hz, H-6b), 3.62 (dd, 1H, J = 10.3, 8.4 Hz, H-2), 3.42-3.47 (m, 2H, H-3 + OCH_aCH_b), 3.30 (t, 1H, J = 9.1 Hz, H-4), 3.24 (ddd, 1H, J = 9.7, 5.7, 2.3 Hz, H-5), 1.96 (s, 3H, Ac), 1.50-1.56 (m, 2H, CH₂), 1.25-1.36 (m, 10H, octyl), 0.89 (t, 3H, J = 6.8 Hz, octyl); ¹³C NMR (125 MHz, CD₃OD): δ 173.62, 102.73, 77.70, 76.10, 70.61, 62.85, 57.49, 49.53, 33.03, 30.70, 30.51, 30.50, 27.16, 23.74, 23.01, 14.43; HRMS m/z calcd. for C₁₆H₃₁NO₆Na (M+Na⁺) 356.2044, found 356.2045.

n-Pentenyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (846 mg) was prepared according to the general procedure, and purified by washing with dichloromethane, yield 64%. If purified by chromatography, the yield was 76%. $[\alpha]_D^{25}$ -27° (c 0.9, CH₃OH); ¹H NMR (600 MHz, D₂O): δ 5.90 (m, 1H, alkene), 5.08 (m, 1H, alkene), 5.02 (m, 1H, alkene), 4.51 (d, 1H, J = 8.5 Hz, H-1), 3.88-3.94 (m, 2H, H-6a + OCH_aCH_b), 3.74 (high order m, 1H, H-6b), 3.68 (dd, 1H, J = 10.4, 8.5 Hz, H-2), 3.61 (ddd, 1H, J = 10.3, 6.4, 6.4 Hz, OCH_aCH_b), 3.54 (high order m, 1H, H-3), 3.41-3.46 (m, 2H, H-4 + H-5), 2.05-2.12 (m, 2H, pent), 2.04 (s, 3H, Ac), 1.65 (m, 2H, pent); ¹³C NMR (125 MHz, D₂O): δ 175.36, 139.61, 115.68, 101.97, 76.72, 74.70, 70.81, 70.53, 61.63, 56.49, 30.14, 28.75, 23.06; HRMS m/z calcd. for C₁₃H₂₃NO₆Na (M+Na⁺) 312.1417, found 312.1416.

2-Hydroxyethyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (216 mg) was prepared according to the general procedure and purified by

(6) Arita, H.; Fukukawa, K.; Matsushima, Y. *Bull. Chem. Soc. Jpn.* **1972**, *45*, 3614-3619.

(7) Boullanger, P.; Chevalier, Y.; Croizier, M.-C.; Lafont D.; Sancho, M. R. *Carbohydr. Res.* **1995**, *278*, 91-102.

chromatography, yield 78%. $[\alpha]_D^{25} -15.6^\circ$ (c 1.0, CH_3OH); ^1H NMR (600 MHz, D_2O): δ 4.55 (d, 1H, $J = 8.5$ Hz, H-1), 3.90-3.94 (m, 2H, H-6a + OCH_aCH_b), 3.69-3.76 (m, 5H, H-2 + H-6b + OCH_aCH_b + $\text{OCH}_a\text{CH}_b\text{CH}_c\text{H}_d\text{OH}$ + $\text{OCH}_a\text{CH}_b\text{CH}_c\text{H}_d\text{OH}$), 3.54 (high order m, 1H, H-3), 3.42-4.38 (m, 2H, H-4 + H-5), 2.12 (s, 3H, Ac); ^{13}C NMR (125 MHz, D_2O): δ 175.67, 102.09, 76.73, 74.74, 71.97, 70.79, 61.60, 61.50, 56.45, 23.04; HRMS m/z calcd. for $\text{C}_{10}\text{H}_{19}\text{NO}_7\text{Na}$ ($\text{M}+\text{Na}^+$) 288.1054, found 288.1053.

4-Hydroxy-(Z)-but-2-enyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (214 mg) was prepared according to the general procedure and purified by chromatography, yield 77%. $[\alpha]_D^{25} -19.5^\circ$ (c 1.0, CH_3OH); ^1H NMR (500 MHz, D_2O): δ 5.84 (m, 1H, alkene), 5.67 (m, 1H, alkene), 4.54 (d, 1H, $J = 8.5$ Hz, H-1), 4.35 (m, 1H, $\text{OCH}_a\text{H}_b\text{CH}=\text{CH}$), 4.30 (m, 1H, $\text{OCH}_a\text{H}_b\text{CH}=\text{CH}$), 4.15-4.20 (m, 2H, $\text{CH}=\text{CHCH}_2\text{OH}$), 3.92 (dd, 1H, $J = 12.4$, 1.3 Hz, H-6a), 3.74 (high order m, 1H, H-6b), 3.68 (dd, 1H, $J = 10.3$, 8.6 Hz, H-2), 3.53 (high order m, 1H, H-3), 3.42-3.47 (m, 2H, H-4 + H-5), 2.03 (s, 3H, Ac); ^{13}C NMR (125 MHz, D_2O): δ 175.46, 133.71, 127.53, 100.67, 76.81, 74.76, 70.77, 65.31, 61.61, 57.99, 56.37, 23.01; HRMS m/z calcd. for $\text{C}_{12}\text{H}_{21}\text{NO}_7\text{Na}$ ($\text{M}+\text{Na}^+$) 314.1210, found 314.1209.

2-Allyloxyethyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (226 mg) was prepared according to the general procedure and purified by chromatography, yield 74%. $[\alpha]_D^{25} -24^\circ$ (c 1.0, CH_3OH); ^1H NMR (500 MHz, D_2O): δ 5.93 (m, 1H, allyl), 5.32 (m, 1H, allyl), 5.26 (m, 1H, allyl), 4.55 (d, 1H, $J = 8.5$ Hz, H-1), 4.03-4.06 (m, 2H, allyl), 4.00 (ddd, 1H, $J = 11.8$, 5.7, 2.9 Hz, $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{O}$), 3.92 (dd, 1H, $J = 12.2$, 1.5 Hz, H-6a), 3.62-3.78 (m, 5H, H-2 + H-6b + $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{O}$ + $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{O}$ + $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{O}$), 3.53 (high order m, 1H, H-3), 3.41-3.46 (m, 2H, H-4 + H-5), 2.03 (s, 3H, Ac); ^{13}C NMR (125 MHz, D_2O): δ 175.38, 134.62, 119.05, 101.92, 76.72, 74.72, 72.60, 70.79, 69.82, 69.61, 56.40, 23.11; HRMS m/z calcd. for $\text{C}_{13}\text{H}_{23}\text{NO}_7\text{Na}$ ($\text{M}+\text{Na}^+$) 328.1367, found 328.1369.

2-Chloroethyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (168 mg) was prepared according to the general procedure and purified by chromatography, yield 65%. $[\alpha]_D^{25} -22^\circ$ (c 0.9, CH_3OH); ^1H NMR (500 MHz, D_2O): δ 4.58 (d, 1H, $J = 8.5$ Hz, H-1), 4.12 (ddd, 1H, $J = 12.0$, 5.8, 3.7 Hz, $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{Cl}$), 3.92 (dd, 1H, $J =$

12.0, 1.4 Hz, H-6a), 3.88 (ddd, 1H, $J = 12, 7.0, 3.7$ Hz, $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{Cl}$), 3.66-3.76 (m, 4H, H-2 + H-6b + $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{Cl}$ + $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{Cl}$), 3.55 (high order m, 1H, H-3), 3.42-3.48 (m, 2H, H-4 + H-5), 2.02 (s, 3H, Ac); ^{13}C NMR (500 MHz, D_2O): δ 175.69, 102.22, 76.80, 71.13, 70.75, 61.58, 56.36, 44.28, 23.13; HRMS m/z calcd. for $\text{C}_{10}\text{H}_{18}\text{NO}_6\text{ClNa}$ ($\text{M}+\text{Na}^+$) 306.0715, found 306.0715.

2-Bromoethyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (181 mg) was prepared according to the general procedure and purified by chromatography, yield 63%. $[\alpha]_{\text{D}}^{25} -9.4^\circ$ (c 1.1, CH_3OH); ^1H NMR (500 MHz, D_2O): δ 4.57 (d, 1H, $J = 8.5$ Hz, H-1), 4.18 (ddd, 1H, $J = 12.2, 6.0, 4.0$ Hz, $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{Br}$), 3.88-3.94 (m, 2H, $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{Br}$ + H-6a), 3.73 (high order m, 1H, H-6b), 3.70 (dd, 1H, $J = 10.4, 8.5$ Hz, H-2), 3.49-3.60 (m, 3H, H-3 + $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{Br}$ + $\text{OCH}_a\text{H}_b\text{CH}_c\text{H}_d\text{Br}$), 3.41-3.47 (m, 2H, H-4 + H-5), 2.04 (s, 3H, Ac); ^{13}C NMR (125 MHz, D_2O): δ 175.64, 102.08, 76.79, 74.53, 70.86, 70.75, 61.58, 56.33, 32.27, 23.28; HRMS m/z calcd. for $\text{C}_{10}\text{H}_{18}\text{NO}_6\text{NaBr}$ ($\text{M}+\text{Na}^+$) 350.0210, found 350.0206.

6-Chlorohexyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (216 mg) was prepared according to the general procedure and purified by chromatography, yield 72%. $[\alpha]_{\text{D}}^{25} -20^\circ$ (c 0.7, $\text{CH}_3\text{OH}:\text{CHCl}_3 = 1:1$, v/v); ^1H NMR (600 MHz, D_2O): δ 4.50 (d, 1H, $J = 8.5$ Hz, H-1), 3.92 (dd, 1H, $J = <1.0, 13.4$ Hz, H-6a), 3.90 (ddd, 1H, $J = 10.5, 6.0, 6.0$ Hz, OCH_aH_b), 3.74 (high order m, 1H, H-6b), 3.67 (dd, 1H, $J = 10.4, 8.8$ Hz, H-2), 3.63 (t, 2H, $J = 6.7$ Hz, CH_2Cl), 3.60 (ddd, 1H, $J = 10.2, 6.4, 6.4$ Hz, OCH_aH_b), 3.54 (high order m, 1H, H-3), 3.41-3.44 (m, 2H, H-4 + H-5), 2.04 (s, 3H, Ac), 1.74-1.80 (m, 2H, CH_2), 1.54-1.60 (m, 2H, CH_2), 1.31-1.43 (m, 4H, $2 \times \text{CH}_2$); ^{13}C NMR (125 MHz, D_2O): δ 175.31, 102.00, 76.73, 74.72, 71.19, 70.82, 61.66, 56.51, 46.57, 32.71, 29.28, 26.50, 25.23, 23.08; HRMS m/z calcd. for $\text{C}_{14}\text{H}_{26}\text{NO}_6\text{ClNa}$ ($\text{M}+\text{Na}^+$) 362.1341, found 362.1342.

6-Azidohexyl 2-acetamido-2-deoxy-b-D-glucopyranoside

The title compound (204 mg) was prepared according to the general procedure and purified by chromatography, yield 70%. $[\alpha]_{\text{D}}^{25} -15^\circ$ (c 0.8, CH_3OH); ^1H NMR (600 MHz, D_2O): δ 4.50 (d, 1H, $J = 8.5$ Hz, H-1), 3.92 (dd, 1H, $J = 12.1, 1.8$ Hz, H-6a), 3.90 (ddd, 1H, $J = 12.0, 6.0, 6.0$ Hz, OCH_aH_b), 3.74 (high order m, 1H, H-6b), 3.67 (dd, 1H, $J = 10.3, 8.5$ Hz, H-2), 3.60 (ddd, 1H, $J = 10.2, 6.5, 6.5$ Hz, OCH_aH_b), 3.53 (high order m, 1H, H-3), 3.41-3.47 (m, 2H, H-4 + H-5), 3.32

(t, 2H, $J = 6.9$ Hz, CH_2N_3), 1.54-1.6m (m, 4H, $2 \times \text{CH}_2$), 1.31-1.41 (m, 4H, $2 \times \text{CH}_2$), 2.04 (s, 3H, Ac); ^{13}C NMR (125 MHz, D_2O): δ 175.31, 102.00, 76.74, 74.72, 71.21, 70.82, 61.66, 56.51, 52.03, 29.31, 28.85, 26.46, 25.53, 23.06; HRMS m/z calcd. for $\text{C}_{14}\text{H}_{26}\text{N}_4\text{O}_6\text{Na}$ ($\text{M}+\text{Na}^+$) 369.1745, found 369.1747.

2-(Trimethylsilyl)ethyl 2-acetamido-2-deoxy- β -D-glucopyranoside

The title compound (196 mg) was prepared according to the general procedure and purified by chromatography, yield 61%. $[\alpha]_{\text{D}}^{25} -23.4^\circ$ (c 0.9, CH_3OH); ^1H NMR (500 MHz, D_2O): δ 4.70 (d, 1H, $J = 8.5$ Hz, H-1), 4.05 (ddd, 1H, $J = 10.3, 10.3, 4.9$ Hz, OCH_2H_b), 3.91 (dd, 1H, $J = 12.5, 1.8$ Hz, H-6a), 3.75 (high order m, 1H, H-6b), 3.65-3.73 (m, 2H, $\text{OCH}_2\text{H}_b + \text{H-2}$), 3.52 (high order m, 1H, H-3), 3.42-3.48 (m, 2H, H-4 + H-5), 2.03 (s, 3H, Ac), 0.98 (ddd, 1H, $J = 13.8, 10.6, 7.3$ Hz, $\text{OCH}_2\text{H}_b\text{CH}_2\text{H}_d\text{Si}$), 0.88 (ddd, 1H, $J = 13.8, 9.8, 4.9$ Hz, $\text{OCH}_2\text{H}_b\text{CH}_2\text{H}_d\text{Si}$), 0.3 (s, 9H, SiMe_3); ^{13}C NMR (125 MHz, D_2O): δ 175.28, 101.21, 76.77, 75.04, 70.80, 69.12, 61.66, 56.48, 23.14, 18.02, -1.52; HRMS m/z calcd. for $\text{C}_{13}\text{H}_{27}\text{NO}_6\text{SiNa}$ ($\text{M}+\text{Na}^+$) 344.1500, found 344.1498.

Cyclohexyl 2-acetamido-2-deoxy- β -D-glucopyranoside

The title compound (299 mg) was prepared according to the general procedure and purified by chromatography, yield 73%. $[\alpha]_{\text{D}}^{25} -37.5^\circ$ (c 1.0, CH_3OH); lit.⁸: $[\alpha]_{\text{D}}^{25} -37^\circ$ (c 0.67, MeOH); ^1H NMR (500 MHz, D_2O): δ 4.62 (d, 1H, $J = 8.5$ Hz, H-1), 3.91 (dd, 1H, $J = 12.3, 1.6$ Hz, H-6a), 3.72-3.80 (m, 2H, H-6b + OCH), 3.65 (dd, 1H, $J = 10.4, 8.4$ Hz, H-2), 3.55 (high order m, 1H, H-3), 3.41-3.47 (m, 2H, H-4 + H-5), 2.02 (s, 3H, Ac), 1.20-1.90 (m, 10H, cyclohexyl); ^{13}C NMR (125 MHz, D_2O): δ 175.69, 102.22, 76.80, 74.54, 71.13, 70.75, 61.58, 56.36, 44.28, 23.13; HRMS m/z calcd. for $\text{C}_{14}\text{H}_{25}\text{NO}_6\text{Na}$ ($\text{M}+\text{Na}^+$) 326.1574, found 326.1573.

Isopropyl 2-acetamido-2-deoxy- β -D-glucopyranoside

The title compound (908 mg) was obtained as a mixture contaminated with α -isomer (α/β : 1/6) according to the general procedure and purified by washing with dichloromethane, yield 66%. If purified by chromatography, the yield was 77%. ^1H NMR (500 MHz, D_2O): δ 4.59 (d, 1H, $J = 8.4$ Hz, H-1), 4.01 (sept, 1H, $J = 6.2$ Hz, OCHMe_2), 3.91 (dd, 1H, $J = 12.2, 2.0$ Hz, H-6a), 3.73

(8) Zemlyakov, A. E.; Tsikalov, V. V.; Kalyuzhin, O. V.; Kur'yanov, V. O.; Chirva, V. Ya. *Russ. J. Bioorg. Chem (Eng. Transl.)*. **2003**, 29, 286-292.

(dd, 1H, $J = 12.2, 5.6$ Hz, H-6b), 3.62 (dd, 1H, $J = 10.4, 8.4$ Hz, H-2), 3.53 (high order m, 1H, H-3), 3.39-3.46 (m, 2H, H-4 + H-5), 2.03 (s, 3H, Ac), 1.19 (d, 3H, $J = 6.2$ Hz, isopropyl), 1.12 (d, 3H, $J = 6.2$ Hz, isopropyl); ^{13}C NMR (125 MHz, D_2O): δ 175.42, 100.52, 76.72, 74.77, 74.52, 70.83, 61.68, 56.73, 23.16, 22.99, 21.99; HRMS m/z Calcd. for $\text{C}_{11}\text{H}_{21}\text{NO}_6\text{Na}$ ($\text{M}+\text{Na}^+$) 286.1261, found, 286.1264.

***t*-Butyl 2-acetamido-2-deoxy-*b*-D-glucopyranoside**

The title compound (134 mg) was obtained by chromatography as a mixture contaminated with α -isomer (α/β : 1/4) according to the general procedure, yield 51%. ^1H NMR (500 MHz, D_2O): δ 4.66 (d, 1H, $J = 8.0$ Hz, H-1), 3.87 (dd, 1H, $J = 12.3, 2.2$ Hz, H-6a), 3.69 (dd, 1H, $J = 12.3, 5.8$ Hz, H-6b), 3.60 (dd, 1H, $J = 10.4, 8.0$ Hz, H-2), 3.55 (dd, 1H, $J = 10.4, 8.1$ Hz, H-3), 3.43 (ddd, 1H, $J = 9.9, 5.9, 2.3$ Hz, H-5), 3.39 (dd, 1H, $J = 9.9, 8.1$ Hz, H-4), 2.03 (s, 3H, Ac), 1.22 (s, 9H, $3 \times \text{CH}_3$, *t*-butyl); ^{13}C NMR (125 MHz, D_2O): δ 175.33, 96.43, 78.16, 76.45, 74.73, 71.00, 61.83, 56.95, 28.43, 23.06; HRMS m/z calcd. for $\text{C}_{12}\text{H}_{23}\text{NO}_6\text{Na}$ ($\text{M}+\text{Na}^+$) 300.1418, found 300.1419.

600 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm), temp 28.0 C -> actual temp = 27.0 C, id600 probe

Pulse Sequence: s2pul

Crude

125 MHz APT in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apt

500 MHz 1D in D₂O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.24 ppm), temp 29.4 C -> actual temp = 27.0 C, sw500u probe

Pulse Sequence: s2pul

125 MHz APT in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apt

500 MHz 1D in CD3OD (ref. to CD3OD @ 3.30 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in CD3OD (ref. to CD3OD @ 49.0 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH2 same, CH & CH3 opposite side of solvent signal

Pulse Sequence: apt

500 MHz 1D in D₂O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

600 MHz 1D in D₂O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 28.0 C -> actual temp = 27.0 C, id600 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in CD3OD (ref. to CD3OD @ 3.30 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in CD3OD (ref. to CD3OD @ 49.0 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH2 same, CH & CH3 opposite side of solvent signal

Pulse Sequence: apt

600 MHz 1D in D2O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 28.0 C -> actual temp = 27.0 C, id600 probe

Pulse Sequence: s2pul

125 MHz APT in D2O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH2 opposite side of CH & CH3

Pulse Sequence: apt

600 MHz 1D in D₂O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 28.0 C -> actual temp = 27.0 C, id600 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in D₂O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D2O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH2 opposite side of CH & CH3

Pulse Sequence: apt

500 MHz 1D in D₂O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in D₂O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in D2O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

600 MHz 1D in D2O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 28.0 C -> actual temp = 27.0 C, id600 probe

Pulse Sequence: s2pul

Pulse Sequence: apt

600 MHz 1D in D2O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 28.0 C -> actual temp = 27.0 C, id600 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in D₂O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in D2O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in D2O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt

500 MHz 1D in D2O (ref. to 0.1 % ext. acetone @ 2.225 ppm), temp 27.2 C -> actual temp = 27.0 C, sw500 probe

Pulse Sequence: s2pul

125 MHz APT in D₂O (ref. to 1% acetone @ 31.07 ppm), temp 27.2 C -> actual temp = 27.0 C, sw probe
C & CH₂ opposite side of CH & CH₃

Pulse Sequence: apt