Supporting Information

Facile Construction of the Benzofuran and Chromene Ring Systems via Pd^{II}-catalyzed Oxidative Cyclization

So Won Youn* and Jeong Im Eom

Department of Chemistry, Pukyong National University, Busan, 608-737, Korea

General

Nuclear Magnetic Resonance spectra were recorded on JEOL 400 Fourier transform NMR spectrometers. Spectra were recorded in CDCl₃ solutions referenced to TMS or solvent residual peak. IR spectra were taken as neat for liquids on NaCl plates using a Perkin-Elmer Spectrum 2000 FT-IR Spectrophotometer. High Resolution Mass Spectra were obtained on a JEOL JMS-700 mass spectrometer. Flash chromatography was performed on MERCK silica gel 60(230-400 mesh). All catalysts bought from Strem or Aldrich and used as received. Unless otherwise noted, all commercially obtained reagents were used as received. THF was distilled from sodium benzophenone ketyl immediately prior to use. Toluene and CH₂Cl₂ were all distilled from CaH₂ immediately prior to use. Reaction temperatures were controlled by an IKAmag temperature modulator. TLC was conducted with E. Merck silica gel 60 F254 precoated plates (0.25 mm) and visualized via UV and anisaldehyde staining.

Systematic Screen

Reactions were conducted at 0.015 M concentration in sealed vials under argon. To a solution of **1** (or **20**) in solvent were added catalyst, base(1 equiv.), and oxidant (1 equiv.). The resulting mixture was heated with stirring for the reported time at the reported temperature. After the reaction was completed (by TLC), the mixture was cooled, quenched with distilled water, extracted with CH₂Cl₂ (three times), dried over MgSO₄, and concentrated in vacuo. Yields were determined by ¹H NMR using trichloroethylene as an internal standard.

Table 1. Optimization Studies for the Cyclization of Compound **1**.

Entry	Catalyst (mol%)	Oxidant	Base	Solvent	Temp. (°C)	Time (h)	Yield (%)
1	Pd(MeCN) ₂ Cl ₂ (100)	-	-	dioxane	rt	4	30
2	Pd(MeCN) ₂ Cl ₂ (100)	-	Cs ₂ CO ₃	dioxane	rt	4	54
3	Pd(MeCN) ₂ Cl ₂ (100)	-	Cs ₂ CO ₃	MeCN	rt	4	3
4	Pd(MeCN) ₂ Cl ₂ (100)	-	Cs ₂ CO ₃	THF	rt	4	13
5	Pd(MeCN) ₂ Cl ₂ (100)	-	Cs ₂ CO ₃	Toluene	rt	4	-
6	Pd(MeCN) ₂ Cl ₂ (100)	-	Cs ₂ CO ₃	CH ₂ Cl ₂	rt	4	43
7	Pd(MeCN) ₂ Cl ₂ (20)	-	-	dioxane	rt	24	18
8	Pd(MeCN) ₂ Cl ₂ (20)	BQ	-	dioxane	rt	5	54
9	Pd(OAc) ₂ (20)	BQ	-	dioxane	rt	5	-
10	Pd(PhCN) ₂ Cl ₂ (20)	BQ	-	dioxane	rt	5	50
11	Pd(PPh ₃) ₂ Cl ₂ (20)	BQ	-	dioxane	rt	5	-
12	PdCl ₂ (20)	BQ	-	dioxane	rt	5	trace
13	PtCl ₂ (20)	BQ	-	dioxane	rt	5	-
14	RuCl ₃ (20)	BQ	-	dioxane	rt	5	-
15	Pd(MeCN) ₂ Cl ₂ (20)	BQ	NaOAc	dioxane	rt	5	trace
16	Pd(MeCN) ₂ Cl ₂ (20)	BQ	Li ₂ CO ₃	dioxane	rt	5	21
17	Pd(MeCN) ₂ Cl ₂ (20)	BQ	Na ₂ CO ₃	dioxane	rt	5	60
18	Pd(MeCN) ₂ Cl ₂ (20)	BQ	K ₂ CO ₃	dioxane	rt	5	48
19	Pd(MeCN) ₂ Cl ₂ (20)	BQ	Cs ₂ CO ₃	dioxane	rt	5	-
20	Pd(MeCN) ₂ Cl ₂ (20)	BQ	NEt ₃	dioxane	rt	5	-
21	Pd(MeCN) ₂ Cl ₂ (20)	CuCl ₂	Na ₂ CO ₃	dioxane	rt	5	trace
22	Pd(MeCN) ₂ Cl ₂ (20)	Cu(OAc) ₂	Na ₂ CO ₃	dioxane	rt	5	5
23	Pd(MeCN) ₂ Cl ₂ (10)	BQ	Na ₂ CO ₃	dioxane	rt	24	37
24	Pd(MeCN) ₂ Cl ₂ (10)	BQ	Na ₂ CO ₃	dioxane	80	0.5	52
25	Pd(MeCN) ₂ Cl ₂ (5)	BQ	Na ₂ CO ₃	dioxane	80	1	53
26	Pd(MeCN) ₂ Cl ₂ (5)	BQ	Na ₂ CO ₃	dioxane	65	5	66
27	Pd(MeCN) ₂ Cl ₂ (2)	BQ	Na ₂ CO ₃	dioxane	80	12	trace

Table 2. Optimization Studies for the Cyclization of Compound **20**.^a

Entry	Catalyst	Oxidant	Base	Time (h)	Yield (%) ^b
1	Pd(MeCN) ₂ Cl ₂	BQ	-	0.5	16
2	Pd(MeCN) ₂ Cl ₂	-	Na ₂ CO ₃	24	-
3	-	BQ	Na ₂ CO ₃	24	-
4	Pd(MeCN) ₂ Cl ₂	BQ	Na ₂ CO ₃	3	77
5	Pd(OAc) ₂	BQ	Na ₂ CO ₃	24	-
6	PdCl ₂	BQ	Na ₂ CO ₃	24	28

^a All reactions were carried out with catalyst (5 mol%), Na₂CO₃ (1 equiv), and BQ (1 equiv) in dioxane (0.015 M) at rt. ^b Determined by ¹H NMR using trichloroethylene as an internal standard.

General procedure for the preparation of allyl aryl ethers

To a solution of the corresponding phenol in DMF were added K₂CO₃ (2 equiv.) and allyl bromide (1.1 equiv.) at 0 °C. The resulting homogeneous solution was warmed up to rt. After the reaction was completed, the reaction mixture was quenched with distilled water, extracted with ether (three times), washed with brine, dried over MgSO₄, and concentrated in vacuo. The residue was purified by column chromatography on silica gel by eluting EtOAc-Hexane (1:3 for **1**, 1:5 for **13**, 1:20 ~ 1:50 for others) to afford the corresponding product (31-88%).

Spectral data for 3-allyloxyphenol (**1**),¹ 3-allyloxylanisole (**3**),^{1,2} 3,5-dimethoxyallyloxybenzene (**5**),² 3,5-dimethylallyloxybenzene (**7**),² 2-allyloxynaphthalene (**9**),¹ 3,4-methylenedioxyallyloxybenzene (**11**),^{2,3} and 3,4,5-trimethoxyallyloxybenzene (**15**)² were consistent with data reported in the literature.

3-Allyloxy-2-methylphenol (**13**)

an orange oil (43%, EtOAc:Hexane = 1:5).

δ_{H} (CDCl₃, 400MHz) 2.15 (s, 3H), 4.53 (td, J = 1.5, 5.1, 2H), 4.65 (s, 1H), 5.27 (qd, J = 1.4, 10.6, 1H), 5.42 (qd, J = 1.7, 17.3, 1H), 6.07 (m, 1H), 6.45 (d, J = 4.1, 1H), 6.47 (d, J = 4.1, 1H), 7.00 (t, J = 8.2,

¹ Gozzo, F. C.; Fernandes, S. A.; Rodrigues, D. C.; Eberlin, M. N.; Marsaioli, A. J. *J. Org. Chem.* **2003**, 68, 5493.

² Taskinen, E. *J. Chem. Soc., Perkin Trans. 2* **2001**, 1824.

³ Baxendale, I. R.; Lee, A.-L.; Ley, S. V. *J. Chem. Soc., Perkin Trans. 1* **2002**, 1850.

1H). δ_C (CDCl₃, 100MHz) 8.07, 69.09, 104.42, 108.12, 112.53, 116.92, 126.33, 133.56, 154.47, 157.63. $\nu_{\max}$ (NaCl)/cm⁻¹ 3419, 3084, 2926, 1595, 1471, 1362, 1096, 925, 769. HREIMS m/z 164.0840 (M)⁺, calcd for C₁₀H₁₂O₂ 164.0837.

(3,5-Dimethoxyphenoxy)-2-butene (17)

a pale brown oil (31%, olefin isomer mixture 3:1, EtOAc:Hexane = 1:30).

Signals corresponding to the major isomer: δ_H (CDCl₃, 400MHz) 1.75 (d, J = 6.5, 3H), 3.76 (s, 6H), 4.41 (d, J = 5.8, 2H), 5.72 (m, 1H), 5.85 (m, 1H), 6.10 (m, 3H). Representative signals corresponding to the minor isomer: δ_H (CDCl₃, 400MHz) 1.60 (d, J = 6.5, 3H), 3.78 (s, 3H), 3.79 (s, 3H), 4.55 (d, J = 6.5, 2H), 5.43 (m, 1H), 5.53 (m, 1H). δ_C (CDCl₃, 100MHz) 17.84, 55.29, 68.74, 92.98, 93.56, 125.89, 130.73, 160.57, 161.45. $\nu_{\max}$ (NaCl)/cm⁻¹ 2938, 1606, 1457, 1379, 1204, 1064, 967, 818. HREIMS m/z 208.1096 (M)⁺, calcd for C₁₂H₁₆O₃ 208.1099.

General procedure for the preparation of aryl homoallyl ethers.

A solution of the 3-buten-1-ol, the corresponding phenol (3 equiv.) and PPh₃ (1.3 equiv.) in THF (0.3 M) was treated with DEAD (1.3 equiv.) at rt. The resulting homogeneous solution was heated at reflux for 1-3 hours, cooled at rt, and concentrated in vacuo. The residue was purified by column chromatography on silica gel by eluting EtOAc-Hexane (1 : 5 ~ 1 : 30) to afford the corresponding product. (47~86%)

Spectral data for 1-(3-methoxyphenoxy)-3-butene (**21**),⁴ 1-(3-hydroxyphenoxy)-3-butene (**23**),⁴ 1-(3,4-methylenedioxyphenoxy)-3-butene (**25**)⁴ and 1-(2-naphthoxy)-3-butene (**27**)⁵ were consistent with data reported in the literature.

1-(3,5-Dimethoxyphenoxy)-3-butene (20)

a colorless oil (70%, EtOAc:Hexane = 1:30).

δ_H (CDCl₃, 400MHz) 2.53 (qt, J = 1.4, 6.8, 2H), 3.77 (s, 6H), 3.98 (t, J = 6.7, 2H), 5.10 (dq, J = 10.3, 1.4, 1H), 5.17 (dq, J = 17.2, 1.7, 1H), 5.90 (qt, J = 6.8, 10.3, 1H), 6.09 (s, 3H). δ_C (CDCl₃, 100MHz) 33.54, 55.29, 67.19, 92.99, 93.41, 116.99, 134.40, 160.79, 161.47. $\nu_{\max}$ (NaCl)/cm⁻¹ 3078, 2937, 1601, 1471, 1387, 1204, 1152, 1066, 918, 818. HREIMS m/z 208.1100 (M)⁺, calcd for C₁₂H₁₆O₃ 208.1099.

1-(3-Hydroxy-2-methylphenoxy)-3-butene (29)

⁴ Hoffmann, N.; Pete, J.-P. *J. Org. Chem.* **1997**, 62, 6952.

⁵ Youn, S. W.; Pastine, S. J.; Sames, D. *Org. Lett.* **2004**, 6, 581.

a yellow oil (47%, EtOAc:Hexane = 1:10).

δ_H (CDCl₃, 400MHz) 2.12 (s, 3H), 2.55 (qt, J = 1.4, 6.6, 2H), 4.00 (t, J = 6.7, 2H), 4.64 (s, 1H), 5.10 (qt, J = 1.5, 10.2, 1H), 5.17 (qt, J = 1.6, 17.2, 1H), 5.92 (m, 1H), 6.44 (d, J = 6.8, 1H), 6.46 (d, J = 7.1, 1H), 7.00 (t, J = 8.2, 1H). δ_C (CDCl₃, 100MHz) 7.96, 33.82, 67.63, 104.18, 107.98, 112.46, 116.86, 126.35, 134.69, 154.45, 157.90. ν_{max} (NaCl)/cm⁻¹ 3433, 3079, 2926, 1595, 1466, 1380, 1272, 1098, 917, 769, 707. HREIMS m/z 178.0996 (M)⁺, calcd for C₁₁H₁₄O₂ 178.0994.

General procedure for the conversion of allyl aryl ethers to 2-methylbenzofurans

To a solution of allyl aryl ether in 1,4-dioxane (0.015 M) were added Pd(MeCN)₂Cl₂, Na₂CO₃ (1 equiv.) and BQ (1 equiv.). The resulting mixture was heated with stirring for the reported temperature. After the reaction was completed,⁶ the mixture was cooled, quenched with distilled water, extracted with CH₂Cl₂ (three times), dried over MgSO₄, and concentrated in vacuo. The residue was purified by column chromatography on silica gel (EtOAc:Hexane = 1:10 ~ 1:50) to afford the corresponding product.

6-Hydroxy-2-methylbenzofurans & 4-Hydroxy-2-methylbenzofurans (2)

5 mol % Pd(MeCN)₂Cl₂ at 65 °C.

The mixture of isomers was obtained as a brown oil (6- : 4- = 84:16, EtOAc : Hexane = 1:10).

Spectral data were consistent with data reported in the literature.⁷

⁶ In TLC diagram, the spots of both starting material and corresponding product are very close in every reaction. After developing TLC with nonpolar solvent (such as EtOAc:Hexane = 1:10~1:50) a few times, visualization with anisaldehyde stain shows color differences between starting material and product.

For examples, TLC diagrams with regards to reactions of **1** and **20** are as follows.

⁷ (a) For spectral data of *p*-isomer, see: Dupont, R.; Cotelle, P. *Synthesis* **1999**, 1651. (b) For spectral data of *o*-isomer, see: Demerseman, P.; Lechartier, J.-P.; Pène, C.; Cheutin, A.; Royer, R. *Bull. Soc. Chim. Fr.* **1965**, 1473.

6-Methoxy-2-methylbenzofurans & 4-Methoxy-2-methylbenzofurans (2)

20 mol % Pd(MeCN)₂Cl₂ at 80 °C.

The mixture of isomers was obtained as a pale yellow oil (6- : 4- = 88:12, EtOAc:Hexane = 1:20).

Spectral data were consistent with data reported in the literature.⁸

4,6-Dimethoxy-2-methylbenzofuran (6)

5 mol % Pd(MeCN)₂Cl₂ at 65 °C.

a white solid (EtOAc : Hexane = 1:20).

δ_{H} (CDCl₃, 400MHz) 2.40 (s, 3H), 3.82 (s, 3H), 3.88 (s, 3H), 6.29 (d, J = 1.7, 1H), 6.35 (s, 1H), 6.59 (d, J = 1.7, 1H). δ_{C} (CDCl₃, 100MHz) 13.91, 55.51, 55.75, 88.16, 93.85, 99.55, 112.63, 152.67, 152.73, 156.29, 158.15. ν_{max} (NaCl)/cm⁻¹ 3119, 2955, 1601, 1501, 1457, 1370, 1216, 1155, 1104, 1043, 949, 814. HREIMS m/z 192.0786 (M)⁺, calcd for C₁₁H₁₂O₃ 192.0786.

2,4,6-Trimethylbenzofuran (8)

25 mol % Pd(MeCN)₂Cl₂ at 80 °C.

a pale yellow oil (EtOAc : Hexane = 1:20).

Spectral data were consistent with data reported in the literature.⁹

2-Methylnaphtho[2,1-b]furan (10)

20 mol % Pd(MeCN)₂Cl₂ at 80 °C.

⁸ Alemagna, A.; Baldoli, C.; Del Buttero, P.; Licandro, E.; Maiorana, S. *Synthesis* **1987**, 192.

⁹ Choi, H.-D.; Seo, P.-J.; Son, B.-W. *J. Kor. Chem. Soc.* **1999**, 43, 237.

a pale yellow solid (EtOAc : Hexane = 1:30).

Spectral data were consistent with data reported in the literature.¹⁰

2-Methyl-5,6-methylenedioxybenzofuran (12)

10 mol % Pd(MeCN)₂Cl₂ at 80 °C.

a white solid (EtOAc : Hexane = 1:20).

δ_{H} (CDCl₃, 400MHz) 2.39 (d, J = 1.0, 3H), 5.95 (s, 2H), 6.24 (m, 1H), 6.84 (s, 1H), 6.91 (s, 1H). δ_{C} (CDCl₃, 100MHz) 14.03, 93.20, 98.85, 101.00, 102.82, 122.18, 144.06, 145.01, 149.58, 154.69. ν_{max} (NaCl)/cm⁻¹ 3110, 2962, 2894, 1600, 1463, 1367, 1318, 1279, 1159, 1038, 942, 850. HREIMS m/z 176.0475 (M)⁺, calcd for C₁₀H₈O₃ 176.0473.

2,7-Dimethyl-6-hydroxybenzofuran (14)

10 mol % Pd(MeCN)₂Cl₂ at 65 °C.

a pale yellow solid (EtOAc : Hexane = 1:10).

δ_{H} (CDCl₃, 400MHz) 2.39 (s, 3H), 2.43 (s, 3H), 4.66 (s, 1H), 6.26 (s, 1H), 6.69 (d, J = 8.2, 1H), 7.12 (d, J = 8.2, 1H). δ_{C} (CDCl₃, 100MHz) 8.31, 14.04, 102.52, 107.12, 111.00, 116.84, 122.27, 150.33, 154.41, 154.58. ν_{max} (NaCl)/cm⁻¹ 3285, 2922, 1605, 1508, 1427, 1277, 1154, 1080, 1038, 935, 810. HREIMS m/z 162.0677 (M)⁺, calcd for C₁₀H₁₀O₂ 162.0681.

2-Methyl-4,5,6-trimethoxybenzofuran (16)

5 mol % Pd(MeCN)₂Cl₂ at 65 °C.

an orange oil (EtOAc : Hexane = 1:5).

δ_{H} (CDCl₃, 400MHz) 2.39 (d, J = 1.4, 3H), 3.85 (s, 3H), 3.87 (s, 3H), 4.06 (s, 3H), 6.42 (m, 1H), 6.73 (s, 1H). δ_{C} (CDCl₃, 100MHz) 13.86, 56.32, 60.31, 61.35, 90.28, 100.45, 114.01, 136.78, 145.28, 151.17, 151.63, 153.32. ν_{max} (NaCl)/cm⁻¹ 3117, 2936, 1621, 1469, 1383, 1200, 1039, 933, 896, 795. HREIMS

¹⁰ (a) Chow, Y. L.; Zhou, X. M.; Gaitan, T. J.; Wu, Z. Z. *J. Am. Chem. Soc.* **1989**, *111*, 3813. (b) Black, M.; Cadogan, J. I. G.; McNab, H.; MacPherson, A. D.; Roddam, V. P.; Smith, C.; Swenson, H. R. *J. Chem. Soc., Perkin Trans. 1* **1997**, 2483.

m/z 222.0892 (M)⁺, calcd for C₁₂H₁₄O₄ 222.0892.

4,6-Dimethoxy-2,3-dimethylbenzofuran (18)

25 mol % Pd(MeCN)₂Cl₂ at 80 °C.

a colorless oil (EtOAc : Hexane = 1:50).

δ_H (CDCl₃, 400MHz) 2.23 (s, 3H), 2.29 (s, 3H), 3.81 (s, 3H), 3.84 (s, 3H), 6.25 (d, J = 1.7, 1H), 6.53 (d, J = 2.0, 1H). δ_C (CDCl₃, 100MHz) 9.72, 11.36, 55.33, 55.68, 87.93, 93.50, 93.78, 109.42, 147.54, 154.31, 155.49, 157.96. ν_{max} (NaCl)/cm⁻¹ 2927, 1606, 1501, 1455, 1214, 1148, 1112, 815. HREIMS m/z 206.0946 (M)⁺, calcd for C₁₂H₁₄O₃ 206.0943.

General procedure for a direct oxidative cyclization of aryl homoallyl ethers to afford chromenes

To a solution of aryl homoallyl ether in 1,4-dioxane (0.015 M) were added Pd(MeCN)₂Cl₂, Na₂CO₃ (1 equiv.) and BQ (1 equiv.). The resulting mixture was heated with stirring for the reported temperature. After the reaction was completed,⁶ the mixture was cooled, quenched with distilled water, extracted with CH₂Cl₂ (three times), dried over MgSO₄, and concentrated in vacuo. The residue was purified by column chromatography on silica gel (EtOAc : Hexane = 1:10 ~ 1:30) to afford the corresponding product.

5,7-Dimethoxy-4-methyl-2H-chromene (19)

5 mol % Pd(MeCN)₂Cl₂ at rt.

a yellowish oil (EtOAc : Hexane = 1:20).

δ_H (CDCl₃, 400MHz) 2.15 (dt, J = 1.7, 3H), 3.77 (s, 3H), 3.78 (s, 3H), 4.49 (dq, J = 4.4, 1.5, 2H), 5.40 (tq, J = 4.3, 1.5, 1H), 6.08 (d, J = 2.4, 1H), 6.11 (d, J = 2.4, 1H). δ_C (CDCl₃, 100MHz) 21.73, 55.28, 55.38, 64.97, 92.92, 93.78, 107.99, 115.21, 131.64, 157.22, 158.23, 160.53. ν_{max} (NaCl)/cm⁻¹ 2962, 1615, 1470, 1213, 1046, 949, 820, 740. HREIMS m/z 206.0942 (M)⁺, calcd for C₁₂H₁₄O₃ 206.0943.

7-Methoxy-4-methyl-2H-chromene & 5-Methoxy-4-methyl-2H-chromene (22)

25 mol % Pd(MeCN)₂Cl₂ at 80 °C.

The mixture of isomers was obtained as a yellow oil (7- : 5- = >99:1, EtOAc : Hexane = 1:30).

Spectral data were consistent with data reported in the literature.¹¹

7-Hydroxy-4-methyl-2H-chromene & 5-Hydroxy-4-methyl-2H-chromene (24)

20 mol % Pd(MeCN)₂Cl₂ at 80 °C.

The mixture of isomers was obtained as a colorless oil (7- : 5- = 94:6, EtOAc : Hexane = 1:10).¹²

Signals corresponding to the major isomer: δ_H (CDCl₃, 400MHz) 1.98 (dt, J = 1.7, 3H), 4.71 (dq, J = 3.6, 1.7, 2H), 5.24 (br, 1H), 5.43 (tq, J = 3.6, 1.6, 1H), 6.33 (d, J = 2.7, 1H), 6.39 (dd, J = 8.3, 2.6, 1H), 7.00 (d, J = 8.2, 1H). δ_C (CDCl₃, 100MHz) 17.95, 65.68, 103.14, 107.93, 115.45, 117.73, 124.57, 130.02, 155.39, 156.29. Representative signals corresponding to the minor isomer: δ_H (CDCl₃, 400MHz) 2.24 (dt, J = 1.7, 3H), 4.53 (dq, J = 4.2, 1.7, 2H), 5.12 (br, 1H), 5.56 (tq, J = 4.1, 1.7, 1H), 6.49 (d, J = 8.2, 1H), 6.95 (t, J = 7.9, 1H). δ_C (CDCl₃, 100MHz) 21.61, 64.78, 108.96, 109.68, 118.36, 128.82. ν_{max} (NaCl)/cm⁻¹ 3376, 2969, 1616, 1506, 1466, 1381, 1160, 1066, 1011, 942, 813. HREIMS m/z 162.0680 (M)⁺, calcd for C₁₀H₁₀O₂ 162.0681.

4-Methyl-6,7-methylenedioxy-2H-chromene (26)

20 mol % Pd(MeCN)₂Cl₂ at 80 °C.

a pale yellow oil (EtOAc : Hexane = 1:30).

δ_H (CDCl₃, 400MHz) 1.97 (dt, J = 1.6, 1.7, 3H), 4.64 (dq, J = 3.8, 1.7, 2H), 5.47 (tq, J = 3.8, 1.7, 1H), 5.90 (s, 2H), 6.41 (s, 1H), 6.66 (s, 1H). δ_C (CDCl₃, 100MHz) 18.27, 65.41, 98.44, 100.99, 103.34,

¹¹ (a) Engler, T. A.; LaTessa, K. O.; Iyengar, R.; Chai, W.; Agrios, K. *Bioorg. Med. Chem.* **1996**, *4*, 1755. (b) Anderson, W. K.; LaVoie, E. J.; Whitkop, P. G. *J. Org. Chem.* **1974**, *39*, 881.

¹² This compound is too unstable to store even for 1 day in the freezer. Therefore, spectral data for this compound should be obtained immediately after purification.

115.65, 117.63, 130.54, 141.82, 147.24, 149.64. $\nu_{\max}$ (NaCl)/ cm^{-1} 2923, 1624, 1485, 1404, 1341, 1265, 1166, 1038, 937, 858, 753. HREIMS m/z 190.0631 (M)⁺, calcd for C₁₁H₁₀O₃ 190.0630.

1-Methyl-3*H*-benzo[*f*]chromene (28) & 1-Methylene-2,3-dihydro-1*H*-benzo[*f*]chromene (28')

25 mol % Pd(MeCN)₂Cl₂ at 80 °C.

The mixture of **28** and **28'** was obtained as an orange oil (**28**:**28'** = 2:1, EtOAc : Hexane = 1:30).

Signals corresponding to **28**¹³: δ_{H} (CDCl₃, 400MHz) 2.42 (dt, J = 1.5, 3H), 4.55 (dq, J = 6.2, 1.4, 2H), 5.76 (tq, J = 6.2, 1.6, 1H), 7.15 (d, J = 8.9, 1H), 7.35 (m, 1H), 7.45 (m, 1H), 7.68 (d, J = 8.9, 1H), 7.78 (d, J = 8.2, 1H), 8.11 (d, J = 8.6, 1H). δ_{C} (CDCl₃, 100MHz) 22.52, 64.41, 114.03, 117.78, 118.07, 123.26, 125.38, 125.89, 128.83, 129.57, 129.99, 130.70, 132.59, 154.07. Representative signals corresponding to **28'**: δ_{H} (CDCl₃, 400MHz) 2.75 (t, J = 6.0, 2H), 4.47 (t, J = 5.8, 2H), 5.38 (m, 1H), 5.66 (s, 1H), 7.02 (d, J = 8.9, 1H), 7.35 (m, 1H), 7.45 (m, 1H), 7.64 (d, J = 8.5, 1H), 7.78 (d, J = 8.2, 1H), 8.45 (d, J = 8.6, 1H). δ_{C} (CDCl₃, 100MHz) 33.24, 68.44, 118.49, 123.16, 124.01, 126.54, 128.52, 129.35, 130.28, 130.96, 139.48, 152.31. $\nu_{\max}$ (NaCl)/ cm^{-1} 3056, 2926, 1633, 1595, 1513, 1466, 1395, 1340, 1231, 1214, 1079, 1028, 996, 816, 747. HREIMS m/z 196.0884 (M)⁺, calcd for C₁₄H₁₂O 196.0888.

4,8-Dimethyl-7-hydroxy-2*H*-chromene (30)

20 mol % Pd(MeCN)₂Cl₂ at 80 °C.

a white solid (EtOAc : Hexane = 1:10).¹²

δ_{H} (CDCl₃, 400MHz) 1.98 (d, J = 1.0, 3H), 2.09 (s, 3H), 4.72 (m, 2H), 4.79 (s, 1H), 5.45 (m, 1H), 6.37 (d, J = 8.2, 1H), 6.89 (d, J = 8.2, 1H). δ_{C} (CDCl₃, 100MHz) 7.99, 18.06, 65.58, 107.14, 111.23, 115.25, 117.58, 121.24, 130.56, 153.22, 154.41. $\nu_{\max}$ (NaCl)/ cm^{-1} 3430, 2968, 1615, 1505, 1456, 1379, 1101, 808. HREIMS m/z 176.0832 (M)⁺, calcd for C₁₁H₁₂O₂ 176.0837.

¹³ For spectral data of **28**, see: (a) Muljani, Z.; Tilak, B. D. *Indian J. Chem.* **1969**, 7, 28 & 30. (b) Iwai, I.; Ide, J. *Chem. Pharm. Bull.* **1962**, 926.