Facile Regiocontrolled Synthesis of Trialkylsubstituted Pyrazines

Tarek Abou Elmaaty and Lyle W. Castle

Department of Chemistry, Idaho State University, Campus Box 8023, Pocatello Idaho, 83209 USA.

Experimental:

¹H NMR were recorded in CDCl₃ on a Varian Mercury NMR spectrometer operating at 300 MHz. Chemical Shifts are expressed in ppm relative to TMS as an internal standard. *J* values are given in Hertz. Final products were purified by Kugelrohr distillation using BUCHI Glass Oven B-580. Starting materials were purchased from Aldrich and used without further purification. Octyl viologen was purchased from KASEI KOGYO CO., LTD (TCI) D1854. The addition procedure was performed using Orion M 361 Sage Pump.

Formation of Pyrazines (6a-d):

A 100 ml round bottom flask is charged with the appropriate α -nitroketone (0.00025 mol), octylviologen (0.00025 mol), 4 mL acetonitrile, 4 mL water and a magnetic stir bar. The round bottomed flask is sealed with suba seal, flushed with nitrogen and placed in an oil bath that is kept at 35°C. To the stirred reaction mixture was added simultaneously a solution of hydrogen sulfite (0.01 mol), potassium carbonate (0.0075 mol) in water (4 mL) and a solution of the appropriate α -aminoketone hydrochloride salt (0.0015 mol) in water (4 mL) over a period of 25 minutes. The α -aminoketone addition should be started one minute earlier than the hydrogen sulfite/potassium carbonate solution. After the addition is complete, the reaction mixture is then left to stir at 35 °C for three hours at which time the reaction flask is removed from the oil bath to reveal a deep blue solution. The reaction mixture was extracted with petroleum ether (2 x 30 mL). The organic solution is then washed with 2% sodium hydroxide (2 x 50 mL) followed by water and dried with sodium sulfate. The solution is filtered and the solvent removed *in vacuo* to give a yellow oil, which may then be purified by distillation using Kugelrohr apparatus.

2-Hexyl-3,5-dimethylpyrazine (6a): ¹H NMR (300 MHz, CDCl₃): δ H, ppm (J, Hz): 0.88 (t, 3H, CH₃), 1.3 (m, 8H, 4 CH₂), 2.49 (s, 3H, CH₃), 2.53 (s, 3H, CH₃), 2.77 (t,

2H, CH₂), 8.19 (s, 1H, CH-pyrazine). ¹³C NMR (300 MHz, CDCl₃): δ C, ppm: 152.93, 150.98, 149.92, 141.12 (pyrazine carbons), 34.83, 31.91, 29.49, 28.56, 22.80, 21.83, 21.24, 14.29 (aliphatic carbons).

2-Heptyl-3,5-dimethylpyrazine (6b): ¹H NMR (300 MHz, CDCl₃): δ H, ppm (J, Hz): 0.96 (t, 3H, CH₃), 1.21 (m, 10H, 5 CH₂), 1.95 (s, 3H, CH₃), 1.97 (s, 3H, CH₃), 2.45 (t, 2H, CH₂), 8.19 (s, 1H, CH-pyrazine). ¹³C NMR (300 MHz, CDCl₃): δ C, ppm: 151.70, 150.8, 146.7, 139.9 (pyrazine carbons), 33.8, 30.9, 30.7, 28.6, 28.3, 27.9, 21.6, 20.0, 13.1 (aliphatic carbons).

3-Ethyl-5-methyl-2-propylpyrazine (6c): ¹H NMR (300 MHz, CDCl₃): δ H, ppm (J, Hz): 1.00 (t, 3H, CH₃), 1.3 (t, 3H, CH₃), 1.59 (m, 2H, CH₂), 1.70-1.79 (q, 2H, CH₂), 2.69 (s, 3H, CH₃), 2.78 (t, 2H, CH₂), 8.19 (s, 1H, CH-pyrazine). ¹³C NMR (300 MHz, CDCl₃): δ C ppm: 151.8, 151.6, 146.9, 141.1 (pyrazine carbons), 36.8, 28.7, 22.0, 14.9, 14.4, 14.0 (aliphatic carbons).

3-Ethyl-5-methyl-2-hexylpyrazine (6d): ¹H NMR (300 MHz, CDCl₃): δ H, ppm (J, Hz): 1.00 (t, 3H, CH₃), 1.25 (t, 3H, CH₃), 1.29-1.6 (m, 8H, 4 CH₂), 2.54 (s, 3H, CH₃), 2.65 (q, 2H, CH₂), 2.75 (t, 2H, CH₂), 8.19 (s, 1H, CH-pyrazine). ¹³C NMR (300 MHz, CDCl₃): δ C, ppm: 154.2, 153.0, 152.0, 140.0 (pyrazine carbons), 35.0, 32.0, 30.0, 27.9, 25.0, 23.0, 19.0, 14.0, 11.0 (aliphatic carbons).

6a

6a

6a

6a

6b

6c

6c

general_31May2005-15:07:47
Data saved to:
ISUNMR/export/home/general/vnmrsvs/data
Archive directory: /export/home/general/vnmrsvs/data
Sample directory: general_31May2005-15:07:47
File: gCOSY
Pulse Sequence: gCOSY
Solvent: CDCl₃
Ambient temperature
Mercury-3000 130MMR
Relax. delay 1.000 sec
Acq. time 0.213 sec
Width 4887.7 Hz
2D Width 4887.7 Hz
Single scan
128 increments
OBSERVE F1: 300.0673615 MHz
DATA PROCESSING
F1 DATA PROCESSING
Sf, sine bell 0.107 sec
Sf, sine bell 0.027 sec
FT size 2048 x 2048
Total time 3 min, 53 sec

6c

6d