Facile Synthesis of β -Organotellurobutenolides via Electrophilic Tellurolactonization of α -Allenic Acids

Qing Xu,[†] Xian Huang,^{* †‡} and Jingqi Yuan[†]

[†] *Department of Chemistry, Zhejiang University (Xixi Campus), Hangzhou 310028, P. R. China*

[‡] *State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu, Shanghai 200032, P. R. China*

huangx@mail.hz.zj.cn

Supporting Information

Table of Contents

1. Experimental Procedure.....	S2
2. ¹ H NMR and ¹³ C NMR spectra of 2 , 4 , 5	S8

General Procedure for the Electrophilic Tellurolactonization of α -Allenic Acids.

The mixture of diphenyl ditelluride (0.205g, 0.50 mmol) and iodine (0.130g, 0.51 mmol) was stirred in dry MeCN (2 mL) at 0 °C under N₂ for 1 h to form PhTeI (1.0 mmol). To the solution of diaryl ditelluride (0.50 mmol) in dry MeCN (1 mL) under N₂ was added dropwise the solution of dry bromine (0.85g, 0.53 mmol) in MeCN (1 mL) and the mixture was stirred at 0 °C for 1 h to form ArTeBr (1.0 mmol). To the solution of diaryl ditelluride (0.50 mmol) in dry MeCN (2 mL) under N₂ was added dropwise SO₂Cl₂ (0.081g, 0.60 mmol) and the mixture was stirred at room temperature for 1 h to form ArTeCl (1.0 mmol). α -Allenic acid (0.50 mmol) in dry MeCN (2 mL) was then added to above prepared ArTeX solution and the mixture stirred for the time as indicated in the Tables. The reaction mixture was concentrated and the residue subjected to flash chromatography or preparative TLC to afford β -organotellurobutenolides **2**. Reactions carried out under different conditions were conducted in a similar manner.

3-Benzyl-4-phenyltelluro-5-(*o*-chloro)phenyl-2(5H)-butenolide (2a).

Pale yellow crystal, mp 118-20 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.36-7.25 (m, 8H), 7.20-7.12 (m, 3H), 7.06-7.02 (m, 2H), 6.97-6.96 (d, *J* = 5.6 Hz, 1H), 6.03 (s, 1H), 3.91-3.87 (d, *J* = 15.3 Hz, 1H), 3.71-3.67 (d, *J* = 15.3 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃): δ 170.7, 144.7, 141.0, 137.1, 136.2, 135.4, 132.3, 130.5, 129.7, 129.6, 129.5, 129.3, 128.9, 127.2, 127.0, 109.0, 33.4. MS (*m/z*): 490 (13, M⁺), 455 (5), 282 (7), 265 (13), 239 (8), 207 (6), 139 (53), 115 (43), 91 (35), 77 (100). IR (KBr): ν 1739, 1602, 1473, 1435, 1300, 1171, 1065 cm⁻¹. Anal. Calcd. for C₂₃H₁₇ClO₂Te: C, 56.56; H, 3.51. Found: C, 56.32; H, 3.54.

3-Benzyl-4-phenyltelluro-2(5H)-butenolide (2b).

Pale yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 7.82-7.80 (d, J = 6.8 Hz, 2H), 7.46-7.42 (m, 1H), 7.35-7.25 (m, 7H), 4.34 (s, 2H), 3.69 (s, 2H). ^{13}C NMR (100 MHz, CDCl_3): δ 171.7, 141.1, 137.0, 133.3, 130.0, 129.9, 129.6, 128.9, 127.1, 108.5, 75.2, 33.1. MS (m/z): 380 (25, M^+), 207 (5), 129 (100), 115 (41), 91 (28), 77 (74). IR (film): ν 3058, 3027, 2926, 1748, 1608, 1573, 1494, 1474, 1453, 1434, 1339, 1079, 1051, 1018 cm^{-1} . Anal. Calcd. for $\text{C}_{17}\text{H}_{14}\text{O}_2\text{Te}$: C, 54.03; H, 3.73. Found: C, 53.85; H, 3.75.

4-Phenyltelluro-5-butyl-2(5H)-butenolide (2c).

Pale yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 7.88-7.86 (dd, J = 8.0 Hz, J = 0.8 Hz, 2H), 7.50-7.46 (m, 1H), 7.38-7.34 (m, 2H), 5.79-7.78 (d, J = 1.2 Hz, 1H), 5.03-5.00 (m, 1H), 1.88-1.80 (m, 1H), 1.60-1.52 (m, 1H), 1.48-1.29 (m, 4H), 0.92-0.88 (t, J = 7.8 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 171.4, 155.2, 140.6, 130.3, 130.0, 124.0, 109.9, 86.9, 34.2, 26.2, 22.3, 13.8. MS (m/z): 346 (12, M^+), 288 (2), 207 (12), 159 (21), 139 (6), 131 (37), 77 (94), 41 (100). IR (film): ν 3055, 2957, 2871, 1741, 1560, 1473, 1436, 1379, 1331, 1301, 1239, 1162, 1120, 1065, 1017, 998 cm^{-1} . Anal. Calcd. for $\text{C}_{14}\text{H}_{16}\text{O}_2\text{Te}$: C, 48.90; H, 4.69. Found: C, 48.75; H, 4.72.

4-(1-Naphthyl)telluro-5-butyl-2(5H)-butenolide(2d).

Pale yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 8.23-8.21 (d, J = 6.8 Hz, 1H), 8.16-8.14 (d, J = 8.0 Hz, 1H), 7.98-7.96 (d, J = 8.0 Hz, 1H), 7.84-7.81 (dd, J = 6.8 Hz, J = 2.4 Hz, 1H), 7.58-7.51 (m, 2H), 7.38-7.34 (m, 1H), 5.54 (d, J = 1.2 Hz, 1H), 4.96-4.92 (m, 1H), 1.86-1.78 (m, 1H), 1.58-1.51 (m, 1H), 1.46-1.38 (m, 1H), 1.37-1.22 (m, 3H), 0.87-0.84 (t, J = 7.2 Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 171.3, 154.4, 142.2, 135.8, 133.9, 132.0, 131.9, 129.2, 128.1, 127.0, 126.6, 124.4,

113.7, 87.0, 34.2, 26.3, 22.2, 13.8. MS (m/z): 396 (7, M^+), 257 (9), 181 (22), 152 (29), 139 (4), 127 (100). IR (film): ν 3052, 2956, 2870, 1742, 1560, 1499, 1457, 1377, 1332, 1239, 162, 1022 cm^{-1} . Anal. Calcd. for $\text{C}_{18}\text{H}_{18}\text{O}_2\text{Te}$: C, 54.88; H, 4.61. Found: C, 54.62; H, 4.65.

3-Benzyl-4-phenyltelluro-5-butyl-2(5H)-butenolide (2e).

Pale yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 7.78-7.76 (d, $J = 7.2$ Hz, 2H), 7.42-7.38 (m, 1H), 7.31-7.23 (m, 7H), 4.642-4.60 (dd, $J = 7.5$ Hz, $J = 1.9$ Hz, 1H), 3.86-3.82 (d, $J = 15.2$ Hz, 2H), 3.57-3.53 (d, $J = 15.2$ Hz, 2H), 1.74-1.70 (m, 1H), 1.34-1.19 (m, 2H), 1.17-1.05 (m, 3H), 0.78-0.76 (t, $J = 6.8$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 171.0, 146.1, 140.5, 137.5, 135.7, 130.0, 129.6, 129.1, 128.7, 126.8, 110.0, 85.0, 33.7, 33.3, 26.2, 22.0, 13.8. MS (m/z): 436 (8, M^+), 228 (6), 207 (11), 183 (37), 159 (13), 141 (34), 129 (55), 115 (69), 91 (84), 77 (100). IR (film): ν 2956, 1749, 1606, 1573, 1495, 1454, 1435, 1329, 1175, 1061 cm^{-1} . Anal. Calcd. for $\text{C}_{21}\text{H}_{22}\text{O}_2\text{Te}$: C, 58.12; H, 5.11. Found: C, 58.05; H, 5.12.

4-Phenyltelluro-5-octyl-2(5H)-butenolide (2f).

Pale yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 7.88-7.86 (d, $J = 7.6$ Hz, 2H), 7.49-7.45 (m, 1H), 7.37-7.33 (m, 2H), 5.79-5.78 (d, $J = 1.6$ Hz, 1H), 5.02-4.99 (m, 1H), 1.88-1.78 (m, 1H), 1.59-1.50 (m, 1H), 1.47-1.27 (m, 12H), 0.91-0.88 (d, $J = 7.0$ Hz, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 171.2, 154.9, 140.6, 130.3, 130.0, 124.1, 110.0, 86.9, 34.5, 31.8, 29.2, 29.13, 29.11, 24.1, 22.6, 14.1. MS (m/z): 402 (8, M^+), 207 (3), 159 (12), 131 (25), 77 (91), 41 (100). IR (film): ν 3055, 2926, 2855, 1747, 1561, 1465, 1435, 1331, 1238, 1160, 1064, 1017 cm^{-1} . Anal. Calcd. for $\text{C}_{18}\text{H}_{24}\text{O}_2\text{Te}$: C, 54.05; H, 6.05. Found: C, 53.85; H, 6.08.

3-Allyl-4-phenyltelluro-5-phenyl-2(5H)-butenolide (2g).

Pale yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 7.48-7.46 (dd, $J = 7.6$ Hz, $J = 1.2$ Hz, 2H), 7.34-7.32 (m, 1H), 7.27-7.25 (m, 1H), 7.21-7.17 (m, 2H), 7.14-7.10 (m, 2H), 6.83-8.61 (d, $J = 7.2$ Hz, 2H), 5.90-5.82 (m, 1H), 5.51 (s, 1H), 5.18 (dd, $J = 24.0$ Hz, $J = 1.2$ Hz, 1H), 5.15 (dd, $J = 3.2$ Hz, $J = 1.2$ Hz, 1H), 3.31-3.25 (m, 1H), 3.09-3.03 (m, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 170.7, 145.5, 141.0, 134.6, 134.5, 132.6, 129.7, 129.4, 129.2, 128.6, 127.7, 117.7, 109.5, 86.9, 31.3. MS (m/z): 406 (11, M^+), 207 (5), 199 (20), 105 (97), 77 (100). IR (film): ν 3064, 2919, 2253, 1752, 1606, 1431, 1298, 1267, 1200, 1069, 995, 912 cm^{-1} . Anal. Calcd. for $\text{C}_{19}\text{H}_{16}\text{O}_2\text{Te}$: C, 56.50; H, 3.99. Found: C, 56.30; H, 4.02.

3-Benzyl-4-phenyltelluro-5-phenyl-2(5H)-butenolide (2h).

White crystal. Mp 80-2 . ^1H NMR (400 MHz, CDCl_3): δ 7.38-7.36 (d, $J = 7.2$ Hz, 2H), 7.32-7.21 (m, 7H), 7.15-7.11 (m, 2H), 7.07-7.03 (m, 2H), 6.74-6.72 (d, $J = 7.6$ Hz, 2H), 5.47 (s, 1H), 3.90-3.87 (d, $J = 15.2$ Hz, 1H), 3.66-3.61 (d, $J = 19.2$ Hz, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 170.9, 145.7, 141.0, 137.3, 135.7, 134.7, 129.6, 129.4, 129.2, 128.9, 128.6, 127.8, 127.0, 109.6, 86.9, 33.2. MS (m/z): 456 (6, M^+), 248 (5), 207 (4), 105 (82), 77 (100). IR (KBr): ν 3061, 3030, 1749, 1603, 1495, 1455, 1297, 1267, 1175, 1082, 1066, 1017, 998 cm^{-1} . Anal. Calcd. for $\text{C}_{23}\text{H}_{18}\text{O}_2\text{Te}$: C, 60.85; H, 4.00. Found: C, 60.76; H, 4.01.

4-Phenyltelluro-5-(*o*-chlorophenyl)-2(5H)-butenolide (2i).

Pale yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 7.75-7.73 (d, $J = 5.6$ Hz, 2H), 7.43-7.38 (m, 2H), 7.33-7.29 (m, 4H), 7.23-7.22 (m, 1H), 6.48 (d, $J = 2.0$ Hz, 1H), 5.82 (d, $J = 2.0$ Hz, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 171.3, 156.3, 140.7, 133.1,

132.9, 130.6, 130.3, 129.9, 127.8, 127.7, 123.2, 110.4, 84.7. MS (m/z): 400 (12, M^+), 365 (9), 232 (6), 207 (6), 193 (10), 149 (35), 139 (20), 114 (24), 102 (79), 77 (100). IR (film): ν 3067, 2927, 1788, 1750, 1557, 1474, 1436, 1279, 1232, 1156, 1055 cm^{-1} . Anal. Calcd. for $\text{C}_{16}\text{H}_{11}\text{ClO}_2\text{Te}$: C, 48.25; H, 2.78. Found: C, 48.49; H, 2.81.

Substitution Reaction of Organotellurobutenolide **2a** with Diethyl Cuprate Reagent.

Ethyl magnesium bromide (1.50 mmol) in dry THF was added at $0\text{ }^\circ\text{C}$ to the THF suspension of CuI (0.75 mmol) in a tube reactor under nitrogen to form the diethyl copper reagent. The mixture was then cooled to $-78\text{ }^\circ\text{C}$ and tellurobutenolide **2a** (0.50 mmol) in dry THF was added dropwise. After the temperature was gradually returned to room temperature, the mixture was washed with saturated aqueous NH_4Cl and NaCl, extracted with EtOAc, dried over MgSO_4 . The solvent was evaporated and the residue purified by TLC to afford substitution product **4**.

3-Benzyl-4-ethyl-5-(*o*-chloro)phenyl-2(5H)-butenolide (4). Pale yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 7.42-7.39 (dd, $J = 7.8\text{ Hz}$, $J = 1.4\text{ Hz}$, 1H), 7.30-7.20 (m, 7H), 7.01-6.99 (dd, $J = 7.2\text{ Hz}$, $J = 1.60\text{ Hz}$, 1H), 6.35 (s, 1H), 3.74-3.64 (m, 2H), 2.53-2.43 (m, 1H), 2.10-2.01 (1H), 0.96-0.92 (t, $J = 7.6\text{ Hz}$, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 174.3, 165.6, 138.1, 133.8, 132.5, 130.4, 130.0, 128.7, 128.5, 127.9, 127.5, 126.6, 126.4, 79.7, 29.6, 20.1, 12.2. MS (m/z): 312 (2, M^+), 84 (100). IR (film): ν 3029, 1760, 1669, 1602, 1495, 1454, 1299, 1164, 1107, 1044, 1015 cm^{-1} . Anal. Calcd. for $\text{C}_{19}\text{H}_{17}\text{ClO}_2$: C, 72.96; H, 5.48. Found: C, 72.65; H, 5.51.

Cross-coupling Reaction of Organotellurobutenolide **2h** with 1-Hexyne.

The mixture of **2h** (0.50 mmol), 1-hexyne (1.0 mmol), PdCl₂ (10 mol%), CuI (10 mol%), and triethylamine (0.50 mmol) in MeOH (5 mL) was stirred overnight under nitrogen. Conventional workup and preparative TLC purification afforded **5**.

3-Benzyl-4-(1-hexynyl)-5-phenyl-2(5H)-butenolide (5). Pale yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 7.37-7.22 (m, 10H), 5.71 (s, 1H), 3.79-3.71 (m, 2H), 2.38-2.35 (t, *J* = 7.00 Hz, 2H), 1.48-1.42 (m, 2H), 1.33-1.28 (m, 2H), 0.88-0.84 (t, *J* = 7.3 Hz, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 173.5, 115.0, 137.6, 134.7, 133.0, 129.2, 128.9, 128.7, 128.6, 126.7, 110.0, 83.7, 72.1, 31.1, 30.1, 21.8, 19.7, 13.5. MS (*m/z*): 330 (2, M⁺). IR (film): ν 3031, 2958, 2872, 2221, 1761, 1640, 1495, 1455, 1356, 1163, 1084, 1021 cm⁻¹. Anal. Calcd. for C₂₃H₂₂O₂: C, 83.60; H, 6.71. Found: C, 83.34; H, 6.75.

ppm

7.36130
7.35018
7.31217
7.30320
7.28539
7.26843
7.25243
7.19574
7.19223
7.17316
7.15162
7.13986
7.12159
7.05537
7.03648
7.01752
6.03327

3.91088
3.87265
3.71138
3.67321

2a ^1H NMR

2a ^{13}C NMR

2b ¹H NMR

2c ¹H NMR

ppm

- 7.87206
- 7.85468
- 7.85241
- 7.49266
- 7.47426
- 7.45559
- 7.37245
- 7.35335
- 7.33463
- 7.27128

- 5.77695
- 5.77344
- 5.02205
- 5.01859
- 5.01363
- 5.00998
- 5.00342
- 4.99980
- 4.99456

- 1.85187
- 1.84020
- 1.55310
- 1.53382
- 1.45300
- 1.43303
- 1.42765
- 1.39640
- 1.38402
- 1.37435
- 1.35989
- 1.35440
- 1.33822
- 1.32018
- 1.30344
- 0.91351
- 0.89577
- 0.87805
- 0.00174

2d ¹H NMR

2d ^{13}C NMR

ppm

7.78301
7.76481
7.76332
7.41564
7.41317
7.39647
7.37827
7.37589
7.31341
7.29557
7.27706
7.25823
7.25696
7.24943
7.23878
7.23075

4.61867
4.61453
4.60064
4.59513

3.86212
3.82402
3.56900
3.53090

1.71551
1.70774
1.28316
1.26319
1.25381
1.24583
1.23446
1.13493
1.11600
1.11283
1.09574
1.08333
1.07922
1.06340
1.04404
0.76327
0.74566
0.72901

2e ¹H NMR

Integral

ppm

9

8

7

6

5

4

3

2

1

2.0053

1.0095

7.0023

1.0000

1.0061

1.0072

1.0070

2.0027

3.0072

3.0104

4 ¹H NMR

5 ¹H NMR

Integral

ppm

5 ^{13}C NMR