Facile Synthesis of the IndenoTetrahydropyridine Core of Haouamine A

Nicole D.Smith, Joji Hayashida, and Viresh H. Rawal

*Department of Chemistry, The University of Chicago
5735 South Ellis Avenue, Chicago, IL 60637*

SUPPORTING INFORMATION

General

All reagents were commercially obtained (Aldrich, Acros) at the highest commercial quality and used without further purification. Air and moisture sensitive reagents were transferred via syringe or cannula. All moisture sensitive reactions were carried out in oven-dried glassware under an atmosphere of argon with freshly distilled solvents. The reactions were monitored by thin-layer chromatography carried out on 0.25mm Whatman silica gel plates using UV light as the visualizing agent. The NMR spectra were obtained on a Bruker DRX-400 (400 MHz) or a Bruker DRX-500 (500 MHz) spectrometer. Chemical shifts (δ) are reported in parts per million (ppm) relative to deuterated solvent signals in d_6 -DMSO. The following abbreviations were used to explain the multiplicities: s=singlet; d=doublet, t=triplet, q=quartet, dd=doublet of doublets, m=multiplet, b=broad. Mass spectroscopic analyses were carried out at the facilities of The University of Chicago or The Ohio State University. IR spectra were recorded on a Nicolet Nexus 670 FT-IR and values are reported in cm^{-1} units.

3-Oxo-piperidine-1-carboxylic acid methyl ester (6)

To a solution of 3-hydroxypiperidine hydrochloride (4.82 g, 35.0 mmol) in anhydrous CH_2Cl_2 (100 mL) was added TEA (9.76 mL, 70.0 mmol) and the mixture was cooled to 0 °C in an ice bath. A solution of dimethyl dicarbonate (4.12 mL, 38.5 mmol) in anhydrous CH_2Cl_2 (10 mL) was then added dropwise via addition funnel, and the reaction mixture was allowed to warm to room temperature and stir overnight. The CH_2Cl_2 was removed *in vacuo*, and Et_2O was added to precipitate out the ammonium salts. The precipitate was filtered and washed with Et_2O (3X, 20 mL). The crude product was purified by flash silica gel chromatography (9:1, Et_2O :Hex) to afford 3-hydroxypiperidine-1-carboxylic acid methyl ester (5.57 g, 35.0 mmol) in quantitative yield.

Anhydrous CH_2Cl_2 (240 mL) was added to a 500 mL round bottom flask, purged with N_2 and cooled to $-78\text{ }^\circ\text{C}$. Oxalyl chloride (3.45 mL, 39.5 mmol) was added, followed by the dropwise addition of anhydrous DMSO (4.69 mL, 66.0 mmol). This solution was allowed to stir at $-78\text{ }^\circ\text{C}$ for 5 min. Next, 3-hydroxy-piperidine-1-carboxylic acid methyl ester (5.25 g, 33.0 mmol) dissolved in anhydrous CH_2Cl_2 (100 mL) was added dropwise via cannula to the reaction mixture at $-78\text{ }^\circ\text{C}$. After 30 min, TEA (18.4 mL, 132 mmol) was added and the reaction mixture was allowed to warm to rt and stir for 2 h. The reaction mixture was diluted with CH_2Cl_2 (100 mL), washed with sat. NH_4Cl (150 mL), sat. NaHCO_3 (150 mL), brine (150 mL) and dried with MgSO_4 . The organic layer was concentrated *in vacuo* and purified via flash silica gel chromatography (3:2, EtOAc:Hex) to afford 3-oxo-piperidine-1-carboxylic acid methyl ester (4.98 g, 31.7 mmol) in a 96% yield. ^1H NMR (d_6 -DMSO, 400 MHz, 363K) δ 1.61-1.68 (m, 2H), 2.14 (t, 2H, $J=7.0$ Hz), 3.27 (t, 2H, $J=6.0$ Hz), 3.35 (s, 3H), 3.66 (s, 2H); ^{13}C NMR (d_6 -DMSO, 400 MHz, 363K) δ 21.4, 37.1, 41.5, 51.7, 53.2, 154.7, 204.4; IR (NaCl) ν 1226, 1407, 1451, 1700, 2873, 2956; MS (ESI) 158 ($\text{M}+\text{H}^+$).

2-(3-Methoxy-benzyl)-3-oxo-piperidine-1-carboxylic acid methyl ester (7)

To a solution of 3-oxo-piperidine-1-carboxylic acid methyl ester (2.00 g, 12.7 mmol) in anhydrous benzene (24.0 mL) was added pyrrolidine (1.57 mL, 19.1 mmol), and the reaction was allowed to stir for 30 min. Next, the mixture was heated to $110\text{ }^\circ\text{C}$ and H_2O and benzene were collected utilizing a Dean-Stark trap. The remaining benzene was removed *in vacuo* and the enamine was redissolved in CH_3CN (24.0 mL). To this solution was added a solution of 3-methoxybenzyl chloride (2.4 mL, 16.5 mmol) in CH_3CN (10 mL), and the reaction mixture was refluxed overnight. The reaction was quenched with brine (30 mL) and extracted with EtOAc (2X, 50 mL). The organic layer was washed with sat. NH_4Cl (40 mL), dried (MgSO_4), concentrated *in vacuo* and purified via flash silica gel chromatography (3:2, Hex:EtOAc) to afford 2-(3-methoxy-benzyl)-3-oxo-piperidine-1-carboxylic acid methyl ester (2.03 g, 7.32 mmol) in a 58% yield. ^1H NMR (d_6 -DMSO, 400 MHz, 363K) δ 1.83-1.90 (m, 2H), 2.35-2.42 (m, 1H), 2.49-2.59 (m, 1H), 2.97-3.02 (m, 2H), 3.18-3.25 (m, 1H), 3.47 (s, 3H), 3.74 (s, 3H), 3.85-3.91 (m, 1H), 4.59 (t, 1H, $J=7.0$ Hz), 6.73-6.79 (m, 3H), 7.17 (t, 1H, $J=8.0$ Hz); ^{13}C NMR (d_6 -DMSO, 400 MHz, 363K) δ 21.8, 35.8, 35.9, 38.5, 51.5, 54.5, 63.8, 111.7, 114.5, 120.9, 128.5, 138.0, 154.7, 158.9, 205.9; IR (NaCl) ν 1448, 1601, 1702, 2954; MS (ESI) 278 ($\text{M}+\text{H}^+$).

2-(3-Methoxy-benzyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (8)

To a solution of 2-(3-methoxy-benzyl)-3-oxo-piperidine-1-carboxylic acid methyl ester (0.30 g, 1.08 mmol) in anhydrous DMSO (3 mL) was added IBX (0.30 g, 1.08 mmol), and the mixture was heated to 70 °C. The reaction was allowed to stir at 70 °C for 4 h then an additional equivalent of IBX was added, and after 4 h, a third equivalent of IBX was added, and the reaction mixture was allowed to stir at 70 °C overnight. The mixture was filtered and the filtrate was washed with ether (3X, 5 mL). The organic extracts were combined and washed with sat. NaHCO₃ (5 mL), brine (5 mL), dried with MgSO₄, filtered and concentrated to afford 2-(3-methoxy-benzyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (0.27 g, 0.98 mmol) in a 90% yield. ¹H NMR (d₆-DMSO, 400 MHz, 363K) δ 2.89 (dd, 2H, J=6.8 Hz, J=6.8 Hz), 3.46 (bs, 3H), 3.75 (s, 3H), 3.93 (ddd, 1H, J=2.5 Hz, J=2.5 Hz, J=21.0 Hz), 4.52 (ddd, 1H, J=2.0 Hz, J=5.0 Hz, J=21.0 Hz), 4.70 (dd, 1H, J=6.0 Hz, J=8.5 Hz), 6.06 (dt, 1H, J=2.5 Hz, J=10.5 Hz), 6.74-6.80 (m, 3H), 7.18-7.13 (m, 1H), 7.18 (t, 1H, J=8.0 Hz); ¹³C NMR (d₆-DMSO, 400 MHz, 363K) δ 35.4, 51.6, 54.4, 60.8, 86.9, 111.7, 114.3, 120.8, 124.4, 128.4, 137.6, 146.7, 154.3, 158.7, 193.4; IR (NaCl) ν 1449, 1684, 1700, 2954; MS (ESI) 274 (M-H⁺).

3-Hydroxy-2-(3-methoxy-benzyl)-3-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (9)

To a solution of 2-(3-methoxy-benzyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (72.0 mg, 0.26 mmol) in anhydrous THF (5 mL), was added 3-methoxyphenylmagnesium bromide (523 μL, 0.52 mmol) dropwise at rt. The reaction was quenched with H₂O (2 mL) after an hour, and extracted with Et₂O (3X, 5 mL). The organic layers were combined, dried with MgSO₄, filtered and concentrated *in vacuo*. The crude product was purified via flash silica gel chromatography (3:2, Et₂O:Hex) to afford 3-hydroxy-2-(3-methoxy-benzyl)-3-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (75.0 mg, 0.20 mmol) in a 75% yield. ¹H NMR (d₆-DMSO, 400 MHz, 363K) δ 2.58 (dd, 1H, J=11.0 Hz, J=14.0 Hz), 2.98 (bs, 3H), 3.29 (dd, 1H, J=3.0 Hz, J=14.5 Hz), 3.61-3.65 (m, 1H), 3.73 (s, 3H), 3.74 (s, 3H), 4.17 (m, 1H), 4.44-4.47 (m, 1H), 5.44 (bs, 1H), 5.77-5.79 (m, 1H), 5.99 (d, 1H, J=10.0 Hz), 6.71-6.73 (m, 1H), 6.76-6.78 (m, 2H), 6.99-7.00 (m, 2H), 7.11-7.20 (m, 2H); ¹³C NMR (d₆-DMSO, 400 MHz, 363K) δ 33.0, 50.9, 51.4, 54.5, 54.5, 72.4, 80.3, 111.1, 111.6, 111.7, 111.7, 114.2, 117.9, 120.9, 122.7, 127.7, 128.1, 129.6, 140.5, 147.6, 158.4, 158.7; IR (NaCl) ν 1455, 1600, 1680, 2923, 2953, 3426; HRMS (EM) [M+H]⁺ calcd for C₂₂H₂₅NO₅Na 406.1624, found 406.1623.

5-Methoxy-13-(3-methoxy-phenyl)-10-aza-tricyclo[7.2.2.0^{2,7}]trideca-2(7),3,5,12-tetraene-10-carboxylic acid methyl ester (10)

A solution of 3-hydroxy-2-(3-methoxy-benzyl)-3-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (31.0 mg, 0.08 mmol) in anhydrous CH₂Cl₂ (1 mL) was added dropwise to a CF₃SO₃H (1 mL) and CH₂Cl₂ (1 mL) mixture at -30 °C. The reaction mixture was allowed to stir at -30 °C for 30 min then poured into H₂O (10 mL) and extracted with CH₂Cl₂ (3X, 25 mL). The organic layers were combined, washed with sat. NaHCO₃ (30 mL), dried with Na₂SO₄ and concentrated *in vacuo* to afford 5-methoxy-13-(3-methoxy-phenyl)-10-aza-tricyclo[7.2.2.0^{2,7}]trideca-2(7),3,5,12-tetraene-10-carboxylic acid methyl ester (16.0 mg, 0.045 mmol) in a 55% yield and 3-methoxy-13-(3-methoxy-phenyl)-10-aza-tricyclo[7.2.2.0^{2,7}]trideca-2(7),3,5,12-tetraene-10-carboxylic acid methyl ester (6.0 mg, 0.02 mmol) in a 21% yield. ¹H NMR (d₆-DMSO, 400 MHz, 363K) δ 2.98-3.03 (m, 1H), 3.31 (dd, 1H, J=4.5 Hz, J=18.0 Hz), 3.45 (dd, 1H, J=4.0 Hz, J=11.0 Hz), 3.61-3.64 (m, 1H), 3.66 (bs, 3H), 3.71, (s, 3H), 3.79 (s, 3H), 5.33 (bs, 1H), 6.65-6.68 (m, 2H), 6.87-6.90 (m, 2H), 6.97-6.98 (m, 1H), 7.03 (dd, 1H, J=1.0 Hz, J=8.0 Hz), 7.30 (t, 1H, J=8.0 Hz); ¹³C NMR (d₆-DMSO, 400 MHz, 363K) δ 27.6, 29.9, 35.9, 50.8, 51.6, 54.6, 54.7, 110.2, 111.1, 112.9, 115.9, 116.7, 128.9, 129.2, 129.6, 131.5, 136.0, 138.4, 154.8, 157.9, 159.3; IR (NaCl) ν 1447, 1498, 1605, 1696, 2952; HRMS (EM) [M+H]⁺ calcd for C₂₂H₂₃NO₄Na 388.1519, found 388.1506.

3-Methoxy-13-(3-methoxy-phenyl)-10-aza-tricyclo[7.2.2.0^{2,7}]trideca-2(7),3,5,12-tetraene-10-carboxylic acid methyl ester (11)

¹H NMR (d₆-DMSO, 400 MHz, 363K) δ 2.99-3.05 (m, 1H), 3.32 (dd, 1H, J=4.5 Hz, J=18.0 Hz), 3.47 (dd, 1H, J=4.0 Hz, J=11.0 Hz), 3.62-3.65 (m, 1H), 3.65 (bs, 3H), 3.79 (s, 3H), 3.82 (s, 3H), 4.45-4.47 (m, 1H), 5.32 (bs, 1H), 6.68 (d, 1H, J=8.0 Hz), 6.80-6.84 (m, 2H), 6.88 (dd, 1H, J=2.0 Hz, J=8.5 Hz), 6.98 (m, 1H), 7.03 (d, 1H, J=8.0 Hz), 7.10 (t, 1H, J=8.0 Hz), 7.30 (t, 1H, J=8.0 Hz); ¹³C NMR (d₆-DMSO, 400 MHz, 363K) δ 20.0, 28.4, 35.8, 50.6, 51.6, 54.7, 55.7, 108.6, 110.1, 112.9, 116.7, 122.9, 126.6, 127.8, 128.9, 129.2, 129.3, 136.3, 138.1, 139.5, 155.1, 159.3; IR (NaCl) ν 1447, 1581, 1697, 2923; HRMS (EM) [M+H]⁺ calcd for C₂₂H₂₃NO₄Na 388.1519, found 388.1519.

2-(2-Bromo-5-methoxy-benzyl)-3-oxo-piperidine-1-carboxylic acid methyl ester (12)

To a solution of 2-(3-methoxy-benzyl)-3-oxo-piperidine-1-carboxylic acid methyl ester (0.55 g, 1.96 mmol) in anhydrous CH_3CN (5 mL) was added a solution of *N*-bromosuccinimide (0.37 g, 2.06 mmol) in anhydrous CH_3CN (5 mL) dropwise via syringe pump. The reaction mixture was allowed to stir for 2 hours, and the CH_3CN was concentrated *in vacuo* and the crude product was redissolved in CH_2Cl_2 (20 mL) and was washed with H_2O (10 mL), brine (10 mL) and dried over Na_2SO_4 . The organic layer was concentrated *in vacuo*, and the crude product was purified via flash silica gel chromatography (7:3, Et_2O :Hex) to afford 2-(2-Bromo-5-methoxy-benzyl)-3-oxo-piperidine-1-carboxylic acid methyl ester (0.70 g, 1.95 mmol) in 99% yield. ^1H NMR (d_6 -DMSO, 400 MHz, 363K) δ 1.91-1.99 (m, 2H), 2.40-2.47 (m, 1H), 2.57-2.65 (m, 1H), 3.12 (dd, 2H, $J=7.5$ Hz, $J=13.5$ Hz), 3.33-3.40 (m, 1H), 3.38 (bs, 3H), 3.75 (s, 3H), 3.91-3.97 (m, 1H), 4.74 (dd, 1H, $J=6.0$ Hz, $J=9.0$ Hz), 6.76 (dd, 1H, $J=3.0$ Hz, $J=9.0$ Hz), 6.84 (d, 1H, $J=3.0$ Hz), 7.44 (d, 1H, $J=9.0$ Hz); ^{13}C NMR (d_6 -DMSO, 400 MHz, 363K) δ 21.7, 35.9, 35.9, 38.3, 51.4, 54.9, 62.2, 114.1, 114.1, 116.9, 132.3, 136.9, 154.1, 158.1, 205.7; IR (NaCl) ν 1446, 1601, 1700, 2953; MS (ESI) 356 ($\text{M}+\text{H}^+$), 358 ($\text{M}+\text{H}^+$).

2-(2-Bromo-5-methoxy-benzyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (13)

To a solution of 2-(2-Bromo-5-methoxy-benzyl)-3-oxo-piperidine-1-carboxylic acid methyl ester (63.5 mg, 0.18 mmol) in anhydrous DMSO (1 mL) was added IBX (50.0 mg, 0.18 mmol), and the mixture was heated to 70 °C. The reaction was allowed to stir at 70 °C for 4 h, then an additional equivalent of IBX was added, and after 4 h, a third equivalent of IBX was added and the reaction mixture was allowed to stir at 70 °C overnight. The mixture was filtered and the filtrate was washed with ether (3X, 3 mL). The organic extracts were combined and washed with sat. NaHCO_3 (3 mL), brine (3 mL), dried with MgSO_4 , filtered and concentrated to afford 2-(2-Bromo-5-methoxy-benzyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (60.0 mg, 0.17 mmol) in a 95% yield. ^1H NMR (d_6 -DMSO, 400 MHz, 363K) δ 2.94-3.04 (m, 2H), 3.39 (bs, 3H), 3.75 (s, 3H), 4.10 (ddd, 1H, $J=2.5$ Hz, $J=2.5$ Hz, $J=11.0$ Hz), 4.55 (ddd, 1H, $J=2.0$ Hz, $J=5.0$ Hz, $J=21.0$ Hz), 4.85 (dd, 1H, $J=5.5$ Hz, $J=9.5$ Hz), 6.10 (dt, 1H, $J=2.5$ Hz, $J=10.5$ Hz), 6.77 (dd, 1H, $J=3.0$ Hz, $J=9.0$ Hz), 6.84 (d, 1H, $J=3.0$ Hz), 7.20 (ddd, 1H, $J=2.0$ Hz, $J=5.0$ Hz, $J=10.0$ Hz), 7.45 (d, 1H, 9.0 Hz); ^{13}C NMR (d_6 -DMSO, 400 MHz, 363K) δ 35.8, 51.6, 54.9, 59.3, 62.2, 114.1, 114.2, 116.9, 124.6, 132.4, 136.4, 146.9, 154.3, 158.1, 193.0; IR (NaCl) ν 1446, 1595, 1688, 1705, 2953; MS (ESI) 354 ($\text{M}+\text{H}^+$), 356 ($\text{M}+\text{H}^+$).

2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (14)

To a solution of 2-(2-Bromo-5-methoxy-benzyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (40.0 mg, 0.11 mmol) in anhydrous THF (3 mL) was added 3-methoxyphenylmagnesium bromide (226 μ L, 0.22 mmol) dropwise at rt. The reaction was quenched with H₂O (2 mL) after an hour and extracted with Et₂O (3X, 5 mL). The organic layers were combined, dried with MgSO₄, filtered and concentrated *in vacuo*. The crude product was purified via flash silica gel chromatography (3:2, Et₂O:Hex) to afford 2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (39.0 mg, 0.08 mmol) in a 76% yield. ¹H NMR (d₆-DMSO, 400 MHz, 363K) δ 2.72 (dd, 1H, J=11.0 Hz, J=14.0 Hz), 2.99 (bs, 3H), 3.47 (dd, 1H, J=3.0 Hz, J=14.0 Hz), 3.72-3.75 (m, 1H), 3.73 (s, 3H), 3.74 (m, 3H), 4.16 (ddd, 1H, J=3.0 Hz, J=3.0 Hz, J=19.0 Hz), 4.63 (d, 1H, J=10.5 Hz), 5.47 (bs, 1H), 5.80-5.86 (m, 1H), 6.01-6.03 (m, 1H), 6.70 (dd, 1H, J=3.0 Hz, J=9.0 Hz), 6.78 (ddd, J=1.0 Hz, J=2.5 Hz, J=8.0 Hz), 6.86 (d, 1H, J=3.0 Hz), 6.99-7.02 (m, 2H), 7.19 (t, 1H, J=8.0 Hz), 7.40 (d, 1H, J=9.0 Hz); ¹³C NMR (d₆-DMSO, 400 MHz, 363K) δ 33.3, 50.9, 54.5, 54.9, 58.5, 58.7, 72.3, 111.6, 111.8, 113.6, 114.3, 116.4, 117.8, 122.7, 127.8, 129.6, 132.2, 139.2, 147.5, 154.6, 158.1, 158.5; IR (NaCl) ν 1456, 1597, 1653, 1684, 2952, 3420; MS (ESI) 444 (M+H⁺), 446 (M+H⁺).

6-Bromo-3-methoxy-13-(3-methoxy-phenyl)-10-aza-tricyclo[7.2.2.0^{2,7}]trideca-2(7),3,5,12-tetraene-10-carboxylic acid methyl ester (15)

A solution of 2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (27.0 mg, 0.06 mmol) in anhydrous CH₂Cl₂ (1 mL) was added dropwise to a CF₃SO₃H (1 mL) and CH₂Cl₂ (1 mL) mixture at -30 °C. The reaction mixture was allowed to stir at -30 °C for 30 min then poured into H₂O (10 mL) and extracted with CH₂Cl₂ (3X, 25 mL). The organic layers were combined, washed with sat. NaHCO₃ (30 mL), dried with Na₂SO₄ and concentrated *in vacuo* to afford 6-Bromo-3-methoxy-13-(3-methoxy-phenyl)-10-aza-tricyclo[7.2.2.0^{2,7}]trideca-2(7),3,5,12-tetraene-10-carboxylic acid methyl ester (22.0 mg, 0.05 mmol) in a 81% yield with only one isomer observed by NMR. ¹H NMR (d₆-DMSO, 400 MHz, 363K) δ 2.85 (dd, 1H, J=2.5 Hz, J=18.0 Hz), 3.21 (dd, 1H, J=4.5 Hz, J=18.0 Hz), 3.48 (dd, 1H, J=4.5 Hz, J=11.0 Hz), 3.65-3.67 (m, 1H), 3.65 (bs, 3H), 3.80 (s, 3H), 3.84 (s, 3H), 4.53 (ddd, 1H, J=1.5 Hz, J=4.0 Hz, J=8.0 Hz), 5.43 (bs, 1H), 6.82 (dd, 1H,

$J=1.0$ Hz, $J=8.0$ Hz), 6.87 (d, 1H, 9.0 Hz), 6.88-6.91 (m, 1H), 7.01 (m, 1H), 7.04-7.07 (m, 1H), 7.31 (t, 1H, $J=8.0$ Hz), 7.43 (d, 1H, $J=9.0$ Hz); ^{13}C NMR ($\text{d}_6\text{-DMSO}$, 400 MHz, 363K) δ 28.7, 38.4, 48.8, 50.2, 51.7, 54.7, 56.0, 110.1, 111.1, 111.2, 113.1, 116.7, 116.73, 128.3, 129.2, 130.4, 130.7, 134.9, 137.6, 139.6, 154.6, 159.3; IR (NaCl) ν 1452, 1577, 1697, 2926; HRMS (EM) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{22}\text{H}_{22}\text{BrNO}_4\text{Na}$ 466.0624, found 466.0609.

2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3-(3-methoxy-phenyl)-piperidine-1-carboxylic acid methyl ester (16)

To a solution of 2-(2-Bromo-5-methoxy-benzyl)-3-oxo-piperidine-1-carboxylic acid methyl ester (18.6 mg, 0.05 mmol) in anhydrous THF (300 μL) was added 3-methoxyphenylmagnesium bromide (61.0 μL , 0.06 mmol) dropwise at rt. The reaction was quenched with H_2O (2 mL) after an hour, and extracted with Et_2O (3X, 5 mL). The organic layers were combined, dried with MgSO_4 , filtered and concentrated *in vacuo*. The crude product was purified via flash silica gel chromatography (3:2, Et_2O :Hex) to afford 2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3-(3-methoxy-phenyl)-piperidine-1-carboxylic acid methyl ester (15.0 mg, 0.03 mmol) in a 65% yield. ^1H NMR ($\text{d}_6\text{-DMSO}$, 400 MHz, 363K) δ 1.25-1.34 (m, 1H), 1.60-1.64 (m, 1H), 1.95-2.12 (m, 2H), 3.09 (dd, 1H, $J=11.0$ Hz, $J=14.0$ Hz), 3.18 (s, 3H), 3.24 (dd, 1H, $J=4.0$ Hz, $J=13.0$ Hz), 3.83 (dd, 1H, $J=6.0$ Hz, $J=13.0$ Hz), 4.95 (dd, $J=3.0$ Hz, $J=11.0$ Hz), 5.07 (bs, 1H), 6.70 (dd, 1H, $J=3.0$ Hz, $J=9.0$ Hz), 6.80 (ddd, 1H, $J=1.0$ Hz, $J=2.5$ Hz, $J=8.0$ Hz), 6.88 (d, 1H, $J=3.0$ Hz), 7.12-7.14 (m, 2H), 7.22 (t, 1H, $J=8.0$ Hz), 7.39 (d, 1H, $J=9.0$ Hz); ^{13}C NMR ($\text{d}_6\text{-DMSO}$, 400 MHz, 363K) δ 21.1, 31.6, 32.3, 37.4, 51.0, 54.5, 54.9, 58.2, 71.8, 111.8, 113.5, 114.5, 116.9, 118.0, 128.1, 132.0, 139.5, 147.7, 154.6, 158.0, 158.6; IR (NaCl) ν 1456, 1652, 1684, 2361, 2951, 3446; MS (ESI) 498 ($\text{M}+\text{Cl}^-$), 500 ($\text{M}+\text{Cl}^-$).

8-Bromo-5-methoxy-4 α -(3-methoxy-phenyl)-2,3,4,4 α ,9,9 α -hexahydro-indeno[2,1- β]pyridine-1-carboxylic acid methyl ester (17)

A solution of 2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3-(3-methoxy-phenyl)-piperidine-1-carboxylic acid methyl ester (22.0 mg, 0.05 mmol) in anhydrous CH_2Cl_2 (1 mL) was added dropwise to a $\text{CF}_3\text{SO}_3\text{H}$ (1 mL) and CH_2Cl_2 (1 mL) mixture at -30°C . The reaction mixture was allowed to stir at -30°C for 30 min, then allowed to warm to rt and stir 2 hours. The reaction mixture was then poured into H_2O (10 mL) and extracted with CH_2Cl_2 (3X, 25 mL). The organic layers were combined, washed with sat. NaHCO_3 (30 mL), dried with Na_2SO_4 and concentrated *in vacuo* to afford 8-Bromo-5-methoxy-4 α -(3-methoxy-phenyl)-2,3,4,4 α ,9,9 α -hexahydro-indeno[2,1- β]pyridine-1-carboxylic acid methyl ester (16.0 mg, 0.04 mmol) in a 77% yield. ^1H NMR (d_6 -DMSO, 400 MHz, 363K) δ 1.38-1.47 (m, 1H), 1.73-1.83 (m, 1H), 2.01-2.08 (m, 1H), 2.38-2.44 (m, 1H), 2.94-3.00 (m, 1H), 3.21 (dd, 1H, $J=8.5$ Hz, $J=17.0$ Hz), 3.46 (s, 3H), 3.57 (s, 3H), 3.72 (s, 3H), 3.83-3.90 (m, 1H), 4.99 (t, 1H, $J=8.0$ Hz), 6.74-6.81 (m, 4H), 7.19 (t, 1H, $J=8.0$ Hz), 7.40 (d, 1H, $J=9.0$ Hz); ^{13}C NMR (d_6 -DMSO, 400 MHz, 363K) δ 19.1, 27.1, 35.7, 37.6, 51.6, 53.9, 54.5, 55.1, 61.7, 109.4, 110.6, 112.6, 112.7, 118.5, 128.2, 130.7, 136.9, 141.0, 147.0, 154.9, 155.2, 158.7; IR (NaCl) ν 1446, 1471, 1600, 1695, 2927; MS (ESI) 446 ($\text{M}+\text{H}^+$), 448 ($\text{M}+\text{H}^+$).

2-(2-Bromo-5-methoxy-benzyl)-5-(3-methoxy-phenyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (18)

To a solution of CuBrSMe_2 (6.0 mg, 0.03 mmol) in anhydrous THF (350 μL) at -78°C was added 3-methoxyphenylmagnesium bromide (207 μL , 0.21 mmol) dropwise, and allowed to stir at -78°C for 30 min. Next, 2-(2-Bromo-5-methoxy-benzyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (49.0 mg, 0.14 mmol) in anhydrous THF (500 μL) was added dropwise via syringe, and the reaction was allowed to stir at -78°C for 30 min. Next, phenylselenenyl chloride (45.0 mg, 0.24 mmol) and diphenyl diselenide (few crystals) in anhydrous THF (500 μL) was added dropwise, and the reaction mixture was allowed to stir at -78°C for 30 min. The reaction mixture was then allowed to warm to 0°C , and a mixture of acetic acid (31.0 μL) in H_2O (156 μL) was added dropwise followed by a 30% H_2O_2 (156 μL). The mixture was then allowed to warm to rt, and after 10 min the complete reaction mixture was added to sat. NaHCO_3 (5 mL) and extracted with Et_2O (3X, 10 mL). The organic layers were combined and

washed with sat. NH_4Cl (15 mL) and brine (15 mL), dried with MgSO_4 , filtered, and concentrated *in vacuo*. The crude product was purified via flash silica gel chromatography (1:1, Et_2O :Hex) to afford 2-(2-Bromo-5-methoxy-benzyl)-5-(3-methoxy-phenyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (36.0 mg, 0.08 mmol) in a 57% yield. ^1H NMR (d_6 -DMSO, 400 MHz, 363K) δ 3.10 (dd, 1H, $J=8.0$ Hz, $J=8.0$ Hz), 3.42 (bs, 3H), 3.76 (s, 3H), 3.86 (s, 3H), 4.52 (dd, 1H, $J=2.5$ Hz, $J=20.0$ Hz), 4.92 (dd, 1H, $J=5.5$ Hz, $J=9.5$ Hz), 4.98 (dd, 1H, $J=1.0$ Hz, $J=20.0$ Hz), 6.52-6.53 (m, 1H), 6.78 (dd, 1H, $J=3.0$ Hz, $J=9.0$ Hz), 6.94 (d, 1H, $J=3.0$ Hz), 7.10 (dd, 1H, $J=2.5$ Hz, $J=8.0$ Hz), 7.21-7.22 (m, 1H), 7.27 (ddd, 1H, $J=1.0$ Hz, $J=1.5$ Hz, $J=8.0$ Hz), 7.42 (t, 1H, $J=8.0$ Hz), 7.46 (d, 1H, $J=9.0$ Hz); ^{13}C NMR (d_6 -DMSO, 400 MHz, 363K) δ 35.4, 41.3, 51.8, 54.9, 55.0, 58.5, 111.4, 114.2, 114.2, 116.2, 117.1, 118.1, 120.9, 129.5, 132.4, 136.1, 136.6, 154.5, 155.2, 158.2, 159.3, 193.2; IR (NaCl) ν 1448, 1576, 1670, 1704, 2953; MS (ESI) 460 ($\text{M}+\text{H}^+$), 462 ($\text{M}+\text{H}^+$).

2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3,5-bis-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (19)

To a solution of 2-(2-Bromo-5-methoxy-benzyl)-5-(3-methoxy-phenyl)-3-oxo-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (30.0 mg, 0.06 mmol) in anhydrous THF (500 μL) was added 3-methoxyphenylmagnesium bromide (130.0 μL , 0.13 mmol) dropwise at rt. The reaction was quenched with H_2O (2 mL) after an hour, and extracted with Et_2O (3X, 5 mL). The organic layers were combined, dried with MgSO_4 , filtered and concentrated *in vacuo*. The crude product was purified via flash silica gel chromatography (1:1, Et_2O :Hex) to afford 2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3,5-bis-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (23.0 mg, 0.04 mmol) in a 62% yield. ^1H NMR (d_6 -DMSO, 400 MHz, 363K) δ 2.81 (dd, 1H, $J=11.0$ Hz, $J=14.5$ Hz), 3.08 (bs, 3H), 3.51 (dd, 1H, $J=3.5$ Hz, $J=14.5$ Hz), 3.74 (s, 3H), 3.84 (s, 3H), 4.07 (dd, 1H, $J=1.5$ Hz, $J=18.5$ Hz), 4.55 (dd, 1H, $J=0.5$ Hz, $J=19.0$ Hz), 4.72 (d, 1H, $J=9.5$ Hz), 6.20 (m, 1H), 6.72 (dd, 1H, $J=3.0$ Hz, $J=9.0$ Hz), 6.80 (m, 1H), 6.93 (d, 1H, $J=3.0$ Hz), 6.96 (dd, 1H, $J=2.5$ Hz, $J=8.0$ Hz), 7.01-7.04 (m, 2H), 7.09-7.10 (m, 1H), 7.15 (d, 1H, $J=8.0$ Hz), 7.20 (t, 1H, $J=8.0$ Hz), 7.36 (t, 1H, $J=8.0$ Hz), 7.42 (d, 1H, $J=9.0$ Hz); ^{13}C NMR (d_6 -DMSO, 400 MHz, 363K) δ 33.2, 41.2, 51.1, 54.5, 54.8, 54.9, 58.5, 72.6, 110.8, 111.6, 111.9, 113.2, 113.7, 114.3, 116.5, 117.2, 117.9, 127.1, 127.9, 129.1, 132.2, 132.5, 138.5, 139.0, 147.6, 158.1, 158.5, 159.2; IR (NaCl) ν 1450, 1481, 1598, 1697, 2953, 3394; HRMS (EM) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{29}\text{H}_{30}\text{BrNO}_6\text{Na}$ 590.1148, found 590.1145.

8-Bromo-5-methoxy-3,4 α -bis-(3-methoxy-phenyl)-2,4 α ,9,9 α -tetrahydro-indeno[2,1- β]pyridine-1-carboxylic acid methyl ester (20)

A solution of 2-(2-Bromo-5-methoxy-benzyl)-3-hydroxy-3,5-bis-(3-methoxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid methyl ester (16.0 mg, 0.03 mmol) in anhydrous CH_2Cl_2 (1 mL) was added dropwise to a $\text{CF}_3\text{SO}_3\text{H}$ (1 mL) and CH_2Cl_2 (1 mL) mixture at $-30\text{ }^\circ\text{C}$. The reaction mixture was allowed to stir at $-30\text{ }^\circ\text{C}$ for 30 min then allowed to warm to rt, and stir 2 hours. The reaction mixture was then poured into H_2O (10 mL) and extracted with CH_2Cl_2 (3X, 25 mL). The organic layers were combined, washed with sat. NaHCO_3 (30 mL), dried with Na_2SO_4 , and concentrated *in vacuo* to afford 8-Bromo-5-methoxy-3,4 α -bis-(3-methoxy-phenyl)-2,4 α ,9,9 α -tetrahydro-indeno[2,1- β]pyridine-1-carboxylic acid methyl ester (11.0 mg, 0.02 mmol) in a 71% yield. ^1H NMR (d_6 -DMSO, 400 MHz, 363K) δ 3.00-3.19 (m, 2H), 3.52 (s, 3H), 3.54 (s, 3H), 3.70 (s, 3H), 3.79 (s, 3H), 4.24 (d, 1H, $J=18.0\text{ Hz}$), 4.66 (d, 1H, $J=18.0\text{ Hz}$), 4.93 (t, 1H, $J=9.0\text{ Hz}$), 6.18 (s, 1H), 6.66 (m, 1H), 6.72 (d, 1H, $J=8.0\text{ Hz}$), 6.80 (d, 1H, $J=8.0\text{ Hz}$), 6.87-6.96 (m, 3H), 7.03 (d, 1H, $J=8.0\text{ Hz}$), 7.21 (t, 1H, $J=8.0\text{ Hz}$), 7.30 (t, 1H, $J=8.0\text{ Hz}$), 7.44 (d, 1H, $J=9.0\text{ Hz}$); ^{13}C NMR (d_6 -DMSO, 400 MHz, 363K) δ 33.6, 40.4, 51.8, 54.5, 54.74, 55.13, 55.84, 60.93, 109.61, 110.68, 111.04, 111.85, 112.82, 112.92, 117.06, 117.9, 122.8, 128.4, 129.1, 131.3, 132.8, 133.1, 139.3, 140.9, 146.3, 154.8, 154.9, 158.7, 159.2; IR (NaCl) ν 1447, 1703, 2850, 2920; HRMS (EM) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{29}\text{H}_{28}\text{BrNO}_5\text{Na}$ 572.1043, found 572.1063.

10

11

15

19