Supplementary Information For:

Ferrocenyl Monophosphine Ligands: Synthesis and Applications in Suzuki Coupling of Aryl Chlorides

Colin Baillie, Lixin Zhang, and Jianliang Xiao*

Department of Chemistry, University of Liverpool, Liverpool L69 7ZD, UK
j.xiao@liv.ac.uk

Contents

1. Preparation of **2**S2
2. Preparation of **3**S4
3. Preparation of **4**S5
4. Preparation of **1**.....S6
5. Suzuki couplingS8

1. Preparation of 2

Diphenylphosphinylferrocene (2a). To a degassed 500-mL Schlenk flask was charged ferrocene (10.0 g, 53.8 mmol). AlCl_3 (7.89 g, 59.2 mmol) was weighed out under an atmosphere of N_2 and added to the reaction vessel, and a second cycle of degassing was performed. Heptane (150 mL) was added, followed by Ph_2PCl (11.6 mL, 64.6 mmol) and heating at $105\text{ }^\circ\text{C}$ for 24 h. The reaction was allowed to cool to rt, and the heptane soln decanted. The remaining dark solid mass was extracted with hot heptane (3×100 mL) to remove unreacted ferrocene. Further extraction with hot toluene (5×100 mL) followed by concentration of the toluene layers yielded the crude product. Recrystallisation from EtOH gave diphenylphosphinoferrocene as bright orange crystals; yield: 11.7 g (59%). ^1H NMR (400 MHz, CDCl_3) δ 7.75-7.28 (m, 10H), 4.37 (m, 2H), 4.09 (m, 2H), 4.07 (s, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ 139.5 (d, $J_{\text{CP}} = 14.4$ Hz), 133.8 (d, $J_{\text{CP}} = 19.2$ Hz), 128.8, 128.5 (d, $J_{\text{CP}} = 6.4$ Hz), 73.3 (d, $J_{\text{CP}} = 15.2$ Hz), 71.1 (d, $J_{\text{CP}} = 4.0$ Hz), 70.1, 69.5; ^{31}P NMR (162 MHz, CDCl_3) δ -15.4. Anal. Calcd for $\text{C}_{22}\text{H}_{19}\text{PFe}$: C, 71.36; H, 5.18. Found: 71.54; H, 5.22.

To a soln of diphenylphosphinoferrocene (11.7 g, 31.6 mmol) in MeOH (50 mL) was added a couple of drops of H_2O_2 at $0\text{ }^\circ\text{C}$. The mixture was stirred until oxidation was complete (as determined by TLC, EtOAc/hexane 2:1). The mixture was treated with aq $\text{Na}_2\text{S}_2\text{O}_3$, and extracted with EtOAc (3×75 mL). The organic layers were concentrated to give the crude product as an orange brown oil. Flash chromatography (EtOAc/hexane 2:1) gave the title product as an orange precipitate; yield: 11.0 g (90%). ^1H NMR (400 MHz, CDCl_3) δ 7.75-7.65 (m, 4H), 7.54-7.42 (m, 6H), 4.48 (m, 2H), 4.37 (m, 2H), 4.21 (s, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ 134.7 (d, $J_{\text{CP}} = 106.3$ Hz), 131.9, 131.8, 128.6 (d, $J_{\text{CP}} = 12.0$ Hz), 72.7 (d, $J_{\text{CP}} = 12.8$ Hz), 72.0 (d, $J_{\text{CP}} = 11.2$ Hz), 70.9, 70.1; ^{31}P NMR (162 MHz, CDCl_3) δ 30.3. Anal. Calcd for $\text{C}_{22}\text{H}_{19}\text{POFe}$: C, 68.41; H, 4.97. Found: 68.39; H, 4.95.

Dicyclohexylphosphinylferrocene (2b). An oven-dried 250-mL Schlenk flask containing Mg (3.38 g, 140 mmol) was degassed, and charged with Et_2O (75 mL). CyCl (15.0 mL, 130 mmol) was added dropwise and the mixture heated at $45\text{ }^\circ\text{C}$ overnight to form the Grignard reagent, CyMgCl . Separately, an oven-dried, degassed 250-mL Schlenk flask charged with Et_2O (100 mL) and PCl_3 (4.0 mL, 46 mmol) was prepared, to which the Grignard reagent was added via cannula at $-40\text{ }^\circ\text{C}$ (CO_2 /acetone bath). The mixture was stirred for 2 h at this temperature during which time a white solid (MgCl_2) precipitated and the reaction was warmed to rt overnight. The white precipitate was filtered under an argon atmosphere by Schlenk filtration and concentrated in vacuo to give a colourless oil of Cy_2PCl . In a third, degassed 250-mL Schlenk flask was charged ferrocene (4.28 g, 23.0 mmol), KO^tBu (0.39 g, 3.5 mmol) and THF (100 mL). Maintaining a temperature of $-78\text{ }^\circ\text{C}$, 1.7 M $^t\text{BuLi}$ in pentane (20.3 mL, 34.5 mmol) was added over 15 min, and stirred for 1 h, resulting in a bright orange suspension. The previously prepared Cy_2PCl in THF (10 mL) was added dropwise and allowed to warm to rt overnight. The mixture was treated with aq NH_4Cl and extracted with Et_2O (3×50 mL). The organic layers were dried (MgSO_4) and concentrated to give a yellow/orange oil. Flash chromatography (hexane/EtOAc 4:1) gave dicyclohexylphosphinoferrocene as

an orange solid; yield: 6.20 g (70%). ^1H NMR (400 MHz, CDCl_3) δ 4.28 (m, 2H), 4.16 (m, 2H), 4.14 (s, 5H), 2.01-1.60 (m, 11H), 1.35-1.05 (m, 11H); ^{13}C NMR (100 MHz, CDCl_3) δ 71.9 (d, $J_{\text{CP}} = 11.2$ Hz), 70.0, 69.5, 68.3, 33.9 (d, $J_{\text{CP}} = 11.2$ Hz), 30.7 (d, $J_{\text{CP}} = 4.0$ Hz), 30.7, 27.8 (d, $J_{\text{CP}} = 18.4$ Hz), 27.7, 26.8; ^{31}P NMR (162 MHz, CDCl_3) δ -6.0. Anal. Calcd for $\text{C}_{22}\text{H}_{31}\text{PFe}$: C, 69.11; H, 8.19. Found: 69.30; H, 8.11.

To a soln of dicyclohexylphosphinoferrocene (6.20 g, 16.2 mmol) in EtOAc (50 mL), H_2O_2 was added dropwise at 0 °C. The soln darkened slightly, and the reaction was monitored by TLC (hexane/EtOAc 4:1). When oxidation was complete the soln was treated with aq $\text{Na}_2\text{S}_2\text{O}_3$, and extracted with Et_2O (3×25 mL). The organic layers were dried (MgSO_4), concentrated, and purified by flash chromatography (EtOAc) to give the title compound as an orange solid; yield: 5.03 g (78%). ^1H NMR (400 MHz, CDCl_3) δ 4.41 (m, 2H), 4.37 (m, 2H), 4.32 (s, 5H), 2.05-1.65 (m, 11H), 1.50-1.16 (m, 11H); ^{13}C NMR (100 MHz, CDCl_3) δ 72.8 (d, $J_{\text{CP}} = 96.7$ Hz), 71.8 (d, $J_{\text{CP}} = 9.6$ Hz), 70.7 (d, $J_{\text{CP}} = 8.8$ Hz), 70.0, 37.6 (d, $J_{\text{CP}} = 69.5$ Hz), 27.2 (d, $J_{\text{CP}} = 2.4$ Hz), 27.0 (d, $J_{\text{CP}} = 2.4$ Hz), 26.7 (d, $J_{\text{CP}} = 3.2$ Hz), 26.4, 26.0 (d, $J_{\text{CP}} = 3.2$ Hz); ^{31}P NMR (162 MHz, CDCl_3) δ 46.8.

Di-*tert*-butylphosphinylferrocene (2c). Method (A). To an oven-dried 100-mL Schlenk flask containing ferrocene (9.36 g, 50.3 mmol) was charged THF (25 mL). At 0 °C, 1.7 M $^t\text{BuLi}$ in pentane (30.0 mL, 50.3 mmol) was added over 5 min. The soln was stirred at this temperature for 25 min, and the solvent concentrated in vacuo. The residue was redissolved in a combination of pentane (100 mL) and THF (5 mL), and $\text{CIP}(^t\text{Bu})_2$ (5.30 mL, 27.7 mmol) added at rt. After stirring for 3 h, degassed MeOH (1 mL) was added and the solvent was again concentrated in vacuo. Unreacted ferrocene was eluted with pentane through a plug of silica under N_2 , followed by the crude product by elution with Et_2O . The Et_2O extract was collected separately and concentrated to give di-*tert*-butylphosphinoferrocene as an orange/red oil; yield: 9.15 g, (61% from $\text{CIP}(^t\text{Bu})_2$). ^1H NMR (400 MHz, CDCl_3) δ 4.34 (m, 2H), 4.26 (m, 2H), 4.16 (s, 5H), 2.01-1.60 (m, 11H), 1.21 (d, $J = 11.2$ Hz, 18H); ^{31}P NMR (162 MHz, CDCl_3) δ 27.9. Anal. Calcd for $\text{C}_{18}\text{H}_{27}\text{PFe}$: C, 65.45; H, 8.26. Found: 65.19; H, 8.11. Method (B). Same procedure as method A only $\text{CIP}(^t\text{Bu})_2$ was prepared from $^t\text{BuMgCl}$ and PCl_3 rather than using a commercial source. This resulted in a lower yield of **52** [45% from $\text{CIP}(^t\text{Bu})_2$].

To a soln of di-*tert*-butylphosphinoferrocene (9.15 g, 27.7 mmol) in EtOAc (100 mL), was added a couple of drops of H_2O_2 at 0 °C, and the stirred soln monitored by TLC (EtOAc). Once oxidation was complete, the soln was treated with aq $\text{Na}_2\text{S}_2\text{O}_3$ and extracted with Et_2O (2×50 mL). The organic extracts were dried (MgSO_4), concentrated and purified by flash chromatography (EtOAc/MeOH 100:1) to give the title compound as an orange precipitate; yield: 6.43 g (67%). ^1H NMR (400 MHz, CDCl_3) δ 4.45 (m, 2H), 4.45 (m, 2H), 4.28 (s, 5H), 1.26 (d, $J = 13.5$ Hz, 18H); ^{13}C NMR (100 MHz, CDCl_3) δ 73.6 (d, $J_{\text{CP}} = 87.9$ Hz), 73.0 (d, $J_{\text{CP}} = 8.8$ Hz), 70.5 (d, $J_{\text{CP}} = 8.0$ Hz), 70.3, 36.5 (d, $J_{\text{CP}} = 63.9$ Hz), 27.7; ^{31}P NMR (162 MHz, CDCl_3) δ 58.6. Anal. Calcd for $\text{C}_{18}\text{H}_{27}\text{POFe}$: C, 62.43; H, 7.87. Found: 61.79; H, 7.86.

2. Preparation of 3

2-Iodo-1-(diphenylphosphinyl)ferrocene 3a. To a degassed 250 mL Schlenk flask charged with THF (60 mL) was added TMEDA (4.30 mL, 28.5 mmol) via syringe. The soln was cooled to $-78\text{ }^{\circ}\text{C}$, and 2.5 M $n\text{BuLi}$ in hexanes (11.4 mL, 28.5 mmol) added over 5 min and stirred for a further 15 min. Maintaining a temperature of $-78\text{ }^{\circ}\text{C}$, a soln of diphenylphosphinylferrocene **2** ($R = \text{Ph}$) (5.00 g, 12.9 mmol) in THF (50 mL) was added over 15 min and stirred for a further 1 h. Again at $-78\text{ }^{\circ}\text{C}$, a soln of I_2 (9.86 g, 38.9 mmol) in THF (50 mL) was slowly added. The mixture was warmed to rt overnight, treated with aq $\text{Na}_2\text{S}_2\text{O}_3$ and extracted with Et_2O ($3 \times 75\text{ mL}$). The organic extracts were dried (MgSO_4), concentrated in vacuo, and purified by flash chromatography (EtOAc /hexane 2:1) to yield the title compound as an orange/brown precipitate (4.10 g, 62%), which was used for the next step without further purification. Analytically pure compound was obtained by recrystallisation (EtOAc /hexane), which gave orange/brown needle shaped crystals. ^1H NMR (400 MHz, CDCl_3) δ 7.79-7.72 (m, 2H), 7.68-7.62 (m, 2H), 7.56-7.40 (m, 6H), 4.76 (m, 1H), 4.39 (s, 5H), 4.35 (m, 1H), 3.89 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 135.5 (d, $J_{\text{CP}} = 107.1\text{ Hz}$), 133.3 (d, $J_{\text{CP}} = 108.6\text{ Hz}$), 132.2, 132.1, 131.7 (d, $J_{\text{CP}} = 10.4\text{ Hz}$), 131.6 (d, $J_{\text{CP}} = 2.4\text{ Hz}$), 128.3 (d, $J_{\text{CP}} = 12.8\text{ Hz}$), 128.2 (d, $J_{\text{CP}} = 12.0\text{ Hz}$), 80.0 (d, $J_{\text{CP}} = 8.8\text{ Hz}$), 75.4 (d, $J_{\text{CP}} = 115.8\text{ Hz}$), 75.0 (d, $J_{\text{CP}} = 15.2\text{ Hz}$), 72.9, 72.7, 71.6 (d, $J_{\text{CP}} = 10.4\text{ Hz}$); ^{31}P NMR (162 MHz, CDCl_3) δ 29.3. Anal. Calcd for $\text{C}_{22}\text{H}_{18}\text{POiFe}$: C, 51.60; H, 3.55. Found: 51.46; H, 3.52.

2-Iodo-1-(dicyclohexylphosphinyl)ferrocene 3b. To an oven-dried, degassed 250 mL Schlenk flask was charged dicyclohexylphosphinylferrocene **2** ($R = \text{Cy}$) (3.00 g, 7.62 mmol) in THF (75 mL). The soln was cooled to $-78\text{ }^{\circ}\text{C}$, and 1.7 M $n\text{BuLi}$ in pentane (5.4 mL, 9.15 mmol) added dropwise and stirred at this temperature for 1 h. The reaction flask was protected against light by aluminium foil, and I_2 (3.00 g, 11.45 mmol) in 15 THF (mL) added slowly and allowed to warm to rt overnight. The soln was treated with aq $\text{Na}_2\text{S}_2\text{O}_3$ and extracted with Et_2O ($2 \times 25\text{ mL}$). The organic layers were concentrated in vacuo to yield a crude orange/brown precipitate, which was protected against light whilst stored, and used without further purification on the same day. ^1H NMR (400 MHz, CDCl_3) δ 4.69 (m, 1H), 4.51 (m, 1H), 4.43 (m, 1H), 4.33 (s, 5H), 2.41-1.00 (m, 22H); ^{31}P NMR (162 MHz, CDCl_3) δ 47.7.

2-Iodo-1-(di-*t*-butylphosphinyl)ferrocene 3c. A 100 mL Schlenk flask was degassed, charged with THF (40 mL), and TMEDA (2.88 mL, 19.1 mmol) added via syringe. The soln was cooled to $-78\text{ }^{\circ}\text{C}$, and 2.5M $n\text{BuLi}$ in hexanes (7.62 mL, 19.1 mmol) added over 2 min and stirred for 15 min. Maintaining a temperature of $-78\text{ }^{\circ}\text{C}$, a soln of di-*t*-

butylphosphinylferrocene **2** (R = ^tBu) (3.00 g, 8.66 mmol) in THF (20 mL) was added over 15 min and stirred for a further 1 h. Again at -78 °C, a soln of I₂ (11.0 g, 43.3 mmol) in THF (20 mL) was slowly added, and upon completion, the reaction vessel was protected against light using aluminium foil. The mixture was warmed to rt overnight, treated with aq Na₂S₂O₃ and extracted with Et₂O (2 × 50 mL). Concentration of the organic layers in vacuo yielded the crude product as a brown oil, which was protected against light whilst stored, and used without further purification on the same day. ¹H NMR (400 MHz, CDCl₃) δ 4.80 (m, 1H), 4.50 (m, 1H), 4.31 (s, 5H), 4.30 (m, 1H), 1.49 (d, *J* = 12.4 Hz, 9H) 1.12 (d, *J* = 13.6 Hz, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 78.1, 73.4, 72.7, 71.0, 70.4, 66.2, 28.0, 27.7, 27.5, 26.9; ³¹P NMR (162 MHz, CDCl₃) δ 59.1.

3. Preparation of **4**

General procedure for the Suzuki coupling of arylboronic acids with 2-iodo-1-(diphenylphosphinyl)ferrocene **3a**

To a Schlenk tube were added **3a** (R = Ph, 2.00 mmol), arylboronic acid (4.00 mmol), PdCl₂(PPh₃)₂ (70.2 mg, 0.10 mmol), and the mixture was degassed. 3M NaOH (1.67 mL, 5.00 mmol), and degassed DME (25 mL) were charged, and the mixture heated at 90 °C for 24 h. After cooling to rt, the mixture was diluted with H₂O (25 mL) and extracted with CHCl₃ (3 × 25 mL). The organic layers were dried (MgSO₄) and concentrated in vacuo. Flash chromatography (2:1 EtOAc/hexane) yielded the coupled products **4a** (R = Ph) in 49-86% yields.

General procedure for the Suzuki coupling of arylboronic acids with 2-iodo-1-(dicyclohexylphosphinyl)ferrocene **3b**

Compound **3b** (R = Cy) was prepared as described above, and divided into three portions. To a degassed Schlenk tube were added **3** (R = Cy, 2.54 mmol) [assuming a 100% conversion of **2** (R = Cy) to **3** (R = Cy)], arylboronic acid (3.81 mmol), PdCl₂(PPh₃)₂ (89 mg, 0.13 mmol), and 3M NaOH (2.1 mL, 6.35 mmol). DME (25 mL) previously degassed was charged, and the mixture heated at 90 °C for 24 h. After cooling to rt, the reaction was diluted with H₂O (25 mL) and extracted with CHCl₃ (3 × 25 mL). The organic layers were dried (MgSO₄) and concentrated in vacuo. Flash chromatography (EtOAc/MeOH 98:2) yielded the coupled products **4b** (R = Cy) in 49-56% yields.

General procedure for the Suzuki coupling of arylboronic acids with 2-iodo-1-(di-*t*-butylphosphinyl)ferrocene **3c**

Compound **3c** (R = *t*Bu) was prepared as described above, and divided into three portions. To a degassed Schlenk tube were added **3** (R = *t*Bu, 2.89 mmol) [assuming a 100% conversion of **2** (R = *t*Bu) to **3** (R = *t*Bu)], arylboronic acid (4.33 mmol), PdCl₂(PPh₃)₂ (0.10 g, 0.14 mmol), and 3M NaOH (2.4 mL, 7.22 mmol). DME (30 mL) previously degassed was charged, and the mixture heated at 90 °C for 24 h. After cooling to rt, the reaction was diluted with H₂O (25 mL) and extracted with CHCl₃ (3 × 25 mL). The organic layers were dried (MgSO₄) and concentrated in vacuo. Flash chromatography (EtOAc/MeOH 98:2) yielded the coupled products **4c** (R = *t*Bu) in 58–74% yields.

4. Preparation of **1**

General procedure for reduction of 2-aryl-1-(diphenylphosphinyl)ferrocene **4a** (R = Ph)

To a degassed 100 mL Schlenk tube containing **4** (R = Ph, 2.00 mmol) was charged toluene (30 mL). NEt₃ (1.53 mL, 11.0 mmol) and HSiCl₃ (0.51 mL, 10.0 mmol) were added in turn via syringe, and the mixture heated at 90 °C for 24 h. The reaction was cooled to rt and cautiously treated with aq NaHCO₃ and stirred for 15 min, before extraction with EtOAc (3 × 25 mL). The organic layers were combined, dried (MgSO₄) and concentrated in vacuo. Purification by flash chromatography (hexane/Et₂O 95:5) yielded the 2-aryl-1-(dicyclohexylphosphino)ferrocene compounds **1a**.

2-Phenyl-1-(diphenylphosphino)ferrocene (1ax). The reduction of 2-phenyl-1-(diphenylphosphinyl)ferrocene **4ax** (R = Ph; Ar = Ph) was conducted on a 1.08 mmol scale according to the general procedure to yield 0.44 g (91%) of the title compound as an orange solid. ¹H NMR (400 MHz, CDCl₃) δ 7.65–7.56 (m, 4H), 7.41–7.37 (m, 3H), 7.24–7.14 (m, 8H), 4.75 (m, 1H), 4.40 (dd, *J* = 2.6, 2.6 Hz, 1H), 4.01 (s, 5H), 3.81 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 140.4 (d, *J*_{CP} = 10.4 Hz), 138.5 (d, *J*_{CP} = 8.0 Hz), 135.8 (d, *J*_{CP} = 21.6 Hz), 132.6 (d, *J*_{CP} = 17.6 Hz), 129.6 (d, *J*_{CP} = 6.4 Hz), 129.5, 128.5, 128.4, 128.4, 128.2, 128.1, 126.8, 94.0, 75.1 (d, *J*_{CP} = 12.0 Hz), 72.7 (d, *J*_{CP} = 7.2 Hz), 72.6 (d, *J*_{CP} = 8.0 Hz), 71.0, 70.2; ³¹P NMR (162 MHz, CDCl₃) δ -19.4. Anal. Calcd for C₂₈H₂₃PFe: C, 75.34; H, 5.20. Found: 75.50; H, 5.25.

2-(2-Methoxyphenyl)-1-(diphenylphosphino)ferrocene (1az). The reduction of 2-(2-methoxyphenyl)-1-(diphenylphosphinyl)ferrocene **4az** (R = Ph; Ar = o-MeOC₆H₄) was conducted on a 1.62 mmol scale according to the general procedure to yield 0.59 g (76%) of the title compound as an orange solid. ¹H NMR (400 MHz, CDCl₃) δ 7.98 (dd, *J* = 7.6, 1.3 Hz, 1H), 7.61-7.55 (m, 2H), 7.40-7.37 (m, 3H), 7.21-7.11 (m, 6H), 6.94 (ddd, *J* = 7.6, 7.6, 1.0 Hz, 1H), 6.72 (d, *J* = 7.8 Hz, 1H), 4.69 (m, 1H), 4.43 (dd, *J* = 2.5, 2.5 Hz, 1H), 4.02 (s, 5H), 3.90 (m, 1H), 3.47 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 157.8, 140.6 (d, *J*_{CP} = 9.6 Hz), 139.1 (d, *J*_{CP} = 9.6 Hz), 135.7 (d, *J*_{CP} = 22.0 Hz), 133.9 (d, *J*_{CP} = 7.3 Hz), 132.6 (d, *J*_{CP} = 17.6 Hz), 129.3, 128.5 (d, *J*_{CP} = 7.3 Hz), 128.3, 128.0 (d, *J*_{CP} = 5.9 Hz), 127.6, 123.5, 120.2, 110.8, 100.5, 77.6, 71.4, 70.6, 70.1, 69.8, 55.3; ³¹P NMR (162 MHz, CDCl₃) δ -19.6. Anal. Calcd for C₂₉H₂₅POFe: C, 73.12; H, 5.30. Found: 73.27; H, 5.69.

General procedure for reduction of 2-aryl-1-(dicyclohexylphosphinyl)ferrocene 4b (R = Cy)

To a degassed 100 mL Schlenk tube containing **4b** (R = Cy) (1.00 mmol) was charged with toluene (20 mL). NEt₃ (0.77 mL, 5.50 mmol) and HSiCl₃ (0.51 mL, 5.00 mmol) were added in turn via syringe, and the mixture heated at 90 °C for 24 h. The reaction was cooled to rt and cautiously treated with aq NaHCO₃ and stirred for 15 min, before extraction with EtOAc (3 × 25 mL). The organic layers were combined, dried (MgSO₄) and concentrated in vacuo. Purification by flash chromatography (hexane/EtOAc 3:1) yielded the 2-aryl-1-(dicyclohexylphosphino)ferrocene compounds **1b**.

2-Phenyl-1-(dicyclohexylphosphino)ferrocene (1bx). The reduction of 2-phenyl-1-(dicyclohexylphosphinyl)ferrocene **4bx** (R = Cy; Ar = Ph) was conducted on a 1.26 mmol scale according to the general procedure to yield 0.39 g (68%) of the title compound as a yellow/orange solid. ¹H NMR (400 MHz, CDCl₃) δ 7.75 (d, *J* = 7.0 Hz, 2H), 7.32-7.18 (m, 3H), 4.58 (m, 1H), 4.42 (dd, *J* = 2.5, 2.5 Hz, 1H), 4.20 (m, 1H), 4.08 (s, 5H), 2.61-2.57 (m, 1H), 2.05-1.76 (m, 5H), 1.58-1.18 (m, 11H), 1.00-0.76 (m, 3H), 0.61-0.46 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 139.6, 130.6 (d, *J*_{CP} = 7.2 Hz), 127.9, 126.6, 92.9 (d, *J*_{CP} = 16.8 Hz), 77.3, 71.2, 70.7 (d, *J*_{CP} = 4.8 Hz), 69.5 (d, *J*_{CP} = 2.4 Hz), 69.1, 36.1 (d, *J*_{CP} = 15.2 Hz), 33.4 (d, *J*_{CP} = 10.4 Hz), 32.7 (d, *J*_{CP} = 18.4 Hz), 30.7 (d, *J*_{CP} = 15.2 Hz), 30.1 (d, *J*_{CP} = 10.4 Hz), 29.2 (d, *J*_{CP} = 4.8 Hz), 28.1, 28.0 (d, *J*_{CP} = 2.4 Hz), 27.8, 27.8, 27.7, 26.8 (d, *J*_{CP} = 30.4 Hz); ³¹P NMR (162 MHz, CDCl₃) δ -12.9. Anal. Calcd for C₂₈H₃₅PFe: C, 73.29; H, 7.71. Found: 73.19; H, 7.80.

2-(2-Methylphenyl)-1-(dicyclohexylphosphino)ferrocene (1by). The reduction of 2-(2-methylphenyl)-1-(dicyclohexylphosphinyl)ferrocene **4by** (R = Cy; Ar = o-MeC₆H₄) was conducted on a 1.25 mmol scale according to the general procedure to yield 0.43 g (73%) of the title compound as a yellow/orange solid. ¹H NMR (400 MHz, CDCl₃) δ 8.13 (d, *J* = 7.6 Hz, 1H), 7.35-7.09 (m, 3H), 4.50 (m, 1H), 4.43 (dd, *J* = 2.4, 2.4 Hz, 1H), 4.19 (m, 1H), 4.16 (s, 5H), 2.59-2.52 (m, 1H), 2.17 (s, 3H), 2.01-1.75 (m, 6H), 1.56-1.07 (m, 9H), 0.92-0.81 (m, 4H), 0.63-0.48 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 137.3, 137.0, 134.6 (d, *J*_{CP} = 8.0 Hz), 130.0, 127.0, 125.3, 93.3, 79.0, 71.9 (d, *J*_{CP} = 2.4 Hz), 71.0, 69.5

(d, $J_{\text{CP}} = 4.0$ Hz), 68.3, 35.7 (d, $J_{\text{CP}} = 15.2$ Hz), 33.5 (d, $J_{\text{CP}} = 10.4$ Hz), 32.7 (d, $J_{\text{CP}} = 19.2$ Hz), 30.2 (d, $J_{\text{CP}} = 10.4$ Hz), 29.7 (d, $J_{\text{CP}} = 11.2$ Hz), 29.7 (d, $J_{\text{CP}} = 16.8$ Hz), 28.0, 27.9, 27.9 (d, $J_{\text{CP}} = 7.2$ Hz), 27.8, 27.7 (d, $J_{\text{CP}} = 8.8$ Hz), 26.8 (d, $J_{\text{CP}} = 20.8$ Hz) 21.3; ^{31}P NMR (162 MHz, CDCl_3) δ -12.0. Anal. Calcd for $\text{C}_{29}\text{H}_{37}\text{PFe}$: C, 73.72; H, 7.91. Found: 73.97; H, 8.04.

2-(2-Methoxyphenyl)-1-(dicyclohexylphosphino)ferrocene (1bz). The reduction of 2-(2-methoxyphenyl)-1-(dicyclohexylphosphinyl)ferrocene **4bz** ($\text{R} = \text{Cy}$; $\text{Ar} = o\text{-MeOC}_6\text{H}_4$) was conducted on a 1.38 mmol scale according to the general procedure to yield 0.49 g (73%) of the title compound as a yellow/orange solid. ^1H NMR (400 MHz, CDCl_3) δ 8.12 (ddd, $J = 7.5, 1.3, 1.3$ Hz, 1H), 7.22 (ddd, 8.2, 7.5, 1.8 Hz, 1H), 7.01 (ddd, 7.5, 7.5, 1.0 Hz, 1H), 6.81 (d, 8.2 Hz, 1H), 4.68 (m, 1H), 4.43 (ddd, $J = 2.5, 2.5, 0.6$ Hz, 1H), 4.18 (m, 1H), 4.12 (s, 5H), 3.70 (s, 3H), 2.59-2.50 (m, 1H), 2.02-1.69 (m, 5H), 1.55-1.06 (m, 10H), 1.01-0.78 (m, 4H), 0.60-0.48 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 158.1, 135.2 (d, $J_{\text{CP}} = 9.6$ Hz), 128.2, 127.1, 120.2, 110.8, 83.1, 77.6, 72.9, 70.9, 69.4 (d, $J_{\text{CP}} = 4.0$ Hz), 68.5, 55.8, 35.6 (d, $J_{\text{CP}} = 14.4$ Hz), 33.3 (d, $J_{\text{CP}} = 10.4$ Hz), 32.7 (d, $J_{\text{CP}} = 19.2$ Hz), 30.3, 30.2, 30.1 (d, $J_{\text{CP}} = 10.4$ Hz), 29.0, 28.1, 28.0, 27.8, 27.8 (d, $J_{\text{CP}} = 13.6$ Hz), 26.9 (d, $J_{\text{CP}} = 26.4$ Hz); ^{31}P NMR (162 MHz, CDCl_3) δ -12.2. Anal. Calcd for $\text{C}_{29}\text{H}_{37}\text{POFe}$: C, 71.30; H, 7.64. Found: 71.52; H, 7.72.

5. Suzuki coupling

General Procedure for the Suzuki Coupling of Aryl Chlorides at 0.1 mol% Catalyst Loading

An oven-dried carousel reaction tube was charged with boronic acid (1.5 mmol), base (3 mmol), and ligand **1** (2×10^{-3} mmol). The reaction vessel was evacuated and backfilled with N_2 . This process was repeated five times. Through a rubber septum, a 1.0 M solution of the aryl chloride (1 mmol) along with 2×10^{-3} M solution of $\text{Pd}(\text{OAc})_2$ (1×10^{-3} mmol, 0.1 mol%) was introduced. Solvent was added to ensure that 5 mL of solution was used in each reaction. The reactions were heated at the stated temperature for the stated time. Conversions were calculated from crude ^1H NMR, and isolated yields were obtained after standard work-up and purification by flash chromatography (100% hexane).