First stereospecific synthesis of (*E*) or (*Z*) α -fluoroenones *via* a kinetically controlled Negishi coupling reaction

Guillaume Dutheuil, Clotilde Paturel, Xinsheng Lei, Samuel Couve-Bonnaire and Xavier Pannecoucke*

IRCOF-ECOFH, UMR CNRS 6014, INSA de ROUEN, rue Tesnière, 76131 Mont-Saint-Aignan, *France*.

Fax: (33) 2.35.52.29.59; Tel: (33) 2.35.52.24.27; E-mail: xavier.pannecoucke@insa-rouen.fr

Table of contents:

S4	General experimental information
S5	Experimental procedure for the obtention of 4a(Z) from <i>E/Z</i> mixture of bromofluoroolefin 2
S6	^1H , ^{13}C and ^{19}F NMR spectra of compound 2₁(Z)
S7	Characterization of compound 4a₁(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S8	^1H , ^{13}C and ^{19}F NMR spectra of compound 2₂(Z)
S9	Characterization of compound 4a₂(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S10	^1H , ^{13}C and ^{19}F NMR spectra of compound 2₃(Z)
S11	Characterization of compound 4a₃(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S12	^1H , ^{13}C and ^{19}F NMR spectra of compound 2₄(Z)
S13	Characterization of compound 4a₄(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S14	^1H , ^{13}C and ^{19}F NMR spectra of compound 2₅(Z)
S15	Characterization of compound 4a₅(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S16	^1H , ^{13}C and ^{19}F NMR spectra of compound 2₆(Z)
S17	Characterization of compound 4a₆(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra

S18	^1H , ^{13}C and ^{19}F NMR spectra of compound 2₇(Z)
S19	Characterization of compound 4a₇(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S20	^1H , ^{13}C and ^{19}F NMR spectra of compound 2₈(Z)
S21	Characterization of compound 4a₈(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S22	Experimental procedure for the obtention of 4b₁(Z) from <i>E/Z</i> mixture of bromofluoroolefin 2
S23	Characterization of compound 4b₁(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S24	Experimental procedure for the obtention of 4b₂(Z) from <i>E/Z</i> mixture of bromofluoroolefin 2
S25	Characterization of compound 4b₂(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S26	Experimental procedure for the obtention of 4c(Z) from <i>E/Z</i> mixture of bromofluoroolefin 2
S27	Characterization of compound 4c₁(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S28	Characterization of compound 4c₂(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S29	Experimental procedure for the obtention of 7(Z) from <i>E/Z</i> mixture of bromofluoroolefin 2
S30	Characterization of compound 7₁(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S31	Characterization of compound 7₂(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S32	Experimental procedure for the obtention of 4d
S33	Characterization of compound 4d₁(Z) ; ^1H , ^{13}C and ^{19}F NMR spectra
S34	Experimental procedure for the obtention of 4a(E) from bromofluoroolefin 2(Z)
S35	Characterization of compound 4a₁(E) ; ^1H , ^{13}C and ^{19}F NMR spectra
S36	Characterization of compound 4a₂(E) ; ^1H , ^{13}C and ^{19}F NMR spectra
S37	Characterization of compound 4a₃(E) ; ^1H , ^{13}C and ^{19}F NMR spectra
S38	Characterization of compound 4a₄(E) ; ^1H , ^{13}C and ^{19}F NMR spectra
S39	Characterization of compound 4a₅(E) ; ^1H , ^{13}C and ^{19}F NMR spectra
S40	Characterization of compound 4a₇(E) ; ^1H , ^{13}C and ^{19}F NMR spectra
S41	Characterization of compound 4a₈(E) ; ^1H , ^{13}C and ^{19}F NMR spectra

S42	Experimental procedure for the obtention of 4b(E) from bromofluoroolefin 2(Z)
S43	Characterization of compound 4b₂(E) ; ¹ H, ¹³ C and ¹⁹ F NMR spectra
S44	Experimental procedure for the obtention of 4c(E) from bromofluoroolefin 2(Z)
S45	Characterization of compound 4c₁(E) ; ¹ H, ¹³ C and ¹⁹ F NMR spectra
S46	Characterization of compound 4c₂(E) ; ¹ H, ¹³ C and ¹⁹ F NMR spectra
S47	Experimental procedure for the obtention of 7(E) from bromofluoroolefin 2(Z)
S48	Characterization of compound 7₁(E) ; ¹ H, ¹³ C and ¹⁹ F NMR spectra
S49	Characterization of compound 7₂(E) ; ¹ H, ¹³ C and ¹⁹ F NMR spectra
S50	Characterization of compound 4d₁(E) ; ¹ H, ¹³ C and ¹⁹ F NMR spectra
S51	Characterization of compound 10 (R, Z) ; ¹ H, ¹³ C and ¹⁹ F NMR spectra
S52	Characterization of compound 10 (R, E) ; ¹ H, ¹³ C and ¹⁹ F NMR spectra
S53	Determination of the enantiomeric purities of 10 (R, Z) and 10 (R, E) ; ¹⁹ F NMR spectra

First stereospecific synthesis of (*E*) or (*Z*) α -fluoroenones *via* a kinetically controlled Negishi coupling reaction

Guillaume Dutheuil, Clotilde Paturel, Xinsheng Lei, Samuel Couve-Bonnaire and Xavier Pannecoucke*

IRCOF-ECOFH, UMR CNRS 6014, INSA de ROUEN, rue Tesnière, 76131 Mont-Saint-Aignan, *France*.

Fax: (33) 2.35.52.29.59; Tel: (33) 2.35.52.24.27; E-mail: xavier.pannecoucke@insa-rouen.fr

Experimental Section.

General: ^{19}F 4 (282.5 MHz), ^{13}C NMR (75.5 MHz) and ^1H NMR (300 MHz) spectra were recorded on a 300 MHz spectrometer. All samples were taken in CDCl_3 solvent unless noted otherwise. All chemical shifts are reported in parts per million downfield (positive) of the standard : TMS for ^1H and ^{13}C , CFCl_3 for ^{19}F NMR. Coupling constants (*J*) are given in cycles per second (Hz). ^1H - and ^{13}C -NMR peaks were assigned using standard COSY, HMQC and HMBC experiments. All reactions were carried out in oven-dried glassware that was placed under vacuum while hot and cooled under argon. Column Chromatographies were performed on silica gel (70-230 mesh). THF was dried by refluxing and distilling under argon from sodium benzophenone dianion. All reagents were obtained from commercial sources, or prepared from the literature. The bromofluorolefins were prepared by our method.^{7a}

Representative procedure for the obtention of (Z) fluorinated methylketones **4a(Z)** from an E/Z mixture of bromofluoroolefins **2**. To a mixture of potassium *tert*-butoxide (0.85 eq.), ethylvinylether (0.85 eq.) in anhydrous THF (5 mL/mmol of *t*-BuOK) at -78°C under argon was added dropwise *n*-BuLi in hexanes (1.6M, 0.85 eq.). The mixture was stirred for 30 min at -78°C and then a solution of dry zincchloride (1.7 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.025 eq.), triphenylphosphine (0.05 eq.) and mixture of bromofluoroolefins **2** (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolefin) at 10°C under argon. The mixture was stirred for 1h. HCl aq. (1N) was added and the solution was stirred for 15 min. The mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on silica gel (eluent: cyclohexane/AcOEt), affording fluorenones **4a(Z)** and unreacted bromofluoroolefin **2(Z)**.

Compound **2₁(Z)**^{7a}

Compound 4a₁(Z)

(Z)-3-Fluoro-4-(4-methoxyphenyl)but-3-en-2-one. Yellow powder (m. p. = 72-74°C), R_f 0.2 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.6 (d, J = 8.9Hz, 2H, H_4), 6.9 (d, J = 8.9Hz, 2H, H_5), 6.7 (d, $^3J_{\text{H-F}}$ = 36.9Hz, 1H, H_2), 3.7 (s, 3H, H_7), 2.3 (d, $^4J_{\text{H-F}}$ = 3.3Hz, 3H, H_9). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -127.1 (dq, $^3J_{\text{H-F}}$ = 36.9Hz, $^4J_{\text{H-F}}$ = 3.3Hz). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.2 (d, $^2J_{\text{C-F}}$ = 32Hz, C_8), 161.2 (d, J = 4Hz, C_6), 153.5 (d, $^1J_{\text{C-F}}$ = 267Hz, C_1), 135.5 (d, J = 6Hz, C_4), 124.0 (d, $^3J_{\text{C-F}}$ = 4Hz, C_3), 116.4 (d, $^2J_{\text{C-F}}$ = 6Hz, C_2), 114.5 (d, J = 18Hz, C_5), 55.7 (C_7), 25.9 (C_9). MS (EI) 194 (M^+), 179 ($\text{M}^+ - \text{CH}_3$), 159, 43 (CH_3CO^+). IR (KBr) 3037, 2970, 2937, 2898, 2837, 1672, 1643, 1605, 1570, 1513, 1424, 1343, 1264, 1185, 1177, 1069, 1022, 869, 808, 539 cm^{-1} . Anal. Calcd for $\text{C}_{11}\text{H}_{11}\text{FO}_2$: C, 68.03; H, 5.71. Found: C, 68.06; H, 5.63.

Compound $2_2(\text{Z})^{7a}$

Compound 4a₂(Z)

(Z)-3-Fluoro-4-(naphthalen-7-yl)but-3-en-2-one. Yellow powder (m. p. = 85-87°C), R_f 0.3 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.0 (s, 1H, H_{12}), 7.8-7.6 (m, 4H, $\text{H}_{7,8,9,10}$), 7.5-7.3 (m, 2H, $\text{H}_{4,5}$), 6.9 (d, $^3J_{\text{H-F}} = 36.6\text{Hz}$, 1H, H_2), 2.3 (d, $^4J_{\text{H-F}} = 3.3\text{Hz}$, 3H, H_{14}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -123.8 (dq, $^3J_{\text{H-F}} = 36.6\text{Hz}$, $^4J_{\text{H-F}} = 3.3\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.5 (d, $^2J_{\text{C-F}} = 33\text{Hz}$, C_{13}), 154.5 (d, $^1J_{\text{C-F}} = 272\text{Hz}$, C_1), 134.1 (d, $J = 2\text{Hz}$, $\text{C}_{\text{ar}}^{\text{q}}$), 133.6 ($\text{C}_{\text{ar}}^{\text{q}}$), 131.8 (d, $J = 8\text{Hz}$, $\text{C}_{\text{ar}}^{\text{H}}$), 129.03 ($\text{C}_{\text{ar}}^{\text{H}}$), 129.0 ($\text{C}_{\text{ar}}^{\text{q}}$), 128.9 ($\text{C}_{\text{ar}}^{\text{H}}$), 128.1 ($\text{C}_{\text{ar}}^{\text{H}}$), 127.8 ($\text{C}_{\text{ar}}^{\text{H}}$), 127.4 (d, $J = 9\text{Hz}$, $\text{C}_{\text{ar}}^{\text{H}}$), 127.0 ($\text{C}_{\text{ar}}^{\text{H}}$), 116.2 (d, $^2J_{\text{C-F}} = 6\text{Hz}$, C_2), 26.1 (C_{14}). MS (EI) 214 (M^+), 199 ($\text{M}^+ - \text{CH}_3$), 179, 171 ($\text{M}^+ - \text{CH}_3\text{CO}$), 43 (CH_3CO^+). IR (KBr) δ 3050, 1702, 1683, 1626, 1365, 1335, 1320, 1243, 1069, 914, 829, 751, 625, 483. Anal. Calcd for $\text{C}_{14}\text{H}_{11}\text{FO}$: C, 78.49; H, 5.18. Found: C, 78.57; H, 5.15.

Compound $2_3(\text{Z})^{7a}$

Compound 4a3(Z)

(Z) 3-Fluoro-4-(4-cyanophenyl)but-3-en-2-one. Yellow powder (m. p. = 119-121°C), R_f 0.3 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.6 (d, J = 8.5Hz, 2H, H_5), 7.1 (d, J = 8.5Hz, 2H, H_4), 6.8 (d, $^3J_{\text{H-F}}$ = 35.3Hz, 1H, H_2), 2.4 (d, $^4J_{\text{H-F}}$ = 3.3Hz, 3H, H_9). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -119.6 (dq, $^3J_{\text{H-F}}$ = 35.3Hz, $^4J_{\text{H-F}}$ = 3.3Hz). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.4 (d, $^2J_{\text{C-F}}$ = 35Hz, C_8), 155.5 (d, $^1J_{\text{C-F}}$ = 278Hz, C_1), 135.8 (d, $^3J_{\text{C-F}}$ = 4Hz, C_3), 132.9 (C_5), 131.2 (d, J = 9Hz, C_4), 118.7 (C_7), 113.4 (d, $^2J_{\text{C-F}}$ = 5Hz, C_2), 113.3 (d, J = 4Hz, C_6), 26.1 (C_9). MS (EI) 189 (M^+), 174 ($\text{M}^+ - \text{CH}_3$), 146 ($\text{M}^+ - \text{CH}_3\text{CO}$), 43 (CH_3CO^+). IR (KBr) 3075, 2987, 2227, 1681, 1648, 1505, 1418, 1382, 1351, 1328, 1293, 1255, 1212, 1081, 968, 834, 616, 561. *Anal. Calcd for $\text{C}_{11}\text{H}_8\text{FNO}$* : C, 69.84; H, 4.26; N, 7.40. *Found* : C, 69.67; H, 4.06; N, 7.24.

Compound **2₄(Z)**^{7a}

Compound 4a₄(Z)

(Z) 3-Fluoro-4-(4-methylbenzoate)but-3-en-2-one. Yellow solid (m. p. = 103–105°C), R_f 0.2 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.0 (d, J = 8.5 Hz, 2H, H_5), 7.6 (d, J = 8.5 Hz, 2H, H_4), 6.8 (d, $^3J_{\text{H-F}}$ = 35.8 Hz, 1H, H_2), 3.9 (s, 3H, H_8), 2.4 (d, $^4J_{\text{H-F}}$ = 3.6 Hz, 3H, H_{10}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -121.1 (dq, $^3J_{\text{H-F}}$ = 35.8 Hz, $^4J_{\text{H-F}}$ = 3.6 Hz). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.7 (d, $^2J_{\text{C-F}}$ = 35 Hz, C_9), 166.8 (C_7), 155.1 (d, $^1J_{\text{C-F}}$ = 277 Hz, C_1), 135.7 (d, J = 4 Hz, C_6), 131.2 (d, $^3J_{\text{C-F}}$ = 3 Hz, C_3), 130.8 (d, J = 9 Hz, C_4), 130.3 (C_5), 114.4 (d, $^2J_{\text{C-F}}$ = 6 Hz, C_2), 52.7 (C_8), 26.2 (C_{10}). MS (EI) 222 (M^+), 207 ($\text{M}^+ - \text{CH}_3$), 191 ($\text{M}^+ - \text{CH}_3\text{O}$), 163 ($\text{M}^+ - \text{CO}_2\text{CH}_3$), 43 (CH_3CO^+). IR (KBr) 3056, 2959, 1721, 1636, 1433, 1364, 1320, 1280, 1192, 1109, 1057, 1019, 968, 904, 832, 769, 697, 571. *Anal. Calcd for $\text{C}_{12}\text{H}_{11}\text{FO}_3$* : C, 64.86; H, 4.99. *Found*: C, 64.98; H, 4.90.

Compound **2₅(Z)**^{7a}

Compound 4a₅(Z)

(Z) 3-Fluoro-4-(4-nitrophenyl)but-3-en-2-one. Yellow solid (m. p. = 139-141°C), R_f 0.1 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.2 (d, J = 8.8Hz, 2H, H_5), 7.8 (d, J = 8.8Hz, 2H, H_4), 6.8 (d, $^3J_{\text{H-F}}$ = 35.0Hz, 1H, H_2), 2.5 (d, $^4J_{\text{H-F}}$ = 3.8Hz, 3H, H_8). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -119.2 (dq, $^3J_{\text{H-F}}$ = 35.5Hz, $^4J_{\text{H-F}}$ = 3.3Hz). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.2 (d, $^2J_{\text{C-F}}$ = 34Hz, C_7), 155.3 (d, $^1J_{\text{C-F}}$ = 280Hz, C_1), 147.9 (d, J = 3Hz, C_6), 137.3 (d, J = 4Hz, C_3), 131.3 (d, J = 13Hz, C_5), 123.3 (C_4), 117.5 (d, $^2J_{\text{C-F}}$ = 5Hz, C_2), 25.9 (C_8). MS (EI) 209 (M^+), 192, 162 ($\text{M}^+ - \text{HNO}_2$), 148. IR (KBr) 3306, 3007, 1707, 1639, 1595, 1511, 1340, 1304, 1268, 1200, 690. Anal. Calcd for $\text{C}_{10}\text{H}_8\text{FNO}_3$: C, 57.42; H, 3.85; N, 6.70. Found: C, 57.78; H, 3.79; N, 6.64.

Compound **2₆(Z)**^{7a}

Compound 4a₆(Z)

(Z) 3-Fluoro-4-(4-bromophenyl)but-3-en-2-one. Yellow solid (m. p. = 88-90°C), R_f 0.3 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.5 (b, 4H, $\text{H}_{4,5}$), 6.7 (d, $^3J_{\text{H-F}} = 35.8\text{Hz}$, 1H, H_2), 2.4 (d, $^4J_{\text{H-F}} = 3.6\text{Hz}$, 3H, H_8). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -122.9 (dq, $^3J_{\text{H-F}} = 35.5\text{Hz}$, $^4J_{\text{H-F}} = 3.2\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.3 (d, $^2J_{\text{C-F}} = 34\text{Hz}$, C_7), 154.2 (d, $^1J_{\text{C-F}} = 274\text{Hz}$, C_1), 132.2 (C_5), 132.0 (d, $J = 9\text{Hz}$, C_4), 130.0 (d, $^3J_{\text{C-F}} = 5\text{Hz}$, C_3), 124.4 (d, $J = 4\text{Hz}$, C_6), 114.3 (d, $^2J_{\text{C-F}} = 6\text{Hz}$, C_2), 25.8 (C_8). MS (EI) 244-242 (M^+), 163 ($\text{M}^+ - \text{Br}$), 148 ($\text{M}^+ - \text{Br} + \text{CH}_3$), 120 ($\text{M}^+ - \text{Br} + \text{COCH}_3$). IR (KBr) 3021, 2959, 1700, 1634, 1360, 1310, 1267, 1197, 1056, 819. Anal. Calcd for $\text{C}_{10}\text{H}_8\text{BrFO}$: C, 49.41; H, 3.32. Found: C, 49.52; H, 3.44.

Compound **2₇(Z)**^{7a}

Compound 4a₇(Z)

(Z) 3-Fluoro-6-phenylhex-3-en-2-one. Yellow oil, R_f 0.3 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.3-7.0 (m, 5H, $\text{H}_{6,7,8}$), 6.0 (dt, $^3J_{\text{H-F}} = 33.8\text{Hz}$, $^3J_{\text{H-H}} = 7.4\text{Hz}$, 1H, H_2), 2.7 (t, $^3J_{\text{H-H}} = 7.4\text{Hz}$, 2H, H_4), 2.6-2.4 (m, 2H, H_3), 2.3 (dd, $^4J_{\text{H-F}} = 2.8\text{Hz}$, $J = 0.8\text{Hz}$, 3H, H_{10}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -127.8 (dq, $^3J_{\text{H-F}} = 33.8\text{Hz}$, $^4J_{\text{H-F}} = 2.8\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 191.9 (d, $^2J_{\text{C-F}} = 32\text{Hz}$, C_9), 155.7 (d, $^1J_{\text{C-F}} = 261\text{Hz}$, C_1), 140.9 (C_5), 128.9 ($\text{C}_{6/7}$), 128.7 ($\text{C}_{6/7}$), 126.7 (C_8), 119.0 (d, $^2J_{\text{C-F}} = 13\text{Hz}$, C_2), 34.8 (d, $^4J_{\text{C-F}} = 2\text{Hz}$, C_4), 26.3 (d, $^3J_{\text{C-F}} = 3\text{Hz}$, C_3), 25.9 (C_{10}). MS (EI) 192 (M^+), 172, 91 (PhCH_2^+), 43 (CH_3CO^+). IR (neat) 3063, 3027, 2929, 2862, 1713, 1693, 1651, 1496, 1454, 1361, 1287, 1099, 750, 700. *Anal. Calcd for $\text{C}_{12}\text{H}_{13}\text{FO}$* : C, 74.98; H, 6.82. *Found* : C, 74.75; H, 6.85.

Compound $2_8(\text{Z})^{7a}$

Compound 4a₈(Z)

(Z) 3-Fluoro-6-(oxytert-butylidiphenylsilyl)hex-3-en-2-one. Yellow oil, R_f 0.2 (2% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.7-7.5 (m, 4H, H_6), 7.4-7.2 (m, 6H, $\text{H}_{7,8}$), 6.1 (dt, $^3J_{\text{H-F}} = 34.3\text{Hz}$, $^3J_{\text{H-H}} = 7.7\text{Hz}$, 1H, H_2), 3.7 (t, $^3J_{\text{H-H}} = 6.4\text{Hz}$, 2H, H_4), 2.5-2.3 (m, 2H, H_3), 2.2 (d, $^4J_{\text{H-F}} = 2.6\text{Hz}$, 3H, H_{12}), 0.9 (s, 9H, H_{10}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -127.4 (m). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 191.7 (d, $^2J_{\text{C-F}} = 31\text{Hz}$, C_{11}), 156.2 (d, $^1J_{\text{C-F}} = 261\text{Hz}$, C_1), 136.0 (C_6), 133.9 (C_5), 130.2 (C_8), 128.2 (C_7), 117.4 (d, $^2J_{\text{C-F}} = 13\text{Hz}$, C_2), 62.4 (d, $^4J_{\text{C-F}} = 2\text{Hz}$, C_4), 28.3 (d, $^3J_{\text{C-F}} = 3\text{Hz}$, C_3), 27.2 (C_{10}), 25.9 (C_{12}), 19.6 (C_9). MS (EI) 313 (M^+ -tBu), 201, 199, 181, 141, 43 (CH_3CO^+). IR (neat) 3410, 3072, 2932, 2858, 1722, 1697, 1471, 1428, 1362, 1114, 821, 701, 612, 509. *Anal. Calcd for $\text{C}_{22}\text{H}_{27}\text{FO}_2\text{Si}$* : C, 71.31; H, 7.34. *Found*: C, 71.55; H, 7.16.

Procedure for the obtention of (Z) 3-Fluoro-4-(4-methoxyphenyl)-1-phenylbut-3-en-2-one 4b₁ from an E/Z mixture of bromofluoroolefin:

With recovering of (Z) starting bromofluoroolefin:

To a mixture of potassium *tert*-butoxide (2 eq.) and freshly synthesized 1-(2-Ethoxyvinyl)benzene (2 eq.) in anhydrous THF (5 mL/mmol of *t*-BuOK) at -78°C under argon was added dropwise *n*-BuLi in hexanes (1.6M, 2 eq.). The mixture was stirred for 30 min at -78°C and then a solution of dry zincchloride (4 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.025 eq.), triphenylphosphine (0.05 eq.) and (Z) & (E) 1-(2-Bromo-2-fluorovinyl)-4-methoxybenzene (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolein) at room temperature under argon. The mixture was stirred for 1h. After the coupling reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, HCl aq. (6M) was added and the mixture was stirred and heated to 70°C for 1h. After cooling to room temperature, the mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on silica gel (eluent: 2% AcOEt in cyclohexane), affording (Z) 1-(2-Bromo-2-fluorovinyl)-4-methoxybenzene (yield = quant.) and (Z) 3-Fluoro-4-(4-methoxyphenyl)-1-phenylbut-3-en-2one (yield = 93%).

With isomerization of (E) enol ether-diene intermediate:

To a mixture of potassium *tert*-butoxide (4 eq.), 1-(2-Ethoxyvinyl)benzene (4 eq.) in anhydrous THF (5 mL/mmol of *t*-BuOK) at -78°C under argon was added dropwise *n*-BuLi in hexanes (1.6M, 4 eq.). The mixture was stirred for 30 min at -78°C and then a solution of dry zincchloride (8 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.05 eq.), triphenylphosphine (0.1 eq.) and (Z) & (E) 1-(2-Bromo-2-fluorovinyl)-4-methoxybenzene (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolein) at 70°C under argon. The mixture was stirred for 1h. After the coupling reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, HCl aq. (6M) was added and the mixture was stirred at 70°C for another hour. After cooling to room temperature, the mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on silica gel (eluent: 2% AcOEt in cyclohexane), affording (Z) 3-Fluoro-4-(4-methoxyphenyl)-1-phenylbut-3-en-2one (yield = 99%).

Compound 4b₁(Z)

(Z) 3-Fluoro-4-(4-methoxyphenyl)-1-phenylbut-3-en-2-one. Yellow solid (m. p. = 90-92°C), R_f 0.2 (2% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.6 (d, J = 8.7Hz, 2H, H_4), 7.3-7.2 (m, 5H, $\text{H}_{11, 12, 13}$), 6.9 (d, J = 8.2Hz, 2H, H_5), 6.8 (d, $^3J_{\text{H-F}}$ = 40.7Hz, 1H, H_2), 3.9 (d, $^4J_{\text{H-F}}$ = 3.1Hz, 2H, H_9), 3.8 (s, 3H, H_7). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -127.9 (dt, $^3J_{\text{H-F}}$ = 38.2Hz, $^4J_{\text{H-F}}$ = 3.1Hz). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.4 (d, $^2J_{\text{C-F}}$ = 33Hz, C_8), 161.4 (d, J = 4Hz, C_6), 153.3 (d, $^1J_{\text{C-F}}$ = 269Hz, C_1), 133.8 (C_{10}), 133.0 (d, J = 9Hz, C_4), 130.0 (C_{12}), 129.1 (C_{11}), 127.5 (C_{13}), 124.2 (d, $^3J_{\text{C-F}}$ = 4Hz, C_3), 116.6 (d, $^2J_{\text{C-F}}$ = 5Hz, C_2), 114.8 (C_5), 55.8 (C_7), 45.2 (C_9). MS (EI) 270 (M^+), 179 ($\text{M}^+ - \text{PhCH}_2$), 151 ($\text{M}^+ - \text{PhCH}_2\text{CO}$). IR (KBr) 3060, 3041, 2968, 2932, 2822, 1703, 1633, 1599, 1513, 1338, 1252, 1167, 1030, 837, 707, 533. Anal. Calcd for $\text{C}_{17}\text{H}_{15}\text{FO}_2$: C, 75.54; H, 5.59. Found: C, 75.57; H, 6.02.

Procedure for the obtention of (Z) fluorinated benzylketones **4b₂(Z) from mixture of bromofluoroolefins **2**.**

To a mixture of potassium *tert*-butoxide (2 eq.), freshly synthesized 1-(2-Ethoxyvinyl)benzene (2 eq.) in anhydrous THF (5 mL/mmol of *t*-BuOK) at -78°C under argon was added dropwise *n*-BuLi in hexanes (1.6M, 2 eq.). The mixture was stirred for 30 min at -78°C and then a solution of dry zincchloride (4 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.025 eq.), triphenylphosphine (0.05 eq.) and mixture of bromofluoroolefins **2** (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolefin) at 10°C under argon. The mixture was stirred for 1h. After the coupling reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, HCl aq. (1N) was added and the mixture was stirred at room temperature for 1h. The mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on silica gel (eluent: 5% AcOEt in cyclohexane), affording (Z) starting bromofluoroolefin **2**(Z) and (Z) benzylketone **4b**(Z).

Compound 4b₂(Z)

(Z) 3-Fluoro-1,6-diphenylhex-3-en-2-one. Orange oil, R_f 0.4 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.3-7.1 (m, 10H, $\text{H}_6, 7, 8, 12, 13, 14$), 6.0 (dt, $^3J_{\text{H-F}} = 34.1\text{Hz}$, $^4J_{\text{H-F}} = 7.7\text{Hz}$, 1H, H_2), 3.8 (d, $^4J_{\text{H-F}} = 2.3\text{Hz}$, 2H, H_{10}), 2.7 (t, $^3J_{\text{H-H}} = 7.5\text{Hz}$, 2H, H_4), 2.5-2.4 (m, 2H, H_3). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -128.3 (dt, $^3J_{\text{H-F}} = 34.4\text{Hz}$, $^4J_{\text{H-F}} = 2.3\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 191.7 (d, $^2J_{\text{C-F}} = 32\text{Hz}$, C_9), 155.4 (d, $^1J_{\text{C-F}} = 262\text{Hz}$, C_1), 140.9 (C_5), 133.6 (C_{11}), 129.9 ($\text{C}_{12/13}$), 129.1 ($\text{C}_{12/13}$), 129.0 ($\text{C}_{6/7}$), 128.7 ($\text{C}_{6/7}$), 127.6 (C_{14}), 126.7 (C_8), 119.5 (d, $^2J_{\text{C-F}} = 12\text{Hz}$, C_2), 45.2 (C_{10}), 34.8 (C_4), 26.5 (C_3). MS (EI) 268 (M^+), 177 ($\text{M}^+ - \text{PhCH}_2$), 157, 91 (PhCH_2^+), 65. IR (neat) 3085, 3062, 3028, 2928, 2860, 1702, 1687, 1650, 1495, 1454, 1297, 1030, 748, 699. Anal. Calcd for $\text{C}_{18}\text{H}_{17}\text{FO}$: C, 80.57; H, 6.39. Found: C, 80.69; H, 6.52.

Procedure for the obtention of (Z) fluorinated propylketones 4c(Z) from mixture of bromofluoroolefins 2.

To a mixture of potassium *tert*-butoxide (2 eq.), butenylethylether (2 eq.) in anhydrous THF (5 mL/mmol of *t*-BuOK) at -78°C under argon was added dropwise *n*-BuLi in hexanes (1.6M, 2 eq.). The mixture was stirred for 30 min at -78°C and then a solution of dry zincchloride (4 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.025 eq.), triphenylphosphine (0.05 eq.) and mixture of bromofluoroolefins **2** (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolefin) at 10°C under argon. The mixture was stirred for 1h. After the reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, HCl aq. (1N) was added and the mixture was stirred for 15 min. The mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on silica gel (eluent: cyclohexane/AcOEt), affording (Z) starting bromofluoroolefin **2(Z)** and (Z) propylketone **4c(Z)**.

Compound 4c₁(Z)

(Z) 2-Fluoro-1-(4-methoxyphenyl)hex-1-en-3-one. White solid (m. p. = 77-79°C), *R_f* 0.3 (5% AcOEt in cyclohexane). ¹H-NMR (300 MHz, CDCl₃) δ 7.6 (d, *J* = 9.0Hz, 2H, H₄), 6.9 (d, *J* = 9.0Hz, 2H, H₅), 6.8 (d, ³*J*_{H-F} = 37.3Hz, 1H, H₂), 3.8 (s, 3H, H₇), 2.7 (dt, ⁴*J*_{H-F} = 2.3Hz, ³*J*_{H-H} = 7.5Hz, 2H, H₉), 1.8-1.6 (m, 2H, H₁₀), 1.0 (t, ³*J*_{H-H} = 7.2Hz, 3H, H₁₁). ¹⁹F-NMR (282.5 MHz, CDCl₃) δ -128.6 (d, ³*J*_{H-F} = 36.7Hz). ¹³C-NMR (75.5 MHz, CDCl₃) δ 195.2 (d, ²*J*_{C-F} = 31Hz, C₈), 161.2 (C₆), 153.6 (d, ¹*J*_{C-F} = 269Hz, C₁), 132.8 (d, *J* = 9Hz, C₄), 124.3 (d, ³*J*_{C-F} = 4Hz, C₃), 115.5 (d, ²*J*_{C-F} = 6Hz, C₂), 114.7 (C₅), 55.7 (C₇), 40.1 (C₉), 17.6 (C₁₀), 14.2 (C₁₁). MS (EI) 222 (M⁺), 179 (M⁺-CH₃CH₂CH₂), 152, 136, 108. IR (KBr) 3019, 2959, 2936, 2873, 2848, 1702, 1637, 1602, 1508, 1464, 1381, 1332, 1250, 1176, 1037, 840, 543. Anal. Calcd for C₁₃H₁₅FO₂: C, 70.25; H, 6.80. Found: C, 69.85; H, 6.71.

Compound 4c₂(Z)

(Z) 4-Fluoro-1-phenyloct-3-en-5-one. Yellow oil, R_f 0.5 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.3-7.1 (m, 5H, $\text{H}_{6,7,8}$), 6.0 (dt, $^3J_{\text{H-F}} = 34.3\text{Hz}$, $^3J_{\text{H-H}} = 7.7\text{Hz}$, 1H, H_2), 2.7 (m, 2H, H_4), 2.5 (m, 4H, $\text{H}_{3,10}$), 1.5 (h, $^3J_{\text{H-H}} = 7.4\text{Hz}$, 2H, H_{11}), 0.9 (t, $^3J_{\text{H-H}} = 7.2\text{Hz}$, 3H, H_{12}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -129.2 (dt, $^3J_{\text{H-F}} = 34.4\text{Hz}$, $^4J_{\text{H-F}} = 2.2\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 194.2 (d, $^2J_{\text{C-F}} = 31\text{Hz}$, C_9), 155.4 (d, $^1J_{\text{C-F}} = 262\text{Hz}$, C_1), 140.6 (C_5), 128.6 ($\text{C}_{6/7}$), 128.4 ($\text{C}_{6/7}$), 126.4 (C_8), 117.8 (d, $^2J_{\text{C-F}} = 12\text{Hz}$, C_2), 39.8 (C_{10}), 34.6 (C_4), 26.0 (C_3), 17.2 (C_{11}), 13.8 (C_{12}). MS (EI) 220 (M^+), 157, 128 ($\text{M}^+ - \text{PhCH}_3$), 91 (PhCH_2^+). IR (neat) 3029, 2964, 2934, 2876, 1732, 1711, 1694, 1656, 1455, 1210, 1169, 995, 749, 700. Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{FO}$: C, 76.33; H, 7.78. Found: C, 76.27; H, 7.77.

Procedure for the obtention of (Z) fluorinated dihydrodioxine 7(Z) from mixture of bromofluoroolefins 2. To 1,4-diox-2-ene (0.85 eq.) in anhydrous THF (5 mL/mmol of enol ether) at -40°C under argon was added dropwise *t*-BuLi in hexanes (1.6M, 0.85 eq.). The mixture was stirred for 30 min at -40°C and then a solution of dry zincchloride (1.7 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise at -78°C. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.05 eq.), triphenylphosphine (0.1 eq.) and mixture of bromofluoroolefins **2** (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolefin) at 10°C under argon. The mixture was stirred for 1h. After the coupling reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, NH₄Cl aq. (sat.) was added and the mixture was stirred for 5 min. The mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on basic alumina (eluent: cyclohexane), affording (Z) starting bromofluoroolefin **2(Z)** and (Z) fluorinated dihydrodioxine **7(Z)**.

Compound 7₁(Z)

(Z) 5-(1-Fluoro-2-(4-methoxyphenyl)vinyl)-2,3-dihydro-1,4-dioxene. White solid (m. p. = 111-113°C), R_f 0.3 (cyclohexane, *Aluminium oxide neutral*). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.4 (d, J = 8.7Hz, 2H, H_4), 6.8 (d, J = 8.6Hz, 2H, H_5), 6.5 (d, $^4J_{\text{H-F}}$ = 2.6Hz, 1H, H_9), 5.7 (d, $^3J_{\text{H-F}}$ = 41.5Hz, 1H, H_2), 4.1-4.0 (m, 4H, $\text{H}_{10, 11}$), 3.7 (s, 3H, H_7). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -133.8 (dd, $^3J_{\text{H-F}}$ = 41.3Hz, $^4J_{\text{H-F}}$ = 2.3Hz). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 158.8 (C_6), 150.8 (d, $^1J_{\text{C-F}}$ = 249Hz, C_1), 129.8 (d, $^2J_{\text{C-F}}$ = 36Hz, C_8), 130.3 (d, J = 7Hz, C_4), 126.7 (d, $^3J_{\text{C-F}}$ = 7Hz, C_9), 125.3 (C_3), 114.3 (C_5), 102.2 (d, $^2J_{\text{C-F}}$ = 6Hz, C_2), 65.0 ($\text{C}_{10/11}$), 64.7 ($\text{C}_{10/11}$), 55.6 (C_7). MS (EI) 236 (M^+), 179 ($\text{M}^+ - \text{CH}_2 - \text{O} - \text{CH} = \text{CH}_2$), 159, 108 (PhOCH_3^+). IR (KBr) 3115, 2941, 2842, 1645, 1643, 1608, 1511, 1315, 1288, 1254, 1165, 1098, 1031, 919, 860, 816, 799, 535. Anal. Calcd for $\text{C}_{13}\text{H}_{13}\text{FO}_3$: C, 66.09; H, 5.55. Found: C, 66.14; H, 5.62.

Compound 7₂(Z)

(Z) 5-(1-Fluoro-4-phenylbut-1-enyl)-2,3-dihydro-1,4-dioxene. Yellow solid (m. p. = 73-75°C), R_f 0.3 (cyclohexane, *Aluminium oxide neutral*). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.3-7.1 (m, 5H, $\text{H}_{6,7,8}$), 6.4 (d, $^4J_{\text{H-F}} = 2.5\text{Hz}$, 1H, H_{10}), 4.9 (dt, $^3J_{\text{H-F}} = 38.7\text{Hz}$, $^3J_{\text{H-H}} = 7.7\text{Hz}$, 1H, H_2), 4.0 (m, 4H, $\text{H}_{11,12}$), 2.6 (m, 2H, H_4), 2.4 (m, 2H, H_3). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -135.6 (d, $^3J_{\text{H-F}} = 38.7\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 151.4 (d, $^1J_{\text{C-F}} = 240\text{Hz}$, C_1), 141.7 (C_5), 130.6 (d, $^3J_{\text{C-F}} = 20\text{Hz}$, C_{10}), 129.5 (d, $^2J_{\text{C-F}} = 38\text{Hz}$, C_9), 128.8 ($\text{C}_{6/7}$), 126.3 ($\text{C}_{6/7}$), 126.1 (C_8), 102.1 (d, $^2J_{\text{C-F}} = 13\text{Hz}$, C_2), 64.6 ($\text{C}_{11/12}$), 64.4 ($\text{C}_{11/12}$), 35.8 (d, $^4J_{\text{C-F}} = 1\text{Hz}$, C_4), 25.4 (d, $^3J_{\text{C-F}} = 5\text{Hz}$, C_3). MS (EI) 234 (M^+), 214, 157 ($\text{M}^+\text{-Ph}$), 143 ($\text{M}^+\text{-CH}_2\text{Ph}$), 128, 99, 91 (PhCH_2^+), 51. IR (KBr) 3113, 3025, 2930, 2862, 1687, 1649, 1453, 1313, 1281, 1163, 1099, 1080, 1036, 920, 815, 752, 699. Anal. Calcd for $\text{C}_{14}\text{H}_{15}\text{FO}_2$: C, 71.78; H, 6.45. Found: C, 71.83; H, 6.41.

Procedure for the obtention of 3-Fluoro-1-hydroxy-4(4-methoxyphenyl)but-3-en-2-one 4d. Representative example **4d(Z)**: To a solution of 5-(1-Fluoro-2-(4-methoxyphenyl)vinyl)-2,3-dihydro-1,4-dioxene **7(Z)-13** in methanol (50 mL/mmol of diene) and CH₂Cl₂ (25 mL/mmol of diene) at 0°C was added dropwise a solution of mCPBA (2 eq.) in MeOH (12 mL/mmol of diene). The solution was stirred at 0°C for 30 min and then neutralized with NaHCO₃ aq. (5%). Solvents were removed under reduced pressure and the product was extracted with CH₂Cl₂ (x 6). The combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was taken up in EtOH (75 mL/mmol of diene) and NaBH₄ (1.32 eq.) was added in three portions at 0°C. The mixture was stirred at room temperature for 5h and EtOH was removed. The residue was then diluted in CH₂Cl₂ and water was added. After extraction, the organic layer was dried over MgSO₄. After filtration and concentration under reduced pressure, the crude diol was taken up in 1% aq. Acetone (50 mL/mmol of diene) and PTSa was added (0.13 eq.). The solution was heated at 55°C for 2h and then neutralized with NaHCO₃ aq. (sat.). The mixture was extracted with AcOEt (3x) and dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on silica gel (eluent: 33% AcOEt in cyclohexane), affording the desired alkoxyketone **4d(Z)**.

Compound 4d₁(Z)

(Z) 3-Fluoro-1-hydroxy-4(4-methoxyphenyl)but-3-en-2-one. Yellow solid (m. p. = 118-120°C), R_f 0.3 (33% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.6 (d, $J = 9.0\text{Hz}$, 2H, H_4), 6.9 (d, $J = 9.0\text{Hz}$, 2H, H_5), 6.8 (d, $^3J_{\text{H-F}} = 37.7\text{Hz}$, 1H, H_2), 4.6 (d, $^4J_{\text{H-F}} = 4.1\text{Hz}$, 2H, H_9), 3.8 (s, 3H, H_7), 3.1 (b, 1H, OH). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -135.6 (d, $^3J_{\text{H-F}} = 37.9\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 193.8 (d, $^2J_{\text{C-F}} = 36\text{Hz}$, C_8), 161.8 (d, $J = 4\text{Hz}$, C_6), 151.6 (d, $^1J_{\text{C-F}} = 264\text{Hz}$, C_1), 133.4 (d, $J = 9\text{Hz}$, C_4), 123.5 (d, $J = 4\text{Hz}$, C_3), 117.3 (d, $^2J_{\text{C-F}} = 4\text{Hz}$, C_2), 114.9 (C_5), 65.8 (C_9), 55.8 (C_7). MS (EI) 210 (M^+), 194, 179 ($\text{M}^+ - \text{OCH}_3$), 161 ($\text{M}^+ - \text{OCH}_3 \& \text{H}_2\text{O}$). IR (KBr) 3440, 3380, 3105, 2966, 1712, 1644, 1604, 1514, 1300, 1260, 1239, 1170, 1100, 1030, 999, 830. Anal. Calcd for $\text{C}_{11}\text{H}_{11}\text{FO}_3$: C, 62.85; H, 5.27. Found: C, 62.62; H, 5.34.

Procedure for the obtention of (*E*) fluorinated methylketones 4a(*E*) from (*Z*) bromofluoroolefins. To a mixture of potassium *tert*-butoxide (2 eq.), ethylvinylether (2 eq.) in anhydrous THF (5 mL/mmol of *t*-BuOK) at -78°C under argon was added dropwise *n*-BuLi in hexanes (1.6M, 2 eq.). The mixture was stirred for 30 min at -78°C and then a solution of dry zincchloride (4 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.05 eq.), triphenylphosphine (0.1 eq.) and bromofluoroolefin **2**(*Z*) (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolefin) at 70°C under argon. The mixture was stirred for 1h. After the reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, the solution was allowed to cool to room temperature. HCl aq. (1N) was added and the mixture was stirred for 15 min. The mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure at low temperature, the residue was purified by chromatography on silica gel (eluent: cyclohexane/AcOEt), affording (*E*) methylketone **4a**(*E*).

Compound 4a₁(*E*)

(*E*) 3-Fluoro-4-(4-methoxyphenyl)but-3-en-2-one. Yellow solid (m. p. < 44°C), R_f 0.2 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.6 (d, $J = 8.9\text{Hz}$, 2H, H_4), 6.9 (d, $J = 8.9\text{Hz}$, 2H, H_5), 6.7 (d, $^3J_{\text{H-F}} = 26.1\text{Hz}$, 1H, H_2), 3.7 (s, 3H, H_7), 2.3 (d, $^4J_{\text{H-F}} = 5.1\text{Hz}$, 3H, H_9). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -115.2 (d, $^3J_{\text{H-F}} = 26.1\text{Hz}$, $^4J_{\text{H-F}} = 5.1\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 193.4 (d, $^2J_{\text{C-F}} = 39\text{Hz}$, C_8), 161.2 (d, $J = 2\text{Hz}$, C_6), 152.6 (d, $^1J_{\text{C-F}} = 252\text{Hz}$, C_1), 132.6 (d, $J = 3\text{Hz}$, C_4), 123.5 (d, $^3J_{\text{C-F}} = 10\text{Hz}$, C_3), 121.0 (d, $^2J_{\text{C-F}} = 29\text{Hz}$, C_2), 114.1 (C_5), 55.7 (C_7), 28.6 (d, $^3J_{\text{C-F}} = 4\text{Hz}$, C_9). MS (EI) 194 (M^+), 179 ($\text{M}^+ - \text{CH}_3$), 159, 43 (CH_3CO^+). IR (KBr) 3005, 2937, 2839, 1729, 1681, 1601, 1513, 1462, 1423, 1371, 1304, 1257, 1177, 1031, 832, 599, 533. *Anal.* Calcd for $\text{C}_{11}\text{H}_{11}\text{FO}_2$: C, 68.03; H, 5.71. Found: C, 68.09; H, 5.54.

Compound 4a₂(*E*)

(*E*) 3-Fluoro-4-(naphthalen-7-yl)but-3-en-2-one. Yellow solid (m. p. = 62-64°C), R_f 0.3 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.1 (s, 1H, H_{12}), 7.9-7.6 (m, 3H, $\text{H}_{7, 8, 9}$), 7.8-7.6 (m, 1H, H_{10}), 7.6-7.4 (m, 2H, $\text{H}_{4, 5}$), 6.9 (d, $^3J_{\text{H-F}} = 24.6\text{Hz}$, 1H, H_2), 2.4 (d, $^4J_{\text{H-F}} = 4.9\text{Hz}$, 3H, H_{14}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -112.8 (dq, $^3J_{\text{H-F}} = 24.6\text{Hz}$, $^4J_{\text{H-F}} = 4.9\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.9 (d, $^2J_{\text{C-F}} = 39\text{Hz}$, C_{13}), 153.3 (d, $^1J_{\text{C-F}} = 258\text{Hz}$, C_1), 133.6 ($\text{C}_{\text{ar}}^{\text{q}}$), 133.1 ($\text{C}_{\text{ar}}^{\text{q}}$), 130.3 (d, $J = 4\text{Hz}$, $\text{C}_{\text{ar}}^{\text{H}}$), 128.6 ($\text{C}_{\text{ar}}^{\text{H}}$), 128.4 ($\text{C}_{\text{ar}}^{\text{q}}$), 127.9 ($\text{C}_{\text{ar}}^{\text{H}}$), 127.7 ($\text{C}_{\text{ar}}^{\text{H}}$), 127.3 (d, $J = 2\text{Hz}$, $\text{C}_{\text{ar}}^{\text{H}}$), 127.1 ($\text{C}_{\text{ar}}^{\text{H}}$), 126.5 ($\text{C}_{\text{ar}}^{\text{H}}$), 120.4 (d, $^2J_{\text{C-F}} = 28\text{Hz}$, C_2), 28.4 (C_{14}). MS (EI) 214 (M^+), 179, 171 ($\text{M}^+ - \text{CH}_3\text{CO}$), 43 (CH_3CO^+). IR (KBr) 3055, 2973, 2879, 1693, 1614, 1392, 1361, 1190, 1183, 1093, 928, 815, 790, 740, 529, 470. Anal. Calcd for $\text{C}_{14}\text{H}_{11}\text{FO}$: C, 78.49; H, 5.18. Found: C, 78.47; H, 5.22.

Compound **4a₃**(*E*)

(*E*) 3-Fluoro-4-(4-cyanophenyl)but-3-en-2-one. Yellow powder (m. p. = 85-87°C), *R_f* 0.3 (10% AcOEt in cyclohexane). ¹H-NMR (300 MHz, CDCl₃) δ 7.7 (dd, *J* = 8.7 Hz, 4H, H_{4,5}), 6.6 (d, ³*J*_{H-F} = 23.3 Hz, 1H, H₂), 2.3 (d, ⁴*J*_{H-F} = 5.1 Hz, 3H, H₉). ¹⁹F-NMR (282.5 MHz, CDCl₃) δ -109.6 (dq, ³*J*_{H-F} = 23.3 Hz, ⁴*J*_{H-F} = 5.1 Hz). ¹³C-NMR (75.5 MHz, CDCl₃) δ 193.2 (d, ²*J*_{C-F} = 40 Hz, C₈), 154.3 (d, ¹*J*_{C-F} = 265 Hz, C₁), 136.0 (d, ³*J*_{C-F} = 11 Hz, C₃), 132.2 (C₅), 130.9 (d, *J* = 3 Hz, C₄), 118.9 (C₇), 118.0 (d, ²*J*_{C-F} = 29 Hz, C₂), 112.8 (C₆), 28.4 (C₉). MS (EI) 189 (M⁺), 174 (M⁺-CH₃), 146 (M⁺-CH₃CO), 43 (CH₃CO⁺). IR (KBr) 3110, 3012, 2960, 2232, 1698, 1616, 1505, 1422, 1368, 1191, 1087, 912, 840, 700, 556, 516. Anal. Calcd for C₁₁H₈FNO: C, 69.84; H, 4.26; N, 7.40. Found: C, 69.85; H, 4.21; N, 7.33.

Compound 4a₄(*E*)

(*E*) 3-Fluoro-4-(4-ethylbenzoate)but-3-en-2-one. Yellow powder (m. p. = 97–99°C), R_f 0.2 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.1 (d, J = 8.2 Hz, 2H, H_5), 7.7 (d, J = 8.3 Hz, 2H, H_4), 6.7 (d, $^3J_{\text{H-F}}$ = 23.7 Hz, 1H, H_2), 3.9 (s, 3H, H_8), 2.3 (d, $^4J_{\text{H-F}}$ = 4.9 Hz, 3H, H_{10}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -111.0 (dq, $^3J_{\text{H-F}}$ = 23.7 Hz, $^4J_{\text{H-F}}$ = 5.4 Hz). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.9 (d, $^2J_{\text{C-F}}$ = 39 Hz, C_9), 166.7 (C_7), 153.7 (d, $^1J_{\text{C-F}}$ = 262 Hz, C_1), 135.6 (d, J = 11 Hz, C_6), 130.6 (d, $^3J_{\text{C-F}}$ = 5 Hz, C_3), 130.0 (d, J = 3 Hz, C_5), 129.5 (C_4), 118.8 (d, $^2J_{\text{C-F}}$ = 28 Hz, C_2), 52.4 (C_8), 28.3 (C_{10}). MS (EI) 222 (M^+), 207 ($\text{M}^+ - \text{CH}_3$), 191 ($\text{M}^+ - \text{CH}_3\text{O}$), 163 ($\text{M}^+ - \text{CO}_2\text{CH}_3$), 43 (CH_3CO^+). IR (KBr) 3066, 2960, 1721, 1637, 1433, 1364, 1320, 1280, 1200, 1109, 1057, 1019, 968, 905, 834, 770, 698, 572. Anal. Calcd for $\text{C}_{12}\text{H}_{11}\text{FO}_3$: C, 64.86; H, 4.99. Found: C, 64.84; H, 4.96.

Compound 4a₅(*E*)

(*E*) 3-Fluoro-4-(4-nitrophenyl)but-3-en-2-one. Yellow solid (m. p. = 89-91°C), R_f 0.2 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.2 (d, $J = 8.8\text{Hz}$, 2H, H_5), 7.7 (d, $J = 8.6\text{Hz}$, 2H, H_4), 6.7 (d, $^3J_{\text{H-F}} = 22.9\text{Hz}$, 1H, H_2), 2.4 (d, $^4J_{\text{H-F}} = 5.2\text{Hz}$, 3H, H_8). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -119.0 (dq, $^3J_{\text{H-F}} = 22.6\text{Hz}$, $^4J_{\text{H-F}} = 5.4\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 192.9 (d, $^2J_{\text{C-F}} = 40\text{Hz}$, C_7), 154.1 (d, $^1J_{\text{C-F}} = 266\text{Hz}$, C_1), 147.8 (C_6), 137.6 (d, $J = 11\text{Hz}$, C_3), 130.9 (d, $J = 3\text{Hz}$, C_5), 123.4 (C_4), 117.2 (d, $^2J_{\text{C-F}} = 29\text{Hz}$, C_2), 28.1 (C_8). MS (EI) 209 (M^+), 192, 162 ($\text{M}^+ - \text{HNO}_2$), 43 (CH_3CO^+). IR (KBr) 3011, 1706, 1595, 1522, 1342, 1199, 1086, 904, 846, 751, 689, 532, 512. Anal. Calcd for $\text{C}_{10}\text{H}_8\text{FNO}_3$: C, 57.42; H, 3.85; N, 6.70. Found: C, 57.43; H, 3.92; N, 6.63

Compound **4a₇**(*E*)

(*E*) 3-Fluoro-6-phenylhex-3-en-2-one. Yellow oil, R_f 0.2 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.3-7.0 (m, 5H, $\text{H}_6, 7, 8$), 5.7 (dt, $^3J_{\text{H-F}} = 22.0\text{Hz}$, $J = 8.0\text{Hz}$, 1H, H_2), 2.9-2.7 (m, 2H, H_4), 2.7-2.5 (m, 2H, H_3), 2.2 (d, $^4J = 4.9\text{Hz}$, 3H, H_{10}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -118.8 (dq, $^3J_{\text{H-F}} = 22.0\text{Hz}$, $^4J = 4.9\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 193.3 (d, $^2J_{\text{C-F}} = 40\text{Hz}$, C_9), 152.4 (d, $^1J_{\text{C-F}} = 256\text{Hz}$, C_1), 139.6 (C_5), 127.3 ($\text{C}_{6/7}$), 127.2 ($\text{C}_{6/7}$), 125.0 (C_8), 119.2 (d, $^2J_{\text{C-F}} = 18\text{Hz}$, C_2), 34.0 (d, $^4J_{\text{C-F}} = 2\text{Hz}$, C_4), 25.8 (C_{10}), 25.7 (C_3). MS (EI) 192 (M^+), 172, 91 (PhCH_2^+), 43 (CH_3CO^+). IR (neat) 3063, 3027, 2927, 2860, 1708, 1640, 1365, 1105, 749, 700. Anal. Calcd for $\text{C}_{12}\text{H}_{13}\text{FO}$: C, 74.98; H, 6.82. Found: C, 75.49; H, 7.17.

Compound 4a₈(*E*)

(*E*) 3-Fluoro-6-(oxytert-butyl)diphenylsilyl)hex-3-en-2-one. Pale yellow oil, R_f 0.4 (2% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.7-7.5 (m, 4H, H_6), 7.4-7.2 (m, 6H, $\text{H}_{7,8}$), 5.8 (dt, $^3J_{\text{H-F}} = 21.8\text{Hz}$, $^3J_{\text{H-H}} = 7.9\text{Hz}$, 1H, H_2), 3.7 (dt, $^5J_{\text{H-F}} = 1.0\text{Hz}$, $^3J_{\text{H-H}} = 6.4\text{Hz}$, 2H, H_4), 2.8-2.6 (m, 2H, H_3), 2.2 (d, $^4J_{\text{H-F}} = 4.9\text{Hz}$, 3H, H_{12}), 0.9 (s, 9H, H_{10}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -118.1 (m). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 194.9 (d, $^2J_{\text{C-F}} = 40\text{Hz}$, C_{11}), 154.3 (d, $^1J_{\text{C-F}} = 255\text{Hz}$, C_1), 136.0 (C_7), 134.0 (C_5), 130.1 (C_8), 128.1 (C_6), 118.9 (d, $^2J_{\text{C-F}} = 20\text{Hz}$, C_2), 63.2 (d, $^4J_{\text{C-F}} = 2\text{Hz}$, C_4), 29.2 (d, $^3J_{\text{C-F}} = 5\text{Hz}$, C_3), 28.1 (d, $^3J_{\text{C-F}} = 3\text{Hz}$, C_{12}), 27.3 (C_{10}), 19.6 (C_9). MS (EI) 313 ($\text{M}^+ - \text{tBu}$), 201, 199, 181, 141, 43 (CH_3CO^+). IR (neat) 3394, 3072, 2959, 2858, 1709, 1643, 1472, 1427, 1365, 1194, 1112, 938, 823, 704, 687, 614, 505. Anal. Calcd for $\text{C}_{22}\text{H}_{27}\text{FO}_2\text{Si}$: C, 71.31; H, 7.34. Found: C, 71.35; H, 7.39.

Procedure for the obtention of (*E*) fluorinated benzylketones **4b(*E*) from bromofluoroolefins **2(*Z*)**.** To a mixture of potassium *tert*-butoxide (4 eq.), freshly synthesized 1-(2-Ethoxyvinyl)benzene (4 eq.) in anhydrous THF (5 mL/mmol of *t*-BuOK) at -78°C under argon was added dropwise *n*-BuLi in hexanes (1.6M, 2 eq.). The mixture was stirred for 30 min at -78°C and then a solution of dry zincchloride (8 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.05 eq.), triphenylphosphine (0.1 eq.) and (*Z*) bromofluoroolefin **2(*Z*)** (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolefin) at 70°C under argon. The mixture was stirred for 1h. After the coupling reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, the solution was allowed to cool to room temperature. HCl aq. (1N) was added and the mixture was stirred for 15 min. The mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on silica gel (eluent: 2% AcOEt in cyclohexane), affording (*E*) benzylketone **4b(*E*)**.

Compound 4b₂(*E*)

(*E*) 3-Fluoro-1,6-diphenylhex-3-en-2-one. Yellow oil, R_f 0.5 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.3-7.1 (m, 10H, $\text{H}_{6,7,8,12,13,14}$), 5.7 (dt, $^3J_{\text{H-F}} = 22.2\text{Hz}$, $^3J_{\text{H-H}} = 7.9\text{Hz}$, 1H, H_2), 3.8 (d, $^4J_{\text{H-F}} = 3.8\text{Hz}$, 2H, H_{10}), 2.8-2.7 (m, 2H, H_4), 2.6-2.5 (m, 2H, H_3). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -120.3 (dt, $^3J_{\text{H-F}} = 23.0\text{Hz}$, $^4J_{\text{H-F}} = 2.3\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 194.6 (d, $^2J_{\text{C-F}} = 39\text{Hz}$, C_9), 154.0 (d, $^1J_{\text{C-F}} = 256\text{Hz}$, C_1), 141.4 (C_5), 133.7 (C_{11}), 130.4 ($\text{C}_{12/13}$), 129.3 ($\text{C}_{12/13}$), 129.2 ($\text{C}_{6/7}$), 129.1 ($\text{C}_{6/7}$), 127.8 (C_{14}), 126.8 (C_8), 122.4 (d, $^2J_{\text{C-F}} = 18\text{Hz}$, C_2), 47.2 (d, $^3J_{\text{C-F}} = 2\text{Hz}$, C_{10}), 35.8 (d, $^4J_{\text{C-F}} = 2\text{Hz}$, C_4), 27.6 (d, $^3J_{\text{C-F}} = 5\text{Hz}$, C_3). MS (EI) 268 (M^+), 177 ($\text{M}^+ - \text{PhCH}_2$), 157, 91 (PhCH_2^+). IR (neat) 3085, 3062, 3028, 2927, 2860, 1704, 1640, 1496, 1454, 1358, 1135, 1080, 749, 729, 698. *Anal. Calcd for* $\text{C}_{18}\text{H}_{17}\text{FO}$: C, 81.57; H, 6.39. *Found*: C, 81.86; H, 6.51.

Procedure for the obtention of (*E*) fluorinated propylketones **4c(*E*) from bromofluoroolefins **2(*Z*)**.** To a mixture of potassium *tert*-butoxide (4 eq.), butenylethylether (4 eq.) in anhydrous THF (5 mL/mmol of *t*-BuOK) at -78°C under argon was added dropwise *n*-BuLi in hexanes (1.6M, 4 eq.). The mixture was stirred for 30 min at -78°C and then a solution of dry zincchloride (8 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.05 eq.), triphenylphosphine (0.1 eq.) and (*Z*) bromofluoroolefin **2(*Z*)** (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolefin) at 55°C under argon. The mixture was stirred for 1h. After the reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, the solution was allowed to cool to room temperature. HCl aq. (1N) was added and the mixture was stirred for 15 min. The mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on silica gel (eluent: cyclohexane/AcOEt), affording (*E*) propylketone **4c(*E*)**.

Compound 4c₁(*E*)

(*E*) 2-Fluoro-1-(4-methoxyphenyl)hex-1-en-3-one. Yellow oil, R_f 0.4 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.7 (d, $J = 9.0\text{Hz}$, 2H, H_4), 6.9 (d, $J = 8.7\text{Hz}$, 2H, H_5), 6.6 (d, $^3J_{\text{H-F}} = 26.6\text{Hz}$, 1H, H_2), 3.8 (s, 3H, H_7), 2.7 (dt, $^4J_{\text{H-F}} = 3.6\text{Hz}$, $^3J_{\text{H-H}} = 7.2\text{Hz}$, 2H, H_9), 1.7 (h, $^3J_{\text{H-H}} = 7.4\text{Hz}$, 2H, H_{10}), 1.0 (t, $^3J_{\text{H-H}} = 7.4\text{Hz}$, 3H, H_{11}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -117.7 (dt, $^3J_{\text{H-F}} = 25.8\text{Hz}$, $^4J_{\text{H-F}} = 3.8\text{Hz}$,). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 195.5 (d, $^2J_{\text{C-F}} = 38\text{Hz}$, C_8), 160.6 (C_6), 152.9 (d, $^1J_{\text{C-F}} = 254\text{Hz}$, C_1), 132.3 (C_4), 123.3 (d, $^3J_{\text{C-F}} = 10\text{Hz}$, C_3), 120.4 (d, $^2J_{\text{C-F}} = 29\text{Hz}$, C_2), 113.7 (C_5), 55.4 (C_7), 42.1 (C_9), 16.8 (C_{10}), 13.8 (C_{11}). MS (EI) 222 (M^+), 193 ($\text{M}^+ - \text{CH}_3\text{CH}_2$), 179 ($\text{M}^+ - \text{CH}_3\text{CH}_2\text{CH}_2$), 136, 108. IR (neat) 3021, 2964, 2936, 2876, 2839, 1701, 1602, 1512, 1464, 1380, 1305, 1258, 1179, 1154, 1033, 875, 831, 598. Anal. Calcd for $\text{C}_{13}\text{H}_{15}\text{FO}_2$: C, 70.25; H, 6.80. Found: C, 70.27; H, 6.89.

Compound **4c₂**(*E*)

(*E*) 4-Fluoro-1-phenyloct-3-en-5-one. Yellow oil, R_f 0.6 (5% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.3-7.1 (m, 5H, $\text{H}_{6,7,8}$), 5.6 (dt, $^3J_{\text{H-F}} = 22.5\text{Hz}$, $^3J_{\text{H-H}} = 7.7\text{Hz}$, 1H, H_2), 2.7 (m, 2H, H_4), 2.5 (m, 2H, H_3), 2.4 (td, $^3J_{\text{H-H}} = 7.2\text{Hz}$, $^4J_{\text{H-F}} = 3.3\text{Hz}$, 2H, H_{10}), 1.6 (m, 2H, H_{11}), 0.9 (t, $^3J_{\text{H-H}} = 7.4\text{Hz}$, 3H, H_{12}). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -121.2 (d, $^3J_{\text{H-F}} = 22.6\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 197.1 (d, $^2J_{\text{C-F}} = 38\text{Hz}$, C_9), 153.8 (d, $^1J_{\text{C-F}} = 256\text{Hz}$, C_1), 140.9 (C_5), 128.6 ($\text{C}_{6/7}$), 128.5 ($\text{C}_{6/7}$), 126.2 (C_8), 120.2 (d, $^2J_{\text{C-F}} = 18\text{Hz}$, C_2), 41.8 (C_{10}), 35.4 (C_4), 26.9 (d, $^3J_{\text{C-F}} = 5\text{Hz}$, C_3), 16.6 (C_{11}), 13.7 (C_{12}). MS (EI) 220 (M^+), 157, 128 ($\text{M}^+ - \text{PhCH}_3$), 104, 91 (PhCH_2^+). IR (neat) 3028, 2964, 2931, 2875, 1706, 1643, 1454, 1378, 1147, 1104, 748, 699. Anal. Calcd for $\text{C}_{14}\text{H}_{17}\text{FO}$: C, 76.33; H, 7.78. Found: C, 76.45; H, 7.96.

Procedure for the obtention of (*E*) fluorinated dihydrodioxine 7(*E*) from bromofluoroolefins 2(*Z*). To 1,4-diox-2-ene (2 eq.) in anhydrous THF (5 mL/mmol of enol ether) at -30°C under argon was added dropwise *t*-BuLi in hexanes (1.6M, 2 eq.). The mixture was stirred for 1h at -20°C and then a solution of dry zincchloride (4 eq.) in THF (4 mL/mmol of ZnCl₂) was added dropwise at -20°C. After 10 min, the cooling bath was removed and the solution was allowed to warm to room temperature for 30 min. The mixture was then added slowly to a solution of palladiumdiacetate (0.05 eq.), triphenylphosphine (0.1 eq.) and (*Z*) bromofluoroolefin 2(*Z*) (1 eq.) in anhydrous THF (10 mL/mmol of bromofluoroolefin) at 70°C under argon. The mixture was stirred for 1h. After the coupling reaction was completed, controlled by monitoring ¹⁹F-NMR signal of the reaction mixture, NH₄Cl aq. (sat.) was added and the mixture was stirred for 5 min. The mixture was then extracted with Et₂O (x 3), washed with brine and the combined organic layers were dried over MgSO₄. After filtration and concentration under reduced pressure, the residue was purified by chromatography on basic alumina (eluent: cyclohexane), affording (*E*) fluorinated dihydrodioxine 7(*E*).

Compound **7₁**(*E*)

(*E*) 5-(1-Fluoro-2-(4-methoxyphenyl)vinyl)-2,3-dihydro-1,4-dioxene. Yellow oil, R_f 0.4 (cyclohexane, *Aluminium oxide neutral*). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.1 (d, $J = 9.1\text{Hz}$, 2H, H_4), 6.7 (d, $J = 8.7\text{Hz}$, 2H, H_5), 6.4 (s, 1H, H_9), 6.1 (d, $^3J_{\text{H-F}} = 22.2\text{Hz}$, 1H, H_2), 4.0-3.9 (m, 4H, $\text{H}_{10,11}$), 3.7 (s, 3H, H_7). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -113.8 (d, $^3J_{\text{H-F}} = 23.0\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 157.5 (C_6), 149.6 (d, $^1J_{\text{C-F}} = 238\text{Hz}$, C_1), 129.7 (C_3), 129.1 (d, $J = 2\text{Hz}$, C_4), 129.0 (d, $^3J_{\text{C-F}} = 6\text{Hz}$, C_9), 124.4 (d, $^2J_{\text{C-F}} = 12\text{Hz}$, C_8), 112.3 (C_5), 108.3 (d, $^2J_{\text{C-F}} = 30\text{Hz}$, C_2), 63.6 ($\text{C}_{10/11}$), 62.8 ($\text{C}_{10/11}$), 54.1 (C_7). MS (EI) 236 (M^+), 216, 179 ($\text{M}^+ - \text{CH}_2 - \text{O} - \text{CH} = \text{CH}_2$), 159. IR (neat) 3107, 2935, 2879, 1656, 1608, 1511, 1300, 1252, 1160, 1100, 1034, 922. *Anal. Calcd for $\text{C}_{13}\text{H}_{13}\text{FO}_3$* : C, 66.09; H, 5.55. *Found*: C, 64.94; H, 5.52.

Compound **7₂(E)**

(E) 5-(1-Fluoro-4-phenylbut-1-enyl)-2,3-dihydro-1,4-dioxene. Colourless oil, R_f 0.3 (cyclohexane, *Aluminium oxide neutral*). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.2–7.1 (m, 5H, $\text{H}_{6,7,8}$), 6.2 (s, 1H, H_{10}), 5.1 (dt, $^3J_{\text{H-F}} = 23.7\text{Hz}$, $^3J_{\text{H-H}} = 7.9\text{Hz}$, 1H, H_2), 4.0 (bs, 4H, $\text{H}_{11,12}$), 2.6–2.4 (m, 4H, $\text{H}_{3,4}$). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -121.5 (d, $^3J_{\text{H-F}} = 23.8\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 151.3 (d, $^1J_{\text{C-F}} = 233\text{Hz}$, C_1), 141.9 (C_5), 131.2 (d, $^2J_{\text{C-F}} = 37\text{Hz}$, C_9), 128.9 ($\text{C}_{6/7}$), 128.8 ($\text{C}_{6/7}$), 128.7 (C_{10}), 126.4 (C_8), 108.3 (d, $^2J_{\text{C-F}} = 23\text{Hz}$, C_2), 64.9 ($\text{C}_{11/12}$), 64.4 ($\text{C}_{11/12}$), 36.9 (d, $^4J_{\text{C-F}} = 2\text{Hz}$, C_4), 27.1 (d, $^3J_{\text{C-F}} = 8\text{Hz}$, C_3). MS (EI) 234 (M^+), 214, 157 ($\text{M}^+\text{-Ph}$), 143 ($\text{M}^+\text{-CH}_2\text{Ph}$), 99, 91 (PhCH_2^+). IR (neat) 3026, 2932, 2879, 1679, 1632, 1455, 1312, 1164, 1110, 924, 700. Anal. Calcd for $\text{C}_{14}\text{H}_{15}\text{FO}_2$: C, 71.78; H, 6.45. Found: C, 71.88; H, 6.41.

Compound 4d₁(*E*)

(*E*) 3-Fluoro-1-hydroxy-4-(4-methoxyphenyl)but-3-en-2-one. Yellow oil, R_f 0.4 (33% AcOEt in cyclohexane). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.8 (d, $J = 8.7\text{Hz}$, 2H, H_4), 6.9 (d, $J = 9.0\text{Hz}$, 2H, H_5), 6.8 (d, $^3J_{\text{H-F}} = 26.9\text{Hz}$, 1H, H_2), 4.5 (m, 2H, H_9), 3.9 (s, 3H, H_7), 3.2 (m, 1H, OH). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -128.6 (d, $^3J_{\text{H-F}} = 26.9\text{Hz}$). $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 193.6 (d, $^2J_{\text{C-F}} = 41\text{Hz}$, C_8), 161.4 (C_6), 150.5 (d, $^1J_{\text{C-F}} = 244\text{Hz}$, C_1), 133.4 (d, $J = 3\text{Hz}$, C_4), 123.9 (d, $^2J_{\text{C-F}} = 28\text{Hz}$, C_2), 122.4 (d, $J = 10\text{Hz}$, C_3), 114.0 (C_5), 67.0 (d, $^3J_{\text{C-F}} = 3\text{Hz}$, C_9), 55.5 (C_7). MS (EI) 210 (M^+), 179 ($\text{M}^+ - \text{OCH}_3$), 161 ($\text{M}^+ - \text{OCH}_3$ & H_2O). IR (neat) 3438, 3392, 3100, 2964, 2840, 1713, 1644, 1602, 1514, 1301, 1259, 1172, 1097, 1030, 831. Anal. Calcd for $\text{C}_{11}\text{H}_{11}\text{FO}_3$: C, 62.85; H, 5.27. Found: C, 62.74; H, 5.36.

Compound 10 (*R,Z*)

(*Z*) (*R*)-3-Fluoro-5-methyl-6-(oxy*tert*-butyl diphenylsilyl)-3-en-2-one.

Colourless oil, *R*_f 0.56 (20% AcOEt in cyclohexane), [α]₂₀^D = -5° (*c* = 0.1, CH₂Cl₂).
¹H-NMR (300 MHz, CDCl₃) δ 7.7-7.6 (m, 4H, H₉), 7.4-7.3 (m, 6H, H₁₀, H₁₁), 5.9-6.0 (dd, ³J_{H-F} = 34.5Hz, ³J_{H-H} = 9.6Hz, 1H, H₂), 3.6-3.5 (m, 2H, H₅), 3.0-2.9 (m, 1H, H₃), 2.3 (d, ⁴J_{H-F} = 2.8Hz, 3H, H₁₃), 1.1 (d, ³J_{H-H} = 6.9Hz, 3H, H₄), 1.0 (s, 9H, H₇).
¹⁹F-NMR (282.5 MHz, CDCl₃) δ -127.9 (s). ¹³C-NMR (75.5 MHz, CDCl₃) δ 191.9 (d, ²J_{C-F} = 31.9Hz, C₁₂), 155.2 (d, ¹J_{C-F} = 260.9Hz, C₁), 135.9 (C₉), 133.8 (C₈), 130.1 (C₁₁), 128.1 (C₁₀), 122.9 (d, ²J_{C-F} = 12.4Hz, C₂), 67.8 (C₅), 33.1 (C₃), 27.0 (d, ³J_{C-F} = 10.9Hz, C₁₃), 25.9 (C₇), 19.7 (C₆), 16.7 (C₄). MS (EI) 327 (M⁺-tBu), 201, 181, 57. IR (neat) 3073, 2968, 2926, 2863, 2357, 1768, 1484, 1368, 1115 cm⁻¹. Anal. Calcd for C₂₃H₂₉FO₂Si : C, 71.83; H, 7.60. Found : C, 71.25; H, 7.32.

Compound 10 (*R,E*)

(*E*) (R)-3-Fluoro-5-methyl-6-(oxy*tert*-butyl diphenylsilyl)hex-3-en-2-one.

Yellow oil, $R_f = 0.69$ (20% AcOEt in cyclohexane), $[\alpha]^{20}_D = -40^\circ$, ($c = 0.057$, CH_2Cl_2). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.7-7.6 (m, 4H, H_9), 7.4-7.3 (m, 6H, H_{10} , H_{11}), 5.7-5.6 (dd, $^3J_{\text{H-F}} = 22.6\text{Hz}$, $^3J_{\text{H-H}} = 9.8\text{Hz}$, 1H, H_2), 3.6-3.5 (m, 3H, $\text{H}_{5,6}$), 2.3 (d, $^5J_{\text{H-F}} = 4.9\text{Hz}$, 1H, H_{13}), 1.1 (d, $^3J_{\text{H-H}} = 2.1\text{Hz}$, 3H, H_4), 1.0 (s, 9H, H_7). $^{19}\text{F-NMR}$ (282.5 MHz, CDCl_3) δ -119.0. $^{13}\text{C-NMR}$ (75.5 MHz, CDCl_3) δ 194.5 (d, $^2J_{\text{C-F}} = 40.0\text{Hz}$, C_1), 153.4 (d, $^1J_{\text{C-F}} = 255.9\text{Hz}$, C_2), 135.8 (C_{10}), 133.7 (C_8), 129.8 (C_{11}), 127.8 (C_9), 124.3 (d, $^2J_{\text{C-F}} = 17.4\text{Hz}$, C_3), 68.1 (C_5), 32.8 (d, $^3J_{\text{C-F}} = 5.1\text{Hz}$, C_4), 27.9 (d, $^3J_{\text{C-F}} = 2.9\text{Hz}$, C_{13}), 26.9 (C_7), 19.4 (C_6), 17.0 (d, $^4J_{\text{C-F}} = 2.2\text{Hz}$, C_{12}). MS (EI) 327 ($\text{M}^+ - \text{tBu}$), 201, 199, 181, 141, 57. IR (neat) 2959, 2856, 1708, 1427, 1363, 1195, 1112, 823, 741 cm^{-1} . Anal. Calcd for $\text{C}_{23}\text{H}_{29}\text{FO}_2\text{Si}$: C, 71.83; H, 7.60. Found: C, 71.15; H, 7.49.

Determination of the diastereoisomeric purities of imines derived from **10 (R, Z)** and **10 (R, E)**; ^{19}F NMR spectra