Formal Total Syntheses of Aspidosperma Alkaloids via a Novel and General Synthetic Pathway Based on an Intramolecular Heck Cyclization

J. Pereira, M. Barlier and C. Guillou *

Institut de Chimie des Substances Naturelles, CNRS Avenue de la Terrasse 91198 Gif-sur-Yvette, France.

guillou@icsn.cnrs-if.fr

Supporting information

Table of Contents

General Procedure for the Synthesis of Acetamide 10a-c	S4
Preparation of compound 10a	S4
Preparation of compound 10b	S4
Preparation of compound 10c	S5
Preparation of compound 10d	S6
Preparation of compound 10e	S7
 General Procedure for the Synthesis of Quinolinone 13a-e	 S8
Preparation of compound 13a	S8
Preparation of compound 13b	S9
Preparation of compound 13c	S10
Preparation of compound 13d	S11
Preparation of compound 13e	S12
 General Procedure for the Synthesis of carbamate derivatives 14a-c	 S13
Preparation of compound 14a	S13
Preparation of compound 14b	S14
Preparation of compound 14c	S15
 General Procedure for the Synthesis of quinolinones 15a-c	 S16
Preparation of compound 15a	S16
Preparation of compound 15b	S17
Preparation of compound 15c	S17
 General Procedure for the Synthesis of carbamates derivatives 9a-c	 S18
Preparation of compound 9a	S18
Preparation of compound 9b	S19
Preparation of compound 9c	S20

General Procedure for the Synthesis of carbazoles 17a-c	S20
Preparation of compound 17a	S21
Preparation of compound 17b	S21
Preparation of compound 17c	S22
General Procedure for the Synthesis of carbazoles 18a-c and 19a-c	S23
Preparation of compound 18a	S23
Preparation of compound 18b	S24
Preparation of compound 18c	S25
Preparation of compound 19a	S26
Preparation of compound 19b	S27
Preparation of compound 19c	S28
General Procedure for the Synthesis of carbazoles 20a-c	S29
Preparation of compound 20a	S29
Preparation of compound 20b	S30
Preparation of compound 20c	S31
General Procedure for the Synthesis of carbazoles 21a-c	S32
Preparation of compound 21a	S33
Preparation of compound 21b	S33
Preparation of compound 21c	S34
General Procedure for the Synthesis of carbazoles 22a-c	S35
Preparation of compound 22a	S35
Preparation of compound 22b	S36
Preparation of compound 22c	S36
General Procedure for the Synthesis of carbazoles 23a-c	S37
Preparation of compound 23a	S38
Preparation of compound 23b	S38
Preparation of compound 23c	S39
General Procedure for the Synthesis of carbazoles 8a-c	S40
Preparation of compound 8a	S40
Preparation of compound 8b	S41
Preparation of compound 8c	S41
Spectral copies	S42-S85

General Procedure for the Synthesis of Acetamide **10a-c**

To a mixture of (1,4-dioxa-spiro[4.5]dec-7-en-8-yl)-acetic acid **11**, the corresponding iodo-aniline and 2-chloro-1-methylpyridinium iodide dissolved in dry CH_2Cl_2 , was added NEt_3 . The reaction mixture was heated for 15 hours. The reaction mixture was extracted with CH_2Cl_2 . The combined organic layer was washed with H_2O , brine and dried (MgSO_4) and evaporated. Silica gel flash-column chromatography of the residue afforded **10a-c** as indicated below for each case.

2-(1',4'-dioxo-spiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-phenyl)-acetamide **10a**

(1,4-dioxo-spiro[4.5]dec-7-en-8-yl)-acetic acid **11** (1 g, 5.05 mmol), 2-iodo-phenylamine **12a** (1.55 g, 7.07 mmol), 2-chloro-1-methylpyridinium iodide (1.94 g, 7.58 mmol), dry CH_2Cl_2 (88 mL) and NEt_3 (2.46 mL, 17.70 mmol) were used to produce **10a**. Purification : Heptane/EtOAc (7:3). White powder (1.83 g, 91%).

H.R.M.S. (E.S.I., m/z) calcd for $\text{C}_{16}\text{H}_{18}\text{INNaO}_3^+$ (MNa^+) : 422.0229, found : 422.0243.

I.R. (CHCl_3) ν (cm^{-1}) : 3322 (N-H), 1681 (C=O (amide)), 1511 (C=C), 1427 (C-Car), 1109 (C-O).

M.S. (E.S.I., m/z) : 422 (MNa^+).

^1H RMN (CDCl_3 , 300 MHz) δ (ppm) : 8.29 (1H, dd, $J = 8.2$ Hz, $J = 1.2$ Hz, H_6), 8.02 (1H, sl, NH), 7.76 (1H, dd, $J = 7.9$ Hz, $J = 1.4$ Hz, H_3), 7.34 (1H, dt, $J = 8.5$ Hz, $J = 1.4$ Hz, H_5), 6.83 (1H, dt, $J = 7.8$ Hz, $J = 1.3$ Hz, H_4), 5.77 (1H, s, $\text{H}_{7'}$), 4.00 (4H, s, $\text{H}_{2'}$ and $\text{H}_{3'}$), 3.16 (2H, s, CH_2), 2.42 (2H, s, $\text{H}_{6'}$), 2.34 (2H, m, $\text{H}_{9'}$), 1.88 (2H, t, $J = 6.4$ Hz, $\text{H}_{10'}$).

^{13}C RMN (CDCl_3 , 75 MHz) δ (ppm) : 169.1 (C (O) NH), 138.9 (C_6), 138.3 (C_1), 132.3 ($\text{C}_{8'}$), 129.3 (C_4), 126.4 ($\text{C}_{7'}$), 125.9 (C_5), 121.6 (C_3), 107.5 ($\text{C}_{5'}$), 89.2 (C_2), 64.5 ($\text{C}_{2'}$ and $\text{C}_{3'}$), 46.8 (CH_2), 36.0 ($\text{C}_{6'}$).

2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-*N*-(2-iodo-5-methoxy-phenyl)-acetamide **10b**

(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-acetic acid **11** (5 g, 25.25 mmol), 2-iodo-5-methoxy-phenylamine **12b** (8.8 g, 35.35 mmol), 2-chloro-1-methylpyridinium iodide (9.68 g, 37.89 mmol), dry CH₂Cl₂ (250 mL), and NEt₃ (12.3 mL, 88.38 mmol) were used to produce **10b**. Purification : Heptane/EtOAc (7:3). White powder (8.55 g, 79%).

H.R.M.S. (E.S.I., *m/z*) calcd for C₁₇H₂₀INNaO₄⁺ (MNa⁺) 452.0335, found : 452.0341.

I.R. (CHCl₃) ν (cm⁻¹) : 3270 (N-H), 1657 (C=O (amide)), 1529 (C=C), 1469 (C-Car), 1309 (Car-O), 1109 (C-O).

M.S. (E.S.I., *m/z*) : 452 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 8.08 (1H, d, *J* = 2.9 Hz, H₆), 8.03 (1H, sl, NH), 7.60 (1H, d, *J* = 8.8 Hz, H₃), 6.47 (1H, dd, *J* = 8.8 Hz, *J* = 3.0 Hz, H₄), 5.77 (1H, s, H_{7'}), 4.01 (4H, s, H_{2'} and H_{3'}), 3.81 (3H, s, OCH₃), 3.16 (2H, s, CH₂), 2.43 (2H, s, H_{6'}), 2.34 (2H, m, H_{9'}), 1.89 (2H, t, *J* = 6.5 Hz, H_{10'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 169.2 (C (O) NH), 160.7 (C_{5'}), 139.1 (C₁), 138.8 (C₃), 132.3 (C_{8'}), 126.6 (C_{7'}), 112.9 (C₄), 107.5 (C_{5'}), 106.7 (C₆), 64.6 (C_{2'} and C_{3'}), 55.6 (OCH₃), 46.9 (CH₂), 36.0 (C_{6'}), 31.2 (C_{10'}), 28.1 (C_{9'}).

2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-*N*-(2-iodo-6-methoxy-phenyl)-acetamide **10c**

(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-acetic acid **11** (1.7 g, 8.60 mmol), 2-iodo-6-methoxy-phenylamine **12c** (3 g, 12.00 mmol), 2-chloro-1-methylpyridinium iodide (3.3 g, 12.90 mmol), dry CH₂Cl₂ (150 mL), and NEt₃ (4.2 mL, 30.10 mmol) were used to

produce **10c**. Purification : Heptane/EtOAc (4:6). Pale yellow amorphous solid (2.95 g, 80 %).

Anal. C 47,39%, H 4,59%, N 3,01%, O 15,16% calcd for C₁₇H₂₀INO₄ C 47,57%, H 4,70%, N 3,26%, O 14,91%.

I.R. (CHCl₃) ν (cm⁻¹) : 3382 (N-H), 1687 (C=O (amide)), 1554 (C=C), 1465 (C-Car), 1261 (Car-O), 1092 (C-O).

M. S. (E.S.I., m/z): 430 (MH⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.44 (1H, d, J = 7.7 Hz, H₃), 7.15 (1H, sl, NH), 6.97 (1H, t, J = 8.2 Hz, J = 7.8 Hz, H₄), 6.89 (1H, d, J = 7.9 Hz, H₅), 5.72 (1H, sl, H_{7'}), 3.99 (4H, s, H_{2'} and H_{3'}), 3.80 (3H, s, OCH₃), 3.14 (2H, s, CH₂), 2.45 (2H, sl, H_{9'}), 2.37 (2H, sl, H_{6'}), 1.85 (2H, t, J = 6.4, H_{10'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 169.1 (C (O) NH), 155.4 (C₆), 133.0 (C₁), 130.8 (C₃), 129.6 (C₄), 128.1 (C_{8'}), 125.0 (C_{7'}), 111.7 (C₅), 107.7 (C_{5'}), 99.8 (C₂), 64.5 (C_{2'} and C_{3'}), 56.0 (OCH₃), 45.7 (CH₂), 35.9 (C_{6'}), 31.1 (C_{10'}), 28.0 (C_{9'}).

Synthesis of the acetamide **10d**

(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-acetyl-(2-iodo-6-methoxy-phenyl)-carboxylic acid *tert*-butyl ester **10d**

10d

To a solution of 2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-6-methoxy-phenyl)-acetamide **10c** (3.01 g, 6.99 mmol) dissolved in dry CH₂Cl₂ (100 mL), NEt₃ (0.97 mL, 6.99 mmol), Boc₂O (3.06 g, 14.00 mmol), DMAP (856.3 mg, 6.99 mmol) were successively added. After being stirred for 16 hours at room temperature, the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. Silica gel flash-column chromatography of the residue (elution with Heptane/EtOAc : 4/6) afforded 3.05 g of (1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-acetyl-(2-iodo-6-methoxy-phenyl)-carboxylic acid *tert*-butyl ester **10d** (81 %) as a white oil.

H.R.M.S. (I.C., m/z) calcd for $C_{22}H_{28}INaO_6^+$ (MNa^+) : 552.0854, found : 552.0877.

I.R. ($CHCl_3$) ν (cm^{-1}) : 1736 (C=O (Boc)), 1699 (C=O (amide)), 1575 (C=C), 1470 (C-Car), 1261 (Car-O), 1152 (C-O (Boc)), 1092 (C-O).

M. S. (I.E., m/z) : 529 (M^{+}).

1H RMN ($CDCl_3$, 300 MHz) δ (ppm) : 7.40 (1H, d, J = 7.9 Hz, H_3), 7.01 (1H, t, J = 7.9 Hz, J = 8.2 Hz, H_4), 6.89 (1H, d, J = 8.3 Hz, H_5), 5.51 (1H, sl, $H_{7'}$), 3.97 (4H, s, $H_{2'}$ and $H_{3'}$), 3.77 (3H, s, OCH_3), 3.70 (2H, d, CH_2), 2.30 (4H, sl, $H_{9'}$ and $H_{6'}$), 1.80 (2H, t, J = 6.3 Hz, $H_{10'}$), 1.37 (9H, s, Boc).

^{13}C RMN ($CDCl_3$, 75 MHz) δ (ppm) : 172.8 (CO (Nvoc)), 155.4 (CO (Boc)), 151.4 (C_6), 131.7 (C_1), 131.1 (C_3), 130.6 (C_4), 130.3 ($C_{8'}$), 123.1 ($C_{7'}$), 111.4 (C_5), 108.0 ($C_{5'}$), 100.7 (C_2), 82.8 (C (Boc)), 64.4 ($C_{2'}$ and $C_{3'}$), 56.0 (OCH_3), 45.5 (CH_2), 35.9 (C_6'), 31.2 ($C_{10'}$), 27.9 (CH_3 (Boc)), 27.9 ($C_{9'}$).

Synthesis of Acetamide 10e

2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-
N-(2-iodo-6-methoxy-phenyl)-*N*-methyl-acetamide **10e**

10e

To a solution of 2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-*N*-(2-iodo-6-methoxy-phenyl)-acetamide **10c** (1.88 g, 4.39 mmol) dissolved in dry THF (150 mL), a solution of NaH (263.0 mg, 10.98 mmol) in dry THF (120 mL) was added dropwise at 0°C. After 15 min, the reaction mixture was let go up at room temperature, and $(MeO)_2SO_2$ was added (1.04 mL, 0.20 mmol). After being stirred for 2 hours at room temperature, $NaHCO_3$ (aq.) was added and the reaction mixture was extracted with Et_2O . The combined organic layer was washed with H_2O , brine, dried ($MgSO_4$) and evaporated. Silica gel flash-column chromatography of the residue (elution with Heptane/ $EtOAc$: 4/6) afforded 1.67 g of 2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-*N*-(2-iodo-6-methoxy-phenyl)-*N*-methyl-acetamide **10e** (86 %) as a colourless oil.

I.R. (CHCl₃) ν (cm⁻¹) : 1652 (C=O (amide)), 1602 (C=C), 1470 (C-Car), 1263 (Car-O), 1090 (C-O).

M. S. (E.S.I., m/z): 444 (MH⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.45 (1H, d, J = 8.0 Hz, H₃), 7.04 (1H, t, J = 8.0 Hz, H₄), 6.93 (1H, d, J = 8.0 Hz, H₅), 5.06 (1H, sl, H₇), 3.94 (4H, s, H₂' and H₃'), 3.82 (3H, s, OCH₃), 3.09 (3H, s, NCH₃), 2.75-2.63 (2H, system AB, J = 15.0 Hz, CH₂), 2.21 (2H, sl, H₉), 2.20 (2H, sl, H₆), 1.79 (2H, m, H₁₀).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 171.3 (C (O) NMe), 156.3 (C₆), 134.8 (C₁), 131.8 (C₃), 131.0 (C₄), 130.7 (C₈'), 122.5 (C₇'), 111.7 (C₅), 107.9 (C₅'), 101.7 (C₂), 64.3 (C₂' and C₃'), 55.9 (OCH₃), 42.3 (CH₂), 35.8 (C₆'), 34.5 (NCH₃), 31.0 (C₁₀'), 27.5 (C₉').

General Procedure for the Synthesis of Quinolinone 13a-e

By Heck condition : A mixture of Pd₂(dba)₃ and dppe in dry DMA was stirred under a oil bath at 110°C for 45 min. A solution of **10a-e** and PMP was added and the reaction mixture was heated at 110°C for 3 hours. After being cooled to room temperature, the reaction mixture was washed by aqueous saturated NaHCO₃ and was extracted with EtOAc. The combined organic layer was washed with H₂O, brine and dried (MgSO₄) and evaporated. Silica gel flash-column chromatography of the residue afforded **13a-e** as indicated below for each case.

By Heck “ligand free” condition : To a mixture of **10a-c** and Pd₂(dba)₃ in dry DMA, NEt₃ was added slowly. The reaction mixture was heated at 140°C for two hours. After cooling, the reaction mixture was extracted with EtOAc. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. Silica gel flash-column chromatography of the residue afforded **13a-c** as indicated below for each case.

3-hydro-1H-quinolin-2-one-4,8'-(1',4'-dioxo-spiro[4.5]dec-6'-enyl) **13a**

By Heck conditions : Pd₂(dba)₃ (5 mg, 5.5 μmol), dppe (4.4 mg, 11.0 μmol), dry DMA (5 mL), 2-(1',4'-dioxo-spiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-phenyl)-acetamide **10a** (43.2 mg, 0.11 mmol), and PMP (0.08 mL, 0.44 mmol) were used to produce **13a**. Purification : Heptane/EtOAc (4:6). White powder (15.4 mg, 52%).

By Heck "ligand free" conditions : 2-(1',4'-dioxo-spiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-phenyl)-acetamide **10a** (3.0 g, 7.52 mmol), Pd₂(dba)₃ (68.8 mg, 0.08 mmol), dry DMA (53 mL), and NEt₃ (3.14 mL, 22.60 mmol) were used to produce **13a**. Purification : Heptane/EtOAc (4:6). White powder (1.86 g, 92%).

H.R.M.S. (E.S.I., m/z) calcd for C₁₆H₁₇NNaO₃⁺ (MNa⁺) 294.1106, found : 294.1096.

I.R. (CHCl₃) ν (cm⁻¹) : 3200 (N-H), 1668 (C=O (amide)), 1600 (C=C), 1481 (C-Car), 1094 (C-O).

M. S. (E.S.I., m/z) : 294 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 8.99 (1H, sl, NH), 7.18 (2H, d, J = 7.7 Hz, H_{ar}), 7.00 (1H, dt, J = 1.2 Hz, J = 7.6 Hz, H_{ar}), 6.85 (1H, dd, J = 7.9 Hz, J = 1.5 Hz, H_{ar}), 5.91 (1H, d, J = 10.0 Hz, H_{6'}), 5.72 (1H, d, J = 10.0 Hz, H_{7'}), 4.04-3.90 (4H, m, H_{2'} and H_{3'}), 2.60 (2H, system AB, J = 16.0 Hz, H_{3a} and H_{3b}), 2.08-1.82 (2H, m, H_{9'}), 1.80-1.65 (2H, m, H_{10'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 170.2 (C₂), 136.2 (C_{4a} and C_{8a}), 135.8 (C_{7'}), 130.7 (C_{6'}), 128.3 (C_{ar}), 127.3 (C_{ar}), 123.42 (C_{ar}), 116.2 (C_{ar}), 105.0 (C_{5'}), 64.8 and 64.7 (C_{2'} and C_{3'}), 42.5 (C₃), 39.3 (C_{8'}), 31.6 (C_{9'}), 29.5 (C_{10'}).

7-methoxy-3-hydroxy-1H-quinolin-2-one-4,8'-(1',4'-dioxo-spiro[4.5]dec-6'-enyl) **13b**

By Heck conditions : Pd₂(dba)₃ (5 mg, 5.5 μmol), dppe (4.4 mg, 11.0 μmol), dry DMA (5 mL), 2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-phenyl)-acetamide **10b** (47.2 mg, 0.11 mmol), and PMP (0.08 mL, 0.44 mmol) were used to produce **13b**. Purification : Heptane/EtOAc (4:6). White powder (29.1 mg, 88%).

By Heck “ligand free” conditions : 2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-5-methoxy-phenyl)-acetamide **10b** (4.05 g, 9.44 mmol), Pd₂(dba)₃ (86.4 mg, 0.09 mmol), dry DMA (65 mL), and NEt₃ (3.9 mL, 28.32 mmol) were used to produce **13b**. Purification : Heptane/EtOAc (4:6). White powder (2.46 g, 87%).

H.R.M.S. (E.S.I., m/z) calcd for C₁₇H₁₉NNaO₄⁺ (MNa⁺) 324.1212, found : 324.1218.

I.R. (CHCl₃) ν (cm⁻¹) : 2923 (N-H), 1681 (C=O lactame), 1591 (C=C), 1265 (Car-O), 1120 (C-O).

M. S. (E.S.I., m/z) : 324 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 8.26 (1H, sl, NH), 7.09 (1H, d, J = 8.5 Hz, H₈), 6.54 (1H, dd, J = 8.5 Hz, J = 2.5 Hz, H₆), 6.36 (1H, d, J = 2.5 Hz, H₅), 5.90 (1H, d, J = 10.0 Hz, H_{6'}), 5.70 (1H, d, J = 10.0 Hz, H_{7'}), 4.06-3.90 (4H, m, H_{2'} and H_{3'}), 3.78 (3H, s, OCH₃), 2.58 (2H, system AB, J = 16.0 Hz, H_{3a} and H_{3b}), 1.77–1.97 (2H, m, H_{9'}), 1.77-1.67 (2H, m, H_{10'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 170.4 (C₂), 159.7 (C₇), 137.2 (C_{4a}), 136.1 (C_{7'}), 130.4 (C_{6'}), 128.3 (C₈), 120.3 (C_{8a}), 108.3 (C₆), 105.0 (C_{5'}), 102.3 (C₅), 64.8 and 64.7 (C_{2'} and C_{3'}), 55.5 (OCH₃), 42.8 (C₃), 38.7 (C_{8'}), 31.8 (C_{9'}), 29.5 (C_{10'}).

8-methoxy-3-hydroxy-1H-quinolin-2-one-4,8'-(1',4'-dioxaspiro[4.5]dec-6'-enyl) **13c**

By Heck conditions : Pd₂(dba)₃ (639.7 mg, 0.70 mmol), dppe (557.2 mg, 1.40 mmol), dry DMA (400 mL), 2-(1',4'-dioxo-spiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-phenyl)-acetamide **10c** (6.0 g, 13.99 mmol), and PMP (10.1 mL, 55.94 mmol) were used to produce **13c**. Purification : Heptane/EtOAc (4:6). Pale yellow amorphous solid (3.9 g, 94%).

By Heck “ligand free” conditions : 2-(1',4'-dioxo-spiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-6-methoxy-phenyl)-acetamide **10c** (3.68 g, 8.58 mmol), Pd₂(dba)₃ (78.5 mg, 0.09 mmol), dry DMA (60 mL), and NEt₃ (3.58 mL, 25.73 mmol) were used to produce **13c**. Purification : Heptane/EtOAc (4:6). Pale yellow amorphous solid (2.21 g, 86%).

H.R.M.S. (I.C., m/z) calcd for C₁₇H₁₉NNaO₄⁺ (MNa⁺) 324.1206, found : 324.1211.

I.R. (CHCl₃) ν (cm⁻¹) : 3404 (N-H), 1679 (C=O (lactame)), 1593 (C=C), 1491 (C-Car), 1260 (Car-O), 1092 (C-O).

M. S. (I.E., m/z) : 301 (M⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 7.88 (1H, sl, NH), 6.97 (1H, t, J = 8.1 Hz, H₆), 6.84 (1H, dd, J = 8.1 Hz, J = 1.1 Hz, H₇), 6.82 (1H, dd, J = 8.1 Hz, J = 1.1 Hz, H₅), 5.90 (1H, d, J = 10.0 Hz, H_{6'}), 5.74 (1H, d, J = 10.0 Hz, H_{7'}), 4.04 (4H, m, H_{2'} and H_{3'}), 3.87 (3H, s, OCH₃), 2.67-2.54 (2H, system AB, J = 15.0 Hz, H_{3a} and H_{3b}), 1.92 (2H, m, H_{9'}), 1.76 (2H, m, H_{10'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 168.4 (C₂), 146.1 (C₈), 135.7 (C_{7'}), 130.4 (C_{6'}), 128.6 (C_{4a}), 125.3 (C_{8a}), 122.7 (C₆), 119.1 (C₅), 109.6 (C₇), 104.9 (C_{5'}), 64.6 (C_{2'} and C_{3'}), 55.9 (OCH₃), 42.3 (C₃), 39.4 (C₄), 31.4 (C_{9'}), 29.5 (C_{10'}).

8-methoxy-3-hydroxy-1H-quinolin-2-one-4,8'-
(1',4'-dioxo-spiro[4.5]dec-6'-enyl)-carboxylic acid *tert*-butyl ester **13d**

13d

By Heck conditions : Pd₂(dba)₃ (45.8 mg, 0.05 mmol), dppe (39.9 mg, 0.10 mmol), dry DMA (40 mL), (1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-acetyl-(2-iodo-6-methoxyphenyl)-carboxylic acid *tert*-butyl ester **10d** (530.2 mg, 1.00 mmol), and PMP (723,4 μL, 4.01 mmol) were used to produce **13d and 13c**. Purification : Heptane/EtOAc (4:6). Pale yellow amorphous solid (219.8 mg and 125.8 mg, 96%).

H.R.M.S. (I.C., m/z) calcd for C₂₂H₂₇NNaO₆⁺ (MNa⁺) 424.1731, found : 424.1710.

I.R. (CHCl₃) ν (cm⁻¹) : 1751 (C=O (Boc)), 1705 (C=O (lactame)), 1589 (C=C), 1479 (C-Car), 1261 (Car-O), 1147 (C-O (Boc)), 1092 (C-O).

M. S. (I.E., m/z) : 401 (M⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.10 (1H, t, J = 8.3 Hz, H₆), 6.86 (1H, dd, J = 8.3 Hz, J = 1.2 Hz, H₇), 6.85 (1H, dd, J = 8.3 Hz, J = 1.3 Hz, H₅), 5.90 (1H, d, J = 10.0 Hz, H_{6'}), 5.67 (1H, d, J = 10.0 Hz, H_{7'}), 4.02 (4H, m, H_{2'} and H_{3'}), 3.83 (3H, s, OCH₃), 2.67-2.48 (2H, system AB, J = 15.0 Hz, H_{3a} and H_{3b}), 1.93 (2H, m, H_{9'}), 1.79 (2H, m, H_{10'}), 1.60 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 167.9 (C₂), 151.9 (CO (Boc)), 149.7 (C₈), 135.3 (C_{7'}), 133.9 (C_{4a}), 130.7 (C_{6'}), 126.4 (C_{8a}), 125.5 (C₆), 119.2 (C₅), 111.4 (C₇), 105.0 (C_{5'}), 84.0 (C (Boc)), 64.7 (C_{2'} and C_{3'}), 56.0 (OCH₃), 44.5 (C₃), 39.5 (C₄), 30.8 (C_{9'}), 29.6 (C_{10'}), 27.8 (CH₃(Boc)).

8-methoxy-1-methyl-3-hydroxy-1H-quinolin-2-one-
4,8'-(1',4'-dioxaspiro[4.5]dec-6'-enyl) **13e**

13e

By Heck conditions : Pd₂(dba)₃ (36.8 mg, 0.04 mmol), dppe (32.1 mg, 0.08 mmol), dry DMA (40 mL), 2-(1',4'-dioxaspiro[4.5]dec-7'-en-8'-yl)-N-(2-iodo-6-methoxy-phenyl)-N-methyl-acetamide **10e** (357.0 mg, 0.81 mmol), and PMP (582.0 μL, 3.22 mmol) were used to produce **13e**. Purification : Heptane/EtOAc (4:6). Pale yellow amorphous solid (203.5 mg, 80%).

H.R.M.S. (I.C., m/z) calcd for C₁₈H₂₂NO₄⁺ (MH⁺) 316.1543, found : 316.1563.

I.R. (CHCl₃) ν (cm⁻¹) : 1659 (C=O (lactame)), 1589 (C=C), 1479 (C-Car), 1261 (Car-O), 1092 (C-O).

M. S. (E.S.I., m/z) : 338 (MNa⁺), 316 (MH⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.05 (1H, t, J=8.0 Hz, H₆), 6.91 (1H, d, J = 8.0 Hz, H₇), 6.84 (1H, d, J = 8.3 Hz, H₅), 5.88 (1H, d, J = 10.0 Hz, H_{6'}), 5.65 (1H, d, J = 10.0 Hz, H_{7'}), 4.03 (4H, m, H_{2'} and H_{3'}), 3.86 (3H, s, OCH₃), 3.40 (3H, s, NCH₃), 2.61-2.46 (2H, system AB, J = 15.0 Hz, H_{3a} and H_{3b}), 1.91-1.75 (4H, m, H_{9'} and H_{10'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 169.9 (C₂), 150.2 (C₈), 136.0 (C_{7'}), 134.7 (C_{4a}), 130.3 (C_{6'}), 129.9 (C_{8a}), 124.6 (C₆), 119.2 (C₅), 112.1 (C₇), 105.1 (C_{5'}), 64.6 (C_{2'} and C_{3'}), 55.9 (OCH₃), 43.7 (C₃), 39.0 (C₄), 34.5 (NCH₃), 30.2 (C_{9'}), 29.7 (C_{10'}).

General Procedure for the Synthesis of Carbamate Derivatives **14a-c**

To **13a-c** in THF were added at 0°C HCl 1N. After being stirred for a night at room temperature, the reaction mixture was extracted with EtOAc. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. The residue was dissolved in dry CH₂Cl₂. Then Boc₂O, DMAP and NEt₃ were added. After being stirred for a night at room temperature, the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. Silica gel flash-column chromatography of the residue afforded **14a-c** as indicated below for each case.

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2'-en-4'-one)-
carboxylic acid *tert*-butyl ester **14a**

14a

3-hydro-1H-quinolin-2-one-4,8'-(1',4'-dioxo-spiro[4.5]dec-6'-enyl) **13a** (4.67 g, 17.23 mmol), THF (400 mL), HCl 1N (200 mL), dry CH₂Cl₂ (200 mL), Boc₂O (7.52 g, 34.46 mmol), DMAP (526.3 mg, 4.31 mmol), and NEt₃ (3.6 mL, 25.85 mmol) were used to produce **14a**. Purification : Heptane/EtOAc (5:5). White amorphous solid (5.55 g, 98%).

H.R.M.S. (E.S.I., m/z) calcd for C₁₉H₂₁NNaO₄⁺ (MNa⁺) 350.1358, found : 350.1368.

I.R. (CHCl₃) ν (cm⁻¹) : 1743 (C=O (Boc)), 1674 (C=O (lactame)), 1600 (C=C), 1492 (C-Car), 1144 (C-O).

M. S. (E.S.I., m/z) : 350 (MNa⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 7.35-7.26 (1H, m, H₆), 7.21-7.13 (2H, m, H₇ and H₈), 7.00 (1H, d, J = 8.2 Hz, H₅), 6.68 (1H, d, J = 10.1 Hz, H_{3'}), 6.28 (1H, d, J = 10.1 Hz, H_{2'}), 2.77 (2H, system AB, J = 15.1 Hz, H_{3a} and H_{3b}), 2.51-2.34 (2H, m, H_{5'}), 2.17 (2H, t, J = 6.4 Hz, H_{6'}), 1.61 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 198.3 (C_{4'}), 166.1 (C₂), 151.3 (CO (Boc)), 151.3 (C_{3'}), 136.1 (C_{8a}), 131.5 (C_{2'}), 128.7 (C₆), 127.9 (C_{4a}), 126.6 and 124.6 (C₇ and C₈), 117.8 (C₅), 85.7 (C (Boc)), 42.6 (C₃), 39.6 (C_{1'}), 33.5 (C_{5'}), 32.4 (C_{6'}), 27.7 (CH₃(Boc)).

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2'-en-4'-one)-
carboxylic acid *tert*-butyl ester **14b**

14b

7-methoxy-3-hydro-1H-quinolin-2-one-4,8'-(1',4'-dioxo-spiro[4.5]dec-6'-enyl) **13b** (4.93 g, 16.38 mmol), THF (400 mL), HCl 1N (200 mL), dry CH₂Cl₂ (200 mL), Boc₂O (7.15 g, 32.76 mmol), DMAP (500.2 mg, 4.09 mmol), and NEt₃ (3.4 mL, 24.57 mmol) were used to produce **14b**. Purification : Heptane/EtOAc (5:5). White amorphous solid (5.67 g, 97%).

H.R.M.S. (E.S.I., m/z) calcd for C₂₀H₂₃NNaO₅⁺ (MNa⁺) 380.1474, found : 380.1448.

I.R. (CHCl₃) ν (cm⁻¹) : 1731(C=O (Boc)), 1682 (C=O (lactame)), 1610 (C=C), 1505 (C-Car), 1300 (Car-O), 1142 (C-O).

M. S. (E.S.I., m/z) : 380 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.06 (1H, d, J = 8.5 Hz, H₅), 6.68-6.63 (2H, m, H_{3'} and H₆), 6.26 (1H, d, J = 10.1 Hz, H_{2'}), 3.79 (3H, s, OCH₃), 2.73 (2H, system AB, J = 15.1 Hz, H_{3a} and H_{3b}), 2.48-2.30 (2H, m, H_{5'}), 2.14 (2H, t, J = 6.6 Hz, H_{6'}), 1.55 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 198.5 (C_{4'}), 166.3 (C₂), 159.8 (C₇), 151.6 (C_{3'}), 151.3 (CO (Boc)), 137.0 (C_{8a}), 131.3 (C_{2'}), 127.6 (C₅), 119.6 (C_{4a}), 109.1 (C₆), 104.3 (C₈), 87.8 (C (Boc)), 55.5 (OCH₃), 42.9 (C₃), 39.1 (C_{1'}), 33.5 (C_{5'}), 32.7 (C_{6'}), 27.6 (CH₃(Boc)).

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2'-en-4'-one)-
carboxylic acid *tert*-butyl ester **14c**

14c

8-methoxy-3-hydro-1H-quinolin-2-one-4,8'-(1',4'-dioxo-spiro[4.5]dec-6'-enyl) **13c** (2.89 g, 9.61 mmol), THF (200 mL), HCl 1N (100 mL), dry CH₂Cl₂ (120 mL), Boc₂O (4.19 g, 19.22 mmol), DMAP (293.5 mg, 2.40 mmol), and NEt₃ (2.0 mL, 14.41 mmol) were used to produce **14c**. Purification : Heptane/EtOAc (5:5). White amorphous solid (3.30 g, 96%).

H.R.M.S. (I.C., m/z) calcd for C₂₀H₂₃NNaO₅⁺ (MNa⁺) 380.1468, found : 380.1483.

I.R. (CHCl₃) ν (cm⁻¹) : 1751 (C=O (Boc)), 1696 (C=O (lactame)), 1590 (C=C), 1480 (C-Car), 1261 (Car-O), 1146 (C-O (Boc)).

M. S. (I.E., m/z) : 357 (M⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.15 (1H, t, J = 8.1 Hz, J = 7.9 Hz, H₆), 6.93 (1H, d, J = 8.3 Hz, H₇), 6.79 (1H, d, J = 7.7 Hz, H₅), 6.67 (1H, d, J = 10.1 Hz, H₂'), 6.27 (1H, d, J = 10.1 Hz, H₃'), 3.86 (3H, s, OCH₃), 2.82-2.60 (2H, system AB, J = 14.4 Hz, H_{3a} and H_{3b}), 2.44 (2H, m, H₅'), 2.19 (2H, m, H₆'), 1.60 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 198.4 (C₄'), 166.9 (C₂'), 151.4 (C₂'), 151.3 (C₈'), 150.0 (CO (Boc)), 131.7 (C_{4a}'), 131.3 (C₃'), 126.5 (C_{8a}'), 125.9 (C₆'), 118.3 (C₅'), 111.9 (C₇'), 84.3 (C (Boc)), 56.0 (OCH₃), 43.6 (C₃'), 40.1 (C₄'), 33.6 (C₅'), 32.0 (C₆'), 27.7 (CH₃ (Boc)).

General Procedure for the Synthesis of Quinolinone **15a-c**

To **14a-c** dissolved in tBuOH was added AcOH, then SeO₂. The reaction mixture was heated at 100°C for 72 hours and every 24 hours SeO₂ was added. After three days of reaction, the mixture was filtered under celite, washed with EtOAc and evaporated. The residue was dissolved in CH₂Cl₂. The combined organic layer was washed with NaOH 1M, H₂O, brine, dried (MgSO₄) and evaporated. Silica gel flash-column chromatography of the residue afforded **15a-c** (and **9c**) as indicated below for each case.

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one) **15a**

15a

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2'-en-4'-one)-carboxylic acid *tert*-butyl ester **14a** (6.18 g, 18.90 mmol), tBuOH (450 mL), AcOH (37.8 mL), and SeO₂ (4.19 g, 37.80 mmol) then (2.10 g, 18.90 mmol) were used to produce **15a**. Purification : Heptane/EtOAc (4:6). Yellow amorphous solid (3.13 g, 74%).

H.R.M.S. (E.S.I., m/z) calcd for C₁₄H₁₁NNaO₂⁺ (MNa⁺) 248.0687, found : 248.0703.

I.R. (CHCl₃) ν (cm⁻¹) : 3219 (N-H), 1665 (C=O lactame), 1614 (C=C), 1462 (C-Car).

M. S. (E.S.I., m/z) : 248 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 9.00 (1H, sl, NH), 7.35-7.28 (1H, m, Har), 7.13-7.03 (2H, m, Har), 7.02 (2H, d, J = J = 10.1, H_{3'} and H_{5'}), 6.99-6.93 (1H, m, Har), 6.42 (2H, d, J = J = 10.1, H_{2'} and H_{6'}), 2.80 (2H, s, H₃).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 184.9 (C_{4'}), 168.2 (C₂), 149.3 (C_{3'} and C_{5'}), 136.6 (C_{8a}), 129.8 (Car), 129.5 (C_{2'} and C_{6'}), 126.8 (Car), 124.5 (Car), 122.2 (C_{4a}), 116.7 (Car), 43.8 (C_{1'}), 40.2 (C₃).

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one) **15b**

15b

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2'-en-4'-one)-carboxylic acid *tert*-butyl ester **14b** (5.0 g, 14.00 mmol), tBuOH (350 mL), AcOH (28.0 mL), and SeO₂ (3.11 g, 28.01 mmol) then (1.55 g, 14.00 mmol) were used to produce **15b**. Purification : Heptane/EtOAc (4:6). Yellow amorphous solid (2.28 g, 64%).

H.R.M.S. (E.S.I., m/z) calcd for C₁₅H₁₃NNaO₃⁺ (MNa⁺) 278.0793, found : 278.0825.

I.R. (CHCl₃) ν (cm⁻¹) : 3209 (N-H), 1656 (C=O lactame), 1613 (C=C), 1485 (C-Car), 1258 (Car-O).

M. S. (E.S.I., m/z) : 278 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 9.10 (1H, sl, NH), 6.99-6.94 (3H, m, H₅, H_{5'} and H_{3'}), 6.57 (1H, dd, J = 2.5 Hz, J = 8.6 Hz, H₆), 6.49 (1H, d, J = 2.5 Hz, H₈), 6.37 (2H, d, J = 10.1 Hz, H_{2'} and H_{6'}), 3.80 (3H, s, OCH₃), 2.75 (2H, s, H₃).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 185.1 (C_{4'}), 168.6 (C₂), 160.8 (C₇), 149.8 (C_{2'} and C_{6'}), 137.9 (C_{8a}), 129.1 (C_{3'} and C_{5'}), 127.7 (C₅), 114.0 (C_{4a}), 109.6 (C₆), 102.7 (C₈), 55.6 (OCH₃), 43.2 (C_{1'}), 40.2 (C₃).

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one) **15c** (and **9c**)

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2'-en-4'-one)-carboxylic acid *tert*-butyl ester **14c** (3.52 g, 9.85 mmol), tBuOH (250 mL), AcOH (19.7 mL), and SeO₂ (2.19 g, 19.71 mmol) then (1.09 g, 9.85 mmol) were used to produce **15c** and **9c**. Purification : Heptane/EtOAc (4:6). Yellow amorphous solid (604.2 mg and 1.80 g, 74%).

H.R.M.S. (I.C., m/z) calcd for C₁₅H₁₄NO₃⁺ (MH⁺) 256.0968, found : 256.0997.

I.R. (CHCl₃) ν (cm⁻¹) : 3400 (N-H), 1669 (C=O (lactame)), 1627 (C=C), 1493 (C-Car), 1263 (Car-O).

M. S. (I.E., m/z) : 255 (M⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.94 (1H, sl, NH), 6.92 (1H, t, J = 8.1 Hz, J = 7.9 Hz, H₆), 6.91 (2H, d, J = 10.1 Hz, H_{2'} and H_{6'}), 6.81 (1H, d, J = 8.2 Hz, H₇), 6.61

(1H, d, $J = 7.7$ Hz, H_5), 6.32 (2H, d, $J = 10.1$ Hz, $H_{3'}$ and $H_{5'}$), 3.84 (3H, s, OCH_3), 2.68 (2H, s, H_3).

^{13}C RMN ($CDCl_3$, 75 MHz) δ (ppm) : 185.0 ($C_{4'}$), 166.6 (C_2), 149.3 ($C_{2'}$ and $C_{6'}$), 146.5 (C_8), 129.4 ($C_{3'}$ and $C_{5'}$), 126.2 (C_{4a}), 123.9 (C_6), 122.3 (C_{8a}), 118.3 (C_5), 110.9 (C_7), 56.1 (OCH_3), 43.9 (C_4), 40.1 (C_3).

General Procedure for the Synthesis of Carbamate Derivatives **9a-c**

To a solution of **15a-c** in dry CH_2Cl_2 , Boc_2O , DMAP and NEt_3 was successfully added. After being stirred for a night at room temperature, the reaction mixture was extracted with CH_2Cl_2 . The combined organic layer was washed with H_2O , brine, dried ($MgSO_4$) and evaporated. Silica gel flash-column chromatography of the residue afforded **9a-c** as indicated below for each case.

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-
carboxylic acid *tert*-butyl ester **9a**

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one) **15a** (1.55 g, 6.89 mmol), dry CH_2Cl_2 (85 mL), Boc_2O (3.0 g, 13.78 mmol), DMAP (210.4 mg, 1.72 mmol), and NEt_3 (1.44 mL, 10.33 mmol) were used to produce **9a**. Purification : Heptane/EtOAc (6:4). Yellow amorphous solid (1.90 g, 85%).

H.R.M.S. (E.S.I., m/z) calcd for $C_{14}H_{11}NNaO_2^+$ (MNa^+) 348.1212, found : 348.1183.

I.R. ($CHCl_3$) ν (cm^{-1}) : 1758 ($C=O$ (Boc)), 1693 ($C=O$ (lactame)), 1629 ($C=C$), 1493 ($C-Car$), 1142 ($C-O$).

M. S. (E.S.I., m/z) : 348 (MNa^+).

1H RMN ($CDCl_3$, 300 MHz) δ (ppm) : : 7.40-7.28 (1H, m, Har), 7.13-7.09 (2H, m, Har), 7.02 (1H, m, Har), 6.96 (2H, d, $J = 10.1$ Hz, $H_{2'}$ and $H_{6'}$), 6.41 (2H, d, $J = 10.1$ Hz, $H_{3'}$ and $H_{5'}$), 2.80 (2H, s, H_3), 1.64 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 184.9 (C_{4'}), 165.4 (C₂), 151.4 (CO (Boc)), 148.7 (C_{2'}), 136.6 (C_{8a}), 129.9 (C_{3'}), 129.5 (Car), 126.7 (Car), 125.3 (Car), 124.4 (C_{4a}), 117.7 (Car), 86.2 (C (Boc)), 43.4 (C_{1'}), 41.3 (C₃), 27.8 (CH₃(Boc)).

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-
carboxylic acid *tert*-butyl ester **9b**

9b

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one) **15b** (1.91 g, 7.48 mmol), dry CH₂Cl₂ (95 mL), Boc₂O (3.27 g, 14.96 mmol), DMAP (228.5 mg, 1.87 mmol), and NEt₃ (1.56 mL, 11.22 mmol) were used to produce **9b**. Purification : Heptane/EtOAc (6:4). Yellow amorphous solid (2.47 g, 93%).

H.R.M.S. (E.S.I., m/z) calcd for C₂₀H₂₁NNaO₅⁺ (MNa⁺) 378.1317, found : 378.1303.

I.R. (CHCl₃) ν (cm⁻¹) : 1764 (C=O (Boc)), 1697 (C=O (lactame)), 1667 (C=C), 1507 (C-Car), 1305 (Car-O), 1143 (C-O).

M. S. (E.S.I., m/z) : 378 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 6.99 (1H, d, J = 8.5 Hz, H₅), 6.93 (2H, d, J = 9.9 Hz, H_{2'} and H_{6'}), 6.63 (1H, dd, J = 2.3 Hz, J = 8.5 Hz, H₆), 6.57 (1H, d, J = 2.4 Hz, H₈), 6.38 (2H, d, J = 9.9 Hz, H_{3'} and H_{5'}), 3.78 (3H, s, OCH₃), 2.77 (2H, s, H₃), 1.64 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 184.9 (C_{4'}), 165.6 (C₂), 160.4 (C₇), 151.0 (CO (Boc)), 149.2 (C_{2'} and C_{6'}), 137.5 (C_{8a}), 129.5 (C_{3'} and C_{5'}), 127.6 (C₅), 116.0 (C_{4a}), 110.0 (C₆), 104.0 (C₈), 86.2 (C (Boc)), 55.6 (OCH₃), 42.9 (C₄), 41.3 (C₃), 27.7 (CH₃(Boc)).

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-
carboxylic acid *tert*-butyl ester **9c**

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one) **15c** (1.16 g, 4.55 mmol), dry CH_2Cl_2 (90 mL), Boc_2O (1.98 g, 9.10 mmol), DMAP (138.9 mg, 1.13 mmol), and NEt_3 (0.95 mL, 6.82 mmol) were used to produce **9c**. Purification : Heptane/EtOAc (6:4). Yellow amorphous solid (1.29 g, 81%).

H.R.M.S. (I.C., m/z) calcd for $\text{C}_{20}\text{H}_{21}\text{NNaO}_5^+$ (MNa^+) 378.1312, found : 378.1320.

I.R. (CHCl_3) ν (cm^{-1}) : 1753 (C=O (Boc)), 1706 (C=O (lactame)), 1628 (C=C), 1480 (C-Car), 1261 (Car-O), 1146 (C-O (Boc)).

M. S. (I.E., m/z) : 355 (M^+).

^1H RMN (CDCl_3 , 500 MHz) δ (ppm) : 7.12 (1H, t, $J = 8.1$ Hz, $J = 7.9$ Hz, H_6), 6.93 (1H, d, $J = 8.3$ Hz, H_7), 6.91 (2H, d, $J = 10.1$ Hz, $\text{H}_{2'}$ and H_6), 6.72 (1H, d, $J = 7.9$ Hz, H_5), 6.43 (2H, d, $J = 10.1$ Hz, $\text{H}_{3'}$ and H_5), 3.86 (3H, s, OCH_3), 2.75 (2H, s, H_3), 1.61 (9H, s, Boc).

^{13}C RMN (CDCl_3 , 75 MHz) δ (ppm) : 184.8 ($\text{C}_{4'}$), 166.2 (C_2), 151.1 (CO (Boc)), 149.9 (C_8), 148.5 ($\text{C}_{2'}$ and C_6), 129.9 ($\text{C}_{3'}$ and C_5), 128.4 (C_{4a}), 127.0 (C_{8a}), 126.5 (C_6), 118.3 (C_5), 112.5 (C_7), 84.7 (C (Boc)), 56.0 (OCH_3), 43.7 (C_4), 42.5 (C_3), 27.7 (CH_3 (Boc)).

General Procedure for the Synthesis of Carbazole **17a-c**

To **9a-c** MeNH_2 (33 % in ethanol) was added. After being stirred for 45 min., the solvent was evaporated. The residue was dissolved in a mixture of THF and NaOH 1M (1/1). After being stirred for 2 hours, the reaction mixture was extracted with CH_2Cl_2 . The combined organic layer was washed with H_2O , brine, dried (MgSO_4) and evaporated. Silica gel flash-column chromatography of the residue afforded **17a-c** as indicated below for each case.

(±)-3-methyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **17a**

17a

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9a** (500.0 mg, 1.54 mmol), MeNH₂ (33 % in ethanol) (60 mL), THF (12 mL), NaOH 1M (12 mL) were used to produce **17a**. Purification : CH₂Cl₂/MeOH (9.9:0.1). Yellow amorphous solid (373.7 mg, 68%).

H.R.M.S. (E.S.I., m/z) calcd for C₂₀H₂₄N₂NaO₄⁺ (MNa⁺) 379.1634, found : 379.1613.

I.R. (CHCl₃) ν (cm⁻¹) : 1686 (C=O), 1479 (C-Car), 1254 (Car-O), 1147 (C-O).

M. S. (E.S.I., m/z) : 379 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.75 (1H, d, J = 7.9 Hz, H₈), 7.30 (1H, dd, J = 7.4 Hz, J = 1.4 Hz, H₉), 7.20 (1H, dd, J = 7.2 Hz, J = 1.4 Hz, H₁₁), 7.06 (1H, dd, J = 7.2 Hz, J = 1.4 Hz, H₁₀), 4.47 (1H, dd, J = 7.2 Hz, J = 5.3 Hz, H_{6a}), 3.97 (1H, dd, J = 7.5 Hz, J = 3.8 Hz, H_{3a}), 2.90 (3H, s, NCH₃), 2.86-2.74 (4H, m, H₁ and H₆), 2.68 (1H, dd, J = 3.8 Hz, J = 17.2 Hz, H₄), 2.49 (1H, dd, J = 7.5 Hz, J = 17.2 Hz, H₄), 1.57 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.2 (C₅), 172.2 (C₂), 151.9 (CO (Boc)), 141.3 (C_{7a}), 133.3 (C_{11a}), 129.7 (C₉), 123.8 (C₁₀), 122.2 (C₁₁), 115.9 (C₈), 82.6 (C (Boc)), 65.1 (C_{6a}), 63.9 (C_{3a}), 47.0 (C_{11b}), 44.9 (C₄), 41.1 and 39.7 (C₁ and C₆), 28.4 (CH₃(Boc)), 27.6 (NCH₃).

(±)-9-methoxy-3-methyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **17b**

17b

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9b** (408.7 mg, 1.15 mmol), MeNH₂ (33 % in ethanol) (45 mL), THF (8 mL), NaOH 1M (8 mL) were used to produce **17b**. Purification : CH₂Cl₂/MeOH (9.9:0.1). Yellow amorphous solid (293.7 mg, 66%).

H.R.M.S. (E.S.I., m/z) calcd for C₂₁H₂₆N₂NaO₅⁺ (MNa⁺) 409.1739, found : 409.1726.

I.R. (CHCl₃) ν (cm⁻¹) : 1692 (C=O), 1497 (C-Car), 1253 (Car-O), 1162 (C-O).

M. S. (E.S.I., m/z) : 409 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.37 (1H, sl, H₈), 7.05 (1H, d, J = 8.3 Hz, H₁₁), 6.58 (1H, dd, J = 8.3 Hz, J = 2.4 Hz, H₁₀), 4.44 (1H, t, J = 6.0 Hz, H_{6a}), 3.90 (1H, dd, J = 3.7 Hz, J = 7.2 Hz, H_{3a}), 3.79 (3H, s, OCH₃), 2.86 (3H, s, NCH₃), 2.79-2.69 (4H, m, H₁ and H₆), 2.63 (1H, dd, J = 3.7 Hz, J = 17.2 Hz, H₄), 2.46 (1H, dd, J = 7.3 Hz, J = 17.2 Hz, H₄), 1.55 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.4 (C₅), 172.3 (C₂), 161.1 (C₉), 151.8 (CO (Boc)), 142.7 (C_{7a}), 125.0 (C_{11a}), 122.7 (C₁₁), 109.9 (C₁₀), 101.8 (C₈), 82.6 (C (Boc)), 65.8 (C_{6a}), 64.1 (C_{3a}), 55.6 (OCH₃), 46.5 (C_{11b}), 45.0 (C₆), 41.2 (C₁), 39.6 (C₄), 28.4 (CH₃(Boc)), 27.6 (NCH₃).

(±)-8-methoxy-3-methyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **17c**

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9c** (297.7 mg, 0.84 mmol), MeNH₂ (33 % in ethanol) (35 mL), THF (7 mL), NaOH 1M (7 mL) were used to produce **17c**. Purification : CH₂Cl₂/MeOH (9.9:0.1). Yellow amorphous solid (240.5 mg, 74%).

H.R.M.S. (I.C., m/z) calcd for C₂₁H₂₆N₂NaO₅⁺ (MNa⁺) 409.1734, found : 409.1728.

I.R. (CHCl₃) ν (cm⁻¹) : 1706 (C=O (Boc)), 1692 (C=O (lactame)), 1491 (C-Car), 1261 (Car-O), 1161 (C-O (Boc)).

M. S. (I.E., m/z) : 386 (M⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.13 (1H, t, J = 7.9 Hz, J = 7.7 Hz, H₁₀), 6.90 (1H, d, J = 8.1 Hz, H₉), 6.78 (1H, d, J = 7.5 Hz, H₁₁), 4.68 (1H, dd, J = 5.1 Hz, J = 11.4 Hz, H_{6a}), 4.02 (1H, dd, J = 9.2 Hz, J = 4.0 Hz, H_{3a}), 3.87 (3H, s, OCH₃), 2.94 (3H, s, NCH₃), 2.80 (1H, d, J = 16.7 Hz, H₁), 2.78 (2H, m, H₄ and H₆), 2.64 (1H, d, J = 16.5 Hz, H₁), 2.40 (2H, dd, J = 10.2 Hz, J = 16.4 Hz, H₄ and H₆), 1.50 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.1 (C₅), 171.7 (C₂), 153.0 (CO (Boc)), 150.5 (C₈), 138.7 (C_{11a}), 129.0 (C_{7a}), 126.9 (C₁₀), 113.6 (C₁₁), 112.9 (C₉), 81.5 (C (Boc)), 67.2 (C_{6a}), 62.0 (C_{3a}), 55.6 (OCH₃), 48.5 (C_{11b}), 43.9 (C₁), 42.8 (C₆), 40.9 (C₄), 28.1 (CH₃ (Boc)), 27.9 (NCH₃).

General Procedure for the Synthesis of Carbazole **18a-c** or **19a-c**

To **9a-c** the corresponding amine (EtNH₂ or BuNH₂) was added. After being stirred for 1h30 at room temperature, the solvent was evaporated. The residue was purified by preparative TLC plates to afford a fraction of the desired product, **18a-c** or **19a-c**, and another fraction of the “intermediate open” **16a-c**, which was engaged thereafter. The “intermediate open” was dissolved in dry THF, and NaH (60 %) was added at 0°C. After being stirred for 1h30 at room temperature, the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. The purification by preparative TLC plates of the residue afforded of the desired product, **18a-c** or **19a-c**, as indicated below for each case.

(±)-3-ethyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **18a**

18a

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9a** (99.7 mg, 0.31 mmol), EtNH₂ (12 mL), dry THF (2 mL), NaH (4.1 mg, 0.10

mmol) were used to produce **18a**. Purification : CH₂Cl₂/MeOH (9.5:0.5). Yellow amorphous solid (40.1 mg, 35%).

H.R.M.S. (I.C., m/z) calcd for C₂₀H₂₄N₂NaO₄⁺ (MNa⁺) 393.1790, found : 393.1811.

I.R. (CHCl₃) ν (cm⁻¹) : 1691 (C=O), 1600 (C=C), 1481 (C-Car).

M. S. (E.S.I., m/z) : 393 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.67 (1H, d, J = 7.4 Hz, H₈), 7.23 (1H, t, J = 7.4 Hz, H₉), 7.15 (1H, d, J = 8.9 Hz, H₁₁), 6.99 (1H, t, J = 7.4 Hz, H₁₀), 4.41 (1H, dd, J = 4.6 Hz, J = 8.1 Hz, H_{6a}), 4.01 (1H, dd, J = 7.8 Hz, J = 3.8 Hz, H_{3a}), 3.69 (1H, dq, J = 7.2 Hz, J = 14.5 Hz, H_{1'}), 2.98 (1H, dq, J = 7.2 Hz, J = 14.0 Hz, H_{1'}), 2.78 (1H, dd, J = 4.4 Hz, J = 17.3 Hz, H₆), 2.76-2.65 (3H, m, H₁ and H₆), 2.60 (1H, dd, J = 3.8 Hz, J = 17.2 Hz, H₄), 2.38 (1H, dd, J = 8.1 Hz, J = 17.2 Hz, H₄), 1.50 (9H, s, Boc), 1.13 (3H, t, J = 7.2 Hz, H_{2'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 204.3 (C₅), 170.7 (C₂), 150.8 (CO (Boc)), 140.0 (C_{7a}), 132.7 (C_{11a}), 128.6 (C₉), 122.7 (C₁₀), 121.1 (C₁₁), 114.8 (C₈), 81.5 (C (Boc)), 63.8 (C_{6a}), 60.1 (C_{3a}), 52.4 (C_{11b}), 44.2 (C₆), 40.1 (C₁), 39.1 (C₄), 34.4 (C_{1'}), 27.4 (CH₃(Boc)), 11.9 (C_{2'}).

(±)-9-methoxy-3-ethyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **18b**

18b

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9b** (92.4 mg, 0.26 mmol), EtNH₂ (10 mL), dry THF (2 mL), NaH (6.7 mg, 0.17 mmol) were used to produce **18b**. Purification : CH₂Cl₂/MeOH (9.5:0.5). Yellow amorphous solid (27.8 mg, 27%).

H.R.M.S. (I.C., m/z) calcd for C₂₂H₂₈N₂NaO₅⁺ (MNa⁺) 423.1896, found : 423.1907.

I.R. (CHCl₃) ν (cm⁻¹) : 1687 (C=O), 1610 (C=C), 1497 (C-Car), 1255 (Car-O).

M. S. (E.S.I., m/z) : 423 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.38 (1H, sl, H₈), 7.10 (1H, d, J = 8.4 Hz, H₁₁), 6.59 (1H, dd, J = 8.3 Hz, J = 2.4 Hz, H₁₀), 4.47 (1H, dd, J = 4.8 Hz, J = 7.7 Hz, H_{6a}), 4.03 (1H, dd, J = 7.8 Hz, J = 3.8 Hz, H_{3a}), 3.81 (3H, s, OCH₃), 3.74 (1H, dq, J = 7.3 Hz, J = 14.6 Hz, H_{1'}), 3.03 (1H, dq, J = 7.1 Hz, J = 14.1 Hz, H_{1''}), 2.83 (1H, dd, J = 4.8 Hz, J = 17.4 Hz, H₆), 2.76 (3H, m, H₁ and H₆), 2.67 (1H, dd, J = 3.8 Hz, J = 17.2 Hz, H₄), 2.45 (1H, dd, J = 7.9 Hz, J = 17.2 Hz, H₄), 1.58 (9H, s, Boc), 1.19 (3H, dq, J = 7.3 Hz, H_{2'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.5 (C₅), 171.9 (C₂), 161.0 (C₉), 151.7 (CO (Boc)), 142.4 (C_{7a}), 125.5 (C_{11a}), 122.6 (C₁₁), 109.9 (C₁₀), 101.7 (C₈), 82.5 (C (Boc)), 65.5 (C_{6a}), 61.4 (C_{3a}), 55.6 (OCH₃), 46.3 (C_{11b}), 45.3 (C₆), 41.2 (C₁), 40.0 (C₄), 35.4 (C_{1'}), 28.4 (CH₃(Boc)), 12.9 (C_{2'}).

(±)-8-methoxy-3-ethyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **18c**

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9c** (87.6 mg, 0.25 mmol), EtNH₂ (10 mL), dry THF (2 mL), NaH (4.3 mg, 0.11 mmol) were used to produce **18c**. Purification : CH₂Cl₂/MeOH (9.5:0.5). Yellow amorphous solid (62.7 mg, 63%).

H.R.M.S. (I.C., m/z) calcd for C₂₂H₂₈N₂NaO₅⁺ (MNa⁺) 423.1890, found : 423.1879.

I.R. (CHCl₃) ν (cm⁻¹) : 1706 (C=O (Boc)), 1676 (C=O (lactame)), 1488 (C-Car), 1255 (Car-O), 1158 (C-O (Boc)).

M. S. (E.S.I., m/z) : 423 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.12 (1H, t, J = 8.2 Hz, J = 7.6 Hz, H₁₀), 6.89 (1H, d, J = 8.2 Hz, H₉), 6.81 (1H, d, J = 7.3 Hz, H₁₁), 4.69 (1H, dd, J = 4.9 Hz, J = 11.6 Hz, H_{6a}), 4.11 (1H, dd, J = 10.1 Hz, J = 4.0 Hz, H_{3a}), 3.87 (3H, s, OCH₃), 3.82 (1H, m, H_{1'}), 3.06 (1H, m, H_{1''}), 2.86 (1H, dd, J = 4.9 Hz, J = 16.8 Hz, H₆), 2.87-2.62 (2H, system AB, J = 16.8 Hz, H₁), 2.80 (1H, dd, J = 4.0 Hz, J = 16.2 Hz, H₄), 2.41 (1H, dd, J

= 11.9 Hz, J = 16.8 Hz, H₆), 2.38 (1H, dd, J = 9.8 Hz, J = 16.5 Hz, H₄), 1.50 (9H, s, Boc), 1.24 (3H, t, J = 7.3 Hz, H₂[•]).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.3 (C₅), 171.2 (C₂), 153.0 (CO (Boc)), 150.5 (C₈), 139.2 (C_{11a}), 128.8 (C_{7a}), 126.9 (C₁₀), 113.6 (C₁₁), 112.8 (C₉), 81.5 (C (Boc)), 67.0 (C_{6a}), 59.3 (C_{3a}), 55.6 (OCH₃), 48.3 (C_{11b}), 44.1 (C₁), 42.8 (C₆), 41.4 (C₄), 35.5 (C₁[•]), 28.1 (CH₃ (Boc)), 12.9 (C₂[•]).

(±)-3-butyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **19a**

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9a** (105.8 mg, 0.32 mmol), BuNH₂ (10 mL), dry THF (2 mL), NaH (5.8 mg, 0.14 mmol) were used to produce **19a**. Purification : CH₂Cl₂/MeOH (9.5:0.5). Yellow amorphous solid (61.9 mg, 48%).

H.R.M.S. (I.C., m/z) calcd for C₂₃H₃₀N₂NaO₅⁺ (MNa⁺) 421.2103, found : 421.2103.

I.R. (CHCl₃) ν (cm⁻¹) : 1688 (C=O), 1480 (C-Car).

M. S. (E.S.I., m/z) : 421 (MNa⁺), 399 (M+H)⁺.

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.74 (1H, d, J = 7.7 Hz, H₈), 7.30 (1H, t, J = 7.7 Hz, H₉), 7.23 (1H, d, J = 7.4 Hz, H₁₁), 7.07 (1H, t, J = 7.4 Hz, H₁₀), 4.47 (1H, dd, J = 7.8 Hz, J = 4.8 Hz, H_{6a}), 4.06 (1H, dd, J = 7.7 Hz, J = 3.8 Hz, H_{3a}), 3.75-3.64 (1H, m, H₁[•]), 3.00-2.90 (1H, m, H₁[•]), 2.88-2.72 (4H, m, H₁ and H₄), 2.67 (1H, dd, J = 17.2 Hz, J = 3.8 Hz, H₆), 2.47 (1H, dd, J = 17.2 Hz, J = 7.8 Hz, H₆), 1.57 (9H, s, Boc), 1.38 (2H, sx, J = 7.4 Hz, H₃[•]), 1.26 (2H, sl, H₂[•]), 0.97 (3H, t, J = 7.4 Hz, H₄[•]).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.4 (C₅), 171.9 (C₂), 151.8 (CO (Boc)), 141.1 (C_{7a}), 135.6 (C_{11a}), 129.6 (C₉), 123.7 (C₁₀), 122.1 (C₁₁), 115.8 (C₈), 82.5 (C (Boc)), 64.9 (C_{6a}), 61.6 (C_{3a}), 46.8 (C_{11b}), 45.1 (C₁), 41.0 (C₆), 40.7 (C₄), 40.0 (C₁[•]), 29.7 (C₂[•]), 28.5 (CH₃ (Boc)), 20.3 (C₃[•]), 13.8 (C₄[•]).

(±)-9-methoxy-3-butyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **19b**

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9b** (89.3 mg, 0.25 mmol), BuNH₂ (10 mL), dry THF (2 mL), NaH (2.6 mg, 0.06 mmol) were used to produce **19b**. Purification : CH₂Cl₂/MeOH (9.5:0.5). Yellow amorphous solid (32.0 mg, 30%).

H.R.M.S. (I.C., m/z) calcd for C₂₄H₃₂N₂NaO₅⁺ (MNa⁺) 451.2209, found : 451.2245.

I.R. (CHCl₃) ν (cm⁻¹) : 1688 (C=O), 1498 (C-Car), 1255 (Car-O).

M. S. (E.S.I., m/z) : 451 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.38 (1H, sl, H₈), 7.09 (1H, d, J = 8.4 Hz, H₁₁), 6.60 (1H, dd, J = 8.4 Hz, J = 2.4 Hz, H₁₀), 4.46 (1H, t, J = 6.0 Hz, H_{6a}), 4.01 (1H, dd, J = 7.6 Hz, J = 3.6 Hz, H_{3a}), 3.81 (3H, s, OCH₃), 3.74-3.63 (1H, m, H_{1'}), 2.98-2.88 (1H, m, H_{1'}), 2.87-2.75 (4H, m, H₁ and H₆), 2.65 (1H, dd, J = 17.3 Hz, J = 3.7 Hz, H₄), 2.46 (1H, dd, J = 17.3 Hz, J = 7.6 Hz, H_{3a}), 1.57 (9H, s, Boc), 1.37 (2H, q, J = 7.3 Hz, H_{3'}), 1.26 (2H, sl, H_{2'}), 0.96 (3H, t, J = 7.3 Hz, H_{4'}).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.6 (C₅), 172.1 (C₂), 161.0 (C₉), 151.7 (CO (Boc)), 142.5 (C_{7a}), 125.3 (C_{11a}), 122.6 (C₁₁), 109.8 (C₁₀), 82.6 (C (Boc)), 65.6 (C_{6a}), 61.8 (C_{3a}), 55.6 (OCH₃), 46.3 (C_{11b}), 45.2 (C₆), 41.1 (C₁), 40.5 (C_{1'}), 39.8 (C₄), 29.7 (C_{2'}), 28.4 (CH₃(Boc)), 20.3 (C_{3'}), 13.8 (C_{4'}).

(±)-8-methoxy-3-butyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **19c**

19c

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9c** (90.4 mg, 0.25 mmol), BuNH₂ (10 mL), dry THF (2 mL), NaH (5.3 mg, 0.11 mmol) were used to produce **19c**. Purification : CH₂Cl₂/MeOH (9.5:0.5). Yellow amorphous solid (69.4 mg, 64%).

H.R.M.S. (I.C., *m/z*) calcd for C₂₄H₃₂N₂NaO₅⁺ (MNa⁺) 451.2203, found : 451.3849.

I.R. (CHCl₃) *v* (cm⁻¹) : 1706 (C=O (Boc)), 1681 (C=O (lactame)), 1490 (C-Car), 1264 (Car-O), 1160 (C-O (Boc)).

M. S. (E.S.I., *m/z*) : 451 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) *δ* (ppm) : 7.13 (1H, t, *J* = 8.2 Hz, *J* = 7.6 Hz, H10), 6.89 (1H, d, *J* = 8.2 Hz, H9), 6.78 (1H, d, *J* = 7.6 Hz, H11), 4.67 (1H, dd, *J* = 4.9 Hz, *J* = 11.6 Hz, H6a), 4.09 (1H, dd, *J* = 10.1 Hz, *J* = 4.3 Hz, H3a), 3.87 (3H, s, OCH₃), 3.74 (1H, m, H1'), 2.97 (1H, m, H1'), 2.85 (1H, dd, *J* = 5.2 Hz, *J* = 16.8 Hz, H6), 2.86-2.63 (2H, system AB, *J* = 17.1 Hz, H1), 2.79 (1H, dd, *J* = 4.0 Hz, *J* = 16.2 Hz, H4), 2.41 (1H, dd, *J* = 11.6 Hz, *J* = 16.8 Hz, H6), 2.40 (1H, dd, *J* = 10.4 Hz, *J* = 16.5 Hz, H4), 1.57 (2H, m, H2'), 1.50 (9H, s, Boc), 1.41 (2H, m, H3'), 0.98 (3H, t, *J* = 7.3 Hz, H4').

¹³C RMN (CDCl₃, 75 MHz) *δ* (ppm) : 205.4 (C5), 171.5 (C2), 153.0 (CO (Boc)), 150.5 (C8), 139.2 (C11a), 128.9 (C7a), 126.9 (C10), 113.6 (C11), 112.8 (C9), 81.5 (C (Boc)), 67.1 (C6a), 59.8 (C3a), 55.6 (OCH₃), 48.4 (C11b), 44.1 (C1), 42.7 (C6), 41.4 (C4), 40.7 (C1'), 29.7 (C2'), 28.2 (CH₃ (Boc)), 20.4 (C3'), 13.8 (C4').

General Procedure for the Synthesis of Carbazole 20a-c

To **9a-c** dissolved in dry MeOH, BnNH₂ was added. After being stirred at room temperature for 4 hours, a solution of HCl 1M was added and the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with a solution of NaOH (1M), H₂O, brine, dried (MgSO₄) and evaporated. The residue was dissolved in a solution of THF/NaOH 1M (1/1). After being stirred at room temperature for 7 hours,

the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. Silica gel flash-column chromatography of the residue afforded **20a-c** as indicated below for each case.

(±)-3-benzyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **20a**

20a

3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9a** (500.0 mg, 1.54 mmol), dry MeOH (8.5 mL), BnNH₂ (34 mL), THF (10 mL), NaOH 1M (10 mL) were used to produce **20a**. Purification : Heptane/EtOAc (5:5). Pale yellow amorphous solid (523.6 mg, 79%).

H.R.M.S. (I.C., *m/z*) calcd for C₂₆H₂₈N₂NaO₅⁺ (MNa⁺) 455.1947, found : 455.1931.

I.R. (CHCl₃) *v* (cm⁻¹) : 1687 (C=O), 1480 (C-Car), 1256 (Car-O).

M. S. (E.S.I., *m/z*) : 455 (MNa⁺).

¹H RMN (CDCl₃, 500 MHz) *δ* (ppm) : 7.69 (1H, d, *J* = 8.4 Hz, H₈), 7.41-7.29 (5H, m, Har (Bn)), 7.25-7.20 (1H, m, H₉), 6.95-6.87 (2H, m, H₁₀ and H₁₁), 5.07 (1H, d, *J* = 14.7 Hz, CH₂(Bn)), 4.47 (1H, dd, *J* = 8.5 Hz, *J* = 4.5 Hz, H_{6a}), 3.97 (1H, d, *J* = 14.7 Hz, CH₂(Bn)), 3.88 (1H, dd, *J* = 8.1 Hz, *J* = 4.0 Hz, H_{3a}), 2.86-2.79 (3H, m, H₁ and H₆), 2.66-2.56 (2H, m, H₄ and H₆), 2.42 (1H, dd, *J* = 8.1 Hz, *J* = 17.2 Hz, H₄), 1.56 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) *δ* (ppm) : 205.3 (C₅), 171.9 (C₂), 151.8 (CO (Boc)), 140.9 (C_{7a}), 135.6 (Car (Bn)), 133.6 (C_{11a}), 129.5 (Car (Bn)), 129.1 (Car (Bn)), 128.6 (Car (Bn)), 128.3 (Car (Bn) and C₉), 123.7 and 122.2 (C₁₀ and C₁₁), 115.8 (C₈), 82.5 (C (Boc)), 64.9 (C_{6a}), 60.8 (C_{3a}), 46.8 (C_{11b}), 45.2 (CH₂(Bn)), 44.8 (C₁), 41.1 (C₆), 40.0 (C₄), 28.4 (CH₃(Boc)).

(±)-9-methoxy-3-benzyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **20b**

7-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9b** (1.0 g, 2.82 mmol), dry MeOH (15 mL), BnNH₂ (75 mL), THF (20 mL), NaOH 1M (20 mL) were used to produce **20b**. Purification : Heptane/EtOAc (5:5). Pale yellow amorphous solid (955.6 mg, 73%).

H.R.M.S. (I.C., m/z) calcd for C₂₇H₃₀N₂NaO₅⁺ (MNa⁺) 485.2052, found : 485.2025.

I.R. (CHCl₃) ν (cm⁻¹) : 1688 (C=O), 1610 (C=O (Boc)), 1496 (C-Car), 1254 (Car-O).

M. S. (E.S.I., m/z) : 485 (MNa⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 7.36-7.26 (6H, m, H₈ and Har (Bn)), 6.78 (1H, d, J = 8.4 Hz, H₁₁), 6.46 (1H, dd, J = 8.4 Hz, J = 2.4 Hz, H₁₀), 5.07 (1H, d, J = 14.7 Hz, CH₂(Bn)), 4.47 (1H, dd, J = 8.2 Hz, J = 4.6 Hz, H_{6a}), 3.97 (1H, d, J = 14.7 Hz, CH₂(Bn)), 3.84 (1H, dd, J = 7.9 Hz, J = 3.9 Hz, H_{3a}), 3.77 (3H, s, OCH₃), 2.85-2.77 (3H, m, H₁ and H₆), 2.64 (1H, dd, J = 8.2 Hz, J = 17.2 Hz, H₆), 2.56 (1H, dd, J = 3.9 Hz, J = 17.2 Hz, H₄), 2.40 (1H, dd, J = 7.9 Hz, J = 17.2 Hz, H₄), 1.56 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.4 (C₅), 172.1 (C₂), 161.0 (C₉), 151.7 (CO (Boc)), 142.3 (C_{7a}), 135.6 (Car (Bn)), 129.1 (Car (Bn)), 128.6 (Car (Bn)), 128.3 (Car (Bn)), 125.3 (C_{11a}), 122.7 (C₁₁), 109.7 (C₁₀), 101.8 (C₈), 82.6 (C (Boc)), 65.6 (C_{6a}), 61.0 (C_{3a}), 55.6 (OCH₃), 46.3 (C_{11b}), 45.3 (C₁), 44.7 (CH₂(Bn)), 41.2 (C₆), 39.9 (C₄), 28.4 (CH₃(Boc)).

(±)-8-methoxy-3-benzyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **20c**

8-methoxy-3-hydro-1H-quinolin-2-one-4,1'-(cyclohex-2',5'-en-4'-one)-carboxylic acid *tert*-butyl ester **9c** (1.0 g, 2.82 mmol), BnNH₂ (55 mL), THF (20 mL), NaOH 1M (20 mL) were used to produce **20c**. Purification : Heptane/EtOAc (5:5). Pale yellow amorphous solid (908.2 mg, 70%).

H.R.M.S. (I.C., m/z) calcd for C₂₇H₃₀N₂NaO₅⁺ (MNa⁺) 485.2053, found : 485.2033.

I.R. (CHCl₃) ν (cm⁻¹) : 1727 (C=O (Boc)), 1679 (C=O (lactame)), 1436 (C-Car), 1266 (Car-O), 1160 (C-O (Boc)).

M. S. (E.S.I., m/z) : 485 (MNa⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 7.36 (5H, m, Har (Bn)), 6.95 (1H, t, J = 7.9 Hz, H₁₀), 6.81 (1H, d, J = 8.0 Hz, H₉), 6.39 (1H, d, J = 7.3 Hz, H₁₁), 5.12 (1H, d, J = 14.6 Hz, CH₂(Bn)), 4.67 (1H, dd, J = 5.2 Hz, J = 11.6 Hz, H_{6a}), 4.00 (1H, d, J = 14.6 Hz, CH₂(Bn)), 3.88 (1H, dd, J = 9.8 Hz, J = 4.0 Hz, H_{3a}), 3.83 (3H, s, OCH₃), 2.91-2.68 (2H, system AB, J = 16.8 Hz, H₁), 2.80 (1H, dd, J = 4.9 Hz, J = 16.8 Hz, H₆), 2.70 (1H, dd, J = 4.0 Hz, J = 16.2 Hz, H₄), 2.35 (1H, dd, J = 10.1 Hz, J = 16.5 Hz, H₄), 2.30 (1H, dd, J = 11.6 Hz, J = 16.8 Hz, H₆), 1.49 (9H, s, Boc).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 205.3 (C₅), 171.2 (C₂), 153.0 (CO (Boc)), 150.2 (C₈), 138.5 (C_{11a}), 135.1 (C_{7a}), 128.7 (Car (Bn)), 128.5 (Car (Bn)), 128.4 (Car (Bn)), 128.1 (Car (Bn)), 126.5 (C₁₀), 113.0 (C₁₁), 112.4 (C₉), 81.3 (C (Boc)), 67.1 (C_{6a}), 58.5 (C_{3a}), 55.4 (OCH₃), 48.0 (C_{11b}), 44.6 (CH₂(Bn)), 43.7 (C₁), 42.2 (C₆), 40.9 (C₄), 27.9 (CH₃ (Boc)).

General Procedure for the Synthesis of Carbazole **21a-c**

To **20a-c** dissolved in CH₂Cl₂, TFA was added. After being stirred at room temperature for 5 hours, NaOH 1M was added and the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. To the residue dissolved in toluene, Na₂SO₄, ethylene glycol and *para*-toluenesulfonic acid were added successively. After being stirred at 140°C for 4 hours, NaOH 1M was added and the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. Silica gel flash-column chromatography of the residue afforded **21a-c** as indicated below for each case.

(±)-3-benzyl-2-oxo-5-(1',4'-dioxaspiro[4.5])-1,3a,4,6,6a,7-hexahydro-pyrrolo[2,3-*d*]carbazole **21a**

21a

(±)-3-benzyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **20a** (350.3 mg, 0.81 mmol), dry CH₂Cl₂ (20 mL), TFA (2 mL), toluene (15 mL), Na₂SO₄ (5 g), ethylene glycol (226.7 μL, 4.05 mmol), and *para*-toluenesulfonic acid (15.4 mg, 0.08 mmol) were used to produce **21a**. Purification : CH₂Cl₂/MeOH (9.9:0.1). White amorphous solid (231.3 mg, 76%).

H.R.M.S. (I.C., *m/z*) calcd for C₂₃H₂₄N₂NaO₃⁺ (MNa⁺) 399.1685, found : 399.1671.

I.R. (CHCl₃) *v* (cm⁻¹) : 3333 (N-H), 1673 (C=O), 1482 (C-Car), 1257 (Car-O), 1049 (C-O).

M. S. (E.S.I., *m/z*) : 399 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) *δ* (ppm) : 7.31-7.25 (5H, m, Har (Bn)), 7.05 (1H, dd, *J* = 7.7 Hz, *J* = 1.2 Hz, H₉), 6.93 (1H, d, *J* = 7.4 Hz, H₁₁), 6.72-6.65 (2H, m, H₈ and H₁₀), 5.07 (1H, d, *J* = 15.0 Hz, CH₂(Bn)), 4.07 (1H, d, *J* = 15.0 Hz, CH₂(Bn)), 3.95-3.88 (5H, m, Hdioxolane and H_{6a}), 3.62 (1H, dd, *J* = 7.5 Hz, *J* = 4.6 Hz, H_{3a}), 2.71 (2H, system AB, *J* = 16.9 Hz, H₁), 2.02-1.87 (3H, m, H₄ and H₆), 1.78 (1H, dd, *J* = 7.5 Hz, *J* = 14.4 Hz, H₄).

¹³C RMN (CDCl₃, 75 MHz) *δ* (ppm) : 172.9 (C₂), 149.3 (C_{7a}), 136.3 (Car (Bn)), 133.4 (C_{11a}), 128.8 (Car (Bn)), 128.6 (Car (Bn)), 128.4 (Car (Bn)), 127.7 (C₉), 122.4 (C₁₁), 119.4 (C₁₀), 111.1 (C₈), 106.9 (C₅), 64.3 and 64.4 (Cdioxolane), 63.2 (C_{6a}), 60.5 (C_{3a}), 47.4 (C_{11b}), 44.5 (CH₂(Bn)), 42.3 (C₁), 35.9 (C₆), 34.3 (C₄).

(±)-9-methoxy-3-benzyl-2-oxo-5-(1',4'-dioxaspiro[4.5])-1,3a,4,6,6a,7-hexahydro-pyrrolo[2,3-*d*]carbazole **21b**

(±)-9-methoxy-3-benzyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **20b** (955.6 mg, 2.07 mmol), dry CH₂Cl₂ (50 mL), TFA (5 mL), toluene (38 mL), Na₂SO₄ (18 g), ethylene glycol (578.3 μL, 10.34 mmol), and *para*-toluenesulfonic acid (39.3 mg, 0.21 mmol) were used to produce **21b**. Purification : CH₂Cl₂/MeOH (9.9:0.1). White amorphous solid (539.7 mg, 64%).

H.R.M.S. (I.C., *m/z*) calcd for C₂₄H₂₆N₂NaO₄⁺ (MNa⁺) 429.1790, found : 429.1776.

I.R. (CHCl₃) *v* (cm⁻¹) : 3333 (N-H), 1680 (C=O), 1498 (C-Car), 1260 (Car-O), 1080 (C-O).

M. S. (E.S.I., *m/z*) : 429 (MNa⁺).

¹H RMN (CDCl₃, 300 MHz) *δ* (ppm) : 7.28-7.22 (5H, m, Har (Bn)), 6.81 (1H, d, *J* = 7.1 Hz, H₁₁), 6.22 (2H, m, H₈ and H₁₀), 5.06 (1H, d, *J* = 15.0 Hz, CH₂(Bn)), 4.06 (1H, d, *J* = 15.0 Hz, CH₂(Bn)), 3.93–3.87 (5H, m, Hdioxolane and H_{6a}), 3.72 (3H, s, OCH₃), 3.59 (1H, dd, *J* = 7.3 Hz, *J* = 4.6 Hz, H_{3a}), 2.68 (2H, system AB, *J* = 17.9 Hz, H₁), 2.01-1.86 (3H, m, H₄ and H₆), 1.77 (1H, dd, *J* = 7.4 Hz, *J* = 14.4 Hz, H₄).

¹³C RMN (CDCl₃, 75 MHz) *δ* (ppm) : 172.9 (C₂), 160.7 (C₉), 150.6 (C_{7a}), 136.3 (Car (Bn)), 128.7 (Car (Bn)), 128.4 (Car (Bn)), 127.7 (Car (Bn)), 125.7 (C_{11a}), 122.8 (C₁₁), 106.9 (C₅), 104.7 (C₈), 97.5 (C₁₀), 64.2 and 64.4 (Cdioxolane), 63.6 (C_{6a}), 60.7 (C_{3a}), 55.4 (OCH₃), 46.7 (C_{11b}), 44.4 (CH₂(Bn)), 42.5 (C₁), 36.0 (C₆), 34.2 (C₄).

(±)-8-methoxy-3-benzyl-2-oxo-5-(1',4'-dioxaspiro[4.5])-1,3a,4,6,6a,7-hexahydro-pyrrolo[2,3-*d*]carbazole **21c**

(±)-8-methoxy-3-benzyl-2,5-dioxo-1,3a,4,6,6a-pentahydro-pyrrolo[2,3-*d*]carbazole-7-carboxylic acid *tert*-butyl ester **20c** (828.4 mg, 1.79 mmol), dry CH₂Cl₂ (45 mL), TFA (4.5 mL), toluene (35 mL), Na₂SO₄ (15 g), ethylene glycol (501.3 µL, 8.96 mmol), and *para*-toluenesulfonic acid (34.1 mg, 0.18 mmol) were used to produce **21c**. Purification : CH₂Cl₂/MeOH (9.9:0.1). White amorphous solid (491.2 mg, 67%).

H.R.M.S. (I.C., m/z) calcd for C₂₄H₂₇N₂O₄⁺ (MH⁺) 407.1971, found : 407.1960.

I.R. (CHCl₃) ν (cm⁻¹) : 3343 (N-H), 1680 (C=O (lactame)), 1490 (C-Car), 1252 (Car-O), 1013 (C-O).

M. S. (E.S.I., m/z) : 429 (MNa⁺), 407 (MH⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.27 (5H, m, Har (Bn)), 6.67 (2H, m, H₉ and H₁₀), 6.57 (1H, dd, J = 6.4 Hz, J = 2.1 Hz, H₁₁), 5.06 (1H, d, J = 14.9 Hz, CH₂(Bn)), 4.05 (1H, d, J = 14.9 Hz, CH₂(Bn)), 3.91 (5H, m, Hdioxolane and H_{6a}), 3.79 (3H, s, OCH₃), 3.68 (1H, dd, J = 4.7 Hz, J = 7.5 Hz, H_{3a}), 2.81-2.58 (2H, system AB, J = 16.8 Hz, H₁), 1.91 (3H, m, H₄ and H₆), 1.78 (1H, dd, J = 7.5 Hz, J = 14.3 Hz, H₄).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 172.8 (C₂), 146.1 (C₈), 138.2 (C_{7a}), 136.2 (Car (Bn)), 134.3 (C_{11a}), 128.7 (Car (Bn)), 128.4 (Car (Bn)), 127.6 (Car (Bn)), 120.5 (C₁₀), 114.5 (C₁₁), 110.2 (C₉), 106.9 (C₅), 64.3 (Cdioxolane), 63.5 (C_{6a}), 60.3 (C_{3a}), 55.2 (OCH₃), 48.2 (C_{11b}), 44.5 (CH₂(Bn)), 42.3 (C₁), 36.0 (C₆), 34.4 (C₄).

General Procedure for the Synthesis of Carbazole **22a-c**

To **21a-c** dissolved in dry THF, LiAlH₄ 1M in THF was added. The reaction mixture was stirred for 5 hours at 60°C. After addition of potassium sodium tartrate at 0°C, the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with H₂O, brine, dried (MgSO₄) and evaporated. Alumina gel flash-column chromatography of the residue afforded **22a-c** as indicated below for each case.

(±)-3-benzyl-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*]carbazole **22a**

22a

(±)-3-benzyl-2-oxo-5-(1',4'-dioxaspiro[4.5])-1,3a,4,6,6a,7-hexahydro-pyrrolo[2,3-*d*]carbazole **21a** (287.3 mg, 0.76 mmol), dry THF (15 mL), and LiAlH₄ 1M in THF (3.8 mL, 3.82 mmol) were used to produce **22a**. Purification : Heptane/EtOAc (7:3). White amorphous solid (216.7 mg, 78%).

H.R.M.S. (I.C., *m/z*) calcd for C₂₃H₂₇N₂O₃⁺ (MH⁺) 363.2073, found : 363.2049.

I.R. (CHCl₃) ν (cm⁻¹) : 3357 (N-H), 1483 (C-Car), 1042 (C-O).

M. S. (E.S.I., *m/z*) : 363 (MH⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.40 (2H, d, *J* = 7.6 Hz, Har (Bn)), 7.30 (2H, t, *J* = 7.6 Hz, Har (Bn)), 7.23 (1H, t, *J* = 7.6 Hz, Har (Bn)), 7.10 (1H, d, *J* = 7.3 Hz, H₁₁), 7.04 (1H, t, *J* = 7.6 Hz, H₉), 6.77 (1H, t, *J* = 7.6 Hz, H₁₀), 6.66 (1H, d, *J* = 7.6 Hz, H₈), 4.05 (1H, d, *J* = 12.8 Hz, CH₂(Bn)), 4.00-3.87 (4H, m, Hdioxolane), 3.78 (1H, t, *J* = 4.3 Hz, H_{6a}), 3.15 (1H, d, *J* = 12.6 Hz, CH₂(Bn)), 2.92 (1H, dt, *J* = 2.1 Hz, *J* = 8.9 Hz, H₂), 2.56 (1H, dd, *J* = 5.2 Hz, H_{3a}), 2.42 (1H, dd, *J* = 14.6 Hz, *J* = 3.7 Hz, H₆), 2.39 (1H, q, *J* = 8.9 Hz, H₂), 2.15 (1H, ddd, *J* = 13.4 Hz, *J* = 8.2 Hz, *J* = 2.1 Hz, H₁), 2.02-1.97 (2H, m, H₄ and H₆), 1.89-1.84 (2H, m, H₄ and H₁).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 150.1 (C_{7a}), 139.7 (Car (Bn)), 136.7 (C_{11a}), 129.0 (Car (Bn)), 128.2 (Car (Bn)), 126.8 (Car (Bn)), 127.8 (C₉), 123.5 (C₁₁), 119.3 (C₁₀), 110.4 (C₈), 108.2 (C₅), 68.5 (C_{3a}), 65.1 (C_{6a}), 64.1 and 63.8 (Cdioxolane), 56.6 (CH₂(Bn)), 53.0 (C_{11b}), 52.2 (C₂), 37.6 (C₁), 35.3 (C₆), 32.1 (C₄).

(±)-9-methoxy-3-benzyl-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*]carbazole **22b**

22b

(±)-9-methoxy-3-benzyl-2-oxo-5-(1',4'-dioxaspiro[4.5])-1,3a,4,6,6a,7-hexahydro-pyrrolo [2,3-*d*] carbazole **21b** (97.5 mg, 0.24 mmol), dry THF (5 mL), and LiAlH₄ 1M in THF (1.2 mL, 1.20 mmol) were used to produce **22b**. Purification : Heptane/EtOAc (7:3). White amorphous solid (54.8 mg, 58%).

H.R.M.S. (I.C., m/z) calcd for C₂₄H₂₉N₂O₃⁺ (MH⁺) 393.2178, found : 393.2165.

I.R. (CHCl₃) ν (cm⁻¹) : 3352 (N-H), 1497 (C-Car), 1196 (Car-O), 1159 (C-O).

M. S. (E.S.I., m/z) : 393 (MH⁺).

¹H RMN (CDCl₃, 300 MHz) δ (ppm) : 7.40 (2H, d, J = 7.6 Hz, Har (Bn)), 7.30 (2H, t, J = 7.6 Hz, Har (Bn)), 7.23 (1H, m, Har (Bn)), 6.97 (1H, d, J = 8.2 Hz, H₁₁), 6.31 (1H, dd, J = 8.2 Hz, J = 2.2 Hz, H₁₀), 6.25 (1H, d, J = 2.2 Hz, H₈), 4.05 (1H, d, J = 12.6 Hz, CH₂(Bn)), 4.00-3.85 (4H, m, Hdioxolane), 3.78 (1H, t, J = 4.2 Hz, H_{6a}), 3.75 (3H, s, OCH₃), 3.13 (1H, d, J = 12.6 Hz, CH₂(Bn)), 2.90 (1H, dt, J = 8.9 Hz, J = 2.0 Hz, H₂), 2.50 (1H, t, J = 3.6 Hz, H_{3a}), 2.43 (1H, dd, J = 3.6 Hz, J = 14.6 Hz, H₄), 2.34 (1H, q, J = 8.9 Hz, H₂), 2.11 (1H, ddd, J = 13.2 Hz, J = 8.2 Hz, J = 2.0 Hz, H₁), 2.00-1.96 (2H, m, H₄ and H₆), 1.87-1.80 (2H, m, H₁ and H₆).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 160.2 (C₉), 151.4 (C_{7a}), 139.8 (Car (Bn)), 129.1 (C_{11a} and Car Bn), 128.2 (Car (Bn)), 126.8 (Car (Bn)), 123.8 (C₁₁), 108.2 (C₅), 104.4 (C₁₀), 96.8 (C₈), 68.7 (C_{3a}), 65.7 (C_{6a}), 64.1 and 63.9 (Cdioxolane), 56.7 (CH₂(Bn)), 55.4 (OCH₃), 52.4 (C_{11b}), 52.2 (C₂), 37.7 (C₁), 35.4 (C₄), 32.1 (C₆).

(±)-8-methoxy-3-benzyl-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*]carbazole **22c**

(±)-8-methoxy-3-benzyl-2-oxo-5-(1',4'-dioxaspiro[4.5])-1,3a,4,6,6a,7-hexahydro-pyrrolo [2,3-*d*] carbazole **21c** (458.4 mg, 1.13 mmol), dry THF (22 mL), and LiAlH₄ 1M in THF (5.6 mL, 5.64 mmol) were used to produce **22c**. Purification : Heptane/EtOAc (7:3). White amorphous solid (339.2 mg, 77%).

H.R.M.S. (I.C., m/z) calcd for C₂₄H₂₉N₂O₃⁺ (MH⁺) 393.2178, found : 393.2148.

I.R. (CHCl₃) ν (cm⁻¹) : 3331 (N-H), 1493 (C-Car), 1240 (Car-O), 1044 (C-O).

M. S. (E.S.I., *m/z*) : 393 (MH⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 7.39 (2H, d, *J* = 7.6 Hz, Har (Bn)), 7.31 (2H, t, *J* = 8.2 Hz, *J* = 7.6 Hz, Har (Bn)), 7.25 (1H, d, *J* = 8.9 Hz, Har (Bn)), 6.78 (2H, m, H₉ and H₁₀), 6.67 (1H, m, H₁₁), 4.06 (1H, d, *J* = 12.2 Hz, CH₂(Bn)), 3.98 (1H, dd, *J* = 3.7 Hz, *J* = 8.2 Hz, H_{6a}), 3.90 (4H, m, Hdioxolane), 3.80 (3H, s, OCH₃), 3.22 (1H, d, *J* = 12.5 Hz, CH₂(Bn)), 2.91 (1H, t, *J* = 7.6 Hz, H₂), 2.65 (1H, sl, H_{3a}), 2.41 (1H, t, *J* = 8.2 Hz, H₂), 2.35 (1H, dd, *J* = 4.0 Hz, *J* = 14.6 Hz, H₆), 2.11 (1H, ddd, *J* = 1.8 Hz, *J* = 8.2 Hz, *J* = 13.1 Hz, H₁), 1.98 (2H, m, H₄ and H₆), 1.87 (2H, m, H₁ and H₄).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 145.8 (C₈), 139.6 (Car (Bn)), 138.9 (C_{7a}), 137.6 (C_{11a}), 129.0 (Car (Bn)), 128.1 (Car (Bn)), 126.7 (Car (Bn)), 119.8 (C₁₀), 115.7 (C₁₁), 109.4 (C₉), 108.1 (C₅), 67.7 (C_{3a}), 65.3 (C_{6a}), 63.9 (Cdioxolane), 56.7 (CH₂(Bn)), 55.2 (OCH₃), 53.9 (C_{11b}), 52.0 (C₂), 37.4 (C₁), 35.6 (C₆), 32.4 (C₄).

General Procedure for the Synthesis of Carbazole **23a-c**

To a spatula of Ni Raney dissolved in EtOH, **22a-c** was added. After being stirred at room temperature for 18 hours under H₂ pressure, the reaction mixture was filtered with celite, washed with MeOH and evaporated. The purification by preparative TLC plates of the residue afforded of the desired product, **23a-c**, and the starting product, **22a-c**, as indicated below for each case.

(±)-5-(1',4'-dioxaspiro[4.5])-1,2,3,3a,4,6,6a,7-octahydro-pyrrolo[2,3-*d*]carbazole **23a**

23a

(±)-3-benzyl-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*]carbazole **22a** (53.7 mg, 0.15 mmol), a spatula of Ni Raney, and EtOH (5 mL) were used to produce **23a**. Purification : CH₂Cl₂/MeOH (9:1). Yellow amorphous solid (22.9 mg, 57%). 7.6 mg of **22a** were recovered.

H.R.M.S. (I.C., m/z) calcd for $C_{16}H_{21}N_2O_3^+$ (MH^+) 273.1603, found : 273.1602.

I.R. ($CHCl_3$) ν (cm^{-1}) : 3345 (N-H), 1483 and 1463 (C-Car), 1046 (C-O).

M. S. (E.S.I., m/z) : 273 (MH^+).

1H RMN ($CDCl_3$, 300 MHz) δ (ppm) : 7.07 (1H, dd, $J = 7.3$ Hz, $J = 7.6$ Hz, H_9 and H_{11}), 6.69 (1H, d, $J = 7.6$ Hz, H_8), 4.03-3.91 (4H, m, Hdioxolane), 3.84 (2H, sl, NH), 3.79-3.74 (2H, m, H_{3a} and H_{6a}), 3.28-3.12 (2H, m, H_2), 2.34 (1H, ddd, $J = 12.9$ Hz, $J = 8.2$ Hz, $J = 4.7$ Hz, H_1), 2.06-2.00 (1H, m, H_4), 1.92-1.84 (3H, m, H_1 , H_4 and H_6), 1.69 (1H, dd, $J = 13.2$ Hz, $J = 9.5$ Hz, H_6).

^{13}C RMN ($CDCl_3$, 75 MHz) δ (ppm) : 149.8 (C_{7a}), 132.5 (C_{11a}), 128.2 (C_9), 122.2 (C_{11}), 119.7 (C_{10}), 110.8 (C_8), 107.9 (C_5), 64.2 and 64.8 (Cdioxolane), 62.7 and 60.7 (C_{3a} and C_{6a}), 53.7 (C_{11b}), 44.2 (C_2), 39.1 (C_1), 38.0 (C_6), 33.3 (C_4).

($\pm$)-9-methoxy-5-(1',4'-dioxaspiro[4.5])-1,2,3,3a,4,6,6a,7-octahydro-pyrrolo[2,3-*d*]carbazole **23b**

23b

($\pm$)-9-methoxy-3-benzyl-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*] carbazole **22b** (63.3 mg, 0.16 mmol), a spatula of Ni Raney, and EtOH (5 mL) were used to produce **23b**. Purification : CH_2Cl_2 /MeOH (9:1). Yellow amorphous solid (29.4 mg, 60%). 17.1 mg of **22b** were recovered.

H.R.M.S. (I.C., m/z) calcd for $C_{17}H_{23}N_2O_3^+$ (MH^+) 303.1709, found : 303.1695.

I.R. ($CHCl_3$) ν (cm^{-1}) : 3341 (N-H), 1498 (C-Car), 1269 (Car-O), 1091 (C-O).

M. S. (E.S.I., m/z) : 303 (MH^+).

1H RMN ($CDCl_3$, 500 MHz) δ (ppm) : 6.94 (1H, d, $J = 8.2$ Hz, H_{11}), 6.31 (1H, d, $J = 8.2$ Hz, H_{10}), 6.27 (1H, s, H_8), 4.00-3.90 (4H, m, Hdioxolane), 3.75 (4H, sl, OCH_3 and H_{6a}), 3.61 (1H, sl, H_{3a}), 3.46 (2H, sl, 2 x NH), 3.19 (1H, m, H_2), 3.11 (1H, m, H_2), 2.28 (1H, ddd, $J = 13.2$ Hz, $J = 8.2$ Hz, $J = 4.8$ Hz, H_1), 1.97 (1H, d, $J = 14.5$ Hz, H_4), 1.91-1.82 (3H, m, H_1 , H_4 and H_6), 1.70 (1H, dd, $J = 13.6$ Hz, $J = 9.1$ Hz, H_6).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 160.4 (C₉), 151.1 (C_{7a}), 125.6 (C_{11a}), 122.4 (C₁₁), 108.0 (C₅), 104.2 (C₁₀), 97.6 (C₈), 64.1 and 64.7 (Cdioxolane), 63.3 (C_{6a}), 60.9 (C_{3a}), 55.4 (OCH₃), 53.1 (C_{11b}), 44.4 (C₂), 39.5 (C₁), 38.0 (C₆), 33.6 (C₄).

(±)-8-methoxy-5-(1',4'-dioxaspiro[4.5])-1,2,3,3a,4,6,6a,7-octahydro-pyrrolo[2,3-*d*]carbazole **23c**

(±)-8-methoxy-3-benzyl-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*] carbazole **22c** (36.6 mg, 0.09 mmol), a spatula of Ni Raney, and EtOH (5 mL) were used to produce **23c**. Purification : CH₂Cl₂/MeOH (9:1). Yellow amorphous solid (16.5 mg, 58%). 14.1 mg of **22c** were recovered.

H.R.M.S. (I.C., m/z) calcd for C₁₇H₂₃N₂O₃⁺ (MH⁺) 303.1708, found : 303.1727.

I.R. (CHCl₃) ν (cm⁻¹) : 3342 (N-H), 1488 (C-Car), 1257 (Car-O), 1035 (C-O).

M. S. (E.S.I., m/z) : 325 (MNa⁺), 303 (MH⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 6.79 (1H, t, J = 7.9 Hz, J = 7.6 Hz, H₁₀), 6.75 (1H, d, J = 7.2 Hz, H₁₁), 6.71 (1H, d, J = 7.9 Hz, H₉), 4.03-3.89 (4H, m, Hdioxolane), 3.82 (3H, s, OCH₃), 3.79 (1H, dd, J = 2.7 Hz, J = 5.2 Hz, H_{3a}), 3.78 (1H, dd, J = 5.5 Hz, J = 10.1 Hz, H_{6a}), 3.64 (2H, sl, NH), 3.21 (2H, m, H₂), 2.39 (1H, ddd, J = 4.3 Hz, J = 8.2 Hz, J = 12.8 Hz, H₁), 2.04 (1H, dd, J = 2.1 Hz, J = 16.5 Hz, H₄), 1.87 (3H, m, H₁, H₄ and H₆), 1.65 (1H, dd, J = 10.1 Hz, J = 13.4 Hz, H₆).

¹³C RMN (CDCl₃, 75 MHz) δ (ppm) : 146.2 (C₈), 138.5 (C_{7a}), 133.0 (C_{11a}), 120.3 (C₁₀), 114.4 (C₁₁), 110.0 (C₉), 107.8 (C₅), 64.4 (Cdioxolane), 62.9 (C_{6a}), 60.4 (C_{3a}), 55.3 (OCH₃), 54.3 (C_{11b}), 43.9 (C₂), 38.9 (C₁), 38.1 (C₆), 33.1 (C₄).

General Procedure for the Synthesis of Carbazole **8a-c**

To **23a-c** dissolved in THF, HCl 35% H₂O was added at 0°C. After being stirred at room temperature for 24 hours, the reaction mixture was extracted with CH₂Cl₂. The combined organic layer was washed with NaOH 1M, H₂O, brine, dried (MgSO₄) and evaporated. The purification by preparative TLC plates of the residue afforded **8a-c** as indicated below for each case.

(±)-5-oxo-1,2,3,3a,4,6,6a,7-octahydro-pyrrolo[2,3-*d*]carbazole **8a**

(±)-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*] carbazole **23a** (20.3 mg, 0.07 mmol), THF (1 mL), and HCl 35% H₂O (1 mL) were used to produce **8a**. Purification : CH₂Cl₂/MeOH (9:1). Yellow amorphous solid (8.8 mg, 52%).

H.R.M.S. (I.C., m/z) calcd for C₁₄H₁₇N₂O₂⁺ (MH⁺) 229.1341, found : 229.1314.

I.R. (CHCl₃) ν (cm⁻¹) : 3350 (N-H), 1708 (C=O), 1483 (C-Car), 1259 (Car-O).

M. S. (E.S.I., m/z) : 229 (MH⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 7.14 (1H, d, J = 7.3 Hz, H₁₁), 7.07 (1H, t, J = 7.3 Hz, H₉), 6.80 (1H, t, J = 7.3 Hz, H₁₀), 6.62 (1H, d, J = 7.3 Hz, H₈), 4.03 (1H, t, J = 3.2 Hz, H_{6a}), 3.29 (1H, t, J = 3.2 Hz, H_{3a}), 3.25 (1H, dd, J = 9.0 Hz, J = 3.5 Hz, H₂), 3.15 (1H, q, J = 9.0 Hz, H₂), 2.96 (1H, dd, J = 17.7 Hz, J = 3.2 Hz, H₆), 2.57 (1H, dd, J = 17.3 Hz, J = 3.2 Hz, H₄), 2.55 (1H, dd, J = 17.7 Hz, J = 3.2 Hz, H₆), 2.51 (1H, dd, J = 8.5 Hz, J = 3.5 Hz, H₁), 2.42 (1H, dd, J = 17.3 Hz, J = 3.2 Hz, H₄), 1.99 (1H, m, H₁).

¹³C RMN (CDCl₃, 125 MHz) δ (ppm) : 210.2 (C₅), 150.3 (C_{7a}), 134.7 (C_{11a}), 128.2 (C₉), 123.2 (C₁₁), 119.6 (C₁₀), 109.6 (C₈), 66.1 (C_{3a}), 65.6 (C_{6a}), 51.4 (C_{11b}), 46.3 (C₂), 41.2 (C₄), 40.5 (C₁), 40.2 (C₆).

(±)-9-methoxy-5-oxo-1,2,3,3a,4,6,6a,7-octahydro-pyrrolo[2,3-*d*]carbazole **8b**

(±)-9-methoxy-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*]carbazole **23b** (27.4 mg, 0.09 mmol), THF (1 mL), and HCl 35% H₂O (1 mL) were used to produce **8b**. Purification : CH₂Cl₂/MeOH (9:1). Yellow amorphous solid (7.5 mg, 32%).

H.R.M.S. (I.C., *m/z*) calcd for C₁₅H₁₉N₂O₂⁺ (MH⁺) 259.1447, found : 259.1434.

I.R. (CHCl₃) ν (cm⁻¹) : 3358 (N-H), 1709 (C=O), 1500 (C-Car), 1218 (Car-O).

M. S. (E.S.I., *m/z*) : 281 (MNa⁺), 259 (MH⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 7.01 (1H, d, *J* = 8.2 Hz, H₁₁), 6.34 (1H, dd, *J* = 8.2 Hz, *J* = 2.2 Hz, H₁₀), 6.20 (1H, d, *J* = 2.2 Hz, H₈), 4.03 (1H, s, H_{6a}), 3.76 (3H, s, OCH₃), 3.26 (1H, s, H_{3a}), 3.23 (1H, d, *J* = 8.2 Hz, *J* = 3.8 Hz, H₂), 3.13 (1H, q, *J* = 8.2 Hz, H₂), 2.95 (1H, dd, *J* = 17.3 Hz, *J* = 3.2 Hz, H₆), 2.55 (1H, dd, *J* = 17.3 Hz, *J* = 3.2 Hz, H₆), 2.53 (1H, dd, *J* = 17.3 Hz, *J* = 2.8 Hz, H₄), 2.47 (1H, tt, *J* = 9.3 Hz, *J* = 4.6 Hz, H₁), 2.41 (1H, dd, *J* = 17.3 Hz, *J* = 2.8 Hz, H₄), 1.97 (1H, m, H₁), 1.79 (2H, sl, 2 x NH).
¹³C RMN (CDCl₃, 125 MHz) δ (ppm) : 210.2 (C₅), 160.6 (C₉), 151.6 (C_{7a}), 126.9 (C_{11a}), 123.6 (C₁₁), 104.7 (C₁₀), 96.1 (C₈), 66.3 and 66.1 (C_{3a} and C_{6a}), 55.5 (OCH₃), 50.8 (C_{11b}), 46.2 (C₂), 41.1 - 40.5 - 40.4 (C₁ - C₄ - C₆).

(±)-8-methoxy-5-oxo-1,2,3,3a,4,6,6a,7-octahydro-pyrrolo[2,3-*d*]carbazole **8c**

(±)-8-methoxy-5-(1',4'-dioxaspiro[4.5])-1,2,3a,4,6,6a,7-heptahydro-pyrrolo[2,3-*d*]carbazole **23c** (32.8 mg, 0.11 mmol), THF (1 mL), and HCl 35% H₂O (1 mL) were

used to produce **8c**. Purification : CH₂Cl₂/MeOH (9:1). Yellow amorphous solid (12.8 mg, 46%).

H.R.M.S. (I.C., m/z) calcd for C₁₅H₁₉N₂O₂⁺ (MH⁺) 259.1441, found : 259.1467.

I.R. (CHCl₃) ν (cm⁻¹) : 3348 (N-H), 1710 (C=O), 1489 (C-Car), 1258 (Car-O).

M. S. (E.S.I., m/z) : 281 (MNa⁺), 259 (MH⁺).

¹H RMN (CDCl₃, 500 MHz) δ (ppm) : 6.73 (2H, m, H₁₀ and H₁₁), 6.63 (1H, m, H₉), 3.97 (1H, sl, H_{6a}), 3.75 (3H, s, OCH₃), 3.28 (1H, t, J = 3.4 Hz, H_{3a}), 3.21 (1H, td, J = 3.4 Hz, J = 8.2 Hz, J = 9.8 Hz, H₂), 3.08 (1H, q, J = 8.9 Hz, H₂), 2.90 (1H, dd, J = 3.4 Hz, J = 17.7 Hz, H₆), 2.55 (1H, dd, J = 3.4 Hz, J = 17.7 Hz, H₄), 2.54 (1H, dd, J = 3.1 Hz, J = 17.7 Hz, H₆), 2.46 (1H, ddd, J = 3.4 Hz, J = 8.2 Hz, J = 13.4 Hz, H₁), 2.39 (1H, dd, J = 3.4 Hz, J = 17.4 Hz, H₄), 1.93 (1H, dt, J = 8.2 Hz, J = 13.7 Hz, H₁).

¹³C RMN (CDCl₃, 125 MHz) δ (ppm) : 209.5 (C₅), 145.2 (C₈), 139.4 (C_{7a}), 135.3 (C_{11a}), 120.3 (C₁₀), 115.4 (C₁₁), 109.9 (C₉), 66.1 (C_{6a}), 65.7 (C_{3a}), 55.4 (OCH₃), 52.1 (C_{11b}), 46.3 (C₂), 41.1 (C₄), 40.2 (C₆), 40.2 (C₁).