SUPPORTING INFORMATION

Formation of Five-Membered Carbocycles by Intramolecular Palladium-Catalyzed Ring-Opening of *tert*-Cyclobutanols

Manivannan Ethirajan, Heong-Sub Oh, and Jin Kun Cha*

Department of Chemistry, Wayne State University, 5101 Cass Ave, Detroit, MI 48202

jcha@chem.wayne.edu

Experimental Section	S1-S4
Characterization Data	S1-S8
¹H and ¹³C NMR Spectra	S9-S52

Mukayama-type Condensation of **1** and **2**.

TMS ether **1** (1.0 g, 4.76 mmol) and acetal **2** (1.25 mL, 6.18 mmol) were dissolved in CH₂Cl₂. A solution of TiCl₄ (0.68 mL, 6.18 mmol) in CH₂Cl₂ was added dropwise at –40 °C. The reaction mixture was stirred at –40 °C for 2 h and then quenched with addition of wet ether. The resulting mixture was allowed to warm to room temperature. The resulting mixture was diluted with ether. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography using 2:98 ethyl acetate–hexane as eluent to afford **3** (1.40 g, 84%) as a 3:1 mixture of two diastereomers.

3 (the major isomer) as a white solid: mp 72 °C; *R*_f 0.7 (1:9 EtOAc–hexane); IR (CH₂Cl₂) 1768 cm^{–1}; ¹H NMR (400 MHz, CDCl₃) δ 1.06 (t, *J* = 7.3 Hz, 3H), 1.08 (m, 1H), 1.24 (m, 1H), 1.41–1.59 (m, 6H), 1.98–2.02 (m, 2H), 2.63 (m, 1H), 2.91–3.06 (m, 2H), 3.16 (m, 1H), 3.27 (m, 1H), 4.71 (d, *J* = 8.1 Hz, 1H), 7.12 (t, *J* = 7.3 Hz, 1H), 7.32 (t, *J* = 7.3 Hz, 1H), 7.42 (d, *J* = 7.3 Hz, 1H), 7.51 (d, *J* = 7.3 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 15.4, 23.1, 24.8, 25.0, 26.9, 34.8, 41.9, 49.0, 64.4, 68.1, 82.7, 124.9, 127.9, 129.0, 132.7, 134.9, 141.7, 214.1; GC-MS *m/z* 352, 350 (M⁺).

The minor isomer: R_f 0.6 (1:9 EtOAc–hexane); IR (CH₂Cl₂) 1776 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.99–1.18 (m, 3H), 1.04 (t, J = 7.3 Hz, 3H), 1.26 (m, 1H), 1.46–1.75 (m, 5H), 2.00 (m, 1H), 2.23 (m, 1H), 2.92–3.03 (m, 2H), 3.03–3.26 (m, 2H), 4.70 (d, J = 9.7 Hz, 1H), 7.11 (t, J = 8.1 Hz, 1H), 7.31 (t, J = 8.1 Hz, 1H), 7.48 (d, J = 8.1 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 14.9, 18.8, 22.2, 24.9, 25.4, 34.1, 41.9, 49.2, 64.1, 67.8, 81.4, 125.6, 128.4, 129.3, 132.3, 134.9, 140.9, 215.1; GC-MS m/z 352, 350 (M⁺), 322, 324, 181 (100).

General Procedure for Grignard Addition of Spirocyclobutanones.

To a solution of a Grignard reagent (10 eq) in THF was added dropwise at –78 °C a THF solution of cyclobutanone **3** (351 mg, 1 mmol). The reaction mixture was stirred at –78 °C for 2 h, allowed to slowly warm up to room temperature, and stirred for an additional hour. The reaction was quenched with addition of sat NH₄Cl. The resulting mixture was extracted with ether. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure. The concentrate was purified by silica gel column chromatography using 2:98 ethyl acetate–hexane as eluent to afford the respective *tert*-cyclobutanols in 79–90% yield.

4a. Obtained as a white solid. mp 107 °C; R_f 0.7 (1:9 EtOAc–hexane); IR (CH₂Cl₂) 3544 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.11 (m, 1H), 1.23 (t, J = 7.3 Hz, 3H), 1.23 (m, 1H), 1.72–1.84 (m, 5H), 1.98–2.05 (m, 2H), 2.17 (m, 2H), 2.53 (q, J = 10.5 Hz, 1H), 2.99 (m, 1H), 3.13 (m, 1H), 3.44 (s, 1H), 4.78 (d, J = 9.7 Hz, 1H), 5.33 (d, J = 7.3 Hz, 1H), 6.83 (d, J = 7.3 Hz, 1H), 6.92 (t, J = 7.3 Hz, 1H), 7.25–7.40 (m, 5H), 7.65 (d, J = 7.3 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 15.6, 20.2, 21.0, 22.3, 26.3, 27.3, 33.9, 46.6, 52.7, 64.0, 79.1, 79.8, 124.6, 125.9, 127.1, 127.5, 128.0, 128.8, 130.1, 131.7, 143.0, 147.9; GC-MS m/z 384, 382 (M⁺–EtOH), 303, 215 (97), 213 (100).

4b. Obtained as a white solid. mp 114 °C; R_f 0.6 (1:9 EtOAc–hexane); IR (CH₂Cl₂) 3465 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.02 (m, 1H), 1.15 (t, J = 7.2 Hz, 3H), 1.13–1.24 (m, 2H), 1.42 (m, 1H), 1.60 (s, 3H), 1.58–1.75 (m, 5H), 1.80 (t, J = 10.0 Hz, 1H), 1.90 (m, 1H), 2.00 (m, 1H), 2.07 (m, 1H), 2.63 (s, 1H), 3.18 (m, 1H), 3.31 (m, 1H),

5.02 (d, $J = 10.4$ Hz, 1H), 7.13 (t, $J = 8.0$ Hz, 1H), 7.39 (t, $J = 8.0$ Hz, 1H), 7.50 (d, $J = 8.0$ Hz, 1H), 7.56 (d, $J = 8.0$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 15.6, 21.0, 21.6, 22.7, 24.5, 24.8, 25.9, 31.3, 48.2, 50.5, 64.0, 75.3, 79.4, 128.3, 129.3 (2C), 129.5, 132.3, 142.9; GC-MS m/z 215, 213 (30), 183 (57), 125 (100).

4c. Obtained as a white solid. mp 122 °C; R_f 0.7 (1:9 EtOAc–hexane); IR (CH_2Cl_2) 3559 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 0.99 (m, 1H), 1.00 (t, $J = 7.3$ Hz, 3H), 1.10–1.22 (m, 2H), 1.11 (t, $J = 7.3$ Hz, 3H), 1.41 (m, 1H), 1.55–1.90 (m, 9H), 2.00 (m, 1H), 2.25 (m, 1H), 3.17 (m, 1H), 3.29 (m, 1H), 5.02 (d, $J = 9.7$ Hz, 1H), 7.12 (m, 1H), 7.35 (t, $J = 8.1$ Hz, 1H), 7.50 (d, $J = 8.1$ Hz, 1H), 7.56 (d, $J = 8.1$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 7.9, 15.5, 21.0, 22.4, 22.6, 24.5, 24.9, 27.1, 28.5, 47.0, 50.7, 63.7, 79.7, 94.7, 125.6, 128.3, 129.3, 129.6, 132.4, 142.9; GC-MS m/z 215, 213 (21), 183 (54), 139 (100).

11. Obtained as a white solid. mp 140 °C; R_f 0.7 (1:9 EtOAc–hexane); IR (CH_2Cl_2) 3563 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.02 (m, 1H), 1.03 (t, $J = 7.0$ Hz, 3H), 1.13 (t, $J = 7.0$ Hz, 3H), 1.17 (m, 2H), 1.38–1.44 (m, 2H), 1.57–1.73 (m, 6H), 1.78–1.92 (m, 4H), 2.02 (m, 1H), 2.22 (t, $J = 8.6$ Hz, 1H), 2.32 (s, 1H), 3.16 (m, 1H), 3.31 (m, 1H), 5.03 (d, $J = 10.1$ Hz, 1H), 7.13 (t, $J = 7.3$ Hz, 1H), 7.35 (t, $J = 7.3$ Hz, 1H), 7.50 (d, $J = 7.3$ Hz, 1H), 7.57 (d, $J = 7.3$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 15.5 (2C's), 16.9, 21.0, 21.4, 22.6, 24.5, 24.8, 27.8, 38.7, 47.4, 50.8, 63.8, 77.5, 79.4, 125.6, 128.3, 129.3, 129.5, 132.5, 143.0.

12. Obtained as a white solid. mp 167 °C; R_f 0.6 (1:9 EtOAc–hexane); IR (CH_2Cl_2) 3461 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 0.89 (m, 1H), 0.91 (t, $J = 7.3$ Hz, 3H), 1.08 (t, $J = 7.3$ Hz, 3H), 1.13–1.30 (m, 3H), 1.39–1.87 (m, 12H), 2.43 (m, 1H), 3.14 (m, 1H), 3.40 (m, 1H), 4.23 (s, 1H), 5.02 (d, $J = 10.3$ Hz, 1H), 7.13 (t, $J = 7.3$ Hz, 1H), 7.35 (t, $J = 7.3$ Hz, 1H), 7.45–7.52 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 15.0, 15.1, 16.2, 21.2, 22.2, 23.8, 27.6, 28.4, 28.7, 38.7, 45.9, 49.2, 63.7, 79.2, 80.5, 125.5, 128.5, 129.4, 129.5, 132.5, 141.0.

Typical Procedure for Palladium-Catalyzed Annulation of *tert*-Cyclobutanols:

To a mixture of Pd(OAc)₂ (0.7 to 2.0 mg, 5 to 15 mol%), dppp (2.4 to 7.2 mg, 10 to 30 mol%), and Cs₂CO₃ (21 mg, 0.064 mmol) in toluene (2 mL) was added cyclobutanol **4a** (25 mg, 0.058 mmol). The reaction mixture was heated at 90 °C for 12 h, cooled to room temperature, and filtered through a short pad of silica gel. The filtrate was concentrated under reduced pressure. The crude product was purified by silica gel column chromatography using 2:98 ethyl acetate–hexane as eluent to afford **5a** (14 mg, 69%).

5a. Obtained as a colorless oil. *R_f* 0.6 (1:9 EtOAc–hexane); IR (CH₂Cl₂) 1684 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.00–1.25 (m, 3H), 1.29 (t, *J* = 7.2 Hz, 3H), 1.47 (m, 1H), 1.57–1.67 (m, 3H), 1.87 (ddd, *J* = 16.2, 11.4, 4.8 Hz, 1H), 1.95 (ddd, *J* = 16.2, 11.4, 4.8 Hz, 1H), 2.27 (m, 1H), 2.45 (m, 1H), 2.74 (ddd, *J* = 16.2, 11.4, 4.8 Hz, 1H), 3.00 (ddd, *J* = 16.2, 11.4, 4.8 Hz, 1H), 3.61–3.69 (m, 2H), 5.08 (d, *J* = 5.7 Hz, 1H), 7.15 (m, 1H), 7.23–7.26 (m, 2H), 7.39–7.42 (m, 3H), 7.51 (t, *J* = 7.3 Hz, 1H), 7.80 (d, *J* = 7.3 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 15.8, 22.6, 23.1, 24.5, 32.3, 34.1, 35.9, 47.8, 50.0, 65.5, 83.7, 123.0, 125.6, 127.1, 127.5, 128.2, 128.8, 133.1, 137.1, 143.8, 145.9, 200.8; GC-MS *m/z* 348 (M⁺), 319, 302, 213, 171 (100); HRMS calcd for C₂₄H₂₈O₂ (M⁺) 348.2089, found 348.2086.

5b. Obtained as a colorless oil. *R_f* 0.6 (1:9 EtOAc–hexane); IR (CH₂Cl₂) 1715 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 0.97–1.08 (m, 2H), 1.16 (m, 1H), 1.29 (t, *J* = 7.1 Hz, 3H), 1.37 (m, 1H), 1.54–1.60 (m, 3H), 1.70 (ddd, *J* = 13.9, 11.6, 4.9 Hz, 1H), 1.79 (ddd, *J* = 13.9, 11.6, 4.9 Hz, 1H), 2.05 (s, 3H), 2.16–2.22 (m, 2H), 2.39 (m, 1H), 2.48 (ddd, *J* = 15.6, 11.4, 5.0 Hz, 1H), 3.61–3.68 (m, 2H), 5.05 (d, *J* = 5.6 Hz, 1H), 7.07 (m, 1H), 7.21–7.23 (m, 2H), 7.37 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 15.8, 22.5, 23.0, 24.5, 30.2, 32.3, 35.1, 39.2, 47.5, 50.0, 65.5, 83.7, 123.0, 125.5, 127.0, 127.5, 143.8, 145.9, 209.3; HRMS calcd for C₁₉H₂₆O₂ (M⁺) 286.1932, found 286.1932.

5c. Obtained as a colorless oil. *R_f* 0.6 (1:9 EtOAc–hexane); IR (CH₂Cl₂) 1715 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 0.97 (t, *J* = 7.6 Hz, 3H), 1.12–1.19 (m, 1H), 1.06–1.19 (m, 2H), 1.28 (t, *J* = 7.0 Hz, 3H), 1.37 (dt, *J* = 12.7, 4.0 Hz, 1H), 1.53–1.60 (m, 3H), 1.70 (ddd, *J* = 16.7, 11.6, 5.0 Hz, 1H), 1.78 (ddd, *J* = 16.7, 11.6, 5.0 Hz, 1H), 2.13–2.17 (m,

2H), 2.32 (m, 2H), 2.38 (m, 1H), 2.45 (ddd, $J = 16.7, 11.6, 5.0$ Hz, 1H), 3.61–3.68 (m, 2H), 5.05 (d, $J = 5.6$ Hz, 1H), 7.07 (m, 1H), 7.21–7.26 (m, 2H), 7.37 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 8.1, 15.8, 22.5, 23.0, 24.5, 32.3, 35.2, 36.2, 37.8, 47.5, 50.0, 65.5, 83.7, 123.0, 125.5, 127.0, 127.4, 143.8, 145.9, 212.0; GC-MS m/z 300 (M^+), 271, 254, 213 (100); HRMS calcd for $\text{C}_{20}\text{H}_{28}\text{O}_2$ (M^+) 300.2089, found 300.2091.

13. Obtained as a colorless oil. R_f 0.6 (1:9 EtOAc–hexane); IR (CH_2Cl_2) 1712 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 0.85 (t, $J = 7.1$ Hz, 3H), 1.03 (m, 1H), 1.28 (t, $J = 7.1$ Hz, 3H), 1.29–1.40 (m, 2H), 1.49–1.71 (m, 5H), 1.76–1.80 (m, 2H), 2.12–2.18 (m, 2H), 2.40 (m, 2H), 2.36–2.47 (m, 3H), 3.61–3.68 (m, 2H), 5.05 (d, $J = 5.6$ Hz, 1H), 7.06 (m, 1H), 7.21–7.25 (m, 2H), 7.37 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 14.0, 15.5, 15.8, 23.0, 23.0, 24.5, 32.3, 35.1, 38.2, 38.5, 47.5, 49.9, 65.5, 83.7, 123.0, 125.5, 127.0, 128.5, 143.0, 143.8, 211.5; GC-MS m/z 314 (M^+), 266, 211, 180, 167 (100); HRMS calcd for $\text{C}_{21}\text{H}_{30}\text{O}_2\text{Na}$ (M^+) 337.2138, found 337.2152.

Reduction product 14. Obtained as a colorless oil. R_f 0.6 (1:9 EtOAc–hexane); IR (CH_2Cl_2) 3446 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 0.83 (m, 1H), 0.91 (t, $J = 7.1$ Hz, 3H), 1.08 (t, $J = 7.1$ Hz, 3H), 1.11–1.29 (m, 4H), 1.39–1.87 (m, 11H), 2.32 (br d, $J = 10.2$ Hz, 1H), 3.18 (m, 1H), 3.25 (m, 1H), 4.29 (s, 1H), 4.39 (d, $J = 10.2$ Hz, 1H), 7.27–7.34 (m, 5H); ^{13}C NMR (125 MHz, CDCl_3) δ 15.0, 15.1, 16.3, 20.5, 22.2, 23.7, 27.4, 28.4, 28.7, 38.7, 45.6, 49.0, 64.0, 80.4, 81.8, 127.7, 127.9, 128.6, 141.5; GC-MS m/z 298 ($\text{M}^+ - \text{H}_2\text{O}$), 270, 252, 209, 184 (100).

15. IR (CH_2Cl_2) 3467 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 0.78 (m, 1H), 0.90 (t, $J = 7.1$ Hz, 3H), 1.10 (m, 1H), 1.18 (t, $J = 7.1$ Hz, 3H), 1.20–1.26 (m, 7H), 1.40–1.74 (m, 7H), 2.28 (s, 1H), 2.40 (s, 3H), 3.49 (m, 2H), 4.28 (s, 1H), 4.70 (d, $J = 10.6$ Hz, 1H), 7.08–7.40 (m, 7H), 7.62 (d, $J = 7.1$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 15.0, 15.4, 16.2, 20.8, 22.1, 22.9, 27.4, 28.3, 28.7, 29.9, 38.7, 48.2, 49.4, 63.9, 76.4, 80.6, 127.7, 127.3, 127.8, 128.2, 128.3, 129.0, 129.8, 129.7, 130.1, 136.8, 140.3, 142.8; MS (ES) m/z 445.20 ($\text{M}^+ + \text{K}$), 429.26 ($\text{M}^+ + \text{Na}$), 407.27 ($\text{M}^+ + \text{H}$).

18. IR (CH₂Cl₂) 3510, 1758 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.19–1.36 (m, 4H), 1.49–1.65 (m, 4H), 1.93 (m, 1H), 2.02 (m, 1H), 2.15 (m, 1H), 2.93 (m, 1H), 3.16 (m, 1H), 3.68 (br s, 1H), 3.96 (d, *J* = 5.1 Hz, 1H), 4.73 (d, *J* = 5.1 Hz, 1H), 4.77 (d, *J* = 8.1 Hz, 1H), 6.84 (d, *J* = 7.1 Hz, 1H), 7.06–7.31 (m, 12H), 7.42 (m, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 14.4, 22.5, 23.5, 24.2, 25.6, 31.3, 42.3, 49.6, 69.6, 77.9, 83.7, 125.0, 126.6, 127.2, 127.9, 128.0, 128.4, 128.6, 128.9, 129.1, 129.3, 132.8, 137.5, 140.2, 141.8, 214.4.

19. IR (CH₂Cl₂) 3446, 1766 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.08–1.11 (m, 3H), 1.34 (m, 1H), 1.52–1.65 (m, 3H), 1.84 (m, 2H), 2.26 (m, 1H), 2.53 (m, 1H), 2.91 (d, *J* = 2.0 Hz, 1H), 3.17 (m, 1H), 3.48 (m, 1H), 4.17 (d, *J* = 8.1 Hz, 1H), 4.75 (dd, *J* = 8.1, 2.0 Hz, 1H), 5.22 (d, *J* = 10.1 Hz, 1H), 6.55 (d, *J* = 7.1 Hz, 2H), 6.82–6.95 (m, 7H), 7.07 (m, 3H), 7.14 (dd, *J* = 7.6, 1.5 Hz, 1H), 7.24 (m, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 18.9, 22.1, 24.6, 25.4, 34.4, 42.3, 49.2, 67.7, 79.5, 82.8, 87.3, 125.4, 127.2, 127.3, 127.4, 127.5, 127.7, 128.1, 128.2, 128.9, 130.4, 131.8, 138.1, 140.4, 140.8, 218.3.

20. IR (CH₂Cl₂) 3485, 1765 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 1.30 (m, 1H), 1.46–1.55 (m, 2H), 1.69 (m, 1H), 1.79 (m, 1H), 1.93 (m, 1H), 2.14 (m, 1H), 2.22 (m, 1H), 2.44 (q, *J* = 8.6 Hz, 1H), 3.02 (m, 1H), 3.11 (m, 1H), 3.64 (d, *J* = 2.0 Hz, 1H), 4.00 (d, *J* = 7.1 Hz, 1H), 4.85 (m, 2H), 6.93–6.98 (m, 4H), 7.10–7.24 (m, 7H), 7.35–7.45 (m, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 23.9, 26.4, 29.6, 31.2, 38.7, 42.9, 56.2, 71.8, 78.2, 83.9, 127.5, 127.6, 127.8, 128.0, 128.2, 128.4, 128.7, 128.9, 129.2, 133.0, 133.1, 136.0, 139.5, 140.0, 215.2; MS (M⁺ – PhCHOH) 397, 399.

21. IR (CH₂Cl₂) 3564, 3461, 1770 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.10 (m, 1H), 1.31 (m, 1H), 1.50–1.71 (m, 2H), 1.86 (m, 1H), 1.96 (m, 1H), 2.06 (m, 1H), 2.62–2.70

(m, 2H), 3.01 (d, $J = 1.6$ Hz, 1H), 3.22 (m, 1H), 3.39 (m, 1H), 4.26 (d, $J = 8.1$ Hz, 1H), 4.72 (dd, $J = 8.1, 1.6$ Hz, 1H), 5.14 (d, $J = 8.9$ Hz, 1H), 6.64 (d, $J = 7.3$ Hz, 2H), 6.80–7.02 (m, 7H), 7.07–7.23 (m, 5H); ^{13}C NMR (100 MHz, CDCl_3) δ 21.4, 22.4, 26.9, 38.5, 43.7, 53.4, 72.5, 79.0, 81.6, 88.0, 127.0, 127.1, 127.8, 127.2, 127.3, 127.7, 127.9, 128.4, 128.7, 130.0, 131.9, 137.5, 139.9, 140.9, 217.8.

22. ^1H NMR (500 MHz, CDCl_3) δ 0.91–1.08 (m, 3H), 1.23 (m, 1H), 1.38–1.52 (m, 3H), 1.61 (s, 3H), 1.65–1.81 (m, 5H), 2.08 (m, 1H), 3.72 (br s, 1H), 4.38 (d, $J = 3.0$ Hz, 1H), 4.71–4.74 (m, 2H), 6.88 (dd, $J = 7.6, 1.5$ Hz, 1H), 6.98–7.06 (m, 2H), 7.18–7.24 (m, 5H), 7.33–7.42 (m, 4H), 7.48 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 21.0, 21.4, 22.6, 24.2, 24.8, 25.6, 32.8, 48.8, 51.4, 75.7, 76.4, 77.1, 83.4, 125.2, 126.3, 127.2, 128.0, 128.1, 128.4, 128.6, 128.9, 129.5, 130.0, 132.2, 138.4, 141.8, 142.4.

23. IR (CH_2Cl_2) 3543, 1715 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.30 (m, 1H), 1.42–1.81 (m, 7H), 1.95–2.10 (m, 4H), 1.97 (s, 3H), 2.25 (m, 1H), 3.47 (s, 1H), 4.67 (d, $J = 8.1$ Hz, 1H), 4.72 (d, $J = 8.1$ Hz, 1H), 4.94 (d, $J = 6.1$ Hz, 1H), 7.02–7.29 (m, 13H), 7.52 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 22.2, 23.5, 24.5, 30.2, 32.7, 34.5, 39.1, 47.2, 48.8, 78.9, 82.4, 87.7, 123.3, 125.3, 127.0, 127.5, 128.1, 128.2, 128.3, 128.4, 128.5, 128.6, 129.0, 138.3, 139.4, 143.2, 209.1.

24. $[\alpha]_{\text{D}} -31$ (c 0.002, CH_2Cl_2); IR (CH_2Cl_2) 1715 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.19–1.28 (m, 2H), 1.30–1.37 (m, 1H), 1.39–1.53 (m, 3H), 1.61–1.68 (m, 1H), 1.75 (ddd, $J = 5.1, 11.6, 14.2$ Hz, 1H), 1.78–1.83 (m, 1H), 1.98 (ddd, $J = 5.1, 11.6, 14.2$ Hz, 1H), 2.06 (s, 3H), 2.14 (m, 1H), 2.23 (ddd, $J = 5.1, 11.6, 17.2$ Hz, 1H), 2.37 (ddd, $J = 5.1, 11.6, 17.2$ Hz, 1H), 2.57 (dd, $J = 5.6, 15.7$ Hz, 1H), 2.98 (dd, $J = 7.1, 15.7$ Hz, 1H), 7.03 (m, 1H), 7.12–7.17 (m, 2H), 7.22 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 22.2, 23.3, 27.7, 30.2, 31.5, 33.8, 36.3, 39.4, 43.9, 48.6, 122.7, 125.9, 126.1, 126.5, 143.4, 149.1,

209.6; MS (GC-MS) m/z 242 (1), 171 (100), 129 (90); HRMS calcd for $C_{17}H_{22}O$ (M^+) 242.1671, found 242.1672.

25. 1H NMR (500 MHz, $CDCl_3$) δ 1.07–1.35 (m, 4H), 1.64 (s, 3H), 1.54–1.71 (m, 2H), 1.77–1.86 (m, 2H), 1.95 (m, 1H), 2.05 (m, 1H), 3.57 (s, 1H), 3.63 (d, $J = 4.9$ Hz, 1H), 4.07 (d, $J = 3.2$ Hz, 1H), 4.31 (d, $J = 9.7$ Hz, 1H), 4.72 (m, 1H), 6.89 (m, 1H), 7.08 (m, 2H), 7.22–7.45 (m, 11H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 19.7, 24.6, 26.7, 27.8, 28.3, 33.0, 53.7, 59.6, 74.0, 77.1, 83.0, 127.1, 127.2, 127.1, 127.9, 128.1, 128.4, 128.5, 128.9, 129.0, 133.0, 132.3, 138.1, 141.4, 141.5.

26. IR (CH_2Cl_2) 3446, 1714 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 1.30–1.38 (m, 1H), 1.51–1.58 (m, 3H), 1.66–1.88 (m, 4H), 1.97 (s, 3H), 1.94–2.24 (m, 3H), 3.68 (d, $J = 1.5$ Hz, 1H), 4.72 (d, $J = 8.1$ Hz, 1H), 4.80 (dd, $J = 8.1, 1.5$ Hz, 1H), 4.97 (d, $J = 7.6$ Hz, 1H), 7.09–7.12 (m, 5H), 7.19–7.30 (m, 8H), 7.52 (m, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 26.1, 27.3, 30.0, 34.3, 40.5, 40.6, 53.8, 58.5, 78.9, 82.5, 89.7, 124.0, 124.3, 127.4, 127.5, 127.9, 128.1, 128.2, 128.3, 128.4, 128.8, 138.5, 139.4, 143.8, 147.8, 209.0.

27. $[\alpha]_D -52$ (c 0.4, CH_2Cl_2); IR (CH_2Cl_2) 1716 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 1.31–1.43 (m, 2H), 1.54–1.61 (m, 1H), 1.72–1.77 (m, 1H), 1.83–1.97 (m, 3H), 2.03 (s, 3H), 2.00–2.15 (m, 2H), 2.32 (ddd, $J = 5.0, 11.7, 16.7$ Hz, 1H), 2.42 (m, 1H), 2.63 (dd, $J = 2.5, 16.7$ Hz, 1H), 3.18 (dd, $J = 9.1, 16.7$ Hz, 1H), 7.07 (d, $J = 7.1$ Hz, 1H), 7.13–7.19 (m, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 26.0, 30.2, 35.3, 35.9, 39.4, 40.6, 41.2, 47.5, 60.1, 123.8, 124.7, 126.8, 126.9, 143.7, 149.5, 209.6; MS (GC-MS) m/z 228 (5), 170 (12), 157 (100); HRMS calcd for $C_{16}H_{20}O$ (M^+) 228.1514, found 228.1520.