Functional Group Scope in the Methylene-Free, Tandem Enyne Metathesis

Brian Peppers, Amol A. Kulkarni and Steven T. Diver*

Department of Chemistry, University at Buffalo, the State University of New York, Amherst, NY, 14260-3000

SUPPORTING INFORMATION

Table of Contents

A. General Information.....	S1
B. General Procedure (Terminal Alkynes).....	S2
C. General Procedure (Internal Alkynes).....	S5
D. Cycloaddition Reactions.....	S8
E. ¹H and ¹³C NMR Spectra for New Compounds.....	S10

General Information

All reactions were conducted under an argon atmosphere unless otherwise noted. Dichloromethane (CH₂Cl₂) (alumina), benzene and toluene (alumina + Q5) were drawn from a solvent purifier (alumina) immediately prior to use. Reagent grade methanol was used for trituration and cycloaddition reaction. Ruthenium [1,3-bis-(2,4,6-trimethylphenyl)-2-imidazolidinylidene]dichloro(phenylmethylene)(tricyclohexylphosphine) (Grubbs' second generation catalyst) was obtained from Materia Inc. (Pasadena, CA). 1,5-Cyclooctadiene (COD) was distilled from sodium prior to use. All-*cis* polybutadiene was purchased from Aldrich (18137-4). Terminal alkynes **5**,¹ **7B**,² **7C**,³ **7D**,⁴ **7E**,⁵ **7F**,⁶ **7G**⁷ and internal alkynes **9C**,⁸ **9G**⁹ were synthesized according

¹ Middleton, M. D.; Diver, S. T., *Tetrahedron Lett.* **2005**, *46*, 4039-4043.

² Rodriguez, J. R.; Castedo, L.; Mascarenas, J. L., *J. Org. Chem.* **2000**, *65*, 2528-2531.

³ Posner, G. H.; Weitzberg, M.; Hamill, T. G.; Asirvatham, E.; He, C. H.; Clardy, J., *Tetrahedron* **1986**, *42*, 2919-29.

⁴ Qin, D.-G.; Zha, H.-Y.; Yao, Z.-J., *J. Org. Chem.* **2002**, *67*, 1038-1040.

⁵ Rodriguez-Conesa, S.; Candal, P.; Jimenez, C.; Rodriguez, J., *Tetrahedron Lett.* **2001**, *42*, 6699-6702.

⁶ Short, K. M.; Ziegler, C. B., Jr., *Tetrahedron Lett.* **1995**, *36*, 355-6.

⁷ Luo, F. T.; Wang, R. T., *Tetrahedron Lett.* **1992**, *33*, 6835-8.

to literature procedures. Alkynes **7A** and **9H** are commercially available from Aldrich. Column chromatography was carried out on Merck silica gel 60 (230 – 400 mesh). ¹H-NMR spectra were recorded at 500 MHz and ¹³C-NMR spectra at 125 MHz in CDCl₃. ¹H-NMR spectra were referenced on the TMS signal for CDCl₃. The ¹³C-NMR spectra were referenced at 77.0 ppm for CDCl₃. The syringe pump (kd scientific, model 210) was used for the tandem enyne metathesis studies using a 5 mL glass gas tight syringe.

General Procedure (Terminal Alkynes)

To an oven-dried 50 mL schlenk tube, cooled under argon, was added 1,5-cyclooctadiene (6 or 9 equiv) and 5 mL of solvent (an argon cooled cold-finger was included for all reactions performed at refluxing conditions). A rubber septum was placed on the side arm of the schlenk tube. This solution was purged with a steady stream of argon for 5 minutes via a stainless steel needle. Upon completion of argon purge, the needle was removed and a teflon cannula attached to a 5 mL Hamilton gas-tight syringe was passed through the septum and lowered until it reached the point where the side arm and the main body of the schlenk tube meet. The catalyst was added all at once and allowed to stir for 1 min. The tube was then immersed in an oil bath at the specified temperature. The teflon cannula was then lowered further until resting about 2-3 cm above the solution line. The addition of 0.250 mmole of alkyne was then started via syringe pump at a constant rate for 4 h. At the end of this period, the syringe was manually plunged to dispense the solution in the teflon cannula and the reaction was stirred for an additional hour. The reaction was then transferred to a 25 mL rb flask and concentrated *in vacuo* (rotary evaporator). A stir bar was added and the vessel was placed in an ice bath and triturated with 5 mL of ice-cold methanol. The trituration was performed three times. The methanol fractions were pooled and concentrated *in vacuo* (rotary evaporator). The resulting residue was further purified by column chromatography on silica gel, eluting with TLC solvent, as indicated for individual compounds. Depending of the efficiency of trituration and polarity of individual substrates, a second column was needed to completely remove all traces of 1,5-cyclooctadiene-derived oligomer byproducts.

Dihydrophenylalanine **6**. Solvent= dichloromethane; reaction temperature= 40 °C; 7.5 mol % complex **1**; 2.30 mmol 1,5-COD (9 equiv). According to the general procedure (0.256 mmol scale) to obtain 88 mg **6** as a colorless oil (80% yield). ¹H NMR (500 MHz, CDCl₃): δ 7.76 (d, J = 7.5 Hz, 2H), 7.60, (m, 2H), 7.40 (t, J = 7.0 Hz, 2H), 7.31 (t, J = 7.5 Hz, 2H), 5.83 (m, 2H), 5.55 (s, 1H), 5.27 (d, J = 8.0

⁸ D'Accolti, L.; Fiorentino, M.; Fusco, C.; Crupi, P.; Curci, R., *Tetrahedron Lett.* **2004**, *45*, 8575-8578.

⁹ Henry, J. R.; Marcin, L. R.; McIntosh, M. C.; Scola, P. M.; Harris, G. D., Jr.; Weinreb, S. M., *Tetrahedron Lett.* **1989**, *30*, 5709-12.

Hz, 1H), 4.41 (m, 1H), 4.34 (m, 2H), 4.23 (m, 1H), 2.48 (m, 2H), 2.10 (m, 4H), 1.47 (s, 9H); ^{13}C NMR (125 MHz, CDCl_3): δ 171.1, 155.5, 143.9, 141.3, 130.9, 127.7, 127.3, 127.0, 126.6, 125.2, 124.4, 120.0, 82.1, 66.1, 53.6, 47.2, 38.5, 28.0, 22.3, 22.1; FT-IR (thin film, cm^{-1}) 3326, 2933, 1718, 1507, 1450, 1367, 1224, 1155, 1050; high resolution ESI molecular ion calcd for $\text{C}_{28}\text{H}_{31}\text{O}_4\text{N} + \text{Na}$ 468.2145 found 468.2152; $[\alpha]_D^{25} +10.0$ ($c = 1.6$, CHCl_3).

Table 2, entry 1. Solvent= dichloromethane; reaction temperature= 40 °C; 5 mol % complex **1**; 1.64 mmol 1,5-COD (6 equiv). According to the general procedure (0.274 mmol scale) to obtain 54 mg **8A** as a colorless oil (82% yield). $R_f = 0.4$ (1:3 ethyl acetate-hexanes). ^1H NMR (500 MHz, CDCl_3) δ 5.85 (s, 1H), 5.77 (dd, $J = 1.5, 10.0$ Hz, 1H), 5.55 (s, 1H), 3.73 (s, 6H), 3.52 (t, $J = 8.0$ Hz, 1H), 2.63 (d, $J = 8.0$ Hz, 2H), 2.08 (m, 4H); ^{13}C NMR (125 MHz, CDCl_3) δ 169.4, 131.9, 127.6, 126.0, 122.8, 52.4, 51.2, 34.6, 22.2, 22.0; FT-IR (thin film, cm^{-1}) 3033, 2952, 2816, 1762, 1742, 1450, 1339, 1278, 1237, 1157 1031; high resolution EI molecular ion calcd for 224.1043 $\text{C}_{12}\text{H}_{16}\text{O}_4$ found 224.1044.

Table 2, entry 2. Solvent= benzene; reaction temperature= 65 °C; 10 mol % complex **1**; 2.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 73 mg **8B** as a colorless oil (75% yield). $R_f = 0.23$ (1:3 ethyl acetate-hexanes). ^1H NMR (400 MHz, CDCl_3) δ 7.93 (d, $J = 7.6$ Hz, 4H), 7.69 (t, $J = 7.2$ Hz, 2H), 7.56 (t, $J = 7.6$ Hz, 4H), 5.77 (m, 1H), 5.59 (d, $J = 10.0$ Hz, 1H), 5.28 (s, 1H), 4.53 (t, $J = 5.2$ Hz, 1H), 2.93 (d, $J = 5.2$ Hz, 2H), 1.92 (s, 4H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.3, 134.4, 129.6, 129.5, 129.0, 128.6, 124.4, 124.2 82.2, 31.2, 22.1, 21.6; FT-IR (thin film, cm^{-1}) 3072, 3033, 2931, 2886, 2835, 1451, 1324, 1310, 1175, 1156, 1082; high resolution EI molecular ion calcd for 388.0798 $\text{C}_{20}\text{H}_{20}\text{O}_4\text{S}_2$ found 388.0801

Table 2, entry 3. Solvent= dichloromethane; reaction temperature= 40 °C; 7.5 mol % complex **1**; 1.50 mmol 1,5-COD (6 equiv). According to the general procedure (0.250 mmol scale) to obtain 43 mg **8C** as a colorless oil (72% yield). $R_f = 0.15$ (1:50 ethyl acetate-hexanes). ^1H NMR (500 MHz, CDCl_3) δ 5.82 (m, 2H), 5.52 (s, 1H), 3.64 (t, $J = 7.0$ Hz, 2H), 2.24 (t, $J = 7.0$ Hz, 2H), 2.09 (m, 4H), 0.89 (s, 9H), 0.043 (s, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 132.9, 127.4, 126.5, 122.1, 62.9, 39.2, 26.0, 22.4, 22.2, 18.4, -5.3; FT-IR (thin film, cm^{-1}) 2927, 2851, 1470, 1258, 1106, 845; high resolution EI molecular ion calcd for 238.1747 $\text{C}_{14}\text{H}_{26}\text{O}_1\text{Si}_1$ found 238.1748

Table 2, entry 4. Solvent= dichloromethane; reaction temperature= 40 °C; 7.5 mol % complex **1**; 2.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 37.5 mg **8D** as a colorless oil (70% yield). $R_f = 0.19$ (1:25 ethyl acetate-hexanes). ^1H NMR (500 MHz, CDCl_3) δ 7.33 (m, 4H), 7.27 (m, 1H), 5.83 (m, 2H), 5.56 (s, 1H), 4.52 (s, 2H), 3.53 (t, $J = 7.0$ Hz, 2H), 2.35 (t, $J = 7.0$ Hz, 2H), 2.10 (m, 4H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.6, 132.7, 128.3,

127.6, 127.5, 127.1, 126.7, 121.9, 72.9, 69.5, 35.9, 35.9, 22.4, 22.2; FT-IR (thin film, cm^{-1}) 3036, 2931, 2862, 1109, 737; high resolution EI molecular ion calcd for $214.1352 \text{ C}_{15}\text{H}_{18}\text{O}$ found 214.1355

Table 2, entry 5. Solvent= dichloromethane; reaction temperature= 40°C ; 5.0 mol % complex **1**; 1.50 mmol 1,5-COD (6 equiv). According to the general procedure (0.250 mmol scale) to obtain 50.0 mg **8E** as a colorless oil (72% yield). $R_f = 0.19$ (1:25 ethyl acetate-hexanes). ^1H NMR (500 MHz, CDCl_3) δ 7.78 (d, $J = 8.0$ Hz, 2H), 7.34 (d, $J = 8.0$ Hz, 2H), 5.80 (m, 1H), 5.68 (d, $J = 8.8$ Hz, 1H), 5.49 (s, 1H), 4.04 (t, $J = 6.8$ Hz, 2H), 2.45 (s, 3H), 2.36 (t, $J = 6.8$ Hz, 2H), 2.06 (m, 4H); ^{13}C NMR (125 MHz, CDCl_3) δ 144.6, 133.2, 130.4, 129.7, 127.8, 127.4, 126.1, 123.3, 69.3, 34.9, 22.2, 21.9, 21.6; FT-IR (thin film, cm^{-1}) 2920, 1603, 1365, 1175, 966; high resolution ESI molecular ion calcd for $301.0869 \text{ C}_{15}\text{H}_{18}\text{O}_3\text{NaS}$ found 301.0862

Table 2, entry 6. Solvent= dichloromethane; reaction temperature= 40°C ; 5.00 mol % complex **1**; 2.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 43.3 mg **8F** as a colorless oil (73% yield). $R_f = 0.15$ (1:5 ethyl acetate-hexanes). ^1H NMR (500 MHz, CDCl_3) δ 7.73 (d, $J = 10$ Hz, 2H), 7.30 (d, $J = 10.0$ Hz, 2H), 5.80 (m, 1H), 5.63 (dd, $J = 1.5, 10.0$ Hz, 1H), 5.45 (s, 1H), 4.46 (t, $J = 6.0$ Hz, 1H), 3.00 (q, $J = 6.5$ Hz, 2H), 2.43 (s, 3H), 2.14 (t, $J = 6.5$, 2H), 2.07 (m, 4H); ^{13}C NMR (125 MHz, CDCl_3) δ 143.3, 137.0, 131.6, 129.6, 128.0, 127.1, 125.6, 123.4, 41.3, 35.1, 22.2, 22.1, 21.5; FT-IR (thin film, cm^{-1}) 3282, 2940, 2874, 2822, 1427, 1326, 1159, 1092; high resolution ESI molecular ion calcd for $300.1029 \text{ C}_{15}\text{H}_{19}\text{O}_2\text{NNaS}$ found 300.1032

Table 2, entry 7. Solvent= dichloromethane; reaction temperature= 40°C ; 5.0 mol % complex **1**; 2.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 54.0 mg **8G** as a colorless oil (74% yield). $R_f = 0.19$ (1:5 ethyl acetate-hexanes). ^1H NMR (500 MHz, CDCl_3) δ 7.75 (d, $J = 8.5$ Hz, 2H), 7.30 (d, $J = 8.0$ Hz, 2H), 5.81 (m, 1H), 5.69 (d, $J = 8.5$ Hz, 1H), 5.39 (s, 1H), 4.60 (t, $J = 6.5$ Hz, 1H), 2.92 (q, $J = 6.5$ Hz, 2H), 2.43 (s, 3H), 2.05 (s, 4H), 1.99 (t, $J = 7.0$ Hz, 2H), 1.56 (dt, $J = 7.0, 14.5$ Hz, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 143.3, 137.0, 134.3, 129.6, 127.3, 127.1, 126.5, 121.0, 42.6, 32.4, 27.9, 22.3, 22.2, 21.5; FT-IR (thin film, cm^{-1}) 3280, 3039, 2934, 2834, 2828, 1429, 1325, 1159, 1092; high resolution ESI molecular ion calcd for $292.1366 \text{ C}_{16}\text{H}_{22}\text{O}_2\text{NS}$ found 292.1369.

Alkyne **7H** (substrate for table 1, entry 8). To a solution of 2-hydroxypent-4-ynoic acid¹⁰ (500 mg, 3.9 mmol) and *tert*-butyldimethylsilylchloride (647 mg,

¹⁰ Dragovich, P. S.; Prins, T. J.; Zhou, R.; Johnson, T. O.; Hua, Y.; Luu, H. T.; Sakata, S. K.; Brown, E. L.; Maldonado, F. C.; Tuntland, T.; Lee, C. A.; Fuhrman, S. A.; Zalman, L. S.; Patick, A. K.; Matthews, D. A.; Wu, E. Y.; Guo, M.; Borer, B. C.; Nayyar, N. K.; Moran, T.; Chen, L.; Rejto, P. A.; Rose, P. W.; Guzman, M. C.; Doval Santos, E. Z.; Lee, S.; McGee, K.; Mohajeri, M.; Liese,

4.29 mmol) in dichloromethane (16 mL) was added imidazole (664 mg, 9.75 mmol). After 12 hours the reaction was washed with 20 mL of saturated ammonium chloride; extracted with dichloromethane (2 x 20 mL). The organic layers were combined and washed with 20 mL of brine and dried over anhydrous sodium sulfate. After filtration and concentration *in vacuo* (rotary evaporator) the residue was further purified by column chromatography on silica gel, eluting with 1:20 ethyl acetate-hexanes to give 745 mg of **7H** (79% yield). R_f = 0.18 (1:25 ethyl acetate-hexanes). ^1H NMR (400 MHz, CDCl_3) δ 4.03 (dd, J = 5.0, 2.4 Hz, 1H), 3.71 (s, 3H), 2.62 (ddd, J = Hz, 1H), 2.62 (m, 1H), 2.52 (m, 1H), 1.98 (t, J = 2.8 Hz, 1H), 0.87 (s, 9H), 0.07 (d, J = 2.0 Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 172.3, 79.9, 71.1, 70.5, 52.0, 25.6, 25.5, 18.3, -5.1, -5.3; FT-IR (thin film, cm^{-1}) 3321, 3296, 2963, 2931, 2854, 1760, 1472, 1267, 1146, 941, 839, 788; high resolution ESI molecular ion calcd for $265.1230 \text{ C}_{12}\text{H}_{22}\text{O}_3\text{NaSi}$ found 265.1227.

Table 2, entry 8. Solvent= dichloromethane; reaction temperature= 40 °C; 10.0 mol % complex **1**; 2.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 44.3 mg **8H** as a colorless oil (60% yield). R_f = 0.25 (1:25 ethyl acetate-hexanes). ^1H NMR (400 MHz, CDCl_3) δ 5.81 (m, 2H), 5.54 (s, 1H), 4.19 (dd, J = 4.0, 8.8 Hz 1H), 3.69 (s, 3H) 2.41 (dd, J = 3.2, 13.2 Hz, 1H), 2.30 (dd, J = 8.8, 13.6 Hz, 1H), 2.07 (m, 4H), 0.85 (s, 9H), 0.00 (d, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 173.9, 131.4, 127.1, 126.7, 124.1, 51.7, 41.7, 25.7, 22.4, 22.1, 18.3, -5.2, -5.3; FT-IR (thin film, cm^{-1}) 2931, 2863, 1762, 1738, 1439, 1259, 1141, 842, 780; high resolution ESI molecular ion calcd for $319.1700 \text{ C}_{16}\text{H}_{28}\text{O}_3\text{NaSi}$ found 319.1698

General Procedure (Internal Alkynes)

To an oven-dried 50 mL schlenk tube, cooled under argon, was added 1,5-cyclooctadiene (9 equiv) and 5 mL of solvent (an argon cooled cold-finger was included for all the reactions performed). A rubber septum was placed on the side arm of the schlenk tube. This solution was purged with a steady stream of argon for 5 minutes via a stainless steel needle. Upon completion of argon purge, the needle was removed and a teflon cannula attached to a 5 mL Hamilton gas-tight syringe was passed through the septum and lowered until it reached the point where the side arm and the main body of the schlenk tube meet. The catalyst was added all at once and allowed to stir for 1 min. The tube was then immersed in an oil bath at the specified temperature. The Teflon cannula was then lowered further until resting about 2-3 cm above the solution line. The addition of 0.250 mmole of alkyne was then started via syringe pump at a constant rate for 4 h. At the end of this period, the syringe was manually plunged to dispense the solution in the teflon cannula and the reaction was stirred for an additional hour. The reaction was then transferred to a 25 mL rb flask and concentrated *in vacuo* (rotary evaporator). A stir bar was added and

A.; Tao, J.; Kosa, M. B.; Liu, B.; Batugo, M. R.; Gleeson, J.-P. R.; Wu, Z. P.; Liu, J.; Meador, J. W., III; Ferre, R. A., *J. Med. Chem.* **2003**, *46*, 4572-4585.

the vessel was placed in an ice bath and triturated with 10 mL of ice-cold methanol. The trituration was performed three times. The methanol fractions were pooled and concentrated *in vacuo* (rotary evaporator). The resulting residue was further purified by column chromatography on silica gel, eluting with TLC solvent, as indicated for individual compounds. Depending of the efficiency of trituration and polarity of individual substrates, a second column was needed to completely remove all traces of 1,5-cyclooctadiene-derived oligomer byproducts.

Table 3, entry 1. Solvent= dichloromethane; reaction temperature= 40 °C; 5 mol % complex **1**; 2.25 mmol 1,5-COD (9 equiv). According to the general procedure to obtain 40 mg **10A** as a pale yellow oil (65% yield). R_f 0.31 (10 % ethyl acetate in hexanes). $^1\text{H-NMR}$ (500 MHz, CDCl_3 , ppm) δ 8.05 (d, J = 8.0 Hz, 2H), 7.55 (t, J = 8.0 Hz, 1H), 7.44 (t, J = 8.0 Hz, 2H), 6.02 (s, 1H), 5.64 (s, 1H), 4.88 (s, 2H), 2.20 (q, J = 7.5 Hz, 2H), 2.16 – 2.10 (m, 4H), 1.07 (t, J = 7.5 Hz, 3H); $^{13}\text{C-NMR}$ (125 MHz, CDCl_3 , ppm) δ 166.4, 137.1, 132.9, 132.7, 130.4, 129.6, 128.3, 128.1, 121.2, 65.9, 24.7, 22.6, 22.2, 13.0; FT-IR (thin film, cm^{-1}) 2934, 1718, 1269, 1109, 710; High-resolution MS (EI^+ , m/z) molecular ion calcd for $\text{C}_{16}\text{H}_{18}\text{O}_2$ 242.1301, found 242.1310, error 3.7 ppm.

Table 3, entry 2. **10B** was synthesized according to the general procedure (0.250 mmol scale) to obtain 36 mg **10B** as a pale yellow oil (59% yield). R_f = 0.32 (10 % ethyl acetate in hexanes). $^1\text{H-NMR}$ (500 MHz, CDCl_3 , ppm) δ 8.04 (d, J = 7.5 Hz, 2H), 7.55 (t, J = 7.5 Hz, 1H), 7.43 (t, J = 7.5 Hz, 2H), 5.69 (s, 1H), 5.62 (s, 1H), 4.36 (t, J = 7.5 Hz, 2H), 2.57 (t, J = 7.5 Hz, 2H), 2.06 – 2.03 (m, 4H), 1.85 (s, 3H); $^{13}\text{C-NMR}$ (125 MHz, CDCl_3 , ppm) δ 166.5, 133.6, 132.8, 132.5, 130.4, 129.5, 128.3, 124.0, 123.2, 64.4, 32.1, 22.8, 22.6, 19.6; FT-IR (thin film, cm^{-1}) 2935, 1719, 1272, 1112, 1027, 710; High-resolution MS (EI^+ , m/z) molecular ion calcd for $\text{C}_{16}\text{H}_{18}\text{O}_2$ 242.1301, found 242.1308, error 3.0 ppm.

Table 3, entry 3. Solvent= dichloromethane; reaction temperature= 40 °C; 5 mol % complex **1**; 9x0.50 mmol 1,5-COD (9 equiv). According to the general procedure (0.50 mmol scale) to obtain 71 mg **10C** as a pale yellow oil (63% yield). R_f = 0.34 (25 % ethyl acetate in hexanes). $^1\text{H-NMR}$ (500 MHz, CDCl_3 , ppm) δ 5.89 (s, 2H), 4.55 (s, 4H), 2.11 (d, J = 1.5 Hz, 4H), 1.98 (s, 6H); $^{13}\text{C-NMR}$ (125 MHz, CDCl_3 , ppm) δ 170.4, 130.4, 128.9, 65.0, 21.8, 20.8; FT-IR (thin film, cm^{-1}) 2939, 1740, 1378, 1230, 1025, 964; High-resolution MS (EI^+ , m/z) molecular ion calcd for $\text{C}_{12}\text{H}_{16}\text{O}_4$ 224.1043, found 224.1050, error 3.3 ppm.

Table 3, entry 4. Solvent= dichloromethane; reaction temperature= 40 °C; 10 mol % complex **1**; 9x1.00 mmol 1,5-COD (9 equiv). According to the general procedure (1.00 mmol scale) to obtain 175 mg **10D** as a pale yellow oil (68% yield). R_f = 0.36 (10 % ethyl acetate in hexanes). $^1\text{H-NMR}$ (500 MHz, CDCl_3 , ppm) δ 7.26 (t, J = 7.5 Hz, 2H), 7.22- 7.16 (m, 3H), 5.78 (s, 1H), 5.61 (s, 1H),

5.56 (t, $J = 6.0$ Hz, 1H), 2.94 (dd, $J = 14.0, 6.0$ Hz, 1H), 2.88 (dd, $J = 14.0, 6.0$ Hz, 1H), 2.05-2.01 (m, 4H), 2.00 (s, 3H), 1.85 (s, 3H); ^{13}C -NMR (125 MHz, CDCl_3 , ppm) δ 170.0, 137.5, 136.9, 131.1, 129.5, 128.1, 126.4, 124.0, 123.3, 74.2, 41.4, 22.3, 22.2, 21.2, 19.7; FT-IR (thin film, cm^{-1}) 2929, 1742, 1369, 1235, 1020; High-resolution MS (EI^+ , m/z) molecular ion calcd for $\text{C}_{17}\text{H}_{20}\text{O}_2$ 256.1458, found 256.1459, error 0.4 ppm.

Table 3, entry 5. Solvent= benzene; reaction temperature= 65°C ; 10 mol % complex **1**; 9x0.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 34 mg **10E** as a colorless oil (53% yield). $R_f = 0.54$ (10 % ethyl acetate in hexanes). ^1H -NMR (500 MHz, CDCl_3 , ppm) δ 5.59 – 5.56 (m, 2H), 3.63 (t, $J = 7.5$ Hz, 2H), 2.33 (t, $J = 7.5$ Hz, 2H), 2.05 – 1.98 (m, 4H), 1.79 (s, 3H), 0.90 (s, 9H), 0.05 (s, 6H); ^{13}C -NMR (125 MHz, CDCl_3 , ppm) δ 134.6, 131.1, 123.5, 122.6, 63.6, 36.5, 26.0, 22.8, 22.7, 19.8, 18.4, -5.3; FT-IR (thin film, cm^{-1}) 2930, 2858, 1472, 1255, 1099, 836, 775; High-resolution MS (EI^+ , m/z) molecular ion calcd for $\text{C}_{15}\text{H}_{28}\text{O}_1\text{Si}_1$ 252.1904, found 252.1905, error 0.3 ppm.

Table 3, entry 6. Solvent= benzene; reaction temperature= 65°C ; 10 mol % complex **1**; 9x0.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 28 mg **10F** as a colorless oil (49% yield). $R_f = 0.45$ (10 % ethyl acetate in hexanes). ^1H -NMR (500 MHz, CDCl_3 , ppm) δ 7.34 – 7.31 (m, 4H), 7.28 – 7.24 (m, 1H), 5.86 (s, 1H), 5.60 (s, 1H), 4.49 (s, 2H), 4.05 (s, 2H), 2.20 (q, $J = 7.5$ Hz, 2H), 2.08 (br s, 4H), 1.03 (t, $J = 7.5$ Hz, 3H); ^{13}C -NMR (125 MHz, CDCl_3 , ppm) δ 138.5, 137.8, 134.4, 128.3, 127.9, 127.5, 126.8, 120.8, 71.6, 71.3, 24.5, 22.6, 22.3, 13.0; FT-IR (thin film, cm^{-1}) 2933, 1454, 1356, 1067, 1028, 827, 734; High-resolution MS (ESI^+ , m/z) molecular ion calcd for $\text{C}_{16}\text{H}_{20}\text{O}_1$ 228.1514, found 251.1406, $[\text{M} + \text{Na}]^+$, error 1.8 ppm.

Table 3, entry 7. Solvent= benzene; reaction temperature= 65°C ; 5 mol % complex **1**; 9x0.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 71 mg **10G** as a pale yellow oil (72% yield). $R_f = 0.42$ (30 % ethyl acetate in hexanes). ^1H -NMR (500 MHz, CDCl_3 , ppm) δ 7.78 (d, $J = 8.5$ Hz, 2H), 7.29 (d, $J = 8.5$ Hz, 2H), 5.69 (s, 1H), 5.60 (s, 1H), 3.85 – 3.81 (m, 2H), 3.54 (t, $J = 8.0$ Hz, 2H), 2.43 (s, 3H), 2.06 – 2.01 (m, 4H), 1.88 (d, $J = 1.5$ Hz, 3H), 1.35 (s, 9H); ^{13}C -NMR (125 MHz, CDCl_3 , ppm) δ 150.8, 144.0, 137.5, 134.4, 132.6, 129.2, 127.7, 124.4, 122.9, 84.0, 47.5, 33.9, 27.8, 22.8, 22.5, 21.5, 19.5; FT-IR (thin film, cm^{-1}) 2932, 1728, 1357, 1287, 1157, 1090, 977, 812; High-resolution MS (ESI^+ , m/z) molecular ion calcd for $\text{C}_{21}\text{H}_{29}\text{NO}_4\text{S}$ 391.1817, found 414.1717, $[\text{M} + \text{Na}]^+$, error 1.9 ppm.

Table 3, entry 8. Solvent= benzene; reaction temperature= 65°C ; 10 mol % complex **1**; 9x0.25 mmol 1,5-COD (9 equiv). According to the general procedure (0.250 mmol scale) to obtain 25 mg **10H** as a colorless oil (61% yield). $R_f = 0.39$ (10 % ethyl acetate in hexanes). ^1H -NMR (500 MHz, CDCl_3 , ppm) δ 6.87 (t, $J =$

10.0 Hz, 1H), 5.66 (d, $J = 1.5$ Hz, 1H), 4.21 (q, $J = 7.0$ Hz, 2H), 2.23 – 2.18 (m, 2H), 2.09 – 2.04 (m, 2H), 1.95 (d, $J = 1.5$ Hz, 3H) 1.31 (t, $J = 7.0$ Hz, 3H); ^{13}C -NMR (125 MHz, CDCl_3 , ppm) δ 166.9, 137.4, 132.2, 130.1, 124.3, 60.3, 23.0, 21.5, 20.6, 14.2; FT-IR (thin film, cm^{-1}) 2940, 1719, 1451, 1382, 1262, 1076; High-resolution MS (ESI^+ , m/z) molecular ion calcd for $\text{C}_{10}\text{H}_{14}\text{O}_2$ 166.1059, found 167.1067, $[\text{M}+\text{H}]^+$, error 0.2 ppm.

***N,O*-cycloadduct 11:**

A mixture of cyclohexadiene **8C** (45 mg, 0.2 mmole, 1.0 equiv) and BocNHOH (32 mg, 0.24 mmole, 1.2 equiv) was dissolved in 4.0 mL CH_2Cl_2 and this mixture was cooled to 0 °C using an ice bath. Tetrapropyl ammonium periodate (90 mg, 0.24 mmole, 1.2 equiv) was dissolved in 2.0 mL CH_2Cl_2 and this solution was taken up in a gas tight syringe. This solution of the oxidant was added to the solution of the cyclohexadiene and the hydroxamic acid in CH_2Cl_2 maintained at 0 °C over 2 h using syringe pump. The analysis of the crude reaction mixture revealed complete consumption of the cyclohexadiene. The reaction mixture was concentrated and purified by flash column chromatography using silica gel. Gradient elution with 10 % ethyl acetate in hexanes (100 mL) followed by 20 % ethyl acetate in hexanes afforded the desired hetero-diels alder adduct **11** (53 mg, 75 % yield) as a yellow oil. R_f 0.30 (30 % ethyl acetate in hexanes). ^1H -NMR (500 MHz, CDCl_3 , ppm) δ 4.91 (d, $J = 13.0$ Hz, 1 H), 4.79 – 4.72 (m, 4H), 4.70 (d, $J = 13.0$ Hz, 1 H), 2.20 – 2.14 (m, 1H), 2.12 – 2.08 (m, 1H), 2.06 (s, 3H), 2.03 (s, 3H), 1.49 – 1.43 (m, 10 H), 1.36 – 1.30 (m, 1H); ^{13}C -NMR (125 MHz, CDCl_3 , ppm) δ 170.5, 170.46, 157.3, 136.7, 134.6, 81.9, 72.9, 60.7, 59.7, 52.1, 28.1, 24.0, 21.1, 20.74, 20.72; FT-IR (thin film, cm^{-1}) 2939, 1742, 1701, 1456, 1370, 1225, 1164, 1026; High-resolution MS (ESI^+ , m/z) molecular ion calcd for $\text{C}_{17}\text{H}_{25}\text{NO}_7$ 355.1631, found 378.1523, $[\text{M}+\text{Na}]^+$, error 0.9 ppm.

Cycloadduct 12:

To a solution of **8A** (45.5 mg, 0.203 mmol) in toluene (2.0 mL) was added maleic anhydride (24.5 mg, 0.250 mmol). This solution was heated to 105 °C for 12 h. After concentration and *in vacuo* (rotary evaporator) the residue was further purified by column chromatography on silica gel, eluting with 1:1 ethyl acetate-hexanes to give 55.9 mg of **12** (85% yield). $R_f = 0.26$ (1:1 ethyl acetate-hexanes). ^1H NMR (500 MHz, CDCl_3) δ 5.95 (dd, $J = 1.5, 6.0$ Hz, 1H), 3.73 (s, 3H), 3.71 (s, 3H), 3.57 (t, $J = 7.5$ Hz, 1H), 3.17 (m, 2H), 3.10 (m, 2H), 2.70 (m, 2H), 1.59 (m, 2H), 1.37 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 172.6, 172.4, 168.9, 168.8, 142.3, 126.2, 52.62, 52.60, 49.5, 44.8, 44.7, 36.1, 33.6, 32.1, 23.7, 22.8; FT-IR (thin film, cm^{-1}) 2956, 2881, 1841, 1782, 1743, 1440, 1342, 1289, 1234, 1158, 1086, 951, 911; high resolution ESI molecular ion calcd for 345.0945 $\text{C}_{16}\text{H}_{18}\text{O}_7\text{Na}$ found 345.0945.

^1H and ^{13}C NMR Spectra

Pulse Sequence: s2pul

Solvent: CDCl₃

Ambient temperature

INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec

Pulse 34.6 degrees

Acq. time 1.891 sec

Width 10000.0 Hz

16 repetitions

OBSERVE H1, 499.8984037 MHz

DATA PROCESSING

Line broadening 0.7 Hz

FT size 65536

Total time 1 min, 18 sec

6

S10

bp2262s-cyclohexadieneC13

Pulse Sequence: s2pul

Solvent: CDCl₃

Ambient temperature

GEMINI-300 "roesy.chem.buffalo.edu"

Relax. delay 5.000 sec

Pulse 90.0 degrees

Acq. time 1.706 sec

Width 18761.7 Hz

704 repetitions

OBSERVE C13, 75.4536589 MHz

DECOUPLE H1, 300.0754431 MHz

Power 1023 dB

continuously on

WALTZ-16 modulated

DATA PROCESSING

line broadening 1.0 Hz

ft size 65536

Total time 1 hr, 59 min, 57 sec

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.891 sec
Width 10000.0 Hz
16 repetitions

OBSERVE H1, 499.8983903 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 1 min, 18 sec.

8A

S12

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 GEMINI-300 "roesy.chem.buffalo.edu"

Relax. delay 5.000 sec
 Pulse 90.0 degrees
 Acq. time 1.706 sec
 Width 18761.7 Hz
 288 repetitions
 OBSERVE C13, 75.4536601 MHz
 DECOUPLE H1, 300.0754431 MHz
 Power 1023 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 1.0 Hz
 FT size 65536
 Total time 1 hr, 59 min, 57 sec

8A

Pulse Sequence: s2pu1

Solvent: CDCl3

Ambient temperature

INNOVA-400 "cosy.chem.buffalo.edu"

Relax. delay 1.000 sec

Pulse 46.9 degrees

Acq. time 1.800 sec

Width 6000.6 Hz

16 repetitions

OBSERVE H1, 399.9388996 MHz

DATA PROCESSING

Line broadening 0.9 Hz

FT size 32768

Total time 0 min, 44 sec

8B

S14

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
GEMINI-300 "roesy.chem.buffalo.edu"

Relax. delay 5.000 sec
Pulse 90.0 degrees
Acq. time 1.706 sec
Width 18761.7 Hz
224 repetitions
OBSERVE C13, 75.4536635 MHz
DECOUPLE H1, 300.0754431 MHz
Power 1023 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 65536
Total time 1 hr, 59 min, 57 sec

129.520
129.034
124.421

134.437
129.596

138.345

128.556
124.208

82.175

31.234

22.120
21.566

ppm

180

160

140

120

100

80

60

40

20

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.891 sec
Width 10000.0 Hz
16 Repetitions
OBSERVE H1, 499.8984025 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 1 min, 18 sec

8C

8C

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 GEMINI-300 "roesy.chem.buffalo.edu"

Relax. delay 5.000 sec
 Pulse 90.0 degrees
 Acq. time 1.706 sec
 Width 18761.7 Hz
 352 repetitions
 OBSERVE C13, 75.4536578 MHz
 DECOUPLE H1, 300.0754431 MHz
 Power 1023 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 1.0 Hz
 FT size 65536
 Total time 1 hr, 59 min, 57 sec

Pulse Sequence: s2put
Solvent: CDCl3
Ambient temperature
INOVA-500 "tocsy.Chem.buffalo.edu"

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.891 sec
Width 10000.0 Hz
16 repetitions
OBSERVE H1, 499.8984053 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 1 min, 18 sec

8D

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
GEMINI-300 "rossy.chem.buffalo.edu"

Relax. delay 5.000 sec
Pulse 90.0 degrees
Acq. time 1.706 sec
Width 18761.7 Hz
224 repetitions
OBSERVE C13, 75.4536601 MHz
DECOUPLE H1, 300.0754431 MHz
Power 1023 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
F1 size 65536
Total time 1 hr, 59 min, 57 sec

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
INOVA-400 "cosy.chem.buffalo.edu"

Relax. delay 1.000 sec
Pulse 46.9 degrees
Acq. time 1.800 sec
Width 6000.6 Hz
16 repetitions
OBSERVE H1, 399.9389010 MHz
DATA PROCESSING
Line broadening 0.9 Hz
FT size 32768
Total time 0 min, 44 sec

8E

S20

13C OBSERVE

Pulse Sequence: s2pu1
Solvent: CDCl3
Ambient temperature
GENINI-300 "roesy.chem.buffalo.edu"
Relax. delay 5.000 sec
Pulse 90.0 degrees
Acq. time 1.706 sec
Width 18761.7 Hz
64 repetitions
OBSERVE C13, 75.4536624 MHz
DECOUPLE H1, 300.0754431 MHz
Power 1023 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FI size 65536
Total time 1 hr, 59 min, 57 sec

Pulse Sequence: s2pul
Solvent: CDC13
Ambient temperature
INOVA-500 "tocsy.Chem.buffalo.edu"

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.891 sec
Width 10000.0 Hz
16 repetitions
OBSERVE H1, 499.8983998 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 1 min, 18 sec

8F

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 GEMINI-300 "roesy.chem.buffalo.edu"

Relax. delay 5.000 sec
 Pulse 90.0 degrees
 Acq. time 1.706 sec
 Width 18761.7 Hz
 176 repetitions
 OBSERVE C13, 75.4536595 MHz
 DECOUPLE H1, 300.0754431 MHz
 Power 1023 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 1.0 Hz
 FT size 65536
 Total time 1 hr, 59 min, 57 sec

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
INOVA-500 "tocsy.Chem.buffalo.edu"

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.891 sec
Width 10000.0 Hz
16 repetitions
OBSERVE H1, 499.8984001 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 1 min, 18 sec

S24

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 GEMINI-300 "roesy.chem.buffalo.edu"

Relax. delay 5.000 sec
 Pulse 90.0 degrees
 Acq. time 1.706 sec
 Width 18761.7 Hz
 320 repetitions
 OBSERVE C13, 75.4536601 MHz
 DECOUPLE H1, 300.0754431 MHz
 Power 1023 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 1.0 Hz
 FT size 65536
 Total time 1 hr, 59 min, 57 sec

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
INOVA-400 "cosy.chem.buffalo.edu"

Relax. delay 1.000 sec
Pulse 46.9 degrees
Acq. time 1.800 sec
Width 6000.6 Hz
16 repetitions
OBSERVE H1, 399.9389105 MHz
DATA PROCESSING
Line broadening 0.9 Hz
FT size 32768
Total time 0 min, 44 sec

7H

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
GEMINI-300 "roesy.chem.buffalo.edu"

Relax. delay 5.000 sec
Pulse 90.0 degrees
Acq. time 1.706 sec
Width 18761.7 Hz
128 repetitions
OBSERVE C13, 75.4536595 MHz
DECOUPLE H1, 300.0754431 MHz
Power 1023 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 65536
Total time 0 min, 0 sec

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
INOVA-400 "cosy.chem.buffalo.edu"

Relax. delay 1.000 sec
Pulse 45.9 degrees
Acq. time 1.800 sec
Width 6000.6 Hz
16 repetitions
OBSERVE H1, 399.9389113 MHz
DATA PROCESSING
Line broadening 0.9 Hz
FT size 32768
Total time 0 min, 44 sec

8H

Relax. delay 5.000 sec
Pulse 90.0 degrees
Acq. time 1.706 sec
Width 18761.7 Hz
224 repetitions
NOISEV C13, 75.436584 MHz
DECOUPLE H1, 300.0754431 MHz
Power 1023 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 65536
Total time 1 hr, 59 min, 57 sec

AAK-IV-prop)Bz-intalkyne

Pulse Sequence: s2pul

Solvent: CDCl₃

Ambient temperature

File: AAK-IV-propOBZ-intyne-1H

INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec

Pulse 34.6 degrees

Acq. time 1.891 sec

Width 10000.0 Hz

32 repetitions

OBSERVE H1, 499.8984031 MHz

DATA PROCESSING

Line broadening 0.7 Hz

FT size 65536

Total time 2 min, 20 sec

10A
(¹H-NMR Spectrum)

STANDARD CARBON PARAMETERS

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 User: 1-14-87
 File: AAK-IV-propOBz-intyne-13C
 INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 6.000 sec
 Pulse 81.8 degrees
 Acq. time 1.300 sec
 Width 37037.0 Hz
 768 repetitions
 OBSERVE C13, 125.6995521 MHz
 DECOUPLE H1, 499.9008903 MHz
 Power 40 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.5 Hz
 FT size 131072
 Total time 20 hr, 47 min, 55 sec

10A (13C-NMR Spectrum)

AAK-IV-215-1H

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
File: AAK-IV-215-1H
INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.891 sec
Width 10000.0 Hz
32 repetitions
OBSERVE H1, 499.8984059 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 2 min, 20 sec

10B
(¹H-NMR Spectrum)

Pulse Sequence: s2pul
 Solvent: CDCl₃
 Ambient temperature
 User: 1-14-87
 File: AAK-IV-215-13C
 INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 6.000 sec
 Pulse 81.8 degrees
 Acq. time 1.300 sec
 Width 37037.0 Hz
 192 repetitions
 OBSERVE C13, 125.695572 MHz
 DECOUPLE H1, 499.9008903 MHz
 Power 40 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.5 Hz
 FI size 131072
 Total time 2 hr, 5 min, 13 sec

10B ¹³C-NMR Spectrum)

AAK-IV-bisacetate-1H

Pulse Sequence: s2pul

Solvent: CDCl3

Ambient temperature

File: AAK-IV-bisacetate-1H

INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec

Pulse 34.6 degrees

Acq. time 1.891 sec

Width 10000.0 Hz

32 repetitions

OBSERVE H1, 499.8983983 MHz

DATA PROCESSING

Line broadening 0.7 Hz

F1 size 65536

Total time 2 min, 20 sec

10C

(¹H-NMR Spectrum)

STANDARD CARBON PARAMETERS

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 User: 1-14-87
 File: AAK-IV-bisacetate-13C
 INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 6.000 sec
 Pulse 81.8 degrees
 Acq. time 1.300 sec
 Width 37037.0 Hz
 160 repetitions
 OBSERVE C13, 125.695702 MHz
 DECOUPLE H1, 499.908903 MHz
 Power 40 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.5 Hz
 FT size 131072
 Total time 2 hr, 5 min, 13 sec

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.881 sec
Width 10000.0 Hz
32 repetitions
OBSERVE H1, 499.8984040 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 2 min, 20 sec

10D
(¹H-NMR Spectrum)

STANDARD CARBON PARAMETERS

Pulse Sequence: s2pul
 Solvent: CDCl₃
 Ambient temperature
 User: l-14-87
 File: AAK-IV-213-13C
 INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 6.000 sec
 Pulse 81.8 degrees
 Acq. time 1.300 sec
 Width 37037.0 Hz
 5944 repetitions
 OBSERVE C13, 125.6995516 MHz
 DECOUPLE H1, 499.9008903 MHz
 Power 40 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.5 Hz
 FT size 131072
 Total time 20 hr, 47 min, 55 sec

10D
 (13C-NMR Spectrum)

STANDARD PROTON PARAMETERS

Pulse Sequence: s2pul

Solvent: CDCl₃

Ambient temperature

File: AAK-IV-229-1H

INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec

Pulse 34.6 degrees

Acq. time 1.891 sec

Width 10000.0 Hz

32 repetitions

OBSERVE H1, 499.8984027 MHz

DATA PROCESSING

Line broadening 0.7 Hz

FT size 65536

Total time 2 min, 20 sec

10E
(¹H-NMR Spectrum)

STANDARD CARBON PARAMETERS

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 User: 1-14-87
 File: AAK-IV-229-13C
 INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 6.000 sec
 Pulse 81.8 degrees
 Acq. time 1.300 sec
 Width 37037.0 Hz
 640 repetitions
 OBSERVE C13, 125.6995499 MHz
 DECOUPLE H1, 499.9008903 MHz
 Power 40 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.5 Hz
 FI size 131072
 Total time 4 hr, 9 min, 58 sec

10E (13C-NMR Spectrum)

AAK-IV-232-F1-1H

Pulse Sequence: s2pul

Solvent: CDCl3

Ambient temperature

File: AAK-IV-232-F1-1H

INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec

Pulse 34.6 degrees

Acq. time 1.891 sec

Width 10000.0 Hz

32 repetitions

OBSERVE H1, 499.8984123 MHz

DATA PROCESSING

Line broadening 0.7 Hz

FI size 65536

Total time 2 min, 20 sec

10F

(¹H-NMR Spectrum)

STANDARD CARBON PARAMETERS

Pulse Sequence: s2pu1
 Solvent: CDCl3
 Ambient temperature
 User: 1-14-87
 File: AAK-IV-232-F1-13C
 INOVA-500 "tocsy.Chem.buffalo.edu"

Relax. delay 6.000 sec
 Pulse 81.8 degrees
 Acq. time 1.300 sec
 Width 37037.0 Hz
 2048 repetitions
 OBSERVE C13, 125.6995555 MHz
 DECOUPLE H1, 499.9008903 MHz
 Power 40 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.5 Hz
 FI size 131072
 Total time 4 hr, 9 min, 58 sec

10F
 (13C-NMR Spectrum)

AAK-IV-231-F1

Pulse Sequence: s2pul
Solvent: CDCl3
Ambient temperature
File: AAK-IV-231-1H
INOVA-500 "tocsy.Chem.Buffalo.edu"

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.891 sec
Width 10000.0 Hz
32 repetitions
OBSERVE H1, 499.8983954 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 2 min, 20 sec

10G
(¹H-NMR Spectrum)

STANDARD CARBON PARAMETERS

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 User: 1-14-87
 File: AAK-IV-231-13C
 INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 6.000 sec
 Pulse 81.8 degrees
 Acq. time 1.300 sec
 Width 37037.0 Hz
 464 repetitions
 OBSERVE C13, 125.6995572 MHz
 DECOUPLE H1, 499.9008903 MHz
 Power 40 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.5 Hz
 FT size 131072
 Total time 2 hr, 5 min, 13 sec

10G
 (¹³C-NMR Spectrum)

AAK-IV-232-1H

Pulse Sequence: s2pul

Solvent: CDCl₃

Ambient temperature

File: AAK-IV-2-ethylbutyrate-1H
INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec

Pulse 34.6 degrees

Acq. time 1.851 sec

Width 10000.0 Hz

32 repetitions

OBSERVE H1, 499.8984013 MHz

DATA PROCESSING

Line broadening 0.7 Hz

FT size 65536

Total time 2 min, 20 sec

10H

(¹H-NMR Spectrum)

STANDARD CARBON PARAMETERS

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 User: 1-14-87
 File: AAK-IV-chd-et-2-butyrate-13C
 INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 6.000 sec
 Pulse 81.8 degrees
 Acq. time 1.300 sec
 Width 37037.0 Hz
 1768 repetitions
 OBSERVE C13, 125.6995516 MHz
 DECOUPLE H1, 499.9008903 MHz
 Power 40 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.5 Hz
 FI size 131072
 Total time 10 hr, 24 min, 12 sec

10H (¹³C-NMR Spectrum)

(spectrometer noise gives
 rise to the peak at δ 123.97 ppm)

STANDARD PROTON PARAMETERS

Pulse Sequence: s2pul
Solvent: CDCl₃
Ambient temperature
File: AAK-IV-235-HDA-1H
INOVA-500 "tocsy.Chem.buffalo.edu"

Relax. delay 2.000 sec
Pulse 34.6 degrees
Acq. time 1.831 sec
Width 10000.0 Hz
32 repetitions
OBSERVE H1, 499.8984024 MHz
DATA PROCESSING
Line broadening 0.7 Hz
FT size 65536
Total time 2 min, 20 sec

11
(¹H-NMR Spectrum)

STANDARD CARBON PARAMETERS

Pulse Sequence: s2pul

Solvent: CDCl3

Ambient temperature

User: 1-14-87

File: AAK-IV-235-HDA-13C

INOVA-500 "tocsy.Chem.Buffalo.edu"

Relax. delay 6.000 sec

Pulse 81.8 degrees

Acq. time 1.300 sec

Width 37037.0 Hz

704 repetitions

OBSERVE C13, 125.6395572 MHz

DECOUPLE H1, 499.9008903 MHz

Power 40 dB

continuously on

WALTZ-16 modulated

DATA PROCESSING

line broadening 0.5 Hz

FT size 131072

Total time 4 hr, 9 min, 58 sec

11 (¹³C-NMR Spectrum)

(spectrometer noise gives
rise to the peak at δ 123.97 ppm)

STANDARD PROTON PARAMETERS

Pulse Sequence: s2pul

Solvent: CDCl₃

Ambient temperature

INOVA-500 "tocsy.chem.buffalo.edu"

Relax. delay 2.000 sec

Pulse 34.6 degrees

Acq. time 1.891 sec

Width 10000.0 Hz

16 repetitions

OBSERVE H1, 499.8983918 MHz

DATA PROCESSING

Line broadening 0.7 Hz

FT size 65536

Total time 1 min, 18 sec

12

Pulse Sequence: s2pul
 Solvent: CDCl3
 Ambient temperature
 GEMINI-300 "roesy.Chem.buffalo.edu"

Relax. delay 5.000 sec
 Pulse 90.0 degrees
 Acq. time 1.706 sec
 Width 18761.7 Hz
 160 repetitions
 OBSERVE C13, 75.4536618 MHz
 DECOUPLE H1, 300.0754431 MHz
 Power 1023 dB
 continuously on
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 1.0 Hz
 FI size 65536
 Total time 0 min, 0 sec

12