Functionalization of cyclodextrins via reactions of 2,3-anhydro-cyclodextrins

De-Qi Yuan,^{*,[a]} Tsutomu Tahara,^[b] Wen-Hua Chen,^[a] Yuji Okabe,^[a] Cheng Yang,^[a] Youichi Yagi,^[a] Yasuyoshi Nogami,^[b] Makoto Fukudome,^[a] Kahee Fujita^{*,[a]}

[a] Department of Molecular Medicinal Sciences, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki 852-8521, Japan; [b] Daiichi College of Pharmaceutical Sciences, Tamagawa-cho, Minami-ku, Fukuoka 815-8511, Japan

¹H and ¹³C NMR spectra

3 ^A -Deoxy-3 ^A -imidazolyl- <i>altro</i> -β-CD (9).....	S2	3 ^A -Amino-3 ^A -deoxy-β-CD (26).....	S18
2 ^A -Deoxy-2 ^A -imidazolyl-β-CD (10).....	S3	2 ^A -Deoxy-2 ^A -iodo- <i>altro</i> -β-CD (27).....	S19
3 ^A -Benzylmercapto-3 ^A -deoxy- <i>altro</i> -β-CD (11).....	S4	3 ^A -Deoxy-3 ^A -iodo-β-CD (28).....	S20
2 ^A -Benzylmercapto-2 ^A -deoxy-β-CD (12).....	S5	3 ^A ,3 ^C -Diazido-3 ^A ,3 ^C -dideoxy-β-CD (32).....	S21
3 ^A -Azido-3 ^A -deoxy- <i>altro</i> -β-CD (13).....	S6	3 ^A ,3 ^D -Diazido-3 ^A ,3 ^D -dideoxy-β-CD (33).....	S22
2 ^A -Azido-2 ^A -deoxy-β-CD (14).....	S7	3 ^A ,3 ^C ,3 ^E -Triazido-3 ^A ,3 ^C ,3 ^E -trideoxy-β-CD (34).....	S23
3 ^A -Amino-3 ^A -deoxy- <i>altro</i> -β-CD (15).....	S8	3 ^A ,3 ^C -Diamino-3 ^A ,3 ^C -dideoxy-β-CD (35).....	S24
2 ^A -Amino-2 ^A -deoxy-β-CD (16).....	S9	3 ^A ,3 ^D -Diamino-3 ^A ,3 ^D -dideoxy-β-CD (36).....	S25
3 ^A -Deoxy-3 ^A -iodo- <i>altro</i> -β-CD (17).....	S10	3 ^A ,3 ^C ,3 ^E -Triamino-3 ^A ,3 ^C ,3 ^E -trideoxy-β-CD (37).....	S26
2 ^A -Deoxy-2 ^A -imidazolyl- <i>altro</i> -β-CD (19).....	S11	2 ^A ,3 ^A -didehydro-2 ^A ,3 ^A -dideoxy-β-CD (42).....	S27
3 ^A -Deoxy-3 ^A -imidazolyl-β-CD (20).....	S12	2 ^A ,2 ^B ,3 ^A ,3 ^B -tetrahydro-2 ^A ,2 ^B ,3 ^A ,3 ^B -tetraideoxy-β-CD(43).....	S28
2 ^A -Benzylmercapto-2 ^A -deoxy- <i>altro</i> -β-CD (21).....	S13	2 ^A ,2 ^C ,3 ^A ,3 ^C -tetrahydro-2 ^A ,2 ^C ,3 ^A ,3 ^C -tetraideoxy-β-CD(44).....	S29
3 ^A -Benzylmercapto-3 ^A -deoxy-β-CD (22).....	S14	2 ^A ,2 ^C ,3 ^A ,3 ^C -tetrahydro-2 ^A ,2 ^C ,3 ^A ,3 ^C -tetraideoxy-β-CD(45).....	S30
2 ^A -Azido-2 ^A -deoxy- <i>altro</i> -β-CD (23).....	S15	2 ^A ,2 ^C ,2 ^E ,3 ^A ,3 ^C ,3 ^E -hexadehydro-2 ^A ,2 ^C ,2 ^E ,3 ^A ,3 ^C ,3 ^E -	
3 ^A -Azido-3 ^A -deoxy-β-CD (24).....	S16	hexadeoxy-β-CD(46).....	S31
2 ^A -Amino-2 ^A -deoxy- <i>altro</i> -β-CD (25).....	S17	Per(3-deoxy)-β-cyclomannin (48).....	S32

Fig. S2 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A-deoxy-3^A-imidazolyl-*altro*-β-CD **9** in D₂O (CH₃CN int).

Fig. S3 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A-deoxy-2^A-imidazolyl-β-CD **10** in D₂O (CH₃CN int).

Fig.S4 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A-benzylmercapto-3^A-deoxy-*altro*-β-CD **11** in DMSO-d₆ (TMS int).

Fig. S5 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A-benzylmercapto-2^A-deoxy-β-CD **12** in DMSO-d₆ (TMS int, deuterated).

Fig. S6 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A-azido-3^A-deoxy-*altro*-β-CD **13** in D₂O (CH₃CN int).

Fig. S7 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A-azido-2^A-deoxy-β-CD **14** in DMSO-d₆ (TMS int, deuterated).

Fig. S8 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A-amino-3^A-deoxy-*altro*-β-CD **15** in D₂O (CH₃CN int).

Fig. S9 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A-amino-2^A-deoxy-β-CD **16** in DMSO-d₆ (TMS int, deuterated).

Fig. S10 ¹H (400 MHz) and ¹³C NMR spectra (100 MHz) of 3^A-deoxy-3^A-iodo-*altro*-β-CD **17** in D₂O (CH₃CN int).

Fig. S11 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A-deoxy-2^A-imidazolyl-*altro*-β-CD **19** in DMSO-d₆ (TMS int).

Fig. S12 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A-deoxy-3^A-imidazolyl-β-CD **20** in D₂O (CH₃CN int).

Fig. S13 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A-benzylmercapto-2^A-deoxy-*α*-D-glucopyranoside **21** in DMSO-d₆ (TMS int, deuterated).

Fig. S14 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A-benzylmercapto-3^A-deoxy-β-CD **22** in DMSO-d₆ (TMS int, deuterated)

Fig. S15 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A-azido-2^A-deoxy-*altro*-β-CD **23** in D₂O (CH₃CN int).

Fig. S16 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A-azido-3^A-deoxy-β-CD **24** in D₂O (CH₃CN int).

Fig. S17 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A-amino-2^A-deoxy-altro-β-CD **25** in D₂O (CH₃CN int).

Fig. S18 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A-amino-3^A-deoxy-β-CD **26** in DMSO-d₆ (TMS int, deuterated).

Fig. S19 ^1H (400 MHz) and ^{13}C NMR spectra (100 MHz) of 2^A-deoxy-2^A-iodo-*altro*- β -CD **27** in D_2O (CH_3CN int).

Fig. S20 ¹H (400 MHz) and ¹³C NMR spectra (100 MHz) of 3^A-deoxy-3^A-iodo-β-CD **28** in D₂O (CH₃CN int).

Fig. S21 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A,3^C-diazido-3^A,3^C-dideoxy-β-CD **32** in DMSO-d₆ (TMS int, deuterated).

33

Fig. S22 ^1H (500 MHz) and ^{13}C NMR spectra (125 MHz) of 3^A,3^D-diazido-3^A,3^D-dideoxy- β -CD **33** in DMSO- d_6 (TMS int, deuterated).

Fig. S23 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A,3^C,3^E-triazido-3^A,3^C,3^E-trideoxy-β-CD **34** in DMSO-d₆ (TMS int, deuterated).

Fig. S24 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 3^A,3^C-diamino-3^A,3^C-dideoxy-β-CD **35** in DMSO-d₆ (TMS int, deuterated).

Fig. S25 ^1H (500 MHz) and ^{13}C NMR spectra (125 MHz) of $3^{\text{A}}, 3^{\text{D}}$ -diamino- $3^{\text{A}}, 3^{\text{D}}$ -dideoxy- β -CD **36** in DMSO- d_6 (TMS int, deuterated).

Fig. S26 ^1H (500 MHz) and ^{13}C NMR spectra (125 MHz) of $3^{\text{A}}, 3^{\text{C}}, 3^{\text{E}}$ -triamino- $3^{\text{A}}, 3^{\text{C}}, 3^{\text{E}}$ -trideoxy- β -CD **37** in DMSO-d_6 (TMS int, deuterated).

Fig. S27 ^1H (500 MHz) and ^{13}C NMR spectra (125 MHz) of 2^A,3^A-dideoxy-2^A,3^A-dideoxy- β -CD **42** in DMSO- d_6 (TMS int).

Fig. S28 ^1H (500 MHz) and ^{13}C NMR spectra (125 MHz) of $2^{\text{A}}, 2^{\text{B}}, 3^{\text{A}}, 3^{\text{B}}$ -tetrahydro- $2^{\text{A}}, 2^{\text{B}}, 3^{\text{A}}, 3^{\text{B}}$ -tetraoxo- β -CD **43** in D_2O (CH_3CN int).

Fig. S29 ¹H (500 MHz) and ¹³C NMR spectra (125 MHz) of 2^A,2^C,3^A,3^C-tetrahydro-2^A,2^C,3^A,3^C-tetraoxo-β-CD **44** in D₂O (CH₃CN int).

Fig. S30 ^1H (500 MHz) and ^{13}C NMR spectra (125 MHz) of $2^{\text{A}}, 2^{\text{D}}, 3^{\text{A}}, 3^{\text{D}}$ -tetrahydro- $2^{\text{A}}, 2^{\text{D}}, 3^{\text{A}}, 3^{\text{D}}$ -tetraoxy- β -CD **45** in D_2O (CH_3CN int).

Fig. S31 ^1H (500 MHz) and ^{13}C NMR spectra (125 MHz) of 2^A,2^C,2^E,3^A,3^C,3^E-hexadehydro-2^A,2^C,2^E,3^A,3^C,3^E-hexadeoxy- β -CD **46** in D_2O (CH_3CN int).

Fig. S32 ^1H - ^{13}C COSY NMR spectrum (500 MHz) of per(3-deoxy)- β -cyclomannin **48** in D_2O (CH_3CN int).