“Generation of acylnitroso dienophiles: a study of metal catalysis”

Mauro F. A. Adamo and Simone Bruschi

Table of Contents

S2: General experimental.

S3-S4: Spectroscopic characterisation of compounds **4, 6, 8, 10a/b, 12a/b, 14a/b**.

S5-S10: ¹H-NMR of compounds **4, 6, 8, 10a/b, 12a/b, 14a/b**.

S11: Experiment described in **Table 1 entry 4**: ¹H-NMR of the crude reaction mixture.

General experimental: ^1H and ^{13}C Spectra were recorded on a 400 MHz spectrometers at ambient temperatures. For ^1H NMR recorded in CDCl_3 chemical shifts (δ_{H}) are quoted in parts per million (ppm) and are referenced to the residual solvent peak. The following abbreviations are used: s, singlet, d, doublet, t, triplet, dd, doublet of doublets, dt, doublet of triplets, tt, triplet of triplets, m, multiplet and br, broad. Coupling constants (J) were recorded in Hertz (Hz) to the nearest 0.5Hz. Infrared (IR) spectra were recorded as thin films between NaCl plates. Absorption maximum (ν_{max}) was reported in wave numbers (cm^{-1}) and only selected peaks are reported. The following abbreviations are used: w, weak, m, medium, s, strong and br, broad.

Flash chromatography was carried out using *silica gel 60* (0.040-0.063mm, 230-400 mesh) as the stationary phase. Thin layer chromatography was carried out on aluminium backed plates pre-coated with *silica gel 60*, which were visualized by quenching of u.v. fluorescence ($\lambda_{\text{max}} = 254 \text{ nm}$) or by staining with either 10% w/v ammonium molybdate in 2M sulphuric acid or basic potassium permanganate solution (followed by heat) as appropriate. Retention factors (R_f) are reported to ± 0.5 .

2-Oxa-3-aza-bicyclo[2.2.2]oct-5-ene-3-carboxylic acid tert-butyl ester 4:²³ pale yellow liquid, $R_f = 0.6$ (ethyl acetate : petroleum spirits 1 : 3); δ_H (400 MHz, $CDCl_3$) 6.50-6.48 (m, 2H), 4.69 (app.s, 2H), 2.18-2.02 (m, 2H), 1.42 (s, 9H), 1.35-1.27 (m, 2H); δ_C (100 MHz, $CDCl_3$) 157.3, 131.2, 131.1, 81.1, 70.2, 49.7, 27.7, 23.1, 20.1.

2-Oxa-3-aza-bicyclo[2.2.1]hept-5-ene-3-carboxylic acid tert-butyl ester 6:²⁴ pale yellow liquid, $R_f = 0.6$ (ethyl acetate : petroleum spirits 1 : 3); δ_H ($CDCl_3$, 400MHz) 6.13-6.07 (m, 2H), 4.88 (s, 1H), 4.67 (s, 1H), 1.67-1.63 (m, 1H), 1.41 (d, $J = 8$, 1H), 1.16 (s, 9H); δ_C ($CDCl_3$, 100MHz) 157.8, 133.3, 132.1, 82.7, 81.0, 64.3, 47.3, 27.4.

4,5-Dimethyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 8:²⁴ pale yellow liquid, $R_f = 0.4$ (ethyl acetate : petroleum spirits 1 : 3); δ_H (400 MHz, $CDCl_3$) 4.19 (s, 2H), 3.88 (s, 2H), 1.65 (s, 3H), 1.58 (s, 3H), 1.49 (s, 9H); δ_C (100 Mhz, $CDCl_3$) 154.5, 122.6, 121.4, 80.9, 70.7, 48.0, 27.8, 14.8, 13.3.

5-Methyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 10a and 4-methyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 10b:²⁵ pale yellow liquid, $R_f = 0.4$ (ethyl acetate : petroleum spirits 1 : 3); δ_H ($CDCl_3$, 400MHz) 5.48 (m, 1H, isomer A), 4.60-4.59 (m, 1H, isomer B), 4.33 (m, 2H, isomer A), 4.23 (m, 2H, isomer B), 4.00 (m, 2H, isomer A), 3.91 (m, 2H, isomer B), 1.70 (s, 3H, isomer B), 1.63 (s, 3H, isomer A), 1.49-1.46 (m, 18H, isomer A and B); δ_C ($CDCl_3$, 100MHz) δ : 154.4, 154.3, 130.9, 129.6, 117.5, 115.9, 80.8, 70.4, 67.3, 53.0, 48.0, 44.2, 27.6, 19.1, 17.6.

3,6-Dimethyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 12a/b:²⁶ pale yellow liquid, $R_f = 0.4$ (ethyl acetate : petroleum spirits 1 : 3); δ_H ($CDCl_3$, 400MHz) δ : 5.78-5.74 (m, 1.5H), 5.66 (d, $J = 10$, 1H), 4.63 (br, 1H), 4.46-4.39 (m, 1.5H), 1.49 (s, 9H), 1.33 (d, $J = 7$, 1H), 1.29 (d, $J = 7$, 3.3H), 1.25 (d, $J = 7$, 3.3H); δ_C ($CDCl_3$, 100MHz) δ : 153.8, 128.3, 127.9, 127.4, 126.6, 80.5, 80.3, 74.3, 72.8, 49.5, 49.3, 27.8, 18.2, 18.0, 17.5, 16.7.

6-Methyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 14a and 3-methyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 14b:²⁴ pale

yellow liquid, $R_f = 0.4$ (ethyl acetate : petroleum spirits 1 : 3); δ_H 5.71-5.64 (m, 4H), 4.55-4.46 (m, 2H), 4.34-4.32 (m, 1H), 4.11-4.02 (m, 2H), 3.86-3.81 (m, 1H), 1.42-1.39 (m, 18H), 1.21 (d, $J = 6$, 3H), 1.17 (d, $J = 7$, 3H); δ_H (CDCl₃, 100MHz) 155.3, 154.3, 153.9, 152.9, 128.9, 127.6, 122.9, 121.6, 84.3, 82.0, 80.7, 73.1, 67.8, 50.0, 44.0, 27.8, 27.7, 27.5, 27.0, 18.1, 17.1.

2-Oxa-3-aza-bicyclo[2.2.2]oct-5-ene-3-carboxylic acid tert-butyl ester 4

2-Oxa-3-aza-bicyclo[2.2.1]hept-5-ene-3-carboxylic acid tert-butyl ester 6

4,5-Dimethyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 8

5-Methyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 10a and 4-methyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 10b

3,6-Dimethyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 12a/b

6-Methyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 14a and 3-methyl-3,6-dihydro-[1,2]oxazine-2-carboxylic acid tert-butyl ester 14b

Experiment described in Table 1 entry 4: ^1H -NMR of the crude reaction mixture