Supporting Information for

Generation of Allylic Indium by Hydroindation of 1,3-Dienes and One-Pot Reaction with Carbonyl Compounds

Naoki Hayashi,[†] Hiroyuki Honda,[†] Makoto Yasuda,[†] Ikuya Shibata,^{*,‡} Akio Baba^{*,†}

[‡]*Research Center for Environmental Preservation, Osaka University, 2-4 Yamadaoka, Suita, Osaka, 565-0871, Japan, and* [†]*Department of Applied Chemistry, Graduate School of Engineering, Osaka University, 2-1 Yamadaoka, Suita, Osaka, 565-0871, Japan*

General:

IR spectra were recorded as thin film on a Horiba FT-720 spectrometer. All ¹H and ¹³C-NMR spectra were recorded with a JEOL JNM-GSX-270 (270 and 67 MHz, respectively) or JEOL JMT-400/54/SS (400 and 100 MHz, respectively) in deuteriochloroform (CDCl₃) containing 0.03% (w/v) of tetramethylsilane. Mass spectra were recorded on a JEOL JMS-DS-303 spectrometer. Column chromatography was performed by using Fuji Davison silica gel FL-100DX. GPC was performed by JAPAN ANALYTICAL INDUSTRY CO., LTD. LC-908.

Unless otherwise noted, materials obtained from commercial supplier were used without further purification. Bu₃SnH was obtained by distillation under reduced pressure from the mixture of Bu₃SnOMe and PMHS.

Typical experimental procedures:

The reaction of 2,3-dimethyl-1,3-butadiene or isoprene with electrophiles

A 10 mL of round bottom flask charged with InCl₃ (0.442 g, 2.0 mmol) was dried by heating at 110 °C under reduced pressure (1 mmHg) for 1 h. After nitrogen was filled, THF (2 mL) was added to dissolve InCl₃. To the mixture was added Bu₃SnH (2.0 mmol) at -78 °C. The mixture was stirred for 10 min to prepare dichloroindium hydride (HInCl₂). After the solution was warmed up to rt, 1,3-diene (4.0 mmol) was added to the mixture and the resulting mixture was stirred for 5 h. Then electrophile (1.0 mmol) was added and stirred for 3 h. The resulting mixture was quenched by 1N HCl aq (10 mL). (When imine was used as electrophile the reaction was quenched by water.) The reaction mixture was extracted with ether (10 mL x 3). The combined organic layer was treated with NH₄F aq and then precipitate was filtered to remove tin compound. The filtrate was extracted with ether and dried over MgSO₄. Concentration followed by silica gel column chromatography eluting with hexane/AcOEt = (9/1) afforded product. Further Purification was performed by distillation under reduced pressure or GPC. Product was determined by ¹H NMR.

The reaction of 1,3-butadiene with electrophiles

A 10 mL of round bottom flask charged with InCl_3 (0.442 g, 2.0 mmol) was dried by heating at 110°C under reduced pressure (1 mmHg) for 1 h. After nitrogen was filled, THF (2 mL) was added to dissolve InCl_3 . To the mixture was added Bu_3SnH (2.0 mmol) at -78°C . The mixture was stirred for 10 min to prepare dichloroindium hydride (HInCl_2). Then 1,3-butadiene (4 mmol, measured by volume using syringe) was added to the mixture. The solution was warmed up to rt and stirred for 5 h. Then electrophile (1.0 mmol) was added and stirred for 3 h. The resulting mixture was quenched by 1N HCl aq (10 mL). (When imine was used as electrophile the reaction was quenched by water.) The reaction mixture was extracted with ether (10 mL x 3). The combined organic layer was treated with NH_4F aq and then precipitate was filtered to remove tin compound. The filtrate was extracted with ether and dried over MgSO_4 . Concentration followed by silica gel column chromatography eluting with hexane/ AcOEt = (9/1) afforded product. Further Purification was performed by distillation under reduced pressure or GPC. Product was determined by ^1H NMR.

(In the case of product **3m**, 1,3-butadiene (ca. 1 mL, 12 mmol) liquefied in another flask at -30°C was added to the mixture after generation of HInCl_2 .)

Physical and spectral data:

Products **3a**, **3g**, **3k**, **3l**, **3m**, **3n**, **3o**, **3p**, **3s** are known compounds.

2,2,3-Trimethyl-1-phenyl-but-3-en-1-ol (**3a**)

Reference : Kobayashi, S.; Nishio, K. *J. Org. Chem.* **1994**, 59, 6620-6628.

2,2,3-Trimethyl-1-*p*-tolyl-but-3-en-1-ol (**3b**)

Shape: colorless liquid

bp 120°C / 0.04 mmHg

IR (neat) 3467 (O-H), 1631 (C=C) cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 7.21 (d, J = 8.0 Hz, 2H), 7.12 (d, J = 8.0 Hz, 2H), 5.02 (s, 1H), 4.96 (s, 1H), 4.60 (s, 1H), 2.34 (s, 3H), 2.04 (s, 1H), 1.86 (s, 3H), 0.98 (s, 3H), 0.93 (s, 3H)

^{13}C NMR (CDCl_3 , 100 MHz) δ 150.68, 137.24, 136.73, 128.04, 127.80, 112.94, 77.29, 44.42, 24.19, 21.04, 20.06, 19.79

MS (CI, 70 eV) m/z 205 ($\text{M}^+ + 1$, 0.58), 188 ($\text{M}^+ + 1 - \text{OH}$, 16), 187 ($\text{M}^+ - \text{OH}$, 100), 121 (33), 84 ($\text{M}^+ - \text{CH}_3\text{C}_6\text{H}_8\text{CHOH}$, 5)

HRMS calcd for $\text{C}_{14}\text{H}_{20}\text{O} + \text{H}$: 205.1592, found: m/z 205.1601 (CI, ($\text{M}^+ + \text{H}$), + 0.9 mmu)

1-(4-Chloro-phenyl)-2,2,3-trimethyl-but-3-en-1-ol (3c)

Shape: colorless liquid

bp 120 °C/ 0.04 mmHg

IR (neat) 3467 (O-H), 1631 (aliphatic C=C), 1184 (C-O) cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 7.30-7.26 (m, 4H), 5.05 (s, 1H), 4.96 (s, 1H), 4.60 (d, J = 1.7 Hz, 1H), 2.10 (d, J = 1.7 Hz, 1H), 1.86 (s, 3H), 0.96 (s, 3H), 0.93 (s, 3H)

^{13}C NMR (CDCl_3 , 100 MHz) δ 150.19, 138.71, 132.97, 129.25, 127.54, 113.54, 76.78, 44.48, 24.04, 19.97, 19.76

MS (CI, 70 eV) m/z 225 ($\text{M}^+ + 1$, 0.22), 210 ($\text{M}^+ + 1 - \text{CH}_3$, 4), 209 ($\text{M}^+ - \text{CH}_3$, 33), 208 (15), 207 ($\text{M}^+ - \text{OH}$, 100), 141 ($\text{M}^+ - \text{C}(\text{CH}_3)_2\text{C}(\text{CH}_3)=\text{CH}_2$, 21), 84 ($\text{M}^+ + 1 - \text{ClC}_6\text{H}_4\text{CHOH}$, 8)

HRMS calcd for $\text{C}_{13}\text{H}_{17}\text{ClO}$: 224.0968, found: m/z 225.1030 (CI, ($\text{M}^+ + \text{H}$), -1.6 mmu)

2,2,3-Trimethyl-1-(4-nitro-phenyl)-but-3-en-1-ol (3d)

Shape: brown liquid

bp 170 °C/ 0.04 mmHg

IR (neat) 3437 (O-H) cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 8.18 (d, J = 8.9 Hz, 2H), 7.52 (d, J = 8.9 Hz, 2H), 5.09 (s, 1H), 4.98 (s, 1H), 4.72 (s, 1H), 2.29 (s, 1H), 1.88 (s, 3H), 0.97 (s, 3H), 0.96 (s, 3H)

^{13}C NMR (CDCl_3 , 100 MHz) δ 149.52, 147.93, 147.19, 128.67, 122.54, 114.16, 76.63, 44.61, 23.84, 20.11, 19.69

MS (CI, 70eV) m/z 236 ($\text{M}^+ + 1$, 100), 218 ($\text{M}^+ - \text{OH}$, 16), 204 ($\text{M}^+ + 1 - \text{OH} - \text{CH}_3$, 100), 154 (79), 84 ($\text{M}^+ + 1 - \text{CHOHC}_6\text{H}_4\text{NO}_2$, 12)

HRMS calcd for $\text{C}_{13}\text{H}_{17}\text{NO}_3 + \text{H}$: 236.1287, found: m/z 236.1284 (CI, ($\text{M}^+ + \text{H}$), - 0.3 mmu)

1-(4-Methoxy-phenyl)-2,2,3-trimethyl-but-3-en-1-ol (3e)

Shape: colorless liquid

bp 155 °C/ 0.7 mmHg

IR (neat) 3537 (O-H) cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 7.24 (d, J = 8.7 Hz, 2H), 6.84 (d, J = 8.7 Hz, 2H), 5.00 (s, 1H), 4.94 (s, 1H), 4.57 (s, 1H), 3.79 (s, 3H), 1.85 (s, 3H), 0.98 (s, 3H), 0.91 (s, 3H)

^{13}C NMR (CDCl_3 , 100 MHz) δ 158.75, 150.69, 132.33, 128.92, 112.97, 112.74, 77.04, 55.11, 44.52, 24.15, 20.02, 19.81

MS (CI, 70 eV) m/z 221 ($\text{M}^+ + 1$, 6), 203 ($\text{M}^+ - \text{OH}$, 100), 137 ($\text{M}^+ - \text{C}(\text{CH}_3)_2\text{C}(\text{CH}_3)=\text{CH}_2$, 31)

HRMS calcd for $\text{C}_{14}\text{H}_{20}\text{O}_2 + \text{H}$: 221.1542, found: m/z 221.1552 (CI, ($\text{M}^+ + \text{H}$), + 1.0 mmu)

methyl 4-(1-Hydroxy-2,2,3-trimethyl-but-3-enyl)-benzoate (3f)

Shape: white solid

mp 63-64 °C

bp 155 °C/ 0.3 mmHg

IR (KBr) 3525 (O-H), 1705 (C=O), 1281 (C-O-C), 1188 (C-OH), 1111 cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 7.99 (d, J = 8.2 Hz, 2H), 7.42 (d, J = 8.2 Hz, 2H), 5.06 (s, 1H), 4.97 (s, 1H), 4.67 (s, 1H), 3.92 (s, 3H), 2.17 (s, 1H), 1.87 (s, 3H), 0.96 (s, 3H), 0.95 (s, 3H)

^{13}C NMR (CDCl_3 , 100 MHz) δ 167.06, 150.09, 145.60, 129.14, 128.67, 127.94, 113.72, 77.07, 52.04, 44.59, 24.07, 20.10, 19.77

MS (CI, 70 eV) m/z 249 ($\text{M}^+ + 1$, 100), 231 ($\text{M}^+ - \text{OH}$, 39), 165 ($\text{M}^+ - \text{C}(\text{CH}_3)_2\text{C}(\text{CH}_3)=\text{CH}_2$, 19), 84 ($\text{M}^+ + 1 - \text{CHOHC}_6\text{H}_4\text{COOCH}_3$, 7)

HRMS calcd for $\text{C}_{15}\text{H}_{20}\text{O}_3 + \text{H}$: 249.1491, found: m/z 249.1493 (CI, ($\text{M}^+ + \text{H}$), + 0.2 mmu)

Anal. calcd for $\text{C}_{15}\text{H}_{20}\text{O}_3$: C, 72.55; H, 8.12, found: C, 72.29; H, 8.04.

4,4,5-Trimethyl-1-phenyl-hex-5-en-3-ol (3g)

Reference : Kobayashi, S.; Nishio, K. *J. Org. Chem.* **1994**, 59, 6620-6628.

3,3,4-Trimethyl-2-phenyl-pent-4-en-2-ol (3h)

Shape: colorless liquid

bp 140 °C/ 0.5 mmHg

IR (neat) 3541 (O-H) cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 7.43-7.40 (m, 2H), 7.40-7.22 (m, 3H), 5.05 (s, 1H), 4.90 (s, 1H), 2.41(s, 1H), 1.59 (s, 3H), 1.57 (s, 3H), 1.12 (s, 3H), 1.06 (s, 3H)

^{13}C NMR (CDCl_3 , 100 MHz) δ 151.47, 145.64, 127.04, 126.98, 126.37, 114.09, 77.10, 46.54, 25.46, 24.29, 23.88, 23.82

MS (CI, 70 eV) m/z 205 ($\text{M}^+ + 1$, 0.92), 188 (15), 187 ($\text{M}^+ - \text{OH}$, 100), 121 ($\text{M}^+ - \text{C}(\text{CH}_3)_2\text{C}(\text{CH}_3)=\text{CH}_2$, 62)

HRMS calcd for $\text{C}_{14}\text{H}_{20}\text{O} + \text{H}$: 205.1592, found: m/z 205.1595 (CI, ($\text{M}^+ + \text{H}$), + 0.3 mmu)

Anal. calcd for $\text{C}_{14}\text{H}_{20}\text{O}$: C, 82.30; H, 9.87, found: C, 82.13; H, 9.74.

3,4,4,5-Tetramethyl-1-phenyl-hex-5-en-3-ol (3i)

Shape: colorless liquid

IR (neat) 3516 (O-H) cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 7.30-7.16 (m, 5H), 5.01 (s, 1H), 4.86 (s, 1H), 2.79 (ddd, $J=13.4$ and 10.7 and 6.3 Hz, 1H), 2.66 (ddd, $J=13.4$ and 10.8 and 6.8 Hz, 1H), 1.87 (s, 3H), 1.84-1.70 (m, 2H), 1.65 (s, 1H), 1.22 (s, 3H), 1.14 (s, 3H), 1.12 (s, 3H)

^{13}C NMR (CDCl_3 , 100 MHz) δ 151.57, 143.18, 128.41, 128.34, 125.61, 113.65, 75.78, 46.45, 38.96, 30.48, 23.74, 23.62, 23.49, 21.46

MS (CI, 70 eV) m/z 215 ($\text{M}^+ + 1 - \text{H}_2\text{O}$, 100), 159 (20), 149 ($\text{M}^+ - \text{C}(\text{CH}_3)_2\text{C}(\text{CH}_3)=\text{CH}_2$, 58), 145 (25), 131 ($\text{M}^+ - \text{H}_2\text{O} - \text{C}(\text{CH}_3)_2\text{C}(\text{CH}_3)=\text{CH}$, 37), 111 (20), 84 (15)

HRMS calcd for $\text{C}_{16}\text{H}_{24}\text{O} + \text{H} - \text{H}_2\text{O}$: 215.1800, found: m/z 215.1789 (CI, ($\text{M}^+ + \text{H} - \text{H}_2\text{O}$), -1.1 mmu)

Phenyl-(2,2,3-trimethyl-1-phenyl-but-3-enyl)-amine (3j)

Reference : Sell, M. S.; Klein, W. R.; Rieke, R. D. *J. Org. Chem.* **1995**, *60*, 1077-1080.

Shape: colorless liquid

IR (neat) 3390 (N-H) cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 7.38-7.37 (m, 2H), 7.30-7.17 (m, 3H), 7.02-6.98 (m, 2H), 6.58 (t, $J=7.2$ Hz, 1H), 6.38 (dt, $J=3.9$ and 1.0 Hz, 2H), 4.99 (s, 2H), 4.17 (s, 1H), 4.11 (s, 1H), 1.75 (s, 3H), 1.03 (s, 3H), 0.99 (s, 3H)

^{13}C NMR (CDCl_3 , 100 MHz) δ 151.35, 147.92, 140.20, 128.95, 128.85, 127.71, 126.93, 117.11, 113.18, 112.20, 63.21, 43.55, 25.90, 20.62, 19.09

MS (EI, 70 eV) m/z 265 (M^+ , 0.19), 182(M, 100), 104($\text{M}^+ - \text{C}(\text{CH}_3)\text{C}(\text{CH}_3)=\text{CH}_2 - \text{C}_6\text{H}_5 - \text{H}$, 7), 77(12)

HRMS calcd for $\text{C}_{19}\text{H}_{23}\text{N}$: 265.1830, found: m/z 265.1832 (EI, (M^+), + 0.2 mmu)

2-Methyl-1-phenyl-but-3-en-1-ol (3k)

Reference : Kobayashi, S.; Nishio, K. *J. Org. Chem.* **1994**, *59*, 6620-6628.

4-Methyl-1-phenyl-hex-5-en-3-ol (3l)

Reference : Tan, K. T.; Chng, S. S.; Cheng, H. S.; Loh, T. P. *J. Am. Chem. Soc.* **2003**, *125*, 2958-2963.

3-Methyl-2-phenyl-pent-4-en-2-ol (3m)

Reference : Wada, R.; Oisaki, K.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2004**, *126*, 8910-8911.

3,4-Dimethyl-1-phenyl-hex-5-en-3-ol (3n)

Reference : Wada, R.; Oisaki, K.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2004**, *126*, 8910-8911.

(2-Methyl-1-phenyl-but-3-enyl)-phenyl-amine (3o)

Reference : (a) Andrade, C. K. Z.; Oliveira, G. R. *Tetrahedron Lett.* **2002**, 43, 1935-1937. (b) Meltzer, P. C.; Bickford, P. H.; Lambert, G. J. *J. Org. Chem.* **1985**, 50, 3121-3124.

2,3-Dimethyl-1-phenyl-but-3-en-1-ol (3p)

Reference : (a) Ishihara, K.; Mouri, M.; Gao, Q.; Maruyama, T.; Furuta, K.; Yamamoto, H. *J. Am. Chem. Soc.* **1993**, 115, 11490-11495. (b) Takahara, J. P.; Masuyama, Y.; Kurusu, Y. *J. Am. Chem. Soc.* **1992**, 114, 2577-2586.

3,4-Dimethyl-2-phenyl-pent-4-en-2-ol (3q)

Isolated as mixture of diastereomers (*dr* = 74:26)

Shape: colorless liquid

bp 90 °C/ 0.4 mmHg

IR (neat) 3498 (O-H), 1639 (aliphatic C=C), 1165 (C-O) cm⁻¹

¹H NMR (CDCl₃, 400 MHz) δ 7.44-7.40 (m, major+minor, 2H), 7.35-7.29 (m, major+minor, 2H), 7.24-7.19 (m, major+minor, 1H), 4.94 (s, major, 1H), 4.87 (s, minor)*, 4.86 (s, major, 1H), 4.76 (s, minor)*, 2.61 (q, *J*= 7.2 Hz, minor)*, 2.57 (q, *J*= 7.0 Hz, major, 1H), 2.29 (s, minor)*, 2.00 (s, major, 1H), 1.72 (s, major, 3H), 1.54 (s, minor)*, 1.51 (s, major, 3H), 1.39 (s, minor)*, 1.09 (d, *J*= 7.2 Hz, minor)*, 0.88 (d, *J*= 7.0 Hz, major, 3H)

¹³C NMR (CDCl₃, 100 MHz) δ 148.31*, 148.00*, 147.97, 147.46, 127.85, 127.74*, 126.37*, 126.25, 125.20*, 124.97, 113.32 (major+minor), 75.71, 75.08*, 51.44, 51.25*, 29.49, 26.68*, 23.44*, 22.57, 14.85*, 14.43

MS (EI, 70 eV) *m/z* 173 (M⁺ - OH, 0.4), 121 (M⁺ - CH(CH₃)C(CH₃)=CH₂, 100)

HRMS calcd for C₁₃H₁₈O + H: 191.1436, found: *m/z* 191.1453 (CI, (M⁺ + H), +1.7 mmu)

Anal. calcd for C₁₃H₁₈O: C, 82.06; H, 9.53, found: C, 81.77; H, 9.32.

* Mark corresponds to minor isomer.

(2,3-Dimethyl-1-phenyl-but-3-enyl)-phenyl-amine (3r)

Isolated as mixture of diastereomers (*dr* = 52:48)

Shape: white solid

mp 50-56 °C

IR (neat) 3398 (N-H), 1643 (aliphatic C=C) cm^{-1}

^1H NMR (CDCl_3 , 400 MHz) δ 7.40-7.19 (m, major+minor, 5H), 7.08-7.00 (m, major+minor, 2H), 6.64-6.57 (m, major+minor, 1H), 6.47-6.43 (m, major+minor, 2H), 4.96 (s, major, 1H), 4.92 (s, major+minor, 1H), 4.86 (s, minor)*, 4.45 (t, J = 4.3 Hz, minor)*, 4.32 (broad, major, 1H), 3.99 (broad, minor)*, 3.92 (dd, J = 9.9 and 2.9 Hz, major, 1H), 2.58 (qd, J = 7.0 and 4.3 Hz, minor)*, 2.43 (qd, J = 7.0 and 9.9 Hz, major, 1H), 1.75 (s, minor)*, 1.67 (s, major, 3H), 0.97 (d, J = 7.0 Hz, minor)*, 0.89 (d, J = 7.0 Hz, major, 3H)

* Mark corresponds to one isomer.

^{13}C NMR (CDCl_3 , 100 MHz) δ 147.79, 147.77, 147.64, 147.17, 143.11, 142.46, 128.92, 128.87, 128.30, 128.22, 127.45, 127.01, 126.82, 126.64, 117.21, 117.11, 113.44, 113.14, 112.04, 60.75, 59.08, 49.64, 46.97, 21.83, 17.29, 16.86, 12.88

One peak overlaps.

MS (EI, 70 eV) m/z 251 (M^+ , 0.72), 182 (M^+ - $\text{CH}(\text{CH}_3)\text{C}(\text{CH}_3)=\text{CH}_2$, 100)

HRMS calcd for $\text{C}_{18}\text{H}_{21}\text{N}$: 251.1674, found: m/z 251.1676 (EI, (M^+), +0.2 mmu)

Anal. calcd for $\text{C}_{18}\text{H}_{21}\text{N}$: C, 86.01; H, 8.42; N, 5.57, found: C, 86.00; H, 8.49, N, 5.57.

2,2-Dimethyl-1-phenyl-but-3-en-1-ol (3s)

Reference : Kobayashi, S.; Nishio, K. *J. Org. Chem.* **1994**, 59, 6620-6628.

2,2,3-Trimethyl-1-*p*-tolyl-but-3-en-1-ol

^1H NMR spectrum

^{13}C NMR spectrum

3c

1-(4-chlorophenyl)-2,2,3-trimethylbut-3-en-1-ol

^1H NMR spectrum

^{13}C NMR spectrum

2,2,3-Trimethyl-1-(4-nitro-phenyl)-but-3-en-1-ol

¹H NMR spectrum

¹³C NMR spectrum

1-(4-Methoxy-phenyl)-2,2,3-trimethyl-but-3-en-1-ol

^1H NMR spectrum

^{13}C NMR spectrum

4-(1-Hydroxy-2,2,3-trimethyl-but-3-enyl)-benzoic acid methyl ester

¹H NMR spectrum

¹³C NMR spectrum

3h

3,3,4-Trimethyl-2-phenyl-pent-4-en-2-ol

¹H NMR spectrum

¹³C NMR spectrum

2,3,3,4-Tetramethyl-1-phenyl-pent-4-en-2-ol

^1H NMR spectrum

^{13}C NMR spectrum

3j

Phenyl-(2,2,3-trimethyl-1-phenyl-but-3-enyl)-amine

¹H NMR spectrum

¹³C NMR spectrum

3,4-Dimethyl-2-phenyl-pent-4-en-2-ol

^1H NMR spectrum

^{13}C NMR spectrum

(2-Methyl-1-phenyl-but-3-enyl)-phenyl-amine

^1H NMR spectrum

^{13}C NMR spectrum