Joshua J. Kennedy-Smith, Steven T. Staben, and F. Dean Toste*

*Center for New Directions in Organic Synthesis, Department of Chemistry, University of California,
Berkeley, California 94720*

Supporting Information

General Information. Unless otherwise noted all commercial materials were used without purification. Small scale reactions (< 3mL) were carried out in Fisher Scientific disposable scintillation vials. Dichloromethane (CH_2Cl_2) and dichloroethane (DCE) were obtained from EMD. Silver trifluoromethanesulfonate (AgOTf) and chloro(triphenylphosphine)gold ($\text{AuCl(PPh}_3\text{)}$) were obtained from Aldrich Chemical Company and Strem Chemicals respectively. Substrates **1**, **3**, **5**, **7**, **23**, **25**, **27**, and **33** were prepared by alkylation of the corresponding acetoacetates or conjugate addition according to the method of Malacria.¹ Substrates **9** and **13** were prepared via intramolecular C-H insertion of the corresponding diazo compounds according to the method of Curran.² Substrates **11**, **15**, and **19** were prepared by conjugate addition to the appropriate enones according to the method of Malacria.³ Substrate **17** was prepared by addition of lithium dimethyl cuprate to the alkynyl substituted enone⁴ followed by trapping with Mander's reagent.⁵ Substrate **21** was prepared in three steps using the method of Stork and Danheiser according to Mattay,⁴ followed by L-Selectride reduction of the enone, and alkylation with Mander's reagent according to Winkler.⁶ Substrate **29** was prepared by alkylation of methyl acetoacetate according to the method of Malacria.⁷ Substrate **34** was prepared by exchange with D_2O and catalytic K_2CO_3 in THF. TLC analysis of reaction mixtures was performed on Merck silica gel 60 F_{254} TLC plates. Flash chromatography was carried out on Merck 60 silica gel (32-63 μm). ^1H and ^{13}C NMR spectra were recorded with Bruker AVB-400, and AVQ-400 spectrometers and referenced to CDCl_3 unless otherwise noted. Mass spectral and analytical data were obtained via the Micro-Mass/Analytical Facility operated by the College of Chemistry, University of California, Berkeley.

General procedure for Gold(I) catalyzed Conia-ene reaction. To a small screw-cap scintillation vial equipped with a magnetic stir bar and charged with a solution of alkynyl α -ketoester (~150 mg, 1 eq) in CH_2Cl_2 (0.4M) was added $\text{AuCl(PPh}_3\text{)}$ (1 mol %) followed by AgOTf (1 mol%). The cloudy white reaction

¹ Cruciani, P.; Stammler, R.; Aubert, C.; Malacria, M. *J. Org. Chem.* **1996**, *61*, 2699.

² Wang, C.; Gu, X.; Yu, M.S.; Curran, D.P. *Tetrahedron*. **1998**, *54*, 8355.

³ Renaud, J.L.; Petit, M.; Aubert, C.; Malacria, M. *Synlett* **1997**, 931.

⁴ Fagnoni, M.; Schmoldt, P.; Kirschberg, T.; Mattay, J. *Tetrahedron*. **1998**, *54*, 6427.

⁵ Laval, G.; Audran, G.; Galano, J.M.; Monti, H. *J. Org. Chem.* **2000**, *65*, 3551.

⁶ Winkler, J.D.; Axten, J.M. *J. Am. Chem. Soc.* **1998**, *120*, 6425.

⁷ Renaud, J.L.; Aubert, C.; Malacria, M. *Tetrahedron*. **1999**, *55*, 5113.

mixture was then stirred at room temperature and monitored periodically by TLC. Upon completion, the reaction mixture was loaded directly on to a silica gel column and chromatographed with the appropriate mixture of hexanes and EtOAc to give the Conia-ene products described below.

1-Acetyl-2-methylene-cyclopentanecarboxylic acid methyl ester (2). ^1H NMR (400 MHz, CDCl_3) δ 5.29 (t, $J = 2.0$, 1H), 5.21 (t, $J = 2.0$, 1H), 3.74 (s, 3H), 2.42 (m, 3H), 2.21 (s, 3H), 2.17 (m, 1H), 1.71 (m, 2H). The following nOe enhancements were observed:

^{13}C NMR (100 MHz, CDCl_3) δ 203.5, 171.6, 148.6, 112.2, 70.4, 52.6, 35.0, 33.9, 26.6, 24.1; HRMS (EI) calc. for $[\text{C}_9\text{H}_{10}\text{O}_2]^+$ ($[\text{M}-\text{MeOH}]^+$) 150.0681, found 150.0678.

1-Benzoyl-2-methylene-cyclopentanecarboxylic acid ethyl ester (4). ^1H NMR (400 MHz, CDCl_3) δ 7.85 (app d, $J = 7.4$, 2H), 7.54 (td, $J = 7.4$, 1.2, 1H), 7.42 (app t, $J = 7.4$, 2H), 5.38 (t, $J = 2.0$, 1H), 5.26 (t, $J = 2.0$, 1H), 4.15 (m, 2H), 2.87 (dt, $J = 13.6$, 7.0, 1H), 2.52-2.50 (m, 2H), 2.20 (dt, $J = 13.6$, 7.0, 1H), 1.88 (m, 1H), 1.72 (m, 1H), 1.07 (t, $J = 7.1$, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 195.4, 171.8, 149.48, 135.42, 132.66, 128.84, 128.40, 111.79, 67.47, 61.59, 36.80, 34.38, 24.34, 13.73; HRMS (EI) calc. for $[\text{C}_{16}\text{H}_{18}\text{O}_3]^+$ 258.1256, found 258.1254; Anal. Calcd.: C, 74.39; H, 7.02, found: C, 74.24; H, 7.12.

1-Acetyl-2-methylene-cyclopentanecarboxylic acid *tert*-butyl ester (6). ^1H NMR (400 MHz, CDCl_3) δ 5.27 (t, $J = 2.0$, 1H), 5.21 (t, $J = 2.0$, 1H), 2.41 (m, 3H), 2.20 (s, 3H), 2.09 (m, 1H), 1.67 (m, 2H), 1.45 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 203.6, 170.0, 148.8, 111.6, 81.8, 71.0, 34.9, 34.0, 27.7, 26.7, 23.9; Anal. Calcd.: C, 69.61; H, 8.99, found: C, 69.63; H, 9.11.

1-Acetyl-2-methylene-cyclopentanecarboxylic acid prop-2-ynyl ester (8). ^1H NMR (400 MHz, CDCl_3) δ 5.33 (app s, 1H), 5.28 (app s, 1H), 4.76 (m, 2H), 2.52-3.39 (m, 4H), 2.25 (s, 3H), 2.22 (td, $J = 13.0, 6.7$, 1H), 1.75 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 203.00, 170.33, 148.23, 112.55, 77.05, 75.23, 70.26, 52.80, 35.02, 33.96, 26.63, 24.09; Anal. Calcd.: C, 69.88; H, 6.84, found: C, 69.76; H, 6.96.

3-Methylene-4-oxo-hexahydro-pentalene-3a-carboxylic acid methyl ester (10). ^1H NMR (400 MHz, CDCl_3) δ 5.35 (t, $J = 2.4$, 1H), 5.20 (t, $J = 2.0$, 1H), 3.71 (s, 3H), 3.20 (m, 1H), 2.45 (m, 4H), 2.14 (m, 1H), 1.99 (m, 1H), 1.60 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 211.1, 170.3, 146.6, 112.2, 69.6, 52.7, 48.7, 37.5, 32.4, 29.5, 24.4; HRMS (EI) calc. for $[\text{C}_{11}\text{H}_{14}\text{O}_3]^+$ 194.0943, found 194.0943.

4-Methylene-3-oxo-octahydro-indene-3a-carboxylic acid methyl ester (12). ^1H NMR (400 MHz, CDCl_3) δ 5.08 (br s, 1H), 4.67 (br s, 1H), 3.74 (s, 3H), 2.96 (m, 1H), 2.42 (m, 1H), 2.34-2.24 (m, 2H), 2.06 (m, 1H), 1.89-1.70 (m, 3H), 1.68-1.58 (m, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 212.66, 170.66, 140.54, 114.24, 67.49, 52.76, 42.27, 36.66, 33.72, 24.97, 22.78, 22.42; LRMS m/z (M^+) 208.

6a-Acetyl-6-methylene-hexahydro-cyclopenta[c]furan-1-one (14). ^1H NMR (400 MHz, CDCl_3) δ 5.53 (t, $J = 2.0$, 1H), 5.29 (t, $J = 1.9$, 1H), 4.34 (dd, $J = 9.3, 7.9$, 1H), 3.94 (dd, $J = 9.3, 4.9$, 1H), 3.47 (m, 1H), 2.58-2.42 (m, 2H), 2.41 (s, 3H), 2.12 (m, 1H), 1.59 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 201.12, 173.70, 146.98, 112.47, 70.64, 70.26, 44.56, 33.64, 29.26, 25.82; LRMS m/z (M^+) 180; Anal. Calcd.: C, 66.65; H, 6.71, found: C, 66.72; H, 6.77.

3-Methylene-4-oxo-octahydro-indene-3a-carboxylic acid ethyl ester (16). ^1H NMR (400 MHz, CDCl_3) δ 5.21 (t, $J = 2.4$, 1H), 4.96 (t, $J = 2.4$, 1H), 4.20 (m, 2H), 3.02 (app. quintet, $J = 7.2$, 1H), 2.38 (m, 4H), 1.85 (m, 3H), 1.67 (m, 1H), 1.50 (m, 2H), 1.25 (t, $J = 7.0$, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 206.0, 170.6, 148.0, 111.7, 71.7, 61.5, 47.7, 39.5, 29.8, 28.0, 25.8, 23.8, 14.0; HRMS (EI) calc. for $[\text{C}_{13}\text{H}_{18}\text{O}_3]^+$ 222.1256, found 222.1256; Anal. Calcd.: C, 70.24; H, 8.16, found: C, 70.24; H, 8.18.

7a-Methyl-3-methylene-4-oxo-octahydro-indene-3a-carboxylic acid methyl ester (18). ^1H NMR (400 MHz, CDCl_3) δ 5.16 (t, $J = 2.2$, 1H), 4.93 (t, $J = 2.5$, 1H), 3.69 (s, 3H), 2.57 (m, 1H), 2.53 (m, 1H), 2.46-2.32 (m, 2H), 1.86 (m, 2H), 1.65 (m, 2H), 1.52 (m, 2H), 1.11 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 206.22, 170.61, 148.88, 112.17, 74.50, 65.81, 51.87, 49.88, 38.13, 36.38, 33.42, 29.25, 24.31, 20.62; HRMS (EI) calc. for $[\text{C}_{13}\text{H}_{18}\text{O}_3]^+$ 222.1254, found 222.1255; Anal. Calcd.: C, 70.24; H, 8.16, found: C, 70.27; H, 8.34.

3-Methylene-4-oxo-octahydro-azulene-3a-carboxylic acid methyl ester (20). ^1H NMR (400 MHz, CDCl_3) δ 5.34 (t, $J = 2.0$, 1H), 5.12 (t, $J = 2.0$, 1H), 3.76 (s, 3H), 2.89 (m, 1H), 2.44 (m, 4H), 2.06 (m, 1H), 1.85-1.40 (m, 7H); ^{13}C NMR (100 MHz, CDCl_3) δ 205.3, 171.9, 147.7, 112.7, 72.6, 52.7, 45.2, 42.4, 33.1, 32.8, 31.9, 27.3, 25.8; HRMS (EI) calc. for $[\text{C}_{13}\text{H}_{18}\text{O}_3]^+$ 222.1256, found 222.1253.

7-Methylene-2-oxo-bicyclo[3.2.1]octane-1-carboxylic acid methyl ester (22). ^1H NMR (400 MHz, CDCl_3) δ 5.64 (app t, $J = 2$, 1H), 5.16 (bs, 1H), 3.77 (s, 3H), 2.78 (m, 1H), 2.65 (m, 1H), 2.45 (m, 3H), 2.26 (ddd, $J = 16.0$, 6.4, 1.6, 1H), 1.50 (dd, $J = 11.6$, 2.4, 1H), 1.89 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 204.7, 169.6, 145.8, 110.9, 68.1, 52.0, 41.9, 38.1, 34.9, 31.7, 31.6; HRMS (EI) calc. for $[\text{C}_{11}\text{H}_{14}\text{O}_3]^+$ 194.0943, found 194.0936; Anal. Calcd.: C, 68.02; H, 7.27, found: C, 68.25; H, 7.39.

1-Acetyl-2-ethyl-5-methylene-cyclopentanecarboxylic acid methyl ester (24). Obtained as an inseparable 17:1 mixture of diastereomers. ^1H NMR (400 MHz, CDCl_3) Major diastereomer: δ 5.26 (t, J = 2.0, 1H), 5.19 (t, J = 2.2, 1H), 3.73 (s, 3H), 2.65 (m, 1H), 2.52 (m, 1H), 2.34 (m, 1H), 2.28 (s, 3H), 2.01 (m, 1H), 1.50 (m, 1H), 1.41 (m, 1H), 1.15 (m, 1H), 0.93 (t, J = 7.4, 3H). Additionally the following peaks were observed for the minor diastereomer: δ 5.24 (m, 1H), 5.21 (m, 1H), 3.77 (s, 3H), 2.16 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 202.34, 171.27, 149.59, 111.93, 73.66, 51.91, 48.12, 32.59, 28.52, 27.31, 23.61, 12.70; LRMS m/z ($\text{M}^+ - \text{OMe}$) 178; Anal. Calcd.: C, 68.54; H, 8.63, found: C, 68.92; H, 8.72.

1-Acetyl-2-methylene-5-phenyl-cyclopentanecarboxylic acid methyl ester (26). Obtained as an inseparable 4.2:1 mixture of diastereomers. ^1H NMR (400 MHz, CDCl_3) Major diastereomer: δ 7.34-7.15 (m, 5H), 5.42 (t, J = 2.0, 1H), 5.39 (t, J = 2.1, 1H), 4.33 (dd, J = 9.2, 7.0, 1H), 3.19 (s, 3H), 2.74-2.65 (m, 1H), 2.32 (s, 3H), 2.23-2.07 (m, 2H). Minor diastereomer: δ 7.34-7.15 (m, 5H), 5.28 (t, J = 2.1, 1H), 5.21 (t, J = 2.0, 1H), 4.23 (dd, J = 11.8, 6.3, 1H), 3.83 (s, 3H), 2.81-2.70 (m, 1H), 2.23-2.07 (m, 2H), 1.52 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) Major diastereomer: δ 200.93, 170.40, 148.32, 140.93, 128.45, 128.28, 126.83, 113.04, 75.76, 51.71, 50.23, 33.35, 29.91, 26.84. Additionally these peaks could be observed for the minor diastereomer: δ 204.60, 170.40, 149.76, 138.77, 111.67, 74.63, 52.72, 51.88, 32.37, 28.78.

1-Acetyl-4-benzyl-2-methylene-cyclopentanecarboxylic acid methyl ester (28). Obtained as an inseparable 4.0:1 mixture of diastereomers. ^1H NMR (400 MHz, CDCl_3) Major diastereomer: δ 7.26 (m, 2H), 7.15 (m, 3H), 5.24 (br m, 1H), 5.18 (m, 1H), 3.71 (s, 3H), 2.58 (m, 1H), 2.48 (m, 1H), 2.42-2.31 (m, 1H), 2.19 (s, 3H), 2.15-2.05 (m, 1H), 1.75 (dd, J = 12.9, 11.0, 1H). Additionally these peaks were observed for the minor diastereomer: δ 5.26 (m, 1H), 5.22 (app t, J = 1.7, 1H), 3.75 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) Major diastereomer: δ 203.29, 171.48, 147.53, 140.46, 128.62, 128.59, 128.34, 126.03, 112.80, 70.20, 52.72, 40.87, 40.82, 39.98, 26.68. Additionally these peaks were observed for the minor diastereomer: δ 203.03, 171.62, 147.81, 140.56, 112.41, 40.97, 40.34, 39.89, 38.95, 26.68; Anal. Calcd.: C, 74.97; H, 7.40, found: C, 75.04; H, 7.59.

1-Acetyl-2-methylene-3-propyl-cyclopentanecarboxylic acid methyl ester (30). Obtained as an inseparable 2.9:1 mixture of diastereomers. ^1H NMR (400 MHz, CDCl_3) Major diastereomer: δ 5.22 (app d, $J = 2.8$, 1H), 5.19 (app d, 2.4, 1H), 3.70 (s, 3H), 2.42 (m, 1H), 2.23 (m, 1H), 2.17 (s, 3H), 1.91 (m, 2H), 1.59 (m, 1H), 1.41-1.13 (m, 4H), 0.89 (t, $J = 7.1$, 3H). Additionally the following peaks were observed for the minor diastereomer: δ 5.23 (app d, $J = 2.6$, 1H), 5.21 (app d, $J = 2.6$, 1H), 3.71 (s, 3H), 2.18 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) Major diastereomer: δ 203.78, 171.77, 152.63, 111.45, 71.08, 52.61, 44.69, 35.86, 32.69, 30.39, 26.82, 20.72, 14.23. Additionally these peaks were observed for the minor diastereomer: 203.51, 172.01, 152.63, 11.81, 36.61, 32.80, 30.14, 26.50.