Supporting Information

Gold(I)-Catalyzed Intermolecular Addition of Phenols and Carboxylic Acids to Olefins

Cai-Guang Yang and Chuan He*

Department of Chemistry, the University of Chicago,
5735 S. Ellis Avenue, Chicago, IL 60617,
E-mail: chuanhe@uchicago.edu

Experimental Section

General Methods. Catalytic addition reactions were performed in sealed oven-dried glass tubes under an atmosphere of dry nitrogen. Flash column chromatography was performed with EM Science Geduran silica gel 60 (35-75 μ m), and eluting with hexanes/ethyl ether. Thin layer chromatography was performed on EM Science silica gel 60 F254 plates (250 μ m). NMR spectra were recorded on Varian XL-400/500 spectrometer for ^1H NMR and 100/125 MHz for ^{13}C NMR in CDCl_3 unless noted otherwise and were referenced to tetramethylsilane (TMS) at 0.00 ppm. The chemical shifts are expressed in ppm and coupling constants are given in Hz. Data for ^1H NMR are recorded as follows: chemical shift (ppm), multiplicity (s, singlet; d, doublet; t, triplet; m, multiplet), coupling constant (Hz), integration. Data for ^{13}C NMR are reported in terms of chemical shift (δ , ppm). Low-Resolution Mass spectra were obtained from the Mass Spectrometry Facility at the University of Chicago. High-Resolution Mass spectra were performed by Research Resources Center at University of Illinois at Chicago. Elemental analyses were performed by Microanalytical Laboratory at University of Illinois at Urbana-Champaign.

Chloro(triphenylphosphine)gold(I) and Silver triflate were purchased from Aldrich and used as received. Dry toluene was obtained from Across Company, and directly used. THF was freshly distilled over sodium/benzoquinone ketyl under dry nitrogen.

General Procedure for intermolecular addition of phenols and carboxylic acids to olefins: To a mixture of chloro(triphenylphosphine)gold(I) (10.0 or 25.0 mg, 0.02 or 0.05 mmol) and silver triflate (5.0 or 13.0 mg, 0.02 or 0.05 mmol) in dry toluene (2.0 mL) was added the olefin (4.0 mmol) and phenol (1.0 mmol) or carboxyl acid (1.0 mmol) with stirring in a sealed tube. After the reaction mixture was heated at 85 $^\circ\text{C}$ overnight (14 – 18 h), the mixture was directly loaded onto silica gel chromatography to give analytic pure products.

(*p*-Methoxyphenoxy)cyclohexane (3a). Colorless oil; ^1H NMR (500 MHz) δ 6.88 – 6.81 (m, 4H), 4.11 (m, 1H), 3.77 (s, 3H), 2.00 – 1.96 (m, 2H), 1.82 – 1.79 (m, 2H), 1.59 – 1.56 (m, 1H), 1.52 – 1.46 (m, 2H), 1.38 – 1.26 (m, 3H); ^{13}C NMR (125 MHz) δ 153.8, 151.7,

117.6, 114.5, 76.6, 55.6, 31.9, 25.6, 23.8; MS (m/z, EI) 206 (M)⁺; HRMS calcd for C₁₃H₁₈O₂: 206.1307; found: 206.1297.

(*p*-tert-Butylphenoxy)cyclohexane (3b).¹ Colorless oil; ¹H NMR (500 MHz) δ 7.28 (m, 2H), 6.84 (m, 2H), 4.20 (m, 1H), 2.01 – 1.98 (m, 2H), 1.83 – 1.79 (m, 2H), 1.60 – 1.48 (m, 3H), 1.40 – 1.30 (m, 3H), 1.30 (s, 9H); ¹³C NMR (125 MHz) δ 155.4, 143.0, 126.1, 115.3, 75.3, 34.0, 31.9, 31.5, 25.6, 23.8; MS (m/z, EI) 232 (M)⁺.

4-(Cyclohexyloxy)benzoic acid methyl ester (3c). Colorless oil; ¹H NMR (500 MHz) δ 7.96 (m, 2H), 6.89 (m, 2H), 4.33 (m, 1H), 3.87 (s, 3H), 2.00 – 1.97 (m, 2H), 1.84 – 1.79 (m, 2H), 1.61 – 1.51 (m, 3H), 1.43 – 1.30 (m, 3H); ¹³C NMR (125 MHz) δ 166.8, 161.6, 131.5, 121.9, 115.0, 75.2, 51.7, 31.5, 25.4, 23.6; MS (m/z, EI) 234 (M)⁺; Anal. Calcd. for C₁₄H₁₈O₃: C, 71.77; H, 7.74. Found C, 71.67; H, 7.72.

(*p*-Nitrophenoxy)cyclohexane (3d).² Pale yellow oil; ¹H NMR (500 MHz) δ 8.17 (m, 2H), 6.92 (m, 2H), 4.37 (m, 1H), 2.00 – 1.97 (m, 2H), 1.83 – 1.79 (m, 2H), 1.61 – 1.52 (m, 3H), 1.45 – 1.31 (m, 3H); ¹³C NMR (125 MHz) δ 163.1, 140.8, 125.8, 115.2, 76.1, 31.3, 25.3, 23.4; MS (m/z, EI) 221 (M)⁺.

2-(4-Methoxyphenoxy)-1-(4-methoxyphenyl)propane (3e). Colorless oil; ¹H NMR (500 MHz) δ 7.15 (m, 2H), 6.83 (m, 2H), 6.82 (m, 4H), 4.39 (m, 1H), 3.79 (s, 3H), 3.76 (s, 3H), 3.01 (dd, *J* = 13.5 and 5.5 Hz, 1H), 2.73 (dd, *J* = 13.5 and 6.5 Hz, 1H), 1.25 (d, *J* = 6.0 Hz, 3H); ¹³C NMR (125 MHz) δ 158.1, 153.9, 151.9, 130.4, 117.5, 114.6, 113.7, 76.1, 55.6, 55.2, 41.7, 19.4; MS (m/z, EI) 272 (M)⁺; Anal. Calcd. for C₁₇H₂₀O₃: C, 74.97; H, 7.40. Found C, 74.88; H, 7.36.

4-[2-(4-Methoxyphenyl)-1-methylethoxy]benzoic acid methyl ester (3f). Colorless oil; ^1H NMR (500 MHz) δ 7.96 (m, 2H), 7.14 (m, 2H), 6.89 (m, 2H), 6.83 (m, 2H), 4.61 (m, 1H), 3.88 (s, 3H), 3.78 (s, 3H), 3.03 (dd, $J = 13.5$ and 6.0 Hz, 1H), 2.80 (dd, $J = 13.5$ and 6.5 Hz, 1H), 1.31 (d, $J = 6.0$ Hz, 3H); ^{13}C NMR (125 MHz) δ 166.7, 161.7, 158.1, 131.5, 130.3, 129.6, 122.1, 114.9, 113.6, 74.7, 55.0, 51.7, 41.4, 19.0; MS (m/z , EI) 300 (M) $^+$; Anal. Calcd. for $\text{C}_{18}\text{H}_{20}\text{O}_4$: C, 71.98; H, 6.71. Found C, 72.14; H, 6.65.

2-(4-Nitrophenoxy)-1-(4-methoxyphenyl)propane (3g). Pale yellow oil; ^1H NMR (500 MHz) δ 8.17 (m, 2H), 7.14 (m, 2H), 6.90 (m, 2H), 6.84 (m, 2H), 4.65 (m, 1H), 3.78 (s, 3H), 3.03 (dd, $J = 14.0$ and 6.0 Hz, 1H), 2.84 (dd, $J = 14.0$ and 6.0 Hz, 1H), 1.35 (d, $J = 6.0$ Hz, 3H); ^{13}C NMR (125 MHz) δ 163.1, 158.3, 141.0, 130.3, 129.2, 125.9, 115.2, 113.8, 75.7, 55.2, 41.4, 19.1; MS (m/z , EI) 287 (M) $^+$; HRMS calcd for $\text{C}_{16}\text{H}_{17}\text{NO}_4$: 287.1158; found: 287.1154.

1-Methoxy-4-(1-methyl-3-phenylpropoxy)benzene (3h). Colorless oil; ^1H NMR (500 MHz) δ 7.30 – 7.25 (m, 2H), 7.21 – 7.18 (m, 3H), 6.82 (s, 4H), 4.24 (m, 1H), 3.77 (s, 3H), 2.82 (m, 1H), 2.74 (m, 1H), 2.09 – 2.02 (m, 1H), 1.90 – 1.85 (m, 1H), 1.29 (d, $J = 6.0$ Hz, 3H); ^{13}C NMR (125 MHz) δ 153.8, 152.0, 141.9, 128.4, 128.3, 125.7, 117.3, 114.5, 73.8, 55.6, 38.2, 31.7, 19.7; MS (m/z , EI) 256 (M) $^+$; HRMS calcd for $\text{C}_{17}\text{H}_{20}\text{O}_2$: 256.1463; found: 256.1473.

1-Bromo-4-(1-methyl-3-phenylpropoxy)benzene (3i). Colorless oil; ^1H NMR (500 MHz) δ 7.35 (m, 2H), 7.30 – 7.14 (m, 5H), 6.74 (m, 2H), 4.30 (m, 1H), 2.82 – 2.69 (m, 2H), 2.09 – 2.03 (m, 1H), 1.92 – 1.85 (m, 1H), 1.31 (d, J = 6.5 Hz, 3H); ^{13}C NMR (125 MHz) δ 157.2, 141.6, 132.3, 128.5, 128.46, 126.0, 117.6, 112.6, 73.0, 38.1, 31.7, 19.6; MS (m/z , EI) 304/306 (1 : 1) (M) $^+$; HRMS calcd for $\text{C}_{16}\text{H}_{17}\text{BrO}$: 304.0463; found: 304.0469.

1-Methyl -3-(phenylpropoxy)benzene (3j). Colorless oil; ^1H NMR (500 MHz) δ 7.32 – 7.28 (m, 4H), 7.23 – 7.19 (m, 3H), 6.97 – 6.94 (m, 1H), 6.91 – 6.88 (m, 2H), 4.39 (m, 1H), 2.84 (m, 1H), 2.75 (m, 1H), 2.10 (m, 1H), 1.91 (m, 1H), 1.35 (d, J = 6.0 Hz, 3H); ^{13}C NMR (125 MHz) δ 158.0, 141.8, 129.4, 128.5, 128.3, 125.8, 120.5, 115.8, 72.6, 38.2, 31.8, 19.7; MS (m/z , EI) 226 (M) $^+$; HRMS calcd for $\text{C}_{16}\text{H}_{18}\text{O}$: 226.1358; found: 226.1356.

Phenyl-acetic acid cyclohexyl ester (3k).³ Colorless oil; ^1H NMR (500 MHz) δ 7.33 – 7.23 (m, 5H), 4.77 (m, 1H), 3.59 (s, 2H), 1.83 – 1.79 (m, 2H), 1.69 – 1.66 (m, 2H), 1.52 – 1.49 (m, 1H), 1.44 – 1.27 (m, 4H), 1.25 – 1.21 (m, 1H); ^{13}C NMR (125 MHz) δ 170.9, 134.4, 129.1, 128.4, 126.9, 72.9, 41.7, 31.4, 25.3, 23.6; MS (m/z , EI) 218 (M) $^+$.

Phenyl-acetic acid 2-(4-methoxyphenyl)-1-methylentyl ester (3l). Colorless oil; ^1H NMR (500 MHz) δ 7.32 – 7.20 (m, 5H), 7.01 (m, 2H), 6.77 (m, 2H), 5.08 (m, 1H), 3.78 (s, 3H), 3.56 (s, 2H), 2.82 (dd, J = 13.5 and 8.5 Hz, 1H), 2.69 (dd, J = 13.5 and 6.5 Hz, 1H), 1.20 (d, J = 6.0 Hz, 3H); ^{13}C NMR (125 MHz) δ 171.0, 158.1, 134.1, 130.3, 129.4, 129.2, 128.4, 126.9, 113.6, 72.1, 55.1, 41.7, 41.2, 19.3; MS (m/z , ESI) 307.1 ($\text{M}+\text{Na}$) $^+$; HRMS calcd for $\text{C}_{18}\text{H}_{20}\text{O}_3\text{Na}$: 307.1310; found: 307.1308.

Phenyl-acetic acid 1-methyl-3-phenylpropyl ester (3m). Colorless oil; ^1H NMR (500 MHz) δ 7.36 – 7.31 (m, 4H), 7.29 – 7.25 (m, 3H), 7.18 (m, 1H), 7.09 (m, 2H), 4.94 (m, 1H), 3.60 (s, 2H), 2.56 (m, 2H), 1.93 – 1.87 (m, 1H), 1.83 – 1.78 (m, 1H), 1.25 (d, J = 6.0 Hz, 3H); ^{13}C NMR (125 MHz) δ 171.2, 141.5, 134.3, 129.2, 128.5, 128.4, 128.3, 127.0, 125.9, 70.9, 41.8, 37.6, 31.6, 20.0; MS (m/z , EI) 268 (M) $^+$; Anal. Calcd. for $\text{C}_{18}\text{H}_{20}\text{O}_2$: C, 80.56; H, 7.51. Found C, 80.53; H, 7.34.

Phenyl-acetic acid 1-methylhexyl ester (3n) and Phenyl-acetic acid 1-ethylpentyl ester (3n'). Colorless oil; ^1H NMR (500 MHz) δ 7.33 – 7.23 (m, 5H), 4.90 (m, 1H for **3n**), 4.81 (m, 1H for **3n'**), 3.60 (s, 2H for **3n'**), 3.59 (s, 2H for **3n**), 1.60 – 1.41 (m, 2H for **3n** and 4H for **3n'**), 1.30 – 1.16 (m, 6H for **3n** and 4H for **3n'**), 1.19 (d, J = 6.0 Hz, 3H for **3n**), 0.86 (t, J = 7.0 Hz, 3H for **3n**), 0.84 (t, J = 7.5 Hz, 3H for **3n'**), 0.82 (t, J = 7.5 Hz, 3H for **3n'**); ^{13}C NMR (125 MHz) δ [171.4], 171.2, [134.4], 134.3, [129.2], 129.1, 128.4, 126.8, [75.9], 71.4, 41.7, 35.8, [31.2], 31.5, [27.3], [26.9], 24.9, 22.5, 19.9, 13.9, [9.4]; MS (m/z , EI) 234 (M) $^+$; HRMS calcd for $\text{C}_{15}\text{H}_{22}\text{O}_2$: 234.1620; found: 234.1613.

Isobutyric acid 2-(4-methoxyphenyl)-1-methylethyl ester (3o). Colorless oil; ^1H NMR (500 MHz) δ 7.11 (m, 2H), 6.82 (m, 2H), 5.06 (m, 1H), 3.78 (s, 3H), 2.84 (dd, J = 13.5 and 7.0 Hz, 1H), 2.70 (dd, J = 13.5 and 6.0 Hz, 1H), 2.46 (hept, J = 7.0 Hz, 1H), 1.20 (d, J = 6.0 Hz, 3H), 1.11 (d, J = 7.0 Hz, 3H), 1.08 (d, J = 7.0 Hz, 3H); ^{13}C NMR (125 MHz) δ 176.6, 158.2, 130.4, 129.7, 113.6, 71.2, 55.2, 41.3, 34.1, 19.4, 18.9, 18.8; MS (m/z , ESI) 259.1 ($\text{M}+\text{Na}$) $^+$; HRMS calcd for $\text{C}_{14}\text{H}_{20}\text{O}_3\text{Na}$: 259.1310; found: 259.1305.

4-Methoxy-benzoic acid bicyclo[2.2.1]hept-2-yl ester (3p).⁴ White solid; mp 63 – 65 °C; ¹H NMR (500 MHz) δ 7.96 (m, 2H), 6.90 (m, 2H), 4.81 (m, 1H), 3.85 (s, 3H), 2.42 (d, J = 5.0 Hz, 1H), 2.31 (br s, 1H), 1.81 (ddd, J = 13.5, 7.0 and 2.0 Hz, 1H), 1.63 (dt, J = 9.5 and 1.5 Hz, 1H), 1.58 – 1.54 (m, 2H), 1.50 – 1.42 (m, 1H), 1.23 – 1.20 (m, 2H), 1.18 – 1.12 (m, 1H); ¹³C NMR (125 MHz) δ 165.8, 163.1, 131.4, 123.2, 113.4, 77.7, 55.3, 41.5, 39.6, 35.3, 28.1, 24.2; MS (m/z , EI) 246 (M)⁺.

2,2-Diphenyl-4-hexen-1-ol (Table 2, entry 2). This substrate was prepared according to a known procedure.⁴ For *E*: ¹H NMR (500 MHz) δ 7.28 – 7.25 (m, 4H), 7.20 – 7.15 (m, 6H), 5.48 (m, 1H), 5.04 (m, 1H), 4.09 (d, J = 6.5 Hz, 2H), 2.87 (d, J = 7.0 Hz, 2H), 1.55 (d, J = 6.5 Hz, 3H), 1.27 (t, J = 7.0 Hz, 1H); ¹³C NMR (125 MHz) δ 145.5, 128.6, 128.2, 128.1, 128.0, 126.6, 126.2, 67.9, 51.7, 39.7, 17.9; MS (m/z , EI) 235 (M - OH)⁺.

Catalytic intramolecular addition of hydroxyl-olefin:

A mixture of chloro(triphenylphosphine)gold(I) (25.0 mg, 0.05 mmol) and silver triflate (13.0 mg, 0.05 mmol) in dry toluene (2.0 mL) was stirred at r.t. for 20 min, and then the white precipitates were filtered off. To the filtrate was added 2,2-diphenyl-4-hexen-1-ol (252 mg, 1.0 mmol) with stirring in a sealed tube. The mixture was heated at 85 °C for 16 h, and the ratio of the two isomers was determined by crude ¹H NMR analysis. Then, the product mixture was directly loaded onto silica gel chromatography to give the analytic pure product (224 mg, 89% combined yield).

2-Ethyl-4,4-diphenyltetrahydrofuran (Scheme 1).⁵ Colorless oil; ¹H NMR (400 MHz) δ 7.33 – 7.18 (m, 10H), 4.62 (dd, J = 8.7 and 0.8 Hz, 1H), 4.12 (d, J = 8.7 Hz, 1H), 3.96 (m, 1H), 2.60 (ddd, J = 12.0 and 5.7 and 1.0 Hz, 1H), 2.28 (dd, J = 12.0 and 9.6 Hz, 1H), 1.68 (m, 1H), 1.56 (m, 1H), 0.94 (t, J = 7.5 Hz, 3H); ¹³C NMR (100 MHz) δ 146.4, 146.0, 128.3, 128.2, 127.1, 126.3, 126.1, 80.1, 76.7, 55.9, 44.4, 28.9, 10.3; MS (m/z , EI) 252 (M)⁺.

2-Methyl-5,5-diphenyltetrahydropyran (Scheme 1).⁵ Colorless oil; ^1H NMR (500 MHz) δ 7.43 - 7.41 (m, 2H), 7.28 - 7.23 (m, 4H), 7.17 - 7.16 (m, 4H), 4.63 (dd, J = 12.5 and 3.0 Hz, 1H), 3.56 (d, J = 12.5 Hz, 1H), 3.56 (m, 1H), 2.48 - 2.38 (m, 2H), 1.56 - 1.52 (m, 1H), 1.29 - 1.18 (m, 1H), 1.16 (d, J = 6.0 Hz); ^{13}C NMR (125 MHz) δ 146.7, 145.8, 128.9, 128.2, 127.9, 126.9, 126.2, 125.5, 75.0, 73.9, 45.7, 34.7, 29.7, 21.6; MS (m/z , EI) 252 (M)⁺.

Catalytic terminal olefin migration:

To a mixture of chloro(triphenylphosphine)gold(I) (25.0 mg, 0.05 mmol) and silver triflate (13.0 mg, 0.05 mmol) in dry toluene (2.0 mL) was added 4-phenyl-1-butene (528 mg, 4.0 mmol) with stirring in a sealed tube. The reaction was heated for 20 h at 85 °C, and the mixture was purified by silica gel flash chromatography. ^1H NMR analysis indicated that it contained 25% of 4-phenyl-1-butene and 75% of 1-phenyl-2-butene (E/Z = 2.2 : 1). The spectral data were in good agreement with the literature values.⁶ ***E*-isomer:** ^1H NMR (500 MHz) δ 7.35 - 7.23 (m, 5H), 5.66 - 5.55 (m, 2H), 3.37 (d, J = 6.5 Hz, 2H), 1.74 (dd, J = 6.0 and 1.0 Hz, 3H); ***Z*-isomer:** ^1H NMR (500 MHz) δ 7.35 - 7.23 (m, 5H), 5.66 - 5.55 (m, 2H), 3.46 (d, J = 5.5 Hz, 2H), 1.78 (d, J = 5.0 Hz, 3H).

Reference:

1. Dalglish, D. T.; Nonhebel, D. C.; Pauson, P. L. *J. Chem. Soc. C* **1971**, 1174-1176.
2. Carrigan, C. N.; Bartlett, R. D.; Esslinger, C. S.; Cybulski, K. A.; Tongcharoensirikul, P.; Bridges, R. J.; Thompson, C. M. *J. Med. Chem.* **2002**, 45, 2260 -2276.
3. Barluenga, J.; Alonso-Cires, L.; Campos, P. J.; Asensio, G. *Synthesis* **1983**, 649-651.
4. Oe, Y.; Ohta, T.; Ito, Y. *Chem. Commun.* **2004**, 1620-1621.
5. Qian, H.; Han, X.; Widenhoefer, R. A. *J. Am. Chem. Soc.* **2004**, 126, 9536-9537.
6. (a) Arnold, D. R.; Mines, S. A. *Can. J. Chem.* **1989**, 67, 689-698. (b) Crich, D.; Mo, X.-S. *J. Am. Chem. Soc.* **1998**, 120, 8298-8304.