Supporting Information for:

A Gold(I)-Catalyzed Propargyl Claisen Rearrangement

Benjamin D. Sherry and F. Dean Toste*

Center for New Directions in Organic Synthesis, Department of Chemistry, University of California, Berkeley, California 94720

General Information: Unless otherwise noted all commercial materials were used without further purification. Claisen rearrangements were conducted in two dram vials equipped with a magnetic stir bar and fitted with a threaded cap. Dichloromethane (CH_2Cl_2) and methanol (MeOH) were obtained from EMD. Silver tetrafluoroborate (AgBF_4), silver trifluoromethanesulfonate (AgOTf), and sodium borohydride (NaBH_4) were obtained from Aldrich Chemical Company. Triphenylphosphinegold(I) chloride (Ph_3PAuCl) was prepared according to the method of Bruce.¹ Tris(triphenylphosphinegold)-oxonium tetrafluoroborate ($[(\text{Ph}_3\text{PAu})_3\text{O}]\text{BF}_4$) was prepared according to the method of Sharp.² Bis(triphenylphosphine)gold(I) tetrafluoroborate ($[(\text{Ph}_3\text{P})_2\text{Au}]\text{BF}_4$) was prepared according to the method of Westland.³ α -Aurated acetaldehyde ($\text{Ph}_3\text{PAuCH}_2\text{CHO}$) was prepared according to the method of Nesmeyanov.⁴ Vinyl ethers **1**, (**R**)-**4a**, (**R**)-**4j**, and (**R**)-**4n** were prepared by reduction of the corresponding ynone according to the method of Midland,⁵ followed by vinylation of the resulting propargyl alcohol according to the method of Yamamoto.⁶ Vinyl ethers **4a-n** were prepared from the corresponding propargyl alcohol according to the method of Yamamoto.⁶ Thin-layer chromatography (TLC) analysis of reaction mixtures was performed using Merck silica gel 60 F_{254} TLC plates and visualized by UV and permanganate staining. Flash column chromatography was carried out on Merck 60 silica gel (32-63 μm). ^1H and ^{13}C NMR spectra were recorded with Bruker DRX-500, AVB-400, AVQ-400, and AV-300 spectrometers and referenced to CDCl_3 unless otherwise noted. Enantiomeric excess was determined either on an Agilent 1100 Series Chiral HPLC with detection at 210, 254, and 280 nm or a Shimadzu VP Series Chiral HPLC with detection at 210, 254, and 280 nm. Mass spectral and analytical data were obtained via the Micro-Mass/Analytical Facility operated by the College of Chemistry, University of California, Berkeley.

Catalyst Optimization Experiments:

To a solution of **1** (0.100 g, 0.467 mmol, 95% ee) in CH_2Cl_2 (1.3 mL) was added Ph_3PAuCl (2.3 mg, 4.7 μmol) followed by AgOTf (1.2 mg, 4.7 μmol) and the reaction mixture was left to stir at rt for 10 min. The solution was diluted with MeOH (2.6 mL), NaBH_4 (17.7 mg, 0.467 mmol) was added, and the reaction mixture was maintained at rt for 30 min. The red solution was concentrated and a crude ^1H NMR (400 MHz, C_6D_6) spectrum was obtained: A 1.7:1.0 ratio of **2**:**3** was observed. The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford homoallenic alcohol **2** (40 mg, 40% yield, R_f = 0.20 in 9:1 hexanes:EtOAc) as a colorless oil and alcohol **3** (24 mg, 24% yield, R_f = 0.20 in 9:1 hexanes:EtOAc) as a colorless oil. Characterization for **2**: Spectral data were consistent with those reported below; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 97:3 hexanes:isopropanol, 1 mL/min; t_R 9.30 min (minor), 12.76 min (major): 83% ee. Characterization for **3**: ^1H NMR (400 MHz, CDCl_3) δ 7.37 (d, 2H, J = 7.2 Hz), 7.33-7.26 (m, 3H), 3.89-3.80 (m, 2H), 3.77-3.71 (m, 2H), 2.23 (td, 2H, J = 6.8, 2.0 Hz), 2.02-1.87 (m, 2H), 1.80 (br s, 1H), 1.55-1.48 (m, 2H), 1.45-1.31 (m, 2H), 0.91 (t, 2H, J = 7.2 Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 142.3, 128.5, 127.4, 126.7, 84.1, 81.1, 61.1, 41.3, 34.7, 31.1, 22.0, 18.5, 13.7; HRMS (EI^+) calc'd for $[\text{C}_{15}\text{H}_{20}\text{O}]^+$: m/z 216.1514, found 216.1517.

To a suspension of AgBF_4 (0.9 mg, 4.7 μmol) in CH_2Cl_2 (0.4 mL) was added Ph_3PAuCl (2.3 mg, 4.7 μmol) and the resulting cloudy mixture as stirred rapidly at rt for 5 min. The mixture was filtered through a cotton plug into a

stirred solution of **1** (0.100 g, 0.467 mmol, 95% ee) in CH₂Cl₂ (0.9 mL) and the resulting mixture was left to stir at rt for 10 min. The reaction mixture was diluted with MeOH (2.6 mL), NaBH₄ (17.7 mg, 0.467 mmol) was added, and the solution was left to stir at rt for 1 h. The red solution was concentrated and the crude oil was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallenlic alcohol **2** (86 mg, 85% yield, *R_f* = 0.38 in 3:1 hexanes:EtOAc) as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 7.35-7.31 (m, 4H), 7.25-7.19 (m, 1H), 6.23 (quintet, 1H, *J* = 3.2 Hz), 3.80 (t, 2H, *J* = 6.4 Hz), 2.47-2.32 (m, 2H), 2.19-2.13 (m, 2H), 1.79 (br s, 1H), 1.57-1.48 (m, 2H), 1.38 (sextet, 2H, *J* = 7.2 Hz), 0.93 (t, 3H, *J* = 7.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 201.9, 135.4, 128.7, 126.8, 126.5, 105.8, 96.1, 60.9, 35.9, 32.8, 29.8, 22.5, 13.9; HRMS (EI⁺) calc'd for [C₁₅H₂₀O]⁺: *m/z* 216.1514, found 216.1513; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 97:3 hexanes:isopropanol, 1 mL/min; *t_R* 14.53 min (minor), 27.10 min (major): 4% ee.

To a solution of **1** (0.100 g, 0.467 mmol, 95% ee) in CH₂Cl₂ (1.3 mL) was added [(Ph₃PAu)₃O]BF₄ (2.3 mg, 1.6 μ mol) and the resulting solution was left to stir at rt for 10 min. The reaction mixture was diluted with MeOH (2.6 mL), NaBH₄ (17.7 mg, 0.467 mmol) was added, and the solution was left to stir at rt for 1 h. The red solution was concentrated and the crude oil was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallenlic alcohol **2** (92 mg, 91% yield, *R_f* = 0.14 in 9:1 hexanes:EtOAc) as a colorless oil: Spectral data were consistent with those reported for **2**; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 97:3 hexanes:isopropanol, 1 mL/min; *t_R* 14.33 min (minor), 26.92 min (major): 90% ee.

General Procedure for the Au(I)-Catalyzed Claisen Rearrangement: To a solution of propargyl vinyl ether (1 equiv) in CH₂Cl₂ (0.35 M) was added [(Ph₃PAu)₃O]BF₄ (1 mol%) and the resulting mixture was maintained at rt until TLC analysis indicated consumption of the starting material. The reaction mixture was diluted with MeOH (2x the volume of CH₂Cl₂) and NaBH₄ (1.0 equiv) was added. The resulting red solution was maintained at rt for a minimum of 1 h, concentrated, and purified by flash column chromatography. For reactions carried out on a scale greater than 0.500 g the reaction mixture after reduction was diluted with H₂O, extracted with Et₂O (3x), washed with brine (1x), dried over MgSO₄, filtered, and concentrated. The crude residue was purified by flash column chromatography.

5a (Table 1, entry 1)

The crude mixture was purified by flash column chromatography (7:1 hexanes:EtOAc eluent) to afford the desired homoallenlic alcohol (45 mg, 90% yield, *R_f* = 0.24 in 3:1 hexanes:EtOAc) as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 7.33-7.29 (m, 4H), 7.23-7.17 (m, 1H), 6.19 (m, 1H), 5.60 (q, 1H, *J* = 6.8 Hz), 3.77 (t, 2H, *J* = 6.4 Hz), 2.39 (m, 2H), 1.78 (br s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 205.7, 134.4, 128.6, 126.9, 126.6, 95.1, 91.5, 61.9, 32.0.

5b (Table 1, entry 2)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallenlic alcohol (89 mg, 89% yield, *R_f* = 0.12 in 9:1 hexanes:EtOAc) as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 7.28-7.22 (m, 4H), 7.17-7.13 (m, 1H), 6.18 (quintet, 1H, *J* = 3.2 Hz), 3.74 (t, 2H, *J* = 6.4 Hz), 3.61 (t, 2H, *J* = 6.4 Hz), 2.39-2.28 (m, 2H), 2.18-2.12 (m, 2H), 1.84 (br s, 1H), 1.72-1.65 (m, 2H), 0.86 (s, 9H), 0.00 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 201.7, 135.2, 128.6, 126.8, 126.4, 105.5, 96.4, 62.6, 60.8, 36.1, 30.6, 29.1, 25.9, 18.3, -5.3; HRMS (EI⁺) calc'd for [C₂₀H₃₂O₂Si]⁺: *m/z* 332.2171, found 332.2174; Anal calc'd for C₂₀H₃₂O₂Si: C, 72.23; H, 9.70; found: C, 71.76; H, 9.91.

5c (Table 1, entry 3)

The crude mixture was purified by flash column chromatography (4:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (82 mg, 81% yield, $R_f = 0.21$ in 3:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.30-7.26 (m, 4H), 7.23-7.18 (m, 1H), 6.32 (quintet, 1H, $J = 2.4$ Hz), 4.69 (dd, 1H, $J = 16.0, 2.8$ Hz), 4.63 (dd, 1H, $J = 14.4, 2.8$ Hz), 3.81 (t, 3H, $J = 6.0$ Hz), 2.39-2.37 (m, 2H), 2.10 (br s, 1H), 1.15 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 202.4, 178.3, 133.9, 128.7, 127.3, 126.8, 102.2, 97.8, 63.8, 60.9, 38.8, 33.0, 27.1; HRMS (EI^+) calc'd for $[\text{C}_{17}\text{H}_{22}\text{O}_3]^+$: m/z 274.1569, found 274.1562; Anal calc'd for $\text{C}_{17}\text{H}_{22}\text{O}_3$: C, 74.42; H, 8.08; found: C, 74.12; H, 8.11.

5d (Table 1, entry 4)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (90 mg, 89% yield, $R_f = 0.37$ in 3:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.20 (d, 2H, $J = 8.4$ Hz), 6.84 (d, 2H, $J = 8.8$ Hz), 6.15 (quintet, 1H, $J = 2.8$ Hz), 3.78 (s, 3H), 3.75 (t, 2H, $J = 6.4$ Hz), 2.39-2.27 (m, 2H), 2.10-2.07 (m, 2H), 1.83 (br s, 1H), 1.50-1.44 (m, 2H), 1.35 (sextet, 2H, $J = 7.2$ Hz), 0.88 (t, 3H, $J = 7.2$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 201.1, 158.7, 127.7, 127.5, 114.2, 105.7, 95.6, 61.0, 55.3, 36.0, 32.9, 29.8, 22.5, 14.0.

5e (Table 1, entry 5)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (87 mg, 86% yield, $R_f = 0.12$ in 9:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.33 (d, 2H, $J = 8.0$ Hz), 7.07 (d, 2H, $J = 8.0$ Hz), 5.87 (sextet, 1H, $J = 2.8$ Hz), 3.45 (t, 2H, $J = 6.0$ Hz), 2.02 (bt, 2H, $J = 6.4$ Hz), 1.56 (d, 3H, $J = 2.8$ Hz), 1.43 (br s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 203.8, 139.6, 127.7 (t, $J =$), 126.8, 125.5, 125.4, 101.3, 93.4, 60.2, 36.8, 18.2; HRMS (EI^+) calc'd for $[\text{C}_{13}\text{H}_{13}\text{F}_3\text{O}]^+$: m/z 242.0918, found 242.0919; Anal calc'd for $\text{C}_{13}\text{H}_{13}\text{F}_3\text{O}$: C, 64.46; H, 5.4; found: C, 64.29; H, 5.58.

5f (Table 1, entry 6)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (97 mg, 96% yield, $R_f = 0.36$ in 3:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.50 (dd, 1H, $J = 8.0, 1.2$ Hz), 7.42 (dd, 1H, $J = 7.6, 1.6$ Hz), 7.22 (td, 1H, $J = 7.6, 1.2$ Hz), 7.01 (td, 1H, $J = 8.0, 1.6$ Hz), 6.62 (quintet, 1H, $J = 3.2$ Hz), 3.76 (t, 2H, $J = 6.0$ Hz), 2.37-2.32 (m, 2H), 2.13-2.08 (m, 2H), 1.68 (br s, 1H), 1.49-1.42 (m, 2H), 1.39-1.31 (m, 2H), 0.88 (t, 3H, $J = 7.6$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 203.1, 134.7, 133.1, 128.0 (2), 127.5, 122.4, 106.0, 95.0, 60.8, 35.8, 32.6, 29.7, 22.5, 13.9; HRMS (EI^+) calc'd for $[\text{C}_{15}\text{H}_{19}\text{BrO}]^+$: m/z 294.0619, found 294.0615; Anal calc'd for $\text{C}_{15}\text{H}_{19}\text{BrO}$: C, 61.03; H, 6.49; found: C, 60.76; H, 6.55.

5g (Table 1, entry 7)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (94 mg, 93% yield, $R_f = 0.15$ in 9:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, C_6D_6) δ 7.50 (d, 2H, $J = 7.6$ Hz), 7.27-7.24 (m, 2H), 7.13 (t, 1H, $J = 7.6$ Hz), 5.45 (septet, 1H, $J = 2.8$ Hz), 3.81 (t, 2H, $J = 6.8$ Hz), 2.67-2.63 (m, 2H), 2.03 (q, 2H, $J = 7.6$ Hz), 1.72 (br s, 1H), 1.48-1.40 (m, 2H), 1.30-1.22 (m, 4H), 0.92 (t, 3H, $J = 6.8$ Hz); ^{13}C NMR (100 MHz, C_6D_6) δ 203.8, 137.0, 128.4, 126.6, 126.0, 102.7, 94.8, 61.1, 33.4, 31.4, 29.0 (2), 22.5, 14.0; HRMS (EI^+) calc'd for $[\text{C}_{16}\text{H}_{22}\text{O}]^+$: m/z 230.1671, found 230.1667; Anal calc'd for $\text{C}_{16}\text{H}_{22}\text{O}$: C, 83.43; H, 9.63; found: C, 81.69; H, 9.80.

5h (Table 1, entry 8)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (88 mg, 87% yield, R_f = 0.20 in 9:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, C_6D_6) δ 7.50 (d, 2H, J = 7.2 Hz), 7.28-7.23 (m, 2H), 7.13 (t, 1H, J = 7.2 Hz), 5.50-5.46 (m, 1H), 3.79 (t, 2H, 6.6 Hz), 2.63 (td, 2H, J = 6.6, 3.6 Hz), 2.34-2.27 (septet, 1H, J = 6.6 Hz), 1.59 (br s, 1H), 1.05 (d, 6H, J = 6.6 Hz); ^{13}C NMR (100 MHz, C_6D_6) δ 202.4, 137.4, 128.7, 127.0, 126.2, 104.3, 102.4, 61.4, 33.8, 29.0, 22.7; HRMS (EI^+) calc'd for $[\text{C}_{14}\text{H}_{18}\text{O}]^+$: m/z 202.1358, found 202.1359.

5i (Table 1, entry 9)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (77 mg, 76% yield, R_f = 0.42 in 9:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 5.29-5.24 (m, 1H), 4.19-4.09 (m, 2H), 3.76-3.70 (m, 1H), 3.65-3.59 (m, 1H), 2.25-2.17 (m, 1H), 2.16-2.08 (m, 1H), 1.97-1.89 (m, 2H), 1.45-1.26 (m, 4H), 0.88 (t, 3H, J = 7.2 Hz), 0.87 (s, 9H), 0.06 (s, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 199.5, 104.4, 93.4, 60.9, 60.0, 35.7, 32.6, 29.7, 25.8, 22.4, 18.4, 14.0, -5.4; HRMS (FAB) calc'd for $[\text{C}_{16}\text{H}_{32}\text{O}_2\text{SiLi}]^+$: m/z 291.2332, found 291.2327; Anal calc'd for $\text{C}_{16}\text{H}_{32}\text{O}_2\text{Si}$: C, 67.54; H, 11.34; found: C, 65.58; H, 10.78.

5j (Table 1, entry 10)

The crude mixture was purified by flash column chromatography (10:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (84 mg, 84% yield, R_f = 0.40 in 3:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.31-7.27 (m, 2H), 7.21-7.18 (m, 3H), 5.22-5.15 (m, 1H), 3.72 (t, 2H, J = 6.0 Hz), 2.65 (t, 2H, J = 7.6 Hz), 2.23-2.18 (m, 2H), 2.00 (td, 2H, J = 7.2, 2.8 Hz), 1.86 (br s, 1H), 1.76 (quintet, 2H, J = 7.6 Hz), 1.68 (d, 3H, J = 6.8 Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 201.5, 142.5, 128.5, 128.3, 125.8, 100.3, 87.7, 60.9, 35.9, 35.5, 32.3, 29.3, 15.1; HRMS (EI^+) calc'd for $[\text{C}_{15}\text{H}_{20}\text{O}]^+$: m/z 216.1514, found 216.1513; Anal calc'd for $\text{C}_{15}\text{H}_{20}\text{O}$: C, 83.28; H, 9.32; found: C, 83.0; H, 9.60.

5k (Table 1, entry 11)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (91 mg, 90% yield, R_f = 0.24 in 9:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 5.20-5.16 (m, 1H), 3.72 (t, 2H, J = 6.4 Hz), 2.30-2.26 (m, 2H), 1.92 (q, 2H, J = 7.6 Hz), 1.80 (s, 1H), 1.37-1.26 (m, 6H), 1.10-1.06 (m, 1H), 0.86 (t, 3H, J = 6.4 Hz), 0.64-0.60 (m, 2H), 0.38-0.31 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 199.3, 104.1, 94.2, 61.0, 36.0, 31.3, 29.4, 28.9, 22.4, 14.0, 12.6, 6.7, 6.2; HRMS (EI^+) calc'd for $[\text{C}_{13}\text{H}_{22}\text{O}]^+$: m/z 194.1671, found 194.1669.

5l (Table 1, entry 12)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (92 mg, 91% yield, R_f = 0.36 in 3:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.30-7.25 (m, 2H), 7.20-7.16 (m, 3H), 3.60 (t, 2H, J = 6.4 Hz), 2.72 (t, 2H, J = 7.6 Hz), 2.27 (t, 2H, J = 8.4 Hz), 2.12 (t, 3H, J = 6.4 Hz), 1.71 (s, 3H), 1.64 (br s, 1H), 1.61 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 198.8, 142.1, 128.4, 128.3, 125.8, 99.0, 95.9, 60.9, 37.4, 36.0, 34.0, 19.6 (2); Anal calc'd for $\text{C}_{15}\text{H}_{20}\text{O}$: C, 83.28; H, 9.32; found: C, 83.02; H, 9.05.

5m (Table 1, entry 13)

The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallenic alcohol (31 mg, 61% yield, R_f = 0.13 in 9:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.29-7.24 (m, 2H), 7.18-7.16 (m, 3H), 3.70 (t, 2H, J = 6.0 Hz), 2.62 (t, 2H, J = 7.6 Hz), 2.17 (t, 2H, J = 6.0 Hz), 2.11 (t, 4H, J = 5.6 Hz), 1.96 (t, 2H, J = 7.2 Hz), 1.73 (quintet, 2H, J = 7.2 Hz), 1.64-1.49 (m, 7H); ^{13}C NMR (100 MHz, CDCl_3) δ 194.6, 142.6, 128.5, 128.3, 125.7, 104.9, 98.4, 61.1, 36.3, 35.3, 32.6, 32.2, 29.3, 27.8, 26.2; HRMS (EI^+) calc'd for $[\text{C}_{19}\text{H}_{26}\text{O}]^+$: m/z 270.1984, found 270.1983; Anal calc'd for $\text{C}_{19}\text{H}_{26}\text{O}$: C, 84.39; H, 9.69; found: C, 84.91; H, 9.68.

Determination of Allene Absolute Stereochemistry:

To a solution of **(R)-4a** (0.100 g, 0.632 mmol, 92% ee) in CH_2Cl_2 (1.8 mL) was added $[(\text{Ph}_3\text{PAu})\text{O}]\text{BF}_4$ (9.3 mg, 6.3 μmol) and the resulting mixture was left to stir at rt for 3 h. The reaction mixture was concentrated, the residue dissolved in t -BuOH (6 mL), and 2-methyl-2-butene (1.8 mL) was added. To the resulting yellow mixture was added a solution of NaClO_2 (85 mg, 0.95 mmol) and NaH_2PO_4 (131 mg, 0.948 mmol) in H_2O (6 mL), and the biphasic mixture was left to stir at rt for 4 h. Organic solvents were removed *in vacuo*, the resulting mixture was diluted with H_2O (30 mL), and extracted with hexanes (3x, 30 mL). The aqueous layer was cooled to 0 $^\circ\text{C}$ and acidified to pH 1 by dropwise addition of concentrated HCl. Organics were extracted with EtOAc (6x, 30 mL), the combined extracts were dried over MgSO_4 , filtered, and concentrated to afford a pale yellow liquid. The crude residue was purified by flash column chromatography (1% AcOH in 5:1 hexanes:EtOAc eluent) to afford the desired carboxylic acid **(R)-15** (22 mg, 66% yield, R_f = 0.13 in 3:1 hexanes:EtOAc) as a colorless oil: Spectral data were consistent with those previously reported;⁷ $[\alpha]_D^{25}$ = -31 (c 1.0, CHCl_3), (lit.⁷ $[\alpha]_D^{25}$ = +102.2 (c 2.0, CHCl_3)).

Allene Racemization Experiments:

To a solution of **(R)-2** (85 mg, 0.39 mmol) in DMF (0.5 mL) was added imidazole (59 mg, 0.87 mmol) followed by TBSCl (71 mg, 0.47 mmol) and the resulting mixture was left to stir at rt for 3 d. The crude mixture was purified directly by flash column chromatography (50:1 hexanes:EtOAc eluent) to afford the desired silyl ether **(R)-14** (118 mg, 91% yield, R_f = 0.88 in 3:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.28-7.25 (m, 4H), 7.17-7.15 (m, 1H), 6.10 (t, 1H, J = 2.8 Hz), 3.74 (t, 2H, J = 7.2 Hz), 2.31 (t, 2H, J = 7.2 Hz), 2.11-2.08 (m, 2H), 1.47-1.43 (m, 2H), 1.37-1.33 (m, 2H), 0.89-0.86 (m, 12H), 0.03 (s, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 202.5, 136.0, 128.5, 126.5, 126.4, 105.6, 95.3, 62.0, 36.1, 32.8, 29.8, 25.8, 22.5, 18.3, 14.0, -5.2.

To a solution of **(R)-14** (42.2 mg, 0.128 mmol, 83% ee) in CH_2Cl_2 (0.36 mL) was added Ph_3PAuCl (0.6 mg, 1.3 μmol) followed by AgOTf (0.2 mg, 1.3 μmol) and the cloudy mixture was left to stir at rt for 1 h. The resulting

solution was filtered through silica gel washing with CH_2Cl_2 (2 mL) and the solvent removed to afford a yellow oil. The oil was dissolved in THF (0.36 mL), TBAF (0.15 mL, 0.15 mmol, 1.0 M in THF) was added, and the reaction mixture was left to stir at rt for 10 min. The crude mixture was purified directly by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallenenic alcohol (19 mg, 68% yield, $R_f = 0.14$ in 9:1 hexanes:EtOAc) as a colorless oil: Spectral data were consistent with those described for **2**; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 97:3 hexanes:isopropanol, 1 mL/min; t_R 9.14 min (minor), 12.49 min (major): 4% ee.

To a suspension of AgBF_4 (0.3 mg, 1.5 μmol) in CH_2Cl_2 (0.2 mL) was added Ph_3PAuCl (0.7 mg, 1.5 μmol) and the resulting cloudy solution was stirred rapidly at rt for 5 min. The mixture was filtered through a cotton plug into a solution of **(R)-14** (50.0 mg, 0.151 mmol, 90% ee) in CH_2Cl_2 (0.4 mL) and the reaction mixture was left to stir at rt for 1 h. The solution was filtered through a silica gel plug washing with 4:1 hexanes:EtOAc and the solvent removed to afford a yellow oil. The resulting oil was dissolved in THF (0.43 mL), TBAF (0.18 mL, 0.18 mmol, 1.0 M in THF) was added, and the reaction mixture was left to stir at rt for 1 h. The crude mixture was purified directly by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallenenic alcohol (27 mg, 83% yield, $R_f = 0.14$ in 9:1 hexanes:EtOAc) as a colorless oil: Spectral data were consistent with those described for **2**; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 97:3 hexanes:isopropanol, 1 mL/min; t_R 14.98 min, 26.87 min: 0% ee.

To a solution of **(R)-14** (50.0 mg, 0.151 mmol, 90% ee) in CH_2Cl_2 (0.43 mL) was added $[(\text{Ph}_3\text{PAu})_3\text{O}]\text{BF}_4$ (0.7 mg, 0.5 μmol) and the resulting solution was left to stir at rt for 1 h. The reaction mixture was filtered through a silica gel plug washing with 4:1 hexanes:EtOAc and the solution concentrated to afford a yellow oil. The resulting oil was dissolved in THF (0.43 mL), TBAF (0.18 mL, 0.18 mmol, 1.0 M in THF) was added, and the reaction mixture was left to stir at rt for 1 h. The crude mixture was purified directly by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallenenic alcohol (31 mg, 95% yield, $R_f = 0.14$ in 9:1 hexanes:EtOAc) as a colorless oil: Spectral data were consistent with those described for **2**; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 97:3 hexanes:isopropanol, 1 mL/min; t_R 14.34 min (minor), 26.59 min (major): 67% ee.

Chirality Transfer Experiments:

The Claisen rearrangement was carried out following the general procedure described above using **(R)-4a** of 92% ee. The crude mixture was purified by flash column chromatography (7:1 hexanes:EtOAc eluent) to afford the desired homoallenenic alcohol (39 mg, 78% yield, $R_f = 0.21$ in 3:1 hexanes:EtOAc) as a pale yellow oil: Spectral data were consistent with those previously described for **5a**; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 97:3 hexanes:isopropanol, 1 mL/min; t_R 21.42 min (minor), 24.43 min (major): 88% ee.

(R)-5j

The Claisen rearrangement was carried out following the general procedure described above using **(R)-4j** of 79% ee. The crude mixture was purified by flash column chromatography (10:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (84 mg, 84% yield, $R_f = 0.36$ in 3:1 hexanes:EtOAc) as a colorless oil: Spectral data were consistent with those described for **5j**; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 99.2:0.8 hexanes:isopropanol, 1 mL/min; t_R 33.23 min (major), 35.17 min (minor): 77% ee.

(S)-5n

The Claisen rearrangement was carried out following the general procedure described above using **(R)-5n** of 92% ee. The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (49 mg, 98% yield, $R_f = 0.40$ in 3:1 hexanes:EtOAc) as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 7.35-7.26 (m, 4H), 7.21-7.17 (m, 1H), 5.99 (t, 1H, $J = 3.2$ Hz), 3.84 (t, 2H, $J = 6.4$ Hz), 2.46-2.39 (m, 2H), 1.79 (br s, 1H), 0.22 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 205.0, 135.4, 128.7, 126.2, 125.9, 98.1, 90.3, 62.2, 32.5, -1.4; HRMS (EI^+) calc'd for $[\text{C}_{14}\text{H}_{20}\text{OSi}]^+$: m/z 232.1283, found 232.1281; Anal calc'd for $\text{C}_{14}\text{H}_{20}\text{OSi}$: C, 72.36; H, 8.67; found: C, 72.78; H, 8.71; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 97:3 hexanes:isopropanol, 1 mL/min; t_R 9.99 min (minor), 23.12 min (major): 92% ee.

Diastereoselectivity Experiments:

SI-1

SI-2

The preparation of **SI-2** was carried out according to the method of Ireland and Wipf.⁸ The crude red oil was purified by flash column chromatography (15:1 hexanes:EtOAc eluent) to afford the desired vinylogous carbonate **SI-2** (1.84 g, 64% yield, $R_f = 0.58$ in 3:1 hexanes:EtOAc) as a colorless liquid: ^1H NMR (400 MHz, CDCl_3) δ 7.64 (d, 1H, $J = 12.8$ Hz), 7.27 (t, 2H, $J = 7.6$ Hz), 7.18 (t, 3H, $J = 8.0$ Hz), 5.38 (d, 1H, $J = 12.4$ Hz), 4.68 (br q, 1H, $J = 6.4$ Hz), 3.68 (s, 3H), 2.69 (t, 2H, $J = 7.6$ Hz), 2.23 (td, 2H, $J = 6.8, 1.6$ Hz), 1.82 (quintet, 2H, $J = 7.2$ Hz), 1.54 (d, 3H, $J = 6.8$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 168.2, 160.5, 141.4, 128.5, 128.4, 126.0, 98.3, 87.9, 78.2, 68.0, 51.1, 34.7, 30.0, 22.2, 18.1; HRMS (EI^+) calc'd for $[\text{C}_{17}\text{H}_{20}\text{O}_3]^+$: m/z 272.1412, found 272.1409; Anal calc'd for $\text{C}_{17}\text{H}_{20}\text{O}_3$: C, 74.97; H, 7.40; found: C, 71.46; H, 7.14.

SI-2

SI-3

The reduction of **SI-2** was carried out according to the method of Ireland and Wipf.⁸ The crude oil was purified by flash column chromatography (2% TEA in 4:1 hexanes:EtOAc eluent) to afford the desired alcohol **SI-3** (0.994 g, 62% yield, $R_f = 0.20$ in 3:1 hexanes:EtOAc) as a colorless oil and the propargyl alcohol **SI-1** (0.160 g, 13% yield, $R_f = 0.38$ in 3:1 hexanes:EtOAc) as a colorless oil. Characterization for **SI-3**: ^1H NMR (400 MHz, CDCl_3) δ 7.28 (t, 2H, $J = 8.0$ Hz), 7.20-7.16 (m, 2H), 6.51 (d, 1H, $J = 12.8$ Hz), 5.24 (dt, 1H, $J = 12.4, 7.6$ Hz), 4.53 (br q, 1H, $J = 6.8$ Hz), 4.05 (d, 2H, $J = 7.2$ Hz), 2.70 (t, 2H, $J = 7.6$ Hz), 2.22 (td, 2H, $J = 6.8, 1.6$ Hz), 1.82 (quintet, 2H, $J = 7.2$ Hz), 1.49 (d, 3H, $J = 6.4$ Hz), 1.23 (br s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 147.9, 141.5, 128.5, 128.4, 126.0, 105.6, 86.4, 79.5, 66.0, 60.7, 34.7, 30.1, 22.3, 18.1; HRMS (EI^+) calc'd for $[\text{C}_{16}\text{H}_{20}\text{O}_2 - \text{C}_3\text{H}_4\text{O}]^+$: m/z 188.1201, found 188.1200; Anal calc'd for $\text{C}_{16}\text{H}_{20}\text{O}_2$: C, 78.65; H, 8.25; found: C, 78.96; H, 8.40.

To a solution of **SI-3** (0.946 g, 3.87 mmol) in DMF (5.5 mL) was added imidazole (0.605 g, 8.90 mmol) followed by TIPSCl (0.99 mL, 4.64 mmol), and the resulting mixture was left to stir at rt. TLC analysis after 3 h indicated consumption of the starting alcohol and the reaction mixture was diluted with H₂O (15 mL). The solution was extracted with Et₂O (3 x 10 mL), and the combined organic layers were washed with brine (3 x 10 mL), dried over MgSO₄, filtered, and concentrated. The crude residue was purified by flash column chromatography (2% TEA in hexanes eluent) to afford the desired silyl ether **6** (1.43 g, 92% yield, *R_f* = 0.76 in 3:1 hexanes:EtOAc) as a colorless liquid: ¹H NMR (400 MHz, CDCl₃) δ 7.29-7.25 (m, 2H), 7.20-7.16 (m, 3H), 6.47 (d, 1H, *J* = 12.4 Hz), 5.16 (dt, 1H, *J* = 12.4, 6.0 Hz), 4.51 (br q, 1H, *J* = 6.4 Hz), 4.18 (d, 2H, *J* = 6.4 Hz), 2.70 (t, 2H, *J* = 7.6 Hz), 2.21 (td, 2H, *J* = 7.2, 1.6 Hz), 1.82 (quintet, 2H, *J* = 7.2 Hz), 1.48 (d, 3H, *J* = 6.8 Hz), 1.12-1.02 (m, 21H); ¹³C NMR (100 MHz, CDCl₃) δ 145.9, 141.6, 128.5, 128.5, 125.9, 106.3, 86.0, 79.7, 65.8, 61.2, 34.8, 30.2, 22.3, 18.1 (2), 12.1; HRMS (EI⁺) calc'd for [C₂₅H₄₀O₂Si]⁺: *m/z* 400.2797, found 400.2795; Anal calc'd for C₂₅H₄₀O₂Si: C, 74.94; H, 10.06; found: C, 74.68; H, 10.07.

The Claisen rearrangement was carried out at 40 °C following the general procedure described above. The crude mixture was purified by flash column chromatography (14:1 hexanes:EtOAc) to afford unreacted **6** (39 mg, 15% yield, *R_f* = 0.71 in 9:1 hexanes:EtOAc) and the desired homoallenlic alcohol **7** (191 mg, 76% yield, *R_f* = 0.39 in 9:1 hexanes:EtOAc, >20:1 dr) as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 7.29-7.25 (m, 2H), 7.20-7.16 (m, 3H), 5.17-5.15 (m, 1H), 3.90 (dd, 1H, *J* = 9.6, 4.0 Hz), 3.78-3.70 (m, 3H), 2.79 (br s, 1H), 2.63 (t, 2H, *J* = 7.6 Hz), 2.31-2.27 (m, 1H), 2.02-1.99 (m, 2H), 1.73 (quintet, 2H, *J* = 7.6 Hz), 1.64 (d, 3H, *J* = 6.8 Hz), 1.14-1.03 (m, 21H); ¹³C NMR (100 MHz, CDCl₃) δ 201.8, 142.4, 128.5, 128.3, 125.7, 101.0, 88.4, 67.2, 66.0, 46.1, 35.4, 31.7, 29.4, 18.0, 15.0, 12.1; HRMS (EI⁺) calc'd for [C₂₅H₄₂O₂Si - C₃H₆]⁺: *m/z* 360.2485, found 360.2428; Anal calc'd for C₂₅H₄₂O₂Si: C, 74.57; H, 10.51; found: C, 74.73; H, 10.66.

The Ag(I)-catalyzed cyclization was carried out using a slight modification of the literature procedure.⁷ To a solution of homoallenlic alcohol **7** (39 mg, 98 μ mol) in acetone (0.9 mL) was added AgNO₃ (1.6 mg, 9.8 μ mol) and the resulting mixture was left to stir at rt. After 17 h an additional 10 mol% AgNO₃ was added and the reaction mixture was maintained at rt for 19 h. The solution was concentrated and the crude oil purified by flash column chromatography (50:1 to 10:1 hexanes:EtOAc eluent) to afford unreacted **7** (27 mg, 69% yield, *R_f* = 0.39 in 9:1 hexanes:EtOAc, >20:1 dr) and the desired pyran **16** (10 mg, 26% yield, *R_f* = 0.68 in 9:1 hexanes:EtOAc, >20:1 dr) as a colorless oil: ¹H NMR (400 MHz, C₆D₆) δ 7.20-7.15 (m, 2H), 7.10-7.07 (m, 3H), 5.18 (d, 1H, *J* = 0.8 Hz), 4.38 (d, 1H, *J* = 10.8 Hz), 4.08 (br q, 1H, *J* = 6.8 Hz), 3.92 (t, 1H, *J* = 9.2 Hz), 3.80 (dd, 1H, *J* = 9.6, 3.6 Hz), 3.49 (dd, 1H, *J* = 11.2, 2.0 Hz), 2.50-2.44 (m, 2H), 2.04-2.02 (br m, 1H), 1.96 (br t, 2H, *J* = 7.6 Hz), 1.72-1.60 (m, 2H), 1.19 (t, 3H, *J* = 6.4 Hz), 1.14-1.04 (m, 21H); ¹³C NMR (100 MHz, C₆D₆) δ 142.2, 135.6, 128.4, 128.3, 127.6, 125.8, 70.5, 64.8, 63.7, 41.5, 35.6, 34.7, 29.5, 21.7, 18.0, 12.0; HRMS (EI⁺) calc'd for [C₂₅H₄₂O₂Si - C₃H₆]⁺: *m/z* 360.2485, found 360.2433.

Selected nOe cross peaks for pyran **16**

A solution of **6** (50 mg, 0.12 mmol) in *m*-xylene (0.36 mL) was heated to 170 °C and maintained for 3 h. The reaction mixture was cooled to rt, diluted with MeOH (0.72 mL) and NaBH₄ (5.4 mg, 0.12 mmol) was added. After 1 h at rt the reaction mixture was filtered through silica gel washing with Et₂O (5 mL) and concentrated to afford homoallenic alcohol **7** (45 mg, 90% yield, *R_f* = 0.27 in 9:1 hexanes:EtOAc, 1.5:1.0 mixture of diastereomers) as a pale yellow oil: ¹H NMR (500 MHz, CDCl₃) The following were observed for the mixture of diastereomers δ 7.29-7.25 (m, 2H), 7.19-7.16 (m, 3H), 5.16-5.13 (m, 1H), 3.91-3.88 (m, 1H), 3.79-3.69 (m, 3H), 2.62 (t, 2H, *J* = 7.5 Hz), 2.31-2.26 (m, 1H), 2.02-1.99 (m, 2H), 1.73 (quintet, 2H, *J* = 7.5 Hz), 1.64 (t, 3H, *J* = 7.0 Hz), 1.12-1.02 (m, 21H); ¹³C NMR (100 MHz, CDCl₃) The following were observed for the major diastereomer δ 100.9, 66.3, 46.0, 31.8, 15.1; The following were observed for the minor diastereomer δ 100.9, 66.0, 46.1, 31.7, 15.0; The following were observed for the mixture of diastereomers δ 201.8, 142.4, 128.5, 128.3, 125.8, 88.4, 67.2, 35.4, 29.4, 18.0, 11.8.

The Ag(I)-catalyzed cyclization was carried out using the modification of the published procedure described above.⁷ The crude oil was purified by flash column chromatography (100:1 hexanes:EtOAc eluent) to afford the desired pyran **16** (5.2 mg, 12% yield, *R_f* = 0.68 in 9:1 hexanes:EtOAc, 1.4:1.0 mixture of diastereomers) as a colorless oil: ¹H NMR (500 MHz, C₆D₆) The following were observed for the major diastereomer δ 5.23 (s, 1H), 4.19 (br q, 1H, *J* = 7.0 Hz), 3.91-3.89 (m, 2H), 3.70-3.66 (m, 2H), 2.22 (m, 1H); The following were observed for the minor diastereomer δ 5.15 (s, 1H), 4.35 (d, 1H, *J* = 11.0 Hz), 4.05 (br q, 1H, *J* = 7.0 Hz), 3.90 (m, 1H), 3.76 (dd, 1H, *J* = 9.5, 3.5 Hz), 3.46 (dd, 1H, *J* = 11.0, 1.5 Hz); The following were observed for the mixture of diastereomers δ 7.16-7.12 (m, 2H), 7.07-7.04 (m, 3H), 2.51-2.39 (m, 2H), 1.99-1.90 (m, 3H), 1.69-1.57 (m, 2H), 1.15 (t, 3H, *J* = 7.0 Hz), 1.10-0.99 (m, 21H).

The preparation of **SI-5** was carried out according to the method of Ireland and Wipf.⁸ The crude red oil was purified by flash column chromatography (12:1 hexanes:EtOAc eluent) to afford the desired vinylogous carbonate **SI-5** (1.36 g, 94% yield, *R_f* = 0.65 in 3:1 hexanes:EtOAc) as a pale yellow liquid: ¹H NMR (400 MHz, CDCl₃) δ 7.70 (d, 1H, *J* = 12.4 Hz), 7.50 (d, 2H, *J* = 6.4 Hz), 7.41-7.34 (m, 3H), 5.63 (s, 1H), 5.44 (d, 1H, *J* = 12.8 Hz), 3.69 (s, 3H), 2.30 (td, 2H, *J* = 6.8, 1.6 Hz), 1.53 (quintet, 2H, *J* = 6.8 Hz), 1.41 (sextet, 2H, *J* = 7.2 Hz), 0.91 (t, 3H, *J* = 7.6 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 168.1, 160.2, 137.0, 129.1, 128.7, 127.5, 99.1, 91.5, 75.6, 73.8, 51.1, 30.4, 22.0, 18.5, 13.6; HRMS (EI⁺) calc'd for [C₁₇H₂₀O₃]⁺: *m/z* 272.1412, found 272.1408.

The reduction of **SI-5** was carried out according to the method of Ireland and Wipf.⁸ The crude oil was purified by flash column chromatography (2% TEA in 4:1 hexanes:EtOAc eluent) to afford the desired alcohol **SI-6** (0.618 g, 69% yield, *R_f* = 0.26 in 3:1 hexanes:EtOAc) as a pale yellow oil: ¹H NMR (400 MHz, CDCl₃) δ 7.50 (d, 2H, *J* = 7.2 Hz), 7.39-7.31 (m, 3H), 6.56 (d, 1H, *J* = 12.4 Hz), 5.48 (s, 1H), 5.30 (dt, 1H, *J* = 12.4, 7.6 Hz), 4.04 (d, 2H, *J* = 7.2 Hz), 2.28 (td, 2H, *J* = 6.8, 1.6 Hz), 1.53 (quintet, 2H, *J* = 6.8 Hz), 1.41 (sextet, 2H, *J* = 7.2 Hz), 0.91 (t, 3H, *J* = 7.2 Hz), 1.25 (br s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 147.5, 138.0, 128.6, 128.5, 127.3, 106.3, 89.9, 76.7, 71.9, 60.6, 30.5, 21.9, 18.5, 13.5; HRMS (EI⁺) calc'd for [C₁₆H₂₀O₂ - CH₃O]⁺: *m/z* 213.1279, found 213.1280.

To a solution of **SI-6** (0.150 g, 0.614 mmol) in DMF (1.5 mL) was added imidazole (96 mg, 1.4 mmol) followed by TIPSCl (0.16 mL, 0.74 mmol) and the resulting solution was left to stir at rt. TLC of the reaction mixture after 18 h showed consumption of starting alcohol and the reaction mixture was diluted with H₂O (5 mL). The solution was extracted with Et₂O (3 x 5 mL), and the combined organic layers washed with brine (3 x 10 mL), dried over MgSO₄, filtered, and concentrated. The crude residue was purified by flash column chromatography (2% TEA in hexanes eluent) to afford the desired silyl ether **(±)-8** (204 mg, 83% yield, *R_f* = 0.89 in 3:1 hexanes:EtOAc) as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 7.50 (d, 2H, *J* = 6.8 Hz), 7.38-7.31 (m, 3H), 6.53 (d, 1H, *J* = 12.4 Hz), 5.46 (s, 1H), 5.22 (dt, 1H, *J* = 12.4, 6.4 Hz), 4.18 (d, 2H, *J* = 6.4 Hz), 2.27 (td, 2H, *J* = 7.2, 2.0 Hz), 1.55-1.48 (m, 2H), 1.44-1.38 (m, 2H), 1.08-1.02 (m, 21H), 0.90 (t, 3H, *J* = 7.6 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 145.6, 138.3, 128.5, 127.4, 107.0, 89.6, 71.8, 61.1, 30.6, 22.0, 18.6, 18.0, 13.6, 12.1.

The Claisen rearrangement was carried out following the general procedure described above. The crude mixture was purified by flash column chromatography (12:1 hexanes:EtOAc eluent) to afford the desired homoallenyl alcohol (167 mg, 89% yield, *R_f* = 0.36 in 9:1 hexanes:EtOAc) as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 7.29-7.22 (m, 4H), 7.19-7.15 (m, 1H), 6.18 (q, 1H, *J* = 1.6 Hz), 4.02 (dd, 1H, *J* = 10.0, 4.4 Hz), 3.83-3.76 (m, 3H), 2.95 (br s, 1H), 2.49-2.45 (m, 1H), 2.15-2.10 (m, 2H), 1.49-1.41 (m, 2H), 1.38-1.31 (m, 2H), 1.10-1.0 (m, 21H), 0.87 (t, 3H, *J* = 7.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 202.1, 135.2, 128.6, 126.8, 126.4, 106.5, 97.0, 67.1, 66.0, 46.8, 32.3, 29.8, 22.5, 17.9, 12.9, 11.8; HRMS (EI⁺) calc'd for [C₂₅H₄₂O₂Si]⁺: *m/z* 402.2954, found 402.2955.

To a solution of **(±)-9** (30 mg, 74 μ mol) in THF (0.2 mL) was added TBAF (0.1 mL, 0.10 mmol, 1.0M in THF), and the resulting solution was maintained at rt for 10 min. The crude mixture was purified by flash column chromatography (2:1 hexanes:EtOAc eluent) to afford the desired diol (7 mg, 50% yield, *R_f* = in 3:1 hexanes:EtOAc) as a white solid: ¹H NMR (400 MHz, CDCl₃) δ 7.31-7.25 (m, 4H), 7.20-7.17 (m, 1H), 6.23 (br q, 1H, *J* = 1.6 Hz), 3.94-3.87 (td, 2H, *J* = 11.2, 4.4 Hz), 3.83-3.78 (m, 2H), 2.44-2.42 (m, 1H), 2.17-2.11 (m, 4H), 1.50-1.40 (m, 2H), 1.35 (sextet, 2H, *J* = 7.2 Hz), 0.87 (t, 3H, *J* = 7.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 202.1, 134.9, 128.8, 127.0, 126.4, 106.7, 97.1, 64.9, 46.6, 32.0, 29.8, 22.5, 13.9.

The Claisen rearrangement was carried out following the general procedure described above. The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallenyl alcohol (122 mg, 81% yield, *R_f* = 0.26 in 9:1 hexanes:EtOAc, >20:1 dr) as a colorless oil: Spectral data were consistent with those reported for **(±)-9**.

Desilylation was carried out using the procedure described above for the preparation **(±)-SI-7**. The crude mixture was purified by flash column chromatography (2:1 hexanes:EtOAc eluent) to afford the desired diol **SI-7** (66 mg,

90% yield, $R_f = 0.05$ in 3:1 hexanes:EtOAc) as a white solid: Spectral data were consistent with those reported for ($\pm$)-**SI-7**; Enantiomeric purity was determined by chiral HPLC analysis, Chiralcel OD column, 92:8 hexanes:isopropanol, 1 mL/min; t_R 14.99 min (minor), 17.43 min (major): 94% ee.

Allyl / Propargyl Competition Experiment:

13

The Claisen rearrangement was carried out following the general procedure described above. The crude mixture was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (81 mg, 80% yield, $R_f = 0.24$ in 3:1 hexanes:EtOAc) as a yellow oil: ^1H NMR (400 MHz, CDCl_3) δ 7.40 (d, 2H, $J = 8.0$ Hz), 7.33 (t, 2H, $J = 7.2$ Hz), 7.23 (t, 1H, $J = 7.2$ Hz), 6.30-6.20 (m, 2H), 5.35-5.28 (m, 1H), 5.09 (d, 1H, $J = 8.8$ Hz), 3.85 (t, 2H, $J = 6.4$ Hz), 2.79-2.70 (m, 2H), 2.02 (br s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 207.7, 135.7, 132.4, 128.3, 127.3, 126.2, 117.0, 104.2, 98.4, 61.1, 33.3; HRMS (EI^+) calc'd for $[\text{C}_{13}\text{H}_{14}\text{O}]^+$: m/z 186.1045, found 186.1045.

Examination of Au(I)-Species Generated Under the Reaction Conditions:

To a solution of ($\pm$)-**4n** (0.100 g, 0.434 mmol) in CH_2Cl_2 (1.24 mL) was added $(\text{Ph}_3\text{P})_2\text{AuBF}_4$ (3.5 mg, 4.3 μmol) and the resulting solution was left to stir at rt and monitored by TLC. After 18 h the mixture was concentrated, redissolved in MeOH (1.24 mL), and NaBH_4 (15 mg, 0.39 mmol) was added. TLC analysis of the reaction mixture after 1 h indicated consumption of the intermediate aldehyde and the solution was concentrated. The crude red oil was purified by flash column chromatography (9:1 hexanes:EtOAc eluent) to afford the desired homoallylic alcohol (67 mg, 66% yield, $R_f = 0.36$ in 3:1 hexanes:EtOAc) as a colorless oil: Spectral data were consistent with those reported for (**S**)-**5n**.

To a solution of ($\pm$)-**4n** (20 mg, 87 μmol) in CH_2Cl_2 (0.25 mL) was added $(\text{Ph}_3\text{PAu})\text{CH}_2\text{CHO}$ (1.3 mg, 2.6 μmol) and the reaction mixture was left to stir at rt and monitored by TLC. After 20 h the mixture was concentrated and a crude ^1H spectrum indicated only the starting vinyl ether ($\pm$)-**4n** present in solution: ^1H NMR (400 MHz, CDCl_3) δ 7.53 (d, 2H, $J = 6.8$ Hz), 7.41-7.33 (m, 2H), 6.50 (dd, 1H, $J = 14.0, 6.4$ Hz), 5.50 (s, 1H), 4.51 (dd, 1H, $J = 14.0, 1.6$ Hz), 4.18 (dd, 1H, $J = 6.8, 2.0$ Hz), 0.19 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 149.3, 137.4, 128.6, 128.5, 127.4, 101.6, 93.7, 90.6, 71.1, -0.2; HRMS (EI^+) calc'd for $[\text{C}_{14}\text{H}_{18}\text{OSi}]^+$: m/z 230.1127, found 230.1127; Anal calc'd for $\text{C}_{14}\text{H}_{18}\text{OSi}$: C, 72.99; H, 7.88; found: C, 72.86; H, 7.83.

References:

- (1) Bruce, M.I.; Nicholson, B.K.; Bin Shawkataly, O. *Inorg. Syn.* **1989**, 26, 324.
- (2) (a) Yang, Y.; Ramamoorthy, V.; Sharp, P.R. *Inorg. Chem.* **1993**, 32, 1946. (b) Nesmeyanov, A.N.; Perevalova, E.G.; Struchkov, Yu. T.; Antipin, M. Yu.; Grandberg, K.I.; Dyadchenko, V.P. *J. Organomet. Chem.* **1980**, 201, 343.
- (3) Westland, A.D. *Can. J. Chem.* **1969**, 47, 4135.
- (4) Nesmeyanov, A.N.; Perevalova, E.G.; Dyadchenko, V.P.; Grandberg, K.I. *Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya* **1974**, 12, 2872.
- (5) Midland, M.M.; Tramontano, A.; Kazubski, A.; Graham, R.S.; Tsai, D.J.S.; Cardin, D.B. *Tetrahedron* **1984**, 40, 1371.
- (6) Nonoshita, K.; Banno, H.; Maruoka, K.; Yamamoto, H. *J. Am. Chem. Soc.* **1990**, 112, 316.
- (7) Wan, Z.; Nelson, S.G. *J. Am. Chem. Soc.* **2000**, 122, 10470.
- (8) Ireland, R.E.; Wipf, P.; Xiang, J.-N. *J. Org. Chem.* **1991**, 56, 3572.