jo030370g
REVISED

1/26/04

Supporting Information

A Halide-Free Dehydrative Allylation Using Allylic Alcohols Promoted by a Palladium-Triphenyl Phosphite Catalyst

Yoshihito Kayaki, Takashi Koda, and Takao Ikariya*

PRESTO, Japan Science and Technology Agency and Graduate School of Science and
Engineering and Frontier Collaborative Research Center, Tokyo Institute of Technology,
O-okayama, Meguro-ku, Tokyo 152-8552, Japan. E-mail: tikariya@o.cc.titech.ac.jp

Table of Contents

General information and experimental procedures	S2-S5
NMR Spectra of $[\text{Pd}\{\text{P}(\text{OC}_6\text{H}_5)_3\}_4]$	S6-S8
NMR Spectra of dicinnamyl ether	S9-S10
^1H NMR spectrum of allyl cyclohexyl ether	S11
^1H NMR spectrum of allyl neopentyl ether	S12
^1H NMR spectrum of allyl benzyl ether	S13
^1H NMR spectrum of allyl furfuryl ether	S14
NMR Spectra of 5,5-diallyl-2,2-dimethyl-1,3-dioxane-4,6-dione	S15-S16
NMR Spectra of 2,2-diallyl-1,3-cyclohexanedione	S17-S18
NMR Spectra of 2,2-diallyl-1,3-cyclopentanedione	S19-S20
NMR Spectra of <i>N, N</i> -diallylaniline	S21-S22

General information:

All reactions were performed under an atmosphere of argon using standard Schlenk techniques. Solvents were dried by standard procedures and freshly distilled under argon. Allylic alcohols other than cinnamyl alcohol were distilled from magnesium and stored in argon. 2,2-dimethyl-1,3-dioxane-4,6-dione was recrystallized from acetone before use. Other reagents were commercially available and were used as received without further purification.

^1H (300.5 MHz) and ^{13}C (75.5 MHz) NMR spectra were referenced to tetramethylsilane ($\delta = 0$ ppm), and ^{31}P (121.7 MHz) NMR spectra were referenced to 85% H_3PO_4 as an external standard.

Preparation of Pd catalyst 3

To a suspension of 1.04 g (4.0 mmol) of $\text{PdCl}_2(\text{CH}_3\text{CN})_2$ in dichloromethane (15 mL) was added 5.25 mL (20.0 mmol) of $\text{P}(\text{OC}_6\text{H}_5)_3$. After stirring at room temperature for 15 min, $\text{N}(\text{C}_2\text{H}_5)_3$ (1.12 mL, 8.0 mmol) and water (72 μL , 4 mmol) were added, and the mixture became a clear yellow solution. Water (5 mL X 2) was added and stirred for 5 min, and the upper layer was removed via cannula. The residue was evaporated to dryness, and the resulted white powder was washed by acetone. The product was recrystallized from hot acetone. Yield 2.10 g, 39%.

^1H NMR (CD_2Cl_2) $\delta = 6.80\text{--}6.83$ (m, 24H), 7.09–7.15 (m, 12H), 7.19–7.25 (m, 24H); ^{13}C NMR (CD_2Cl_2) $\delta = 141.5, 124.3, 129.6, 151.9$; ^{31}P NMR (CD_2Cl_2) $\delta = 138.0$ (s).

Typical procedure for dehydrative etherification of allyl alcohol. The Pd catalyst 3 (67.4 mg, 5.0×10^{-2} mmol) in a 20 mL glass reactor was added to degassed and distilled alcohol 1 under an argon atmosphere. The solution was stirred at 80 $^\circ\text{C}$ for 6 h. The product ether 2 was isolated from the reaction mixture by distillation. The NMR yields were estimated by using hexamethylbenzene (16.2 mg, 0.1 mmol) as an internal standard.

Experimental procedure for dehydrative etherification of cinnamyl alcohol:

Palladium complex **3** (32.9 mg, 0.024 mmol), cinnamyl alcohol (657 mg, 4.9 mmol), and toluene (2.5 mL) were placed under argon in a Schlenk tube with a magnetic stirring bar. The mixture was stirred at 80 °C for 18 h. After the mixture had cooled to room temperature, the crude products were purified by column chromatography on silica gel (hexane/ethyl acetate) to yield dicinnamyl ether. Yield 586 mg, 96%.

Dicinnamyl ether: ^1H NMR (CDCl_3) δ = 4.1-4.2 (d, 4H), 6.27-6.36 (d, J = 6.1 Hz, 2H), 6.60-6.66 (d, J = 15.9 Hz, 2H), 7.20-7.41 (m, 10H); ^{13}C NMR (CDCl_3) δ = 70.2, 126.0, 126.5, 127.7, 128.5, 128.5, 132.6, 136.7.

General procedure for cross etherification using allyl alcohol. The Pd catalyst **3** (67.4 mg, 5.0×10^{-2} mmol) in a 20 mL glass reactor was added to degassed and distilled allyl alcohol **1** (0.68 mL, 10.0 mmol) and other alcohols (5.0 mmol) under an argon atmosphere. The solution was stirred at 80 °C for 6-42 h. The reaction mixture was cooled to room temperature and purified by column chromatography on silica gel with hexane/ethyl acetate as eluent to afford the product unsymmetrical ethers **4**. The NMR yields were estimated by using hexamethylbenzene (16.2 mg, 0.1 mmol) as an internal standard.

NMR data for cross etherification products 4:

Allyl cyclohexyl ether: ^1H NMR (CDCl_3) δ = 1.03-1.40 (m, 5H), 1.40-1.58 (m, 1H), 1.58-1.78 (m, 2H), 1.78-1.96 (m, 2H), 3.18-3.31 (m, 1H), 3.93-4.02 (m, 2H), 5.01-5.30 (m, 2H), 5.80-6.00 (m, 1H); ^{13}C NMR (CDCl_3) δ = 24.1, 25.8, 32.2, 68.7, 76.9, 116.2, 135.6.

Allyl neopentyl ether: ^1H NMR (CDCl_3) δ = 0.90 (s, 9H), 3.05 (s, 2H), 3.95 (d, J = 5.4 Hz, 2H), 5.07-5.17 (m, 2H), 5.17-5.31 (m, 2H), 5.78-5.93 (m, 1H); ^{13}C NMR (CDCl_3) δ = 26.7, 31.6, 72.3, 80.9, 116.1, 135.4.

Allyl benzyl ether: ^1H NMR (CDCl_3) δ = 3.98-4.05 (m, 2H), 4.52 (s, 2H), 5.13-5.36 (m, 4H), 5.85-6.04 (m, 1H), 7.20-7.40 (m, 5H); ^{13}C NMR (CDCl_3) δ = 71.1, 72.1, 117.1, 127.6, 127.7, 128.4, 134.7, 138.2.

Allyl furfuryl ether: ^1H NMR (CDCl_3) δ = 3.98-4.02 (m, 2H), 4.44 (s, 2H), 5.16-5.32 (m, 2H), 5.82-5.98 (m, 1H), 6.27-6.35 (m, 2H), 7.38 (s, 1H); ^{13}C NMR (CDCl_3) δ = 63.8, 70.9, 109.2, 110.2, 117.5, 134.4, 142.7, 151.7.

Experimental procedure for dehydrative allylation of active methylene compounds:

Palladium complex **3** (13.9 mg, 0.010 mmol) was added to toluene (2.5 mL) followed by active methylene compounds (2.5 mmol) and allyl alcohol **1** (0.35 mL, 5.0 mmol). The stirred mixture was heated to 80 °C for 2 h. After the mixture had cooled to room temperature, the crude products were purified by column chromatography on silica gel using hexane/ethyl acetate as eluent to yield the respective pure compounds.

Experimental procedure for dehydrative allylation of aniline

Palladium complex **3** (23.4 mg, 0.017 mmol) was added to toluene (2.5 mL) followed by aniline (0.24 mL, 1.75 mmol) and allyl alcohol **1** (0.24 mL, 3.5 mmol). The stirred mixture was heated to 80 °C for 4 h. After the mixture had cooled to room temperature, the crude products were purified by column chromatography on alumina (hexane/ethyl acetate) to yield *N,N*-diallylaniline.

NMR data for C-allylation and N-allylation products **6:**

5,5-diallyl-2,2-dimethyl-1,3-dioxane-4,6-dione: ^1H NMR (CDCl_3) δ = 1.66 (s, 6H), 2.70 (d, J = 7.6 Hz, 4H), 5.15-5.16 (d, J = 4.2 Hz, 2H), 5.19-5.22 (d, J = 11.2 Hz, 2H), ^{13}C NMR (CDCl_3) δ = 29.4, 42.3, 54.9, 105.3, 120.7, 130.6, 167.8.

2,2-diallyl-1,3-cyclohexanedione: ^1H NMR (CDCl_3) δ = 1.89 (m, 2H), 2.47 (d, J = 7.3 Hz, 4H), 2.50 (d, J = 6.6 Hz, 4H), 4.98 (m, 2H), 5.02 (m, 2H), 5.41-5.56 (m, 2H); ^{13}C NMR (CDCl_3) δ = 16.4, 39.9, 40.8, 68.2, 119.3, 132.4, 210.3.

2,2-diallyl-1,3-cyclopentanedione: ^1H NMR (CDCl_3) δ = 2.30 (d, J = 7.6 Hz, 4H), 2.56 (d, J = 0.5 Hz, 4H), 4.98-5.03 (m, 4H), 5.44-5.58 (m, 2H); ^{13}C NMR (CDCl_3) δ = 36.4, 39.3, 61.2, 120.0, 131.3, 216.4.

***N,N*-diallylaniline:** ^1H NMR (CDCl_3) δ = 3.90 (d, J = 4.9 Hz, 4H), 5.12-5.20 (m, 4H), 5.79-5.91 (m, 2H), 6.67-6.70 (m, 3H), 7.16-7.21 (m, 2H); ^{13}C NMR (CDCl_3) δ = 52.7, 112.4, 116.0, 116.3, 129.0, 134.0, 148.6.

S6

```

Date . Sat Nov 4 13:36:37 2000
FileName LoadingF10.mcd
Comment Pd[P10C645] 3) 4702012/r
SliceHistory non
ENCODE

POINT 32768 points
SAMP0 32768 points
FREQU 6000.0 Hz
FILTR 4000 Hz
DELAY 50.0 usec
DEADT 72.2 usec
INSTR 125.0 usec
TIMES 8 times
DUMMY 0 times
PR0 1 0000 sec
ACQTH 4096 0000 msec
FREQD 10.00000 msec
INSTR 1600 0000 msec
RESOL 0.24 Hz
PR1 5 60 usec
CORNUC 1H
CORR HQ
CORSET 300 40 MHz
CORATN 130780.00 Hz
SCANS 16
      8 times

CORREL 17 Hz
      217 C

```

Pd[P(Oc6H5)3]4/CD2Cl2/r.t./31P

Date Sat Nov 4 13:44:09 2000

Filename .loadingf10.metadata

Comment Pd[P(Oc6H5)3]4/CD2Cl2/r

Slcchistory

EXMODE

POINT 32768 POINTS

SAMPD 32768 POINTS

FREQU 45045.0 HZ

FILTR 22500 HZ

DELAY 8.9 USEC

DEADT 10.0 USEC

INVL 22.2 USEC

TIMES 512 TIMES

DUMMY 0 TIMES

PD 0.5000 SEC

ACQTM 727.4495 NSEC

PREDL 10.00000 NSEC

INVT 1000.0000 NSEC

RESOL 1.37 HZ

PW 3.65 USEC

31P 121.55 MHz

DESET 106557.48 HZ

RGAIN 26

IRFNUC 300.40 MHz

IRFRD 130282.93 HZ

IRFET 45.0 USEC

IRPM 0

IRPNS 175 TIMES

SCANS

SLVNT CD2Cl2

SPINNING 16 HZ

TEMP 22.6 C

08-03107-025/00C13/r.t./1H

21-21211-117/CDC13/P. 6/13C

Date Thu Dec 12 21 18:58 2002
 FileName LoadingID: noda
 Comment 21-21211-117/CDC13/P. 6/13C
 SliceHistory
 EXMODE 0ch
 POINT 65536 points
 SAMPD 65536 points
 FREQU 25000.1 Hz
 PULPR 12500 Hz
 DELAY 15.0 usec
 DEADT 21.0 usec
 INTVI 40.0 usec
 TIMES 3000 times
 JUMPS 0 times
 DO 5 0000 sec
 ACQTN 2621.4399 msec
 PHEDL 10 00000 msec
 INUM1 1000.0000 msec
 RESOL 0.38 Hz
 SWH 6.00 usec
 13C 75.45 MHz
 126064 52 Hz
 1H 300.40 MHz
 130282.93 Hz
 45.0 usec
 D
 SCANS 103 times
 SYNT
 SPINNING
 TEMP 16 Hz
 23.1 C

Date Thu Oct 24 13:52:04 2002

```

FileShare
Connect
Directory
EXCH02
Exchange
RoutingIC Update
21-21036-261/250033
out

```

POINT	32768	PCNTS	
SAMP	32768	PCNTS	
REC	8000	Hz	
CLIF	4000	Hz	
DELAY	50	USEC	
LEAD	125	USEC	
INVL	125	USEC	
TIMES	16	CMPS	
CONV	1	CMPS	
BD	1	0050	SEC
BCOIM	4036	0000	msec
PRED	15	0060	msec
ININT	1040	0050	msec
RESOL	0	24	Hz
PH	5	50	USEC
GENUC	14		
BBFRO	300	40	MHz
CBSE	133780	50	Hz
PGAIN	13		
GCANS	16	CMPS	
ELVNT	000.3		
SPINNING	16	Hz	
TEMP	25.0	C	

W. J. P. R.