Supporting Information for

Hangman Salen Platforms Containing Two Xanthene Scaffolds

Jenny Y. Yang, Julien Bachmann and Daniel G. Nocera*

Department of Chemistry, 6-335, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, Massachusetts 02139-4301

<i>Index</i>	<i>Page</i>
General Experimental Methods	S2
Synthetic Scheme S1	S3
X-ray Structural Data for 9	S4-S14
X-ray Structural Data for 24	S15-S22
DFT Calculations	S23-S36
¹ H NMR Spectra for 3 , 5-14 , 16 , 22 , 24-26	S37

Materials. Silica gel 60 (70 - 230 and 230 - 400 mesh) was used for column chromatography. Analytical thin layer chromatography was performed using F254 silica gel (pre-coated sheets, 0.2 mm thick). Solvents for synthesis were reagent grade or better and used as received or dried according to standard methods.¹ The following compounds were obtained using published protocols and their purity confirmed by ¹H NMR: 4-hydroxycarbonyl-5-bromo-2,7-di-*tert*-butyl-9,9-dimethylxanthene (1),² 4-methoxycarbonyl-5-bromo-2,7-di-*tert*-butyl-9,9-dimethylxanthene (2),² 4-(5-bromo-2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthen-4-yl)-benzoic acid methyl ester (4),³ and 4-hydroxy-biphenyl-3-carbaldehyde (21).⁴

Physical measurements. ¹H NMR spectra were collected in CDCl₃ or C₄D₈O using a 500 Hz spectrometer at 25 °C. All chemical shifts are reported using the standard δ notation in parts-per-million relative to tetramethylsilane and spectra have been internally calibrated to the monoprotio impurity of the deuterated solvent used. High-resolution mass spectral analyses were carried out using an ESI source.

-
1. Armarego, W. L. F.; Perrin, D. D. *Purification of Laboratory Chemicals*; 4th ed.; Butterworth-Heinmann: Oxford, 1966.
 2. Chang, C. J.; Chng, L. L.; Nocera, D. G. *J. Am. Chem. Soc.* **2003**, *125*, 1866-1876.
 3. Liu, S.-Y.; Nocera, D. G. *J. Am. Chem. Soc.* **2005**, *127*, 5278-5279.
 4. Morris, G. A.; Nguyen, S. T. *Tetrahedron Lett.* **2001**, *42*, 2093-2096.

Scheme S1

(a) H_2SO_4 , methanol, (b) bis(pinacolato)diboron, Pd(dppf)Cl_2 , KOAc, DMSO, (c) 5-bromosalicylaldehyde, $\text{Pd(PPh}_3)_4$, Na_2CO_3 , DMF/ H_2O , (d) BBr_3 , CH_2Cl_2 , (e) benzyl alcohol, DMAP, DCC, CH_2Cl_2 , (g) H_2 on Pd/charcoal, ethyl acetate

Figure S1. Thermal ellipsoid plot of 2,7-di-*tert*-butyl-5-(3-formyl-4-hydroxyphenyl)-9,9-dimethyl-9H-xanthene-4-carboxylic acid (**9**). Thermal ellipsoids are drawn at the 50% probability level.

Table S1.1. Crystal data and structure refinement 2,7-di-*tert*-butyl-5-(3-formyl-4-hydroxy-phenyl)-9,9-dimethyl-9H-xanthene-4-carboxylic acid (**9**).

Identification code	03084a	
Empirical formula	C ₃₁ H ₃₄ O ₅	
Formula weight	486.58	
Temperature	467(2) K	
Wavelength	0.71073 Å	
Crystal system	Triclinic	
Space group	$P\bar{1}$	
Unit cell dimensions	$a = 9.7828(9)$ Å	$\alpha = 78.978(2)^\circ$
	$b = 9.8015(9)$ Å	$\beta = 82.309(2)^\circ$
	$c = 15.2502(14)$ Å	$\gamma = 68.047(2)^\circ$
Volume	1336.8(2) Å ³	
<i>Z</i>	2	
Density (calculated)	1.209 Mg/m ³	
Absorption coefficient	0.081 mm ⁻¹	
<i>F</i> (000)	520	
Crystal size	0.34 × 0.31 × 0.15 mm ³	
θ range for data collection	2.27 to 24.73°	
Index ranges	$-11 \leq h \leq 7, -11 \leq k \leq 9, 18 \leq \ell \leq 18$	
Reflections collected	6974	
Independent reflections	4471 [$R_{\text{int}} = 0.0274$]	
Completeness to $\theta = 24.73^\circ$	97.6 %	
Absorption correction	None	
Refinement method	Full-matrix least-squares on F^2	
Data / restraints / parameters	4471 / 0 / 334	
Goodness-of-fit on F^2	1.021	
Final <i>R</i> indices [$I > 2\sigma(I)$]	$R_I = 0.0720, wR_2 = 0.1568$	
<i>R</i> indices (all data)	$R_I = 0.1084, wR_2 = 0.1738$	
Largest diff. peak and hole	0.476 and -0.284 e/Å ³	

Tables S.1.2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 03084a. U(eq) is defined as one third of the trace of the orthogonalized U_{ij} tensor.

Atom	x	y	z	U(eq)
C(1)	2435(3)	7977(3)	2723(2)	23(1)
C(2)	1525(3)	9389(3)	2921(2)	25(1)
C(3)	1189(3)	10581(3)	2233(2)	28(1)
C(4)	1697(4)	10418(3)	1362(2)	28(1)
C(5)	2612(4)	9002(3)	1201(2)	30(1)
C(6)	3007(3)	7762(3)	1868(2)	25(1)
C(7)	4036(3)	6234(3)	1647(2)	28(1)
C(8)	4104(3)	5033(3)	2457(2)	26(1)
C(9)	4810(3)	3530(3)	2388(2)	29(1)
C(10)	4876(3)	2397(3)	3098(2)	28(1)
C(11)	4176(3)	2826(3)	3907(2)	29(1)
C(12)	3481(3)	4309(3)	4019(2)	26(1)
C(13)	3476(3)	5397(3)	3282(2)	25(1)
C(14)	1276(4)	11770(4)	622(2)	39(1)
C(15)	−408(5)	12490(4)	636(3)	56(1)
C(16)	1904(5)	12911(4)	787(3)	51(1)
C(17)	1875(6)	11353(4)	−301(2)	66(1)
C(18)	3445(4)	5868(4)	878(2)	41(1)
C(19)	5588(4)	6283(4)	1383(3)	45(1)
C(20)	5717(4)	752(3)	3001(2)	36(1)
C(21)	4762(12)	−169(10)	3414(7)	50(3)
C(22)	7055(10)	203(10)	3648(6)	56(3)
C(23)	6347(13)	473(12)	2129(7)	73(3)
C(21A)	5400(10)	−331(9)	3743(6)	57(2)
C(22A)	7343(9)	470(9)	2800(6)	63(3)
C(23A)	5230(8)	426(8)	2126(5)	46(2)
C(24)	2788(3)	4703(3)	4909(2)	29(1)
C(25)	3138(4)	5700(4)	5314(2)	33(1)
C(26)	2585(4)	5994(4)	6165(2)	39(1)
C(27)	1655(4)	5332(4)	6619(2)	34(1)

C(28)	1230(3)	4364(3)	6254(2)	31(1)
C(29)	1829(3)	4056(3)	5388(2)	32(1)
C(30)	233(4)	3667(4)	6728(3)	43(1)
C(31)	881(3)	9638(3)	3836(2)	26(1)
O(1)	2760(2)	6847(2)	3438(1)	29(1)
O(2)	540(3)	8715(2)	4377(1)	37(1)
O(3)	677(3)	10965(2)	4002(2)	42(1)
O(4)	1103(3)	5645(3)	7462(2)	49(1)
O(5)	−278(3)	3828(3)	7496(2)	55(1)

Table S1.3. Bond lengths (Å) for 2,7-di-*tert*-butyl-5-(3-formyl-4-hydroxyphenyl)-9,9-dimethyl-9H-xanthene-4-carboxylic acid (**9**).

Bond	Length	Bond	Length
C(1)-O(1)	1.377(3)	C(13)-O(1)	1.382(3)
C(1)-C(6)	1.380(4)	C(14)-C(17)	1.529(5)
C(1)-C(2)	1.402(4)	C(14)-C(15)	1.529(5)
C(2)-C(3)	1.387(4)	C(14)-C(16)	1.535(5)
C(2)-C(31)	1.487(4)	C(20)-C(23)	1.427(11)
C(3)-C(4)	1.380(4)	C(20)-C(21A)	1.486(8)
C(4)-C(5)	1.390(4)	C(20)-C(22A)	1.508(8)
C(4)-C(14)	1.536(4)	C(20)-C(21)	1.531(9)
C(5)-C(6)	1.397(4)	C(20)-C(23A)	1.607(8)
C(6)-C(7)	1.531(4)	C(20)-C(22)	1.614(10)
C(7)-C(8)	1.529(4)	C(24)-C(29)	1.379(4)
C(7)-C(19)	1.534(5)	C(24)-C(25)	1.413(4)
C(7)-C(18)	1.535(5)	C(25)-C(26)	1.386(5)
C(8)-C(13)	1.382(4)	C(26)-C(27)	1.354(5)
C(8)-C(9)	1.392(4)	C(27)-O(4)	1.379(4)
C(9)-C(10)	1.389(4)	C(27)-C(28)	1.396(5)
C(10)-C(11)	1.393(4)	C(28)-C(29)	1.419(4)
C(10)-C(20)	1.539(4)	C(28)-C(30)	1.437(5)
C(11)-C(12)	1.389(4)	C(30)-O(5)	1.233(4)
C(12)-C(13)	1.397(4)	C(31)-O(2)	1.218(4)
C(12)-C(24)	1.495(4)	C(31)-O(3)	1.310(4)

Table S1.4. Bond angles [°] for 2,7-di-*tert*-butyl-5-(3-formyl-4-hydroxyphenyl)-9,9-dimethyl-9H-xanthene-4-carboxylic acid (**9**).

Bond Angle	Angle	Bond Angle	Angle
O(1)-C(1)-C(6)	122.6(3)	O(1)-C(13)-C(8)	123.1(3)
O(1)-C(1)-C(2)	115.9(3)	O(1)-C(13)-C(12)	114.9(3)
C(6)-C(1)-C(2)	121.4(3)	C(8)-C(13)-C(12)	122.0(3)
C(3)-C(2)-C(1)	118.6(3)	C(17)-C(14)-C(15)	108.9(3)
C(3)-C(2)-C(31)	119.2(3)	C(17)-C(14)-C(16)	108.0(3)
C(1)-C(2)-C(31)	122.2(3)	C(15)-C(14)-C(16)	108.9(3)
C(4)-C(3)-C(2)	122.2(3)	C(17)-C(14)-C(4)	112.4(3)
C(3)-C(4)-C(5)	117.0(3)	C(15)-C(14)-C(4)	109.0(3)
C(3)-C(4)-C(14)	120.2(3)	C(16)-C(14)-C(4)	109.5(3)
C(5)-C(4)-C(14)	122.8(3)	C(23)-C(20)-C(21A)	126.9(6)
C(4)-C(5)-C(6)	123.4(3)	C(23)-C(20)-C(22A)	61.8(6)
C(1)-C(6)-C(5)	117.3(3)	C(21A)-C(20)-C(22A)	113.5(5)
C(1)-C(6)-C(7)	122.0(3)	C(23)-C(20)-C(21)	113.5(6)
C(5)-C(6)-C(7)	120.8(3)	C(21A)-C(20)-C(21)	30.9(4)
C(8)-C(7)-C(6)	110.3(2)	C(22A)-C(20)-C(21)	136.6(5)
C(8)-C(7)-C(19)	109.3(3)	C(23)-C(20)-C(10)	115.7(5)
C(6)-C(7)-C(19)	108.7(3)	C(21A)-C(20)-C(10)	114.8(4)
C(8)-C(7)-C(18)	109.0(3)	C(22A)-C(20)-C(10)	110.0(4)
C(6)-C(7)-C(18)	109.2(3)	C(21)-C(20)-C(10)	109.8(4)
C(19)-C(7)-C(18)	110.4(3)	C(23)-C(20)-C(23A)	42.4(5)
C(13)-C(8)-C(9)	117.6(3)	C(21A)-C(20)-C(23A)	105.3(5)
C(13)-C(8)-C(7)	121.4(3)	C(22A)-C(20)-C(23A)	103.7(5)
C(9)-C(8)-C(7)	121.0(3)	C(21)-C(20)-C(23A)	78.9(5)
C(10)-C(9)-C(8)	123.2(3)	C(10)-C(20)-C(23A)	108.8(3)
C(9)-C(10)-C(11)	116.8(3)	C(23)-C(20)-C(22)	107.7(6)
C(9)-C(10)-C(20)	121.4(3)	C(21A)-C(20)-C(22)	73.6(5)
C(11)-C(10)-C(20)	121.8(3)	C(22A)-C(20)-C(22)	48.7(4)
C(12)-C(11)-C(10)	122.6(3)	C(21)-C(20)-C(22)	104.3(6)
C(11)-C(12)-C(13)	117.9(3)	C(10)-C(20)-C(22)	104.8(4)
C(11)-C(12)-C(24)	120.2(3)	C(23A)-C(20)-C(22)	142.8(5)
C(13)-C(12)-C(24)	121.9(3)	C(29)-C(24)-C(25)	117.2(3)

O(1)-C(1)-C(6)	122.6(3)	O(1)-C(13)-C(8)	123.1(3)
O(1)-C(1)-C(2)	115.9(3)	O(1)-C(13)-C(12)	114.9(3)
C(29)-C(24)-C(12)	121.3(3)	C(27)-C(28)-C(30)	122.3(3)
C(25)-C(24)-C(12)	121.4(3)	C(29)-C(28)-C(30)	119.7(3)
C(26)-C(25)-C(24)	122.1(3)	C(24)-C(29)-C(28)	121.5(3)
C(27)-C(26)-C(25)	119.2(3)	O(5)-C(30)-C(28)	123.4(4)
C(26)-C(27)-O(4)	118.6(3)	O(2)-C(31)-O(3)	122.8(3)
C(26)-C(27)-C(28)	121.9(3)	O(2)-C(31)-C(2)	123.6(3)
O(4)-C(27)-C(28)	119.5(3)	O(3)-C(31)-C(2)	113.6(3)
C(27)-C(28)-C(29)	118.0(3)	C(1)-O(1)-C(13)	119.0(2)

Table S1.5. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 2,7-di-*tert*-butyl-5-(3-formyl-4-hydroxyphenyl)-9,9-dimethyl-9H-xanthene-4-carboxylic acid (**9**). The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U_{11} + \dots + 2hka^*b^*U_{12}]$.

Atom	U ₁₁	U ₂₂	U ₃₃	U ₂₃	U ₁₃	U ₁₂
C(1)	22(2)	19(2)	25(2)	1(1)	-2(1)	-5(1)
C(2)	23(2)	24(2)	27(2)	-2(1)	-1(1)	-7(1)
C(3)	31(2)	20(2)	31(2)	-6(1)	-1(1)	-5(1)
C(4)	34(2)	19(2)	29(2)	-1(1)	-4(1)	-9(1)
C(5)	40(2)	27(2)	21(2)	-3(1)	4(1)	-13(2)
C(6)	26(2)	24(2)	24(2)	-5(1)	2(1)	-9(1)
C(7)	31(2)	23(2)	25(2)	-5(1)	4(1)	-5(1)
C(8)	25(2)	26(2)	25(2)	-5(1)	0(1)	-7(1)
C(9)	28(2)	27(2)	27(2)	-8(1)	1(1)	-4(1)
C(10)	27(2)	22(2)	31(2)	-1(1)	-5(1)	-4(1)
C(11)	30(2)	24(2)	28(2)	3(1)	-1(1)	-7(1)
C(12)	23(2)	23(2)	27(2)	-3(1)	-3(1)	-3(1)
C(13)	21(2)	21(2)	28(2)	-5(1)	-2(1)	-3(1)
C(14)	55(2)	23(2)	31(2)	2(2)	-1(2)	-9(2)
C(15)	66(3)	38(2)	57(3)	10(2)	-25(2)	-13(2)
C(16)	74(3)	33(2)	47(2)	1(2)	4(2)	-26(2)
C(17)	122(4)	38(2)	29(2)	4(2)	-3(2)	-22(2)
C(18)	62(3)	28(2)	28(2)	-5(2)	-8(2)	-8(2)
C(19)	39(2)	34(2)	52(2)	-1(2)	14(2)	-9(2)
C(20)	38(2)	20(2)	42(2)	-6(2)	2(2)	-3(2)
C(24)	26(2)	22(2)	30(2)	4(1)	-3(1)	0(1)
C(25)	34(2)	30(2)	32(2)	-5(2)	-1(2)	-9(2)
C(26)	42(2)	35(2)	39(2)	-6(2)	-8(2)	-12(2)
C(27)	33(2)	27(2)	34(2)	-3(2)	-5(2)	-2(2)
C(28)	27(2)	23(2)	37(2)	2(1)	-3(2)	-5(1)
C(29)	31(2)	25(2)	33(2)	-3(1)	-3(2)	-4(1)
C(30)	34(2)	39(2)	52(3)	-5(2)	1(2)	-10(2)
C(31)	27(2)	18(2)	28(2)	-2(1)	-1(1)	-3(1)
O(1)	35(1)	19(1)	23(1)	-2(1)	0(1)	0(1)

O(2)	52(2)	24(1)	31(1)	−8(1)	11(1)	−11(1)
O(3)	69(2)	25(1)	32(1)	−12(1)	16(1)	−18(1)
O(4)	57(2)	55(2)	32(1)	−14(1)	8(1)	−18(1)
O(5)	59(2)	60(2)	42(2)	−11(1)	19(1)	−22(1)

Table S1.6. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 2,7-di-*tert*-butyl-5-(3-formyl-4-hydroxyphenyl)-9,9-dimethyl-9H-xanthene-4-carboxylic acid (**9**).

Atom	x	y	z	U(eq)
H(3)	602	11522	2362	34
H(5)	2981	8872	620	36
H(9)	5260	3272	1840	34
H(11)	4174	2091	4389	35
H(15A)	−680	13322	165	84
H(15B)	−788	12827	1198	84
H(15C)	−813	11773	555	84
H(16A)	2962	12473	771	76
H(16B)	1527	13204	1360	76
H(16C)	1616	13769	334	76
H(17A)	1474	10659	−425	99
H(17B)	2933	10905	−317	99
H(17C)	1594	12231	−739	99
H(18A)	4055	4888	760	62
H(18B)	3461	6581	356	62
H(18C)	2450	5908	1038	62
H(19A)	5928	6550	1861	68
H(19B)	5556	7007	860	68
H(19C)	6251	5321	1262	68
H(21A)	3894	147	3090	75
H(21B)	4477	−28	4023	75
H(21C)	5316	−1203	3388	75
H(22A)	7580	−847	3662	84
H(22B)	6664	400	4237	84
H(22C)	7716	727	3428	84
H(23A)	6720	−580	2116	109
H(23B)	7140	846	1977	109
H(23C)	5604	964	1709	109
H(21D)	4351	−90	3840	86
H(21E)	5788	−285	4274	86

H(21F)	5855	−1317	3593	86
H(22D)	7851	−544	2706	95
H(22E)	7732	647	3292	95
H(22F)	7482	1127	2275	95
H(23D)	5465	1052	1613	69
H(23E)	4186	629	2182	69
H(23F)	5749	−599	2059	69
H(25)	3760	6173	4999	39
H(26)	2850	6638	6421	47
H(29)	1569	3403	5137	38
H(30)	−41	3064	6441	52
H(3A)	322	11042	4513	64
H(4)	554	5185	7666	73

Figure S2. Thermal ellipsoid plot of 5-(boronic acid)-2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthene-4-carboxylic acid methyl ester (**24**). Thermal ellipsoids are drawn at the 50% probability level.

Table S2.1. Crystal data and structure refinement for 5-(boronic acid)-2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthene-4-carboxylic acid methyl ester (**24**).

Identification code	030780a	
Empirical formula	C ₂₅ H ₃₃ BO ₅	
Formula weight	424.32	
Temperature	467(2) K	
Wavelength	0.71073 Å	
Crystal system	Triclinic	
Space group	$P\bar{1}$	
Unit cell dimensions	$a = 9.157(7)$ Å	$\alpha = 77.035(14)^\circ$
	$b = 11.143(10)$ Å	$\beta = 83.053(17)^\circ$
	$c = 12.500(11)$ Å	$\gamma = 79.421(18)^\circ$
Volume	1217.6(18) Å ³	
<i>Z</i>	2	
Density (calculated)	1.157 Mg/m ³	
Absorption coefficient	0.078 mm ⁻¹	
<i>F</i> (000)	456	
Crystal size	0.20 × 0.17 × 0.04 mm ³	
θ range for data collection	2.26 to 24.69°	
Index ranges	$-8 \leq h \leq 7, -11 \leq k \leq 12, -5 \leq \ell \leq 14$	
Reflections collected	2658	
Independent reflections	2654 [$R_{\text{int}} = 0.0508$]	
Completeness to $\theta = 24.69^\circ$	63.9 %	
Absorption correction	None	
Refinement method	Full-matrix least-squares on F^2	
Data / restraints / parameters	2654 / 0 / 291	
Goodness-of-fit on F^2	1.083	
Final <i>R</i> indices [$I > 2\sigma(I)$]	$R_I = 0.0970, wR_2 = 0.1710$	
<i>R</i> indices (all data)	$R_I = 0.1935, wR_2 = 0.2057$	
Largest diff. peak and hole	0.201 and -0.212 e/Å ³	

Tables S2.2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 5-(boronic acid)-2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthene-4-carboxylic acid methyl ester (**24**). U(eq) is defined as one third of the trace of the orthogonalized U_{ij} tensor.

Atom	x	y	z	U(eq)
B(1)	13069(11)	5899(8)	5728(7)	28(3)
O(1)	10585(5)	5580(4)	1694(4)	49(2)
O(2)	11546(5)	5096(4)	3323(4)	47(2)
O(3)	10351(5)	6702(4)	4621(3)	31(1)
O(4)	13180(5)	5266(5)	4906(4)	41(2)
O(5)	14354(5)	5789(5)	6219(4)	45(2)
C(1)	11852(9)	4767(7)	1304(6)	62(3)
C(2)	10577(9)	5675(6)	2750(6)	33(2)
C(3)	9208(8)	6502(6)	3104(5)	29(2)
C(4)	7976(8)	6787(6)	2448(5)	34(2)
C(5)	6676(7)	7557(6)	2734(5)	29(2)
C(6)	5343(9)	7909(7)	2019(6)	38(2)
C(7)	5169(10)	9299(7)	1528(6)	88(3)
C(8)	5533(9)	7184(9)	1106(7)	97(4)
C(9)	3913(8)	7650(8)	2735(6)	65(3)
C(10)	6663(8)	8017(6)	3663(5)	34(2)
C(11)	7830(7)	7785(6)	4335(5)	27(2)
C(12)	9098(8)	6996(6)	4026(5)	30(2)
C(13)	10349(8)	7162(6)	5572(5)	27(2)
C(14)	9137(7)	7964(6)	5928(5)	27(2)
C(15)	7698(7)	8386(6)	5333(5)	31(2)
C(16)	6373(7)	8017(7)	6123(5)	50(3)
C(17)	7523(7)	9826(5)	4916(5)	44(2)
C(18)	9292(8)	8383(6)	6870(5)	38(2)
C(19)	10531(8)	8038(6)	7464(5)	31(2)
C(20)	10566(9)	8540(7)	8503(5)	39(2)
C(21)	9319(11)	8128(9)	9350(6)	93(3)
C(22)	12056(10)	8103(9)	9019(7)	93(4)
C(23)	10328(10)	9978(7)	8199(6)	77(3)
C(24)	11694(8)	7234(6)	7047(5)	32(2)
C(25)	11651(8)	6775(6)	6108(5)	28(2)

Table S2.3. Bond lengths (Å) for 5-(boronic acid)-2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthene-4-carboxylic acid methyl ester (**24**).

Bond	Length	Bond	Length
B(1)-O(4)	1.356(8)	C(10)-C(11)	1.390(8)
B(1)-O(5)	1.365(9)	C(11)-C(12)	1.392(8)
B(1)-C(25)	1.568(10)	C(11)-C(15)	1.526(8)
O(1)-C(2)	1.346(7)	C(13)-C(14)	1.386(8)
O(1)-C(1)	1.442(7)	C(13)-C(25)	1.390(8)
O(2)-C(2)	1.206(7)	C(14)-C(18)	1.391(8)
O(3)-C(12)	1.391(7)	C(14)-C(15)	1.537(8)
O(3)-C(13)	1.396(7)	C(15)-C(16)	1.525(9)
C(2)-C(3)	1.497(9)	C(15)-C(17)	1.556(8)
C(3)-C(12)	1.371(8)	C(18)-C(19)	1.379(9)
C(3)-C(4)	1.422(8)	C(19)-C(24)	1.390(8)
C(4)-C(5)	1.393(8)	C(19)-C(20)	1.532(8)
C(5)-C(10)	1.369(8)	C(20)-C(21)	1.522(10)
C(5)-C(6)	1.541(9)	C(20)-C(22)	1.530(10)
C(6)-C(8)	1.516(9)	C(20)-C(23)	1.542(9)
C(6)-C(7)	1.519(10)	C(24)-C(25)	1.390(8)
C(6)-C(9)	1.529(9)		

Table S2.4. Bond angles (°) for 5-(boronic acid)-2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthene-4-carboxylic acid methyl ester (**24**).

Bond Angle	Angle	Bond Angle	Angle
O(4)-B(1)-O(5)	115.2(7)	O(3)-C(12)-C(11)	121.5(6)
O(4)-B(1)-C(25)	126.2(8)	C(14)-C(13)-C(25)	123.4(6)
O(5)-B(1)-C(25)	118.5(7)	C(14)-C(13)-O(3)	121.6(6)
C(2)-O(1)-C(1)	114.9(5)	C(25)-C(13)-O(3)	114.9(6)
C(12)-O(3)-C(13)	120.5(5)	C(13)-C(14)-C(18)	116.0(6)
O(2)-C(2)-O(1)	121.8(6)	C(13)-C(14)-C(15)	123.2(6)
O(2)-C(2)-C(3)	126.3(6)	C(18)-C(14)-C(15)	120.9(6)
O(1)-C(2)-C(3)	111.8(6)	C(16)-C(15)-C(11)	110.1(6)
C(12)-C(3)-C(4)	118.9(6)	C(16)-C(15)-C(14)	109.7(5)
C(12)-C(3)-C(2)	122.9(7)	C(11)-C(15)-C(14)	109.8(5)
C(4)-C(3)-C(2)	118.2(6)	C(16)-C(15)-C(17)	111.4(5)
C(5)-C(4)-C(3)	120.7(6)	C(11)-C(15)-C(17)	108.2(5)
C(10)-C(5)-C(4)	116.6(6)	C(14)-C(15)-C(17)	107.7(6)
C(10)-C(5)-C(6)	121.7(6)	C(19)-C(18)-C(14)	124.7(6)
C(4)-C(5)-C(6)	121.7(6)	C(18)-C(19)-C(24)	115.5(6)
C(8)-C(6)-C(7)	109.8(7)	C(18)-C(19)-C(20)	120.2(6)
C(8)-C(6)-C(9)	108.1(7)	C(24)-C(19)-C(20)	124.3(6)
C(7)-C(6)-C(9)	108.1(7)	C(21)-C(20)-C(22)	109.1(7)
C(8)-C(6)-C(5)	112.6(6)	C(21)-C(20)-C(19)	109.7(7)
C(7)-C(6)-C(5)	108.3(6)	C(22)-C(20)-C(19)	112.6(6)
C(9)-C(6)-C(5)	109.9(6)	C(21)-C(20)-C(23)	109.0(7)
C(5)-C(10)-C(11)	125.6(6)	C(22)-C(20)-C(23)	107.5(7)
C(10)-C(11)-C(12)	115.8(6)	C(19)-C(20)-C(23)	108.9(6)
C(10)-C(11)-C(15)	120.8(6)	C(19)-C(24)-C(25)	124.0(6)
C(12)-C(11)-C(15)	123.5(6)	C(13)-C(25)-C(24)	116.4(6)
C(3)-C(12)-O(3)	116.1(6)	C(13)-C(25)-B(1)	125.7(6)
C(3)-C(12)-C(11)	122.4(7)	C(24)-C(25)-B(1)	117.9(7)

Table S2.5. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 5-(boronic acid)-2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthene-4-carboxylic acid methyl ester (**24**). The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U_{11} + \dots + 2hka^*b^*U_{12}]$.

Atom	U ₁₁	U ₂₂	U ₃₃	U ₂₃	U ₁₃	U ₁₂
B(1)	20(7)	35(5)	35(5)	1(4)	-11(5)	-24(5)
O(1)	52(4)	54(3)	37(3)	-21(3)	0(3)	8(3)
O(2)	39(4)	54(3)	50(3)	-23(3)	-13(3)	11(3)
O(3)	25(4)	37(3)	32(3)	-13(2)	-6(2)	4(3)
O(4)	23(4)	52(3)	46(3)	-16(3)	-12(2)	9(3)
O(5)	29(4)	65(4)	50(3)	-29(3)	-18(3)	4(3)
C(1)	69(7)	59(5)	53(5)	-28(4)	9(4)	15(5)
C(2)	32(6)	33(4)	35(4)	-4(4)	-2(4)	-15(4)
C(3)	23(5)	35(4)	29(4)	-4(3)	0(4)	-10(4)
C(4)	39(6)	40(5)	22(4)	-7(3)	4(4)	-6(4)
C(5)	27(5)	40(4)	22(4)	-8(3)	-1(3)	-12(4)
C(6)	23(6)	59(5)	35(4)	-6(4)	-18(4)	-12(5)
C(7)	97(9)	83(7)	73(6)	27(5)	-55(6)	-8(6)
C(8)	50(7)	167(10)	93(7)	-78(7)	-37(6)	17(7)
C(9)	30(6)	113(7)	47(5)	2(5)	-16(4)	-12(6)
C(10)	23(5)	44(5)	30(4)	-3(4)	-7(4)	9(4)
C(11)	15(5)	27(4)	38(4)	-6(3)	-10(4)	-1(4)
C(12)	28(6)	33(4)	32(4)	-5(3)	-13(4)	-8(4)
C(13)	24(5)	38(5)	20(4)	-4(3)	1(3)	-12(4)
C(14)	25(5)	29(4)	22(4)	-2(3)	4(3)	2(4)
C(15)	20(5)	31(4)	37(4)	-8(3)	-14(3)	15(4)
C(16)	25(6)	83(6)	33(4)	-19(4)	5(4)	11(5)
C(17)	45(6)	41(5)	42(4)	-8(4)	-13(4)	10(4)
C(18)	36(6)	39(5)	37(4)	-17(4)	6(4)	0(4)
C(19)	32(6)	34(4)	23(4)	3(3)	-9(3)	-6(4)
C(20)	39(6)	55(5)	30(4)	-15(4)	-9(4)	-16(5)
C(21)	106(9)	142(9)	54(6)	-49(6)	20(6)	-59(8)
C(22)	76(8)	142(9)	79(7)	-68(6)	-42(6)	19(7)
C(23)	113(9)	65(6)	67(6)	-39(5)	-30(5)	-2(6)
C(24)	22(5)	30(4)	40(4)	0(3)	-7(4)	-3(4)
C(25)	17(5)	29(4)	45(5)	-15(4)	-5(4)	-13(4)

Table S2.6. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for 5-(boronic acid)-2,7-di-*tert*-butyl-9,9-dimethyl-9H-xanthene-4-carboxylic acid methyl ester (**24**).

Atom	x	y	z	U(eq)
H(4)	12395	5433	4611	61
H(5)	14177	6123	6753	68
H(1A)	12752	5058	1375	94
H(1B)	11780	4767	544	94
H(1C)	11867	3935	1733	94
H(4A)	8039	6456	1821	41
H(7A)	6050	9481	1065	132
H(7B)	5025	9753	2110	132
H(7C)	4321	9541	1099	132
H(8A)	4699	7461	665	145
H(8B)	5585	6310	1422	145
H(8C)	6435	7323	653	145
H(9A)	3074	7948	2300	98
H(9B)	3794	8071	3338	98
H(9C)	3973	6768	3016	98
H(10)	5798	8530	3862	41
H(16A)	5474	8300	5754	74
H(16B)	6304	8393	6751	74
H(16C)	6507	7125	6360	74
H(17A)	6648	10112	4519	66
H(17B)	8384	10031	4438	66
H(17C)	7430	10222	5533	66
H(18)	8503	8936	7116	45
H(21A)	9481	7234	9564	140
H(21B)	8380	8409	9035	140
H(21C)	9312	8480	9985	140
H(22A)	12039	8470	9646	140
H(22B)	12848	8354	8486	140
H(22C)	12215	7210	9247	140
H(23A)	9401	10281	7869	116

X-tal Structural data for 24

H(23B)	11132	10246	7687	116
H(23C)	10303	10302	8852	116
H(24)	12554	6989	7420	38

Table S3.1. Bond distance parameters from DFT calculated energy minimized structures of the hydroperoxide complexes of Mn(HSX*) and Mn(H_{ph}SX*).

Distance (Å)	Mn(HSX*)OOH	Mn(H _{ph} SX*)OOH
Mn-O	1.89 Å	1.85 Å
peroxide O-O	1.46 Å	1.46 Å
peroxide O-H	0.99	0.99 Å
acid OH- Mn	5.14 Å, 6.77 Å	4.58 Å, 6.03 Å
	peroxide H to acid OH :	peroxide H to acid CO :
peroxide-acid	1.94 Å	3.11 Å
acid dimer (OH-CO)	no dimer	2.57 Å, 2.68 Å

Table S3.2 Total bond energies of calculated compounds.

	Total Energy (kJ/mol)
Mn(HSX*)-OOH	−56695.92
Mn(HphSX*)-OOH (carboxylic acid dimer)	−69749.31
Mn(HphSX*)-OOH (no carboxylic acid dimer)	−69704.88

Table S3.3 Cartesian coordinates from the DFT calculated energy minimized structure of the hydroperoxide complex of Mn(HSX*). The structure is shown in Figure 3(a).

	x	y	z
1.C	10.54703	-3.38522	-3.3624
2.C	9.68632	-2.64592	-2.55441
3.C	8.311687	-2.76836	-2.75099
4.C	7.783106	-3.63145	-3.72707
5.C	8.682754	-4.36068	-4.51001
6.C	10.05356	-4.24241	-4.33893
7.O	7.416122	-2.0485	-1.96717
8.C	7.862076	-0.94236	-1.24597
9.C	9.218721	-0.75202	-0.9971
10.C	10.21893	-1.77446	-1.45312
11.C	6.884345	-0.0688	-0.7487
12.C	7.324951	1.027343	0.001137
13.C	8.675675	1.254335	0.240083
14.C	9.615952	0.364363	-0.26163
15.C	5.427121	-0.25561	-0.94458
16.C	4.814239	-0.16331	-2.2162
17.C	3.44543	-0.20198	-2.36716
18.C	2.57427	-0.34933	-1.24733
19.C	3.203745	-0.47449	0.049393
20.C	4.608608	-0.41023	0.159554
21.O	1.287077	-0.34475	-1.42562
22.Mn	-0.16069	-0.86555	-0.23384
23.O	-0.6646	-2.5896	0.3577
24.O	-2.07419	-2.96625	0.295007
25.C	2.429979	-0.60886	1.24072
26.N	1.137132	-0.75186	1.294269
27.C	0.434907	-0.76124	2.575628
28.C	-0.67066	0.304545	2.506114
29.N	-1.35727	0.138633	1.233124
30.C	-2.63639	0.283297	1.136077
31.C	-3.39444	0.07964	-0.06945

32.C	-2.80265	-0.30302	-1.316
33.C	-3.66507	-0.45679	-2.42489
34.C	-5.03447	-0.30466	-2.30187
35.C	-5.62856	0.015866	-1.0648
36.C	-4.79101	0.222814	0.020031
37.O	-1.51506	-0.5233	-1.49761
38.C	-7.09904	0.10156	-0.94635
39.C	-7.83231	0.931162	-1.80171
40.C	-9.22335	0.979313	-1.7475
41.C	-9.90188	0.188831	-0.82458
42.C	-9.20428	-0.64331	0.053971
43.C	-7.81255	-0.67904	-0.01552
44.O	-7.0592	-1.48972	0.813131
45.C	-7.65478	-2.53137	1.48274
46.C	-9.04792	-2.58636	1.623098
47.C	-9.91558	-1.47852	1.081779
48.C	-6.83248	-3.54502	2.033475
49.C	-7.45473	-4.59552	2.719975
50.C	-8.8336	-4.66286	2.858963
51.C	-9.62136	-3.65903	2.303248
52.C	-5.33755	-3.66437	1.922399
53.O	-4.73639	-2.61443	1.329674
54.C	6.320222	-3.86776	-4.03141
55.O	5.401275	-3.2661	-3.24738
56.O	-4.73514	-4.64188	2.338189
57.O	5.975872	-4.58544	-4.94947
58.H	-3.77078	-2.82908	1.23245
59.H	5.849483	-2.71298	-2.56649
60.H	8.254883	-5.0207	-5.26105
61.H	10.73623	-4.81956	-4.95944
62.H	11.62155	-3.29034	-3.20684
63.H	10.49567	-2.41122	-0.59286
64.H	11.13619	-1.27075	-1.78935
65.H	10.67786	0.520843	-0.07428
66.H	8.991218	2.126241	0.809769
67.H	6.575785	1.725685	0.37136

68.H	-6.8036	-5.36913	3.125449
69.H	-9.29376	-5.49536	3.391576
70.H	-10.7081	-3.69612	2.401853
71.H	-10.992	0.218998	-0.77225
72.H	-9.77516	1.637244	-2.41979
73.H	-7.28691	1.557318	-2.5098
74.H	-10.8272	-1.91062	0.638539
75.H	-10.2388	-0.82912	1.918061
76.H	-3.2148	-0.72904	-3.38048
77.H	-5.6749	-0.47482	-3.1698
78.H	-5.22411	0.485137	0.985485
79.H	2.988855	-0.11889	-3.3517
80.H	5.442582	-0.03236	-3.09736
81.H	5.059577	-0.49095	1.149005
82.H	1.121107	-0.55949	3.414924
83.H	-0.02263	-1.75599	2.701444
84.H	-0.20891	1.308071	2.529538
85.H	-1.35302	0.208945	3.369173
86.H	-2.1853	-3.16188	-0.66379
87.H	2.982576	-0.57186	2.192149
88.H	-3.22249	0.563181	2.027737

Table S3.4 Cartesian coordinates from the DFT calculated energy minimized structure of the hydroperoxide complex of Mn(H_{ph}SX*) with the xanthenes oriented so the carboxylic acids can form a hydrogen bonded dimer. The structure is shown in Figure 3(b).

	x	y	z
1.C	-3.75165	4.546489	14.31793
2.C	-3.38406	3.626824	15.30982
3.C	-2.8434	2.395334	14.92456
4.C	-2.68628	2.083854	13.58282
5.C	-3.06025	3.004904	12.59827
6.C	-3.58758	4.242615	12.97275
7.C	-3.60206	3.958867	16.73365
8.C	-2.56721	3.891009	17.68142
9.C	-2.77102	4.23596	19.01614
10.C	-4.04147	4.653395	19.41208
11.C	-5.08914	4.71473	18.49963
12.C	-4.86443	4.370164	17.17086
13.C	-1.63375	4.113736	19.99311
14.C	-0.29351	4.125105	19.30752
15.C	-0.20586	3.78953	17.95878
16.O	-1.33477	3.477823	17.22283
17.C	1.021914	3.710811	17.27923
18.C	2.176114	3.978908	18.02141
19.C	2.115825	4.326536	19.3671
20.C	0.882194	4.40687	20.00279
21.C	1.11115	3.36242	15.84516
22.C	0.330315	4.025557	14.87198
23.C	0.413865	3.706708	13.53512
24.C	1.281166	2.691984	13.06558
25.C	2.13297	2.065424	14.03725
26.C	2.009736	2.405781	15.40133
27.C	3.137512	1.125374	13.65775
28.N	3.331359	0.699528	12.44032
29.C	4.394219	-0.26124	12.15586
30.C	3.868309	-1.22721	11.08788

31.N	3.242431	-0.423	10.04361
32.C	3.509082	-0.6284	8.784802
33.C	3.007294	0.125546	7.684865
34.C	2.209732	1.311089	7.875835
35.C	1.736861	1.958986	6.698018
36.C	2.071322	1.513461	5.438487
37.C	2.928479	0.399599	5.246521
38.C	3.364036	-0.27635	6.374625
39.C	3.459637	0.040274	3.909948
40.C	2.648788	-0.24499	2.794942
41.C	3.186567	-0.58831	1.553547
42.C	4.57522	-0.60557	1.405997
43.C	5.406733	-0.31751	2.483065
44.C	4.846457	-0.00795	3.720003
45.C	2.279383	-0.99611	0.421165
46.C	0.856353	-0.55775	0.641278
47.C	0.431035	-0.20078	1.921345
48.O	1.284142	-0.19645	3.002532
49.C	-0.89868	0.185909	2.177049
50.C	-1.79941	0.201045	1.107698
51.C	-1.39444	-0.15816	-0.17648
52.C	-0.07203	-0.5304	-0.40197
53.C	-1.31587	0.571328	3.542132
54.C	-1.92146	1.812961	3.783046
55.C	-2.25587	2.200459	5.077877
56.C	-1.98531	1.343799	6.154086
57.C	-1.41963	0.086463	5.90944
58.C	-1.09142	-0.29527	4.619417
59.O	1.932871	1.80829	9.040583
60.Mn	2.017818	0.941913	10.83467
61.O	1.273912	2.407007	11.78954
62.C	-2.24824	1.724205	7.56468
63.O	-2.774	2.917394	7.745202
64.C	-2.88906	2.675209	11.16074
65.O	-2.53684	1.425547	10.95134
66.O	0.535003	-0.20666	11.02656

67.O	0.250044	-0.84851	9.744538
68.O	-1.96951	0.937607	8.494387
69.O	-3.06877	3.51929	10.25959
70.H	-2.83754	0.474442	1.302656
71.H	-2.11176	-0.15753	-0.99779
72.H	0.252183	-0.81884	-1.40393
73.H	2.306334	-2.09848	0.319297
74.H	2.655165	-0.57658	-0.52584
75.H	5.000639	-0.85511	0.432238
76.H	6.4899	-0.31843	2.358736
76.H	6.4899	-0.31843	2.358736
77.H	5.487723	0.252168	4.563406
78.H	-5.67994	4.39356	16.44961
79.H	-6.08157	5.022087	18.8235
80.H	-4.20853	4.91942	20.45552
81.H	0.822128	4.681388	21.05555
82.H	3.03054	4.549584	19.91288
83.H	3.136077	3.953563	17.50816
84.H	-1.68326	4.932077	20.72549
85.H	-1.74167	3.168577	20.55642
86.H	-0.19761	4.225949	12.79949
87.H	-0.35209	4.813256	15.18598
88.H	2.633856	1.880058	16.12387
89.H	1.111855	2.843248	6.830108
90.H	1.697964	2.051326	4.565478
91.H	3.999387	-1.15685	6.253866
92.H	4.684173	-1.8534	10.68829
93.H	3.095807	-1.88001	11.53033
94.H	5.264114	0.286524	11.75112
95.H	4.711448	-0.81151	13.059
96.H	-0.41679	-0.22796	9.366585
97.H	-2.88509	3.120288	8.762245
98.H	-0.6426	-1.2716	4.439241
99.H	-1.23114	-0.58331	6.745625
100.H	-2.0949	2.494781	2.94807
101.H	-2.70507	3.176113	5.260958

102.H	-2.34413	1.271822	9.951154
103.H	-2.26015	1.128916	13.28726
104.H	-2.54897	1.674658	15.68419
105.H	-4.15206	5.516006	14.61272
106.H	-3.86272	4.958224	12.20067
107.H	4.190518	-1.45728	8.5362
108.H	3.779046	0.736605	14.46404

Figure S3.1 Energy minimized structure obtained from DFT of the hydroperoxide complexes of $\text{Mn}(\text{H}_{\text{ph}}\text{SX}^*-\text{COOH})$ with the xanthenes splayed away from each other, precluding formation of the $-(\text{COOH})_2-$ dimer. Each of the carboxylic acids has a hydrogen-bonding interaction with the hydroperoxide.

Table S3.5 Cartesian coordinates from the DFT calculated energy minimized structure of the hydroperoxide complex of Mn(H_{ph}SX*) with the xanthenes splayed away from each other, precluding formation of the $-(\text{COOH})_2$ - dimer. (The structure is shown in Figure S3.1)

	x	y	z
1.C	0.318602	5.619625	15.67877
2.C	-1.03035	5.288744	15.85218
3.C	-1.80647	5.010826	14.72388
4.C	-1.24529	5.050496	13.45124
5.C	0.107431	5.358143	13.2815
6.C	0.878647	5.657568	14.40875
7.C	-1.6201	5.278902	17.21186
8.C	-2.11217	4.100881	17.78755
9.C	-2.64949	4.071659	19.07316
10.C	-2.68154	5.255792	19.80812
11.C	-2.19993	6.44201	19.26403
12.C	-1.67672	6.448512	17.974
13.C	-3.20089	2.774928	19.59977
14.C	-2.52797	1.59271	18.95657
15.C	-1.98332	1.738529	17.68226
16.O	-2.02149	2.951232	17.02309
17.C	-1.41374	0.660886	16.98784
18.C	-1.40223	-0.58576	17.62143
19.C	-1.92735	-0.75141	18.89896
20.C	-2.48304	0.33765	19.56226
21.C	-0.87982	0.807457	15.61754
22.C	0.104325	1.763175	15.2795
23.C	0.607953	1.866599	13.99913
24.C	0.134787	1.043494	12.95146
25.C	-0.84444	0.056638	13.29364
26.C	-1.3209	-0.03865	14.61848
27.C	-1.31766	-0.88833	12.34079
28.N	-1.0586	-0.86453	11.0687
29.C	-1.54578	-1.93995	10.20082

30.C	-1.84123	-1.30742	8.848462
31.N	-0.68379	-0.46505	8.531889
32.C	-0.11339	-0.54823	7.362761
33.C	1.03234	0.181635	6.943407
34.C	1.653	1.157236	7.788614
35.C	2.751387	1.868368	7.240496
36.C	3.238248	1.570288	5.983477
37.C	2.684113	0.54916	5.169879
38.C	1.567746	-0.1046	5.666886
39.C	3.344486	0.149931	3.905987
40.C	2.643765	-0.23746	2.741226
41.C	3.303061	-0.66561	1.587981
42.C	4.699641	-0.68156	1.578923
43.C	5.422227	-0.29293	2.701968
44.C	4.7454	0.106703	3.850936
45.C	2.520279	-1.14983	0.393329
46.C	1.056677	-0.79951	0.466635
47.C	0.508408	-0.3362	1.661119
48.O	1.267409	-0.1736	2.80125
49.C	-0.86574	-0.05478	1.786744
50.C	-1.68432	-0.25771	0.670982
51.C	-1.15558	-0.70771	-0.53681
52.C	0.208347	-0.97313	-0.63097
53.C	-1.43785	0.417976	3.066854
54.C	-0.91968	1.543641	3.720266
55.C	-1.43228	1.948707	4.94659
56.C	-2.47937	1.236602	5.543434
57.C	-3.03863	0.14502	4.869497
58.C	-2.519	-0.26422	3.646006
59.O	1.262347	1.418726	9.005161
60.Mn	-0.09393	0.616229	10.08887
61.O	0.61384	1.212467	11.74124
62.C	-2.91828	1.51337	6.949663
63.O	-2.24575	2.522214	7.507636
64.C	0.784147	5.441983	11.93933
65.O	0.216481	4.806699	10.89687

66.O	-1.60327	1.950353	9.950737
67.O	-1.86347	2.95312	11.00138
68.O	-3.75027	0.810921	7.521208
69.O	1.815217	6.075576	11.79406
70.H	-2.74756	-0.02911	0.75873
71.H	-1.80473	-0.85078	-1.40125
72.H	0.631984	-1.33507	-1.57025
73.H	2.625178	-2.24968	0.321945
74.H	2.954824	-0.72685	-0.52722
75.H	5.218955	-1.01916	0.680064
76.H	6.512325	-0.32059	2.692441
77.H	5.310193	0.362008	4.748086
78.H	-1.3059	7.37239	17.53514
79.H	-2.23651	7.361413	19.84392
80.H	-3.10162	5.24174	20.81458
81.H	-2.90872	0.213146	20.55823
82.H	-1.90116	-1.72979	19.37691
83.H	-0.94288	-1.4274	17.10416
84.H	-3.0875	2.726311	20.69039
85.H	-4.28774	2.729835	19.38454
86.H	1.378574	2.600403	13.76485
87.H	0.485305	2.425654	16.05639
88.H	-2.08101	-0.78558	14.84409
89.H	3.202437	2.663417	7.841351
90.H	4.080114	2.154097	5.592003
91.H	1.090486	-0.88679	5.079501
92.H	-2.00464	-2.06467	8.070056
93.H	-2.72927	-0.65756	8.909303
94.H	-0.73716	-2.68945	10.09072
95.H	-2.43145	-2.42744	10.62344
96.H	-2.09593	2.384333	11.76924
97.H	-2.20382	2.406437	8.528486
98.H	-2.92749	-1.14358	3.143468
99.H	-3.86153	-0.39244	5.341844
100.H	-0.09344	2.093599	3.269636
101.H	-1.00029	2.801417	5.468514

DFT Calculations

102.H	−0.50246	4.160987	11.13767
103.H	−1.88492	4.879932	12.588
104.H	−2.86528	4.782888	14.84809
105.H	0.930679	5.839595	16.5547
106.H	1.921994	5.928019	14.26209
107.H	−1.93624	−1.70839	12.7255
108.H	−0.54179	−1.24444	6.631932

24

25