Supporting Information

Hesseltin A, a novel metabolite from *Penicillium hesseltinei*.

Richard K. Phipps, Bent O. Petersen, Kathrine B. Christensen, Jens Ø. Duus, Jens C. Frisvad, Thomas O. Larsen

[†]*Center for Microbial Biotechnology, BioCentrum-DTU, 2800 Kgs. Lyngby, Denmark*
and [‡]*Carlsberg Laboratory, Gamle Carlsberg Vej 10, DK-2500 Valby, Denmark.*

Experimental procedures

General Experimental Procedures. UV spectra were measured on a Perkin-Elmer UV/VIS spectrometer lambda 2. $[\alpha]_D$ values were measured using a Perkin-Elmer 241 Polarimeter at 589 nm. NMR spectra were recorded at 25°C in DMSO- d_6 on a Varian Unity Inova 800 instrument operating at 799.96 MHz for proton and 201.12 MHz for carbon, and on a Bruker 600 Avance instrument operating at 600.13 MHz for proton and 150.92 MHz for carbon. Spectra were referenced to DMSO at 2.50 ppm for proton and 39.5 ppm for carbon. On the Varian instrument standard programs tndqcosy, tnnoesy (mixing time 400 ms), tntocsy (spinlock time 80 ms), gHSQC and gHMBC were used with digital resolution in F2 dimension <2 Hz/pt. On the Bruker instrument standard programs cosydfprtp, noesyprtp (mixing time 400 ms), mlevprtp (spinlock time 80 ms), inv4gstp and inv4gslplrnd were used with digital resolution in F2 dimension <2 Hz/pt. Mass spectra were acquired using an Agilent HP 1100 liquid chromatograph with a DAD system (Waldbronn, Germany) coupled to an LCT oaTOF mass spectrometer (Micromass, Manchester, UK) with a Z-spray ESI source and a Lock Spray probe. Column chromatography used Waters Sep-Pak Vac 35cc (10g) C₁₈ cartridges. Semi-preparative purification was carried with a Waters 600 system attached to a Waters 996 Photodiode Array detector on a Waters RCM25 C₁₈ column. 12396 DA was gradient elution from 35 % CH₃CN increasing to 50 % CH₃CN over 20 minutes, with a flow rate of 20 mL/min at room temperature. Solvents for extraction and chromatography were of commercial HPLC grade. Culture media YES agar consisted of yeast extract (Difco, 212750) 2%, sucrose 15%, MgSO₄·7H₂O 0.05%, ZnSO₄·7H₂O 0.001%, CuSO₄·5H₂O 0.0005%, agar 2%, H₂O to 1 litre, pH 6.0.

Fermentation and Isolation. *Penicillium hesseltinei* (IBT12396) was three point inoculated onto yeast extract sucrose agar (YES) and incubated for 14 days in the dark at 25°C (for media formulations, see Samson *et al.* 1996¹).

For extraction and purification of the targeted metabolites, the *P. hesseltinei* strain IBT 12396 was 3 point inoculated onto CYA agar (200 x 9 cm Petri dishes) and incubated in darkness at 25°C for 14 days. Following incubation, the agar and fungal colonies were macerated in a stomacher and extracted with an equal volume of EtOAc (solvent: agar). The mixture was left to stand for one day, decanted then filtered by gravity through filter paper (Whatman #4). The solid residue was re-extracted with fresh EtOAc and the extracts combined. The combined extracts were concentrated under vacuum yielding a dark brown residue (998 mg).

The residue was adsorbed to C₁₈ (2 g) and chromatographed on Waters Sep-pak C₁₈ (10 g) disposable cartridges under vacuum in five steps (10:90, 25:75, 50:50, 0:100, 0:100; 0.05% TFA; 200 mL each). The fractions were taken to dryness and evaluated by RP-HPLC-DAD analysis. This showed that the target compounds eluted in the H₂O:MeOH (25:75) fraction (234 mg).

¹ Samson, R.A., Hoekstra, J.C., Frisvad J.C., Filtenborg, O., Introduction to Food- and Air Borne Fungi. 6th Ed., Centraalbureau voor Schimmelcultures: Utrecht, 2002.

Table 1. 1D and 2D NMR data for Hesseltn A (**1**) in DMSO-*d*₆

Atom No.	¹ H (δ ppm, <i>J</i> Hz)	¹³ C/HSQC correlations	HMBC correlations	TOCSY correlations	NOESY correlations
1	1.81 (3H, d, 6.7)	18.4	C2, C3	H2, H3, H4, H5	H2, H3
2	6.16 (1H, dq, 14.3, 6.7)	136.7	C1, C3, C4	H1, H3, H4, H5	H1, H4
3	6.25 (1H, dd, 10.6, 14.3)	130.4	C1, C2, C4, C5	H1, H2, H4, H5	H1, H4, H5
4	6.92 (1H, dd, 10.6, 15.5)	135.1	C2, C3, C5, C6	H1, H2, H3, H5	H2, H3, H5
5	6.22 (1H, d, 15.5)	120.0	C3, C4, C6, C7	H1, H2, H3, H4	H3, H4, H7
6	-	157.6	-	-	-
7	6.10 (1H, s)	111.2	C5, C6, C8, C9	-	H5
8	-	178.2	-	-	-
9	-	102.5	-	-	-
10	-	161.6	-	-	-
11	4.76 (1H, dd, 4.3, 4.8)	57.6	C8, C9, C10, C12, C13, C17	H12, OH11	OH11, H12, H18b, H19b, H25a
12	2.00 (1H, d, 4.3)	46.5	C9, C13, C14, C16, C17, C18, C24, C25	H11, OH11	H11, H23
13	-	85.2	-	-	-
14a	2.06 (1H, m)*	35.0	C12, C13, C15, C16, C24	H14b	H14b, H24
14b	1.70 (1H, m)	-	-	H14a	H14a
15a	2.03 (1H, m)*	26.8	C13, C14, C15	H15b	H15b, H25b
15b	1.63 (1H, m)*	-	-	H15a	H15a, OH16
16	-	73.6	-	-	-
17	-	41.7	-	-	-
18a	2.00 (1H, m)*	27.3	C12, C16, C17, C20, C25	H18b	H18b, H25a
18b	1.63 (1H, m)*	-	-	H18a, H25b†	H18a, H11, OH16
19a	1.99 (1H, m)*	28.9	C17, C18, C20, C21	H19b	H19b
19b	1.56 (1H, m)	-	-	H19a	H11, H19a, OH20
20	-	96.9	-	-	-
21	-	46.2	-	-	-
22	0.96 (3H, s)	21.1	C16, C20, C21, C23	H23	OH20, H23, H25b
23	0.88 (3H, s)	20.2	C16, C20, C21, C22	H22	H12, OH16, H22
24	1.52 (3H, s)	21.5	C12, C13, C14	-	OH11, H14a, H25a, H25b
25a	4.57 (1H, d, 9.7)	66.5	C12, C16, C17, C18, C20	H25a	H11, OH11, H18a, H24, H25b
25b	3.87 (1H, dd, 2.1, 9.7)	-	-	H18b†, H25a	H15a, H22, H24, H25a
11OH	5.00 (1H, d, 4.8)	-	C9, C11, C12, C17	H11, H12	H11, H24, H25a
16OH	4.21 (1H, s)	-	C15, C16, C21	-	H15b, H18b, H23
20OH	5.40 (1H, s)	-	C19, C20, C21	-	H19b, H22

* signals overlapped

† W coupling

Figure S 1. ^1H NMR spectrum (800 MHz) of hesseltin A (**1**) in $\text{DMSO-}d_6$

Figure S 2. ^{13}C NMR spectrum (600 MHz) of hesseltin A (**1**) in $\text{DMSO}-d_6$

Figure S 2. ^1H - ^{13}C HMBC NMR spectrum (800 MHz) of hesseltin A (1) in $\text{DMSO-}d_6$