Supporting Information Available
for the Article entitled

Heteroannulation of Nitroketene *N,S*-
arylaminoacetals with POCl₃: A Novel Highly
Regioselective Synthesis of Unsymmetrical 2,3-
Substituted Quinoxalines

C. Venkatesh,^a B. Singh,^a P. K. Mahata,^a H. Ila^{a,*} and H. Junjappa^b

^a,^aDepartment of Chemistry, Indian Institute of Technology, Kanpur-208016, India

^bBioorganics and Applied Materials Pvt. Ltd., # B-64/1, 3rd Stage, Peenya Industrial Area, Peenya, Bangalore – 560 058

Email: hila@iitk.ac.in

Experimental Section

General. ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) spectra (JEOL JNM LA-400) were recorded in CDCl₃, and TMS was used as an internal reference. Melting points were obtained on Mel-Temp melting point apparatus (capillary method) and were uncorrected. The IR spectra were recorded on Perkin Elmer 1320 spectrophotometer. The FAB mass spectra were recorded on a JEOL SX 102/DA-6000 Mass Spectrometer/Data System. Chromatographic purification was conducted by column chromatography using 100-200 mesh silica gel obtained from Acme Synthetic Chemicals. DMF, NMP and CH₂Cl₂ were distilled over CaH₂ and stored over 4Å molecular sieves. CH₃CN and Cl₂CHCHCl₂ were distilled over P₂O₅. THF was distilled over sodium/benzophenone prior to use. NaH (as 60% suspension in oil), *N*-benzyl piperazine and Fe(acac)₃ were purchased from the standard firms and used as such whereas NiCl₂(PPh₃)₂ and NiCl₂(dppp) catalysts were

prepared according to the literature procedure. *m*CPBA (50-55% suspension in water) was diluted with CH₂Cl₂ and dried over anhydrous Na₂SO₄. All the known **1a-b**, **1f-h**, **1j**, **4** and unknown **1c-e**, **1i**, **1k**, **9**, **11** nitroketene *N,S*-arylaminoacetals were prepared by slight modification of the reported procedure.²³

General Procedure for Preparation of Nitroketene *N,S*-Acetals **1a-k, **4** and Bis(nitroketene-*N,S*-acetals) **9** and **11****

In a typical experiment a solution of nitroketene *S,S*-acetal (5.0 g, 30.30 mmol) and corresponding aniline (30.30 mmol) in 150 mL ethanol was refluxed for 24 h with constant stirring, the reaction being monitored by TLC. The reaction mixture was then cooled by ice-water, filtered and washed with ether to give crystalline nitroketene *N,S*-acetals **1a-k**, **4** in high yields.

The bis(nitroketene-*N,S*-acetals) **9** and **11** were also prepared following the similar procedure from the corresponding diamine (1.1 g, 10 mmol) and nitroketene *S,S*-acetal (3.30 g, 20 mmol).

2-(3'-Methoxyanilino)-2-(methylthio)-1-nitroethylene (1a**)**

Yield 88% (6.41 g); Yellow crystals; m.p. 122-123 °C; *R_f* 0.49 (7:3 hexanes-EtOAc); IR (KBr): 3069, 2999, 1599, 1546, 1330 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 2.30 (s, 3H, SMe), 3.74 (s, 3H, OMe), 6.61 (s, 1H, =CH), 6.75 (t, *J* = 2.20 Hz, 1H, ArH), 6.80 (td, *J* = 8.56, 2.20 Hz, 2H, ArH), 7.24 (t, *J* = 8.04 Hz, 1H, ArH), 11.74 (brs, 1H, NH); ¹³C NMR (100 MHz, CDCl₃): δ 14.7, 55.4, 107.7, 111.4, 113.7, 117.8, 130.1, 137.1, 160.2, 163.4; Anal. Calcd. for C₁₀H₁₂N₂O₃S (240.28): C, 49.99; H, 5.03; N, 11.66 %. Found C, 49.81; H, 5.14; N, 11.45 %.

2-(3',4'-Dimethoxyanilino)-2-(methylthio)-1-nitroethylene (1b**)**

Yield 86% (7.04 g); Yellow solid; m.p. 138-139 °C; *R_f* 0.40 (7:3 hexanes-EtOAc); IR (KBr): 2991, 1558, 1426, cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 2.38 (s, 3H, SMe),

3.89 (s, 3H, OMe), 3.91 (s, 3H, OMe), 6.69 (s, 1H, =CH), 6.79 (d, $J = 1.44$ Hz, 1H, ArH), 6.85 (d, $J = 8.52$ Hz, 1H, ArH), 6.88 (d, $J = 8.76$ Hz, 1H, ArH), 11.71 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.6, 56.0, 56.1, 107.3, 109.6, 110.9, 118.6, 128.9, 149.0, 149.3, 164.4; Anal. Calcd. for $\text{C}_{11}\text{H}_{14}\text{N}_2\text{O}_4\text{S}$ (270.31): C, 48.88; H, 5.22; N, 10.36 %. Found C, 48.72 ; H, 5.44; N, 10.15 %.

2-(3',5'-Dimethoxyanilino)-2-(methylthio)-1-nitroethylene (1c)

Yield 87% (7.12 g); Yellow solid; m.p. 142-143 °C; R_f 0.40 (8.5:1.5 hexanes-EtOAc); IR (KBr): 2998, 1610, 1544, 1426, 1325 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.39 (s, 3H, SMe), 3.80 (s, 6H, OMe), 6.43 (d, $J = 0.96$ Hz, 1H, ArH), 6.44 (brs, 2H, ArH), 6.69 (s, 1H, =CH), 11.82 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.8, 55.6, 100.0, 103.9, 107.8, 137.6, 161.1, 163.3; Anal. Calcd. for $\text{C}_{11}\text{H}_{14}\text{N}_2\text{O}_4\text{S}$ (270.31): C, 48.88; H, 5.22; N, 10.36 %. Found C, 48.71; H, 5.43; N, 10.54 %.

2-(2',5'-Dimethoxyanilino)-2-(methylthio)-1-nitroethylene (1d)

Yield 84% (6.87 g); Orange solid; m.p. 121-123 °C; R_f 0.60 (8.5:1.5 hexanes-EtOAc); IR (KBr): 2996, 1543, 1415, 1322 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.40 (s, 3H, SMe), 3.79 (s, 3H, OMe), 3.84 (s, 3H, OMe), 6.72 (s, 1H, =CH), 6.82 (dd, $J = 9.02$, 2.92 Hz, 1H, ArH), 6.90 (d, $J = 9.04$ Hz, 1H, ArH), 7.01 (d, $J = 2.92$ Hz, 1H, ArH), 11.86 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.8, 55.8, 56.2, 108.2, 112.2, 113.0, 125.8, 147.1, 153.0, 157.8, 162.8; Anal. Calcd. for $\text{C}_{11}\text{H}_{14}\text{N}_2\text{O}_4\text{S}$ (270.31): C, 48.88; H, 5.22; N, 10.36 %. Found C, 48.71; H, 5.38; N, 10.22 %.

2-(4'-Methoxy-2'-methylanilino)-2-(methylthio)-1-nitroethylene (1e)

Yield 89% (6.85 g); Brown solid; m.p. 172-173 °C; R_f 0.48 (8.5:1.5 hexanes-EtOAc); IR (KBr): 2955, 1557, 1418, 1341 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.26 (s, 3H, Me), 2.35 (s, 3H, SMe), 3.82 (s, 3H, OMe), 6.69 (s, 1H, =CH), 6.77 (dd, $J = 8.64$, 2.62 Hz, 1H, ArH), 6.82 (d, $J = 2.44$ Hz, 1H, ArH), 7.14 (d, $J = 8.56$ Hz, 1H, ArH), 11.43 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.4, 18.0, 55.3, 107.0, 111.8, 116.1, 127.6, 128.8, 136.8, 159.7, 165.6; Anal. Calcd. for $\text{C}_{11}\text{H}_{14}\text{N}_2\text{O}_3\text{S}$ (254.31): C, 51.95; H, 5.55; N, 11.02 %. Found C, 51.81; H, 5.36 N, 11.21 %.

2-(Anilino)-2-(methylthio)-1-nitroethylene (1f)

Yield 80% (5.1 g); Yellow crystals; m.p. 147-148 °C; R_f 0.50 (4:1 hexanes-EtOAc); IR (KBr): 1542, 1422, 1336 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.39 (s, 3H, SMe), 6.71 (s, 1H, =CH), 7.30 (d, $J = 7.56$ Hz, 2H, ArH), 7.37 (t, $J = 7.32$ Hz, 1H, ArH), 7.44 (t, $J = 7.08$ Hz, 2H, ArH), 11.83 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.7, 107.8, 126.0, 128.7, 129.4, 136.1, 163.6; Anal. Calcd. for $\text{C}_9\text{H}_{10}\text{N}_2\text{O}_2\text{S}$ (210.25): C, 51.41; H, 4.79; N, 13.32 %. Found C, 51.22; H, 4.65; N, 13.54 %.

2-(4'-Methoxyanilino)-2-(methylthio)-1-nitroethylene (1g)

Yield 85% (6.18 g); Yellow crystals; m.p. 157-158 °C; R_f 0.50 (8.5:1.5 hexanes-EtOAc); IR (KBr): 3002, 1546, 1426, 1332 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.41 (s, 3H, SMe), 3.88 (s, 3H, OMe), 6.73 (s, 1H, =CH), 7.0 (d, $J = 9.50$ Hz, 2H, ArH), 7.25 (d, $J = 8.8$ Hz, 2H, ArH), 11.70 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.6, 55.5,

107.3, 114.5, 127.7, 128.7, 159.5, 164.5; Anal. Calcd. for $C_{10}H_{12}N_2O_3S$ (240.28): C, 49.99; H, 5.03; N, 11.66 %. Found C, 49.87; H, 5.25; N, 11.45 %.

2-(4'-Chloroanilino)-2-(methylthio)-1-nitroethylene (1h)

Yield 87% (6.45 g); White solid; m.p. 165-166 °C; R_f 0.45 (7:3 hexanes-EtOAc); IR (KBr): 3142, 1563, 1357, 1268 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$): δ 2.37 (s, 3H, SMe), 6.66 (s, 1H, =CH), 7.22 (d, J = 8.56 Hz, 2H, ArH), 7.38 (d, J = 8.56 Hz, 2H, ArH), 11.68 (brs, 1H, NH); ^{13}C NMR (100 MHz, $CDCl_3$): δ 14.7, 108.1, 127.3, 127.4, 129.7, 134.7, 163.2; Anal. Calcd. for $C_9H_9ClN_2O_2S$ (244.70): C, 44.18; H, 3.71; N, 11.45 %. Found C, 44.31; H, 3.54; N, 11.58 %.

2-(3',4'-Dichloroanilino)-2-(methylthio)-1-nitroethylene (1i)

Yield 72% (6.08 g); Yellow crystals; m.p. 157-158 °C; R_f 0.35 (6:1 hexanes-EtOAc); IR (KBr): 1565, 1475, 1262, 829 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$): δ 2.41 (s, 3H, SMe), 6.67 (s, 1H, =CH), 7.18 (d, J = 8.56 Hz, 1H, ArH), 7.45 (d, J = 1.80 Hz, 1H, ArH), 7.50 (dd, J = 8.56, 1.80 Hz, 1H, ArH), 11.66 (brs, 1H, NH); ^{13}C NMR (100 MHz, $CDCl_3$): δ 14.7, 108.5, 125.2, 127.8, 131.0, 132.1, 133.3, 135.5, 162.5; Anal. Calcd. for $C_9H_8N_2O_2SCl_2$ (279.14): C, 38.72; H, 2.89; N, 10.04 %. Found C, 38.61; H, 2.75; N, 10.21 %.

2-(3'-Fluoroanilino)-2-(methylthio)-1-nitroethylene (1j)

Yield 80% (5.53 g); Yellow solid; m.p. 126-127 °C; R_f 0.45 (8.5:1.5 hexanes-EtOAc); IR (KBr): 1548, 1416, 1341, cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$): δ 2.38 (s, 3H, SMe), 6.66 (s, 1H, =CH), 7.03-7.08 (m, 3H, ArH), 7.36 (dd, J = 8.8, 7.6 Hz, 1H, ArH),

11.75 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.7, 108.3, 113.3, (d, $J_{\text{F-C}} = 23.9$ Hz), 115.0 (d, $J_{\text{C-F}} = 20.5$ Hz), 121.5, 130.7 (d, $J_{\text{C-F}} = 9.0$ Hz), 137.6 (d, $J_{\text{C-F}} = 9.9$ Hz), 162.9, 163.9; Anal. Calcd. for $\text{C}_9\text{H}_9\text{N}_2\text{O}_2\text{SF}$ (228.24): C, 47.36; H, 3.97; N, 12.27 %. Found C, 47.20; H, 3.84; N, 12.41 %.

2-(Methylthio)-2-(1'-naphthylamino)-1-nitroethylene (1k)

Yield 92% (7.25 g); Yellow solid; m.p. 187-188 °C; R_f 0.40 (9:1 hexanes-EtOAc); IR (KBr): 1557, 1418, 1341 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.33 (s, 3H, SMe), 6.80 (s, 1H, =CH), 7.50-7.53 (m, 2H, ArH), 7.56-7.62 (m, 2H, ArH), 7.90-7.96 (m, 3H, ArH), 12.03 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.6, 107.9, 122.0, 125.0, 125.1, 127.0, 127.6, 128.5, 129.3, 129.4, 132.3, 134.2, 165.2; Anal. Calcd. for $\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$ (260.31): C, 59.98; H, 4.65; N, 10.76 %. Found C, 59.80; H, 4.49; N, 10.54 %.

2-(3'-Aminopyridyl)-2-(methylthio)-1-nitroethylene (4)

Yield 62% (3.96 g); Yellow solid; m.p. 125-126 °C; R_f 0.45 (1:1 hexanes-EtOAc); IR (KBr): 3143, 1557, 1463, 1418, 1349, 1175 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.36 (s, 3H, SMe), 6.66 (s, 1H, =CH), 7.33 (dd, $J = 8.16, 4.64$ Hz, 1H, ArH), 7.63 (dd, $J = 8.18, 1.48$ Hz, 1H, ArH), 8.53-8.55 (m, 2H, ArH), 11.62 (brs, 1H, NH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.5, 108.5, 123.6, 133.0, 133.5, 147.2, 148.9, 163.1; Anal. Calcd. for $\text{C}_8\text{H}_9\text{N}_3\text{O}_2\text{S}$ (211.24): C, 45.49; H, 4.29; N, 19.89 %. Found C, 45.34; H, 4.44; N, 19.62 %.

1,3-Bis[2'-(methylthio)-1'-nitroethenyl]diaminobenzene (9)

Yield 95% (3.25 g); Yellow solid; m.p. 130-132 °C; R_f 0.50 (4:1 hexanes-EtOAc); IR (KBr): 3146, 1552, 1344, 1199 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.39 (s, 6H, $2 \times \text{SMe}$), 6.67 (s, 2H, $2 \times =\text{CH}$), 7.26 (dd, $J = 8.06, 1.96$ Hz, 2H, ArH), 7.30 (s, 1H, ArH), 7.46 (t, $J = 7.80$ Hz, 1H, ArH), 11.76 (brs, 2H, NH); ^{13}C NMR (100 MHz, CDCl_3): 14.76, 108.4, 123.2, 125.2, 130.4, 137.3, 162.6; Anal. Calcd. for $\text{C}_{12}\text{H}_{14}\text{N}_4\text{O}_4\text{S}_2$ (342.40): C, 42.09; H, 4.12; N, 16.36 %. Found C, 42.32; H, 4.34; N, 16.12 %.

1,4-Bis[2'-(methylthio)-1'-nitroethenyl]diaminobenzene (11)

Yield 91% (3.1 g); Yellow solid; m.p. 224-225 °C; R_f 0.20 (7:3 hexanes-EtOAc); IR (KBr): 1565, 1475, 1353, 1275 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 3.30 (s, 6H, $2 \times \text{SMe}$), 6.77 (s, 2H, $2 \times =\text{CH}$), 7.41 (s, 4H, ArH), 11.53 (brs, 2H, NH); Anal. Calcd. for $\text{C}_{12}\text{H}_{14}\text{N}_4\text{O}_4\text{S}_2$ (342.40): C, 42.09; H, 4.12; N, 16.36 %. Found C, 42.21; H, 4.01; N, 16.49 %.

Note: ^{13}C NMR of compound **11** couldn't be recorded because of its highly insoluble nature in most of the deuterated solvents including $\text{DMSO}-d_6$.

General Procedure for Cyclization of Nitroketene *N,S*-acetals **1a-k, **4** to Quinoxalines **3a-k**, **5****

To a suspension of *N,S*-acetals **1a-k**, **4** (8.36 mmol) in 20 mL CH_3CN , POCl_3 (2.34 mL, 25.10 mmol) was added dropwise at 0 °C over a period of 15-minutes with constant stirring. After the completion of addition, the reaction mixture was warmed to 80 °C for 3-4 h, monitored by TLC. It was then cooled and neutralized with ice cold saturated NaHCO_3 solution (100 mL), extracted with CHCl_3 (3×50 mL), washed with H_2O (2×50 mL), brine (1×50 mL) and dried over anhydrous Na_2SO_4 . The solvent was

evaporated under vacuum to give quinoxalines **3a-k**, **5**, which were purified by column chromatography over silica gel using hexane: EtOAc (10:1) as eluent.

3-Chloro-7-methoxy-2-(methylthio)quinoxaline (**3a**)

Yield 70 % (1.41 g); Pale Yellow crystals; m.p. 119-120 °C; R_f 0.85 (10:1 hexanes-EtOAc); IR (KBr): 2925, 1615, 1519, 1494 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.65 (s, 3H, SMe), 3.95 (s, 3H, OCH_3), 7.23-7.27 (m, 2H, ArH), 7.78 (d, J = 8.80 Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.7, 55.7, 105.8, 121.1, 129.0, 134.5, 142.5, 142.8, 155.8, 161.0; MS (m/z , %): 240 (M^+ , 30), 238 ($\text{M}-2$, 100), (204, 88); Anal. Calcd. for $\text{C}_{10}\text{H}_9\text{N}_2\text{OSCl}$ (240.71): C, 49.90; H, 3.77; N, 11.64 %. Found C, 49.78; H, 3.52; N, 11.46%.

3-Chloro-6,7-dimethoxy-2-(methylthio)quinoxaline (**3b**)

Yield 75 % (1.69 g); Pale yellow crystals; m.p. 184-185 °C; R_f 0.60 (9:1 hexanes-EtOAc); IR (KBr): 2934, 1615, 1507 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.66 (s, 3H, SMe), 4.01 (s, 3H, OMe), 4.06 (s, 3H, OMe), 7.32 (s, 1H, ArH), 7.28 (s, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.7, 56.2, 56.3, 105.7, 106.1, 135.5, 138.4, 142.8, 151.6, 152.6, 152.7; MS: (m/z , %): 271 ($\text{M}+1$, 100), 273 ($\text{M}+1$, 43); Anal. Calcd. for $\text{C}_{11}\text{H}_{11}\text{N}_2\text{O}_2\text{SCl}$ (270.74): C, 48.80; H, 4.10; N, 10.35 %. Found C, 48.59; H, 4.32; N, 10.57 %.

3-Chloro-5,7-dimethoxy-2-(methylthio)quinoxaline (3c)

Yield 71 % (1.61 g); Orange solid; m.p. 177-178 °C; R_f 0.82 (9.5:0.5 hexanes-EtOAc); IR (KBr): 2932, 1707, 1604, 1458 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.63 (s, 3H, SMe), 3.92 (s, 3H, OMe), 3.99 (s, 3H, OMe), 6.61 (d, $J = 2.44$ Hz, 1H, ArH), 6.85 (d, $J = 2.20$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.7, 55.8, 56.3, 98.0, 100.6, 127.1, 141.4, 143.6, 155.2, 155.6, 161.6; MS: (m/z, %): 271 ($M+1$, 75), 273 ($M+1$, 30); Anal. Calcd. for $\text{C}_{11}\text{H}_{11}\text{N}_2\text{O}_2\text{SCl}$ (270.74): C, 48.80; H, 4.10; N, 10.35 %. Found C, 48.99; H, 4.22; N, 10.11 %.

3,5-Dichloro-8-methoxy-2-(methylthio)quinoxaline (3d)

Yield 61 % (1.40 g); White solid; m.p. 181-182 °C; R_f 0.75 (9.5:0.5 hexanes-EtOAc); IR (KBr): 2926, 1606, 1466, 1223 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.67 (s, 3H, SMe), 3.96 (s, 3H, OMe), 7.21 (d, $J = 2.68$ Hz, 1H, ArH), 7.39 (d, $J = 2.68$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.9, 56.0, 105.3, 120.9, 129.4, 131.7, 132.6, 143.2, 157.4, 160.4; MS: (m/z, %): 275 (M^+ , 65), 277 (M^+ , 43); Anal. Calcd. for $\text{C}_{10}\text{H}_8\text{N}_2\text{OSCl}_2$ (275.15): C, 43.65; H, 2.93; N, 10.18 %. Found C, 43.45; H, 2.60; N, 10.42 %.

3-Chloro-6-methoxy-8-methyl-2-(methylthio)quinoxaline (3e)

Yield 62 % (1.32 g); Pale yellow crystals; m.p. 137-138 °C; R_f 0.70 (9.5:0.5 hexanes-EtOAc); IR (KBr): 2918, 1615, 1473, 1392 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ

2.63 (s, 3H, Me), 2.66 (s, 3H, SMe), 3.87 (s, 3H, OMe), 7.06 (d, $J = 2.68$ Hz, 1H, ArH), 7.16 (d, $J = 2.68$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.7, 17.1, 55.6, 104.0, 122.5, 136.7, 137.3, 140.6, 145.1, 151.1, 159.5; MS: (m/z , %): 255 ($M+1$, 100), 257 ($M+1$, 43); Anal. Calcd. for $\text{C}_{11}\text{H}_{11}\text{ClN}_2\text{OS}$ (254.74): C, 51.86; H, 4.35; N, 11.00 %. Found C, 51.55; H, 4.60; N, 11.22 %.

3f

3-Chloro-2-(methylthio)quinoxaline (3f)

Yield 50 % (0.88 g); White needles; m.p. 102-103 °C; R_f 0.54 (10:1 hexanes-EtOAc); IR (KBr): 2959, 1525, 1261, 1175, 761 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.68 (s, 3H, SMe), 7.63 (dt, $J = 7.08, 1.00$ Hz, 1H, ArH), 7.70 (dt, $J = 7.08, 0.96$ Hz, 1H, ArH), 7.93 (d, $J = 9.28$ Hz, 1H, ArH), 7.96 (d, $J = 9.04$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.9, 127.4, 128.1, 128.7, 130.1, 139.0, 141.3, 145.2, 155.9; MS: (m/z , %): 210 (M^+ , 100), 212 (M^+ , 40). Anal. Calcd. for $\text{C}_9\text{H}_7\text{N}_2\text{SCl}$ (210.68): C, 51.31; H, 3.35; N, 13.30 %. Found C, 51.18; H, 3.54; N, 13.18 %.

3g

3-Chloro-6-methoxy-2-(methylthio)quinoxaline (3g)

Yield 57 % (1.14 g); Pale yellow crystals; m.p. 137-138 °C; R_f 0.5 (20:1 hexanes-EtOAc); IR (KBr): 2963, 1611, 1523, 1483 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.64 (s, 3H, SMe), 3.92 (s, 3H, OCH_3), 7.24 (d, $J = 2.68$ Hz, 1H, ArH), 7.33 (dd, $J = 9.14, 2.68$ Hz, 1H, ArH), 7.84 (d, $J = 9.04$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.7, 55.8, 106.3, 122.7, 128.4, 137.4, 140.4, 145.5, 152.6, 159.9; MS: (m/z , %): 241 ($M+1$, 100), 243 ($M+1$, 37). Anal. Calcd. for $\text{C}_{10}\text{H}_9\text{N}_2\text{OSCl}$ (240.71): C, 49.90; H, 3.77; N, 11.64 %. Found C, 50.14; H, 3.49; N, 11.45%.

3,6-Dichloro-2-(methylthio)quinoxaline (3h)

Yield 50 % (1.02 g); White solid; m.p. 126-127 °C; R_f 0.77 (9:1 hexanes-EtOAc); IR (KBr): 2926, 1524, 1272, 1240, 1151, 1118, 1012, 821 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.66 (s, 3H, SMe), 7.63 (dd, J = 8.90, 2.44 Hz, 1H, ArH), 7.88 (d, J = 9.54, Hz, 1H, ArH), 7.90 (d, J = 2.44 Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.8, 127.0, 128.4, 130.9, 134.1, 139.0, 139.7, 146.1, 156.3; MS: (m/z , %): 244 (M^+ , 100), 246 (M^+ , 71); Anal. Calcd. for $\text{C}_9\text{H}_6\text{N}_2\text{SCl}_2$ (245.13): C, 44.10; H, 2.47; N, 11.43 %. Found C, 44.23; H, 2.25; N, 11.66 %.

2-(Methylthio)-3,6,7-trichloroquinoxaline (3i)

Yield 42 % (0.98 g); Colorless crystals; m.p. 164-165 °C; R_f 0.95 (20:1 hexanes-EtOAc); IR (KBr): 3080, 1526, 1261, 1138, 1024 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.63 (s, 3H, SMe), 7.98 (s, 1H, ArH), 8.02 (s, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.0, 128.0, 128.7, 132.9, 134.7, 137.5, 139.8, 146.4, 157.8; MS: (m/z , %): 279 (M^+ , 67), 281 (M^+ , 66); Anal. Calcd. for $\text{C}_9\text{H}_5\text{N}_2\text{SCl}_3$ (279.57): C, 38.66; H, 1.80; N, 10.02 %. Found C, 38.47; H, 1.68; N, 10.29 %.

3-Chloro-7-fluoro-2-(methylthio)quinoxaline (3ja)

Yield 40 % (0.76 g); White solid; m.p. 110-111 °C; R_f 0.60 (9:1 hexanes-EtOAc); IR (KBr): 2924, 1611, 1567, 1486, 1363, 1266, 1180, 1010, 825 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.64 (s, 3H, SMe), 7.37 (dt, J = 8.28, 2.44, Hz, 1H, ArH), 7.56 (dd, J = 9.02, 2.68 Hz, 1H, ArH), 7.90 (dd, J = 9.28, 5.72 Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.9, 111.6 (d, $J_{\text{C-F}}$ = 22.2 Hz), 118.5 (d, $J_{\text{C-F}}$ = 25.5 Hz), 130.1 (d, $J_{\text{C-F}}$ = 10.7 Hz), 136.0, 142.0 (d, $J_{\text{C-F}}$ = 10.9 Hz), 157.4, 161.7, 164.2; MS: (m/z , %): 228 (M^+ ,

100), 230 (M^+ , 38). Anal. Calcd. for $C_9H_6N_2SClF$ (228.67): C, 47.27; H, 2.64; N, 12.25 %. Found C, 47.12; H, 2.32; N, 12.47 %.

3-Chloro-5-fluoro-2-(methylthio)quinoxaline (3jb)

Yield 20 % (0.38 g); White solid; m.p. 105-106 °C; R_f 0.60 (9:1 hexanes-EtOAc); IR (KBr): 2924, 1474, 1258, 1179, 1139, 1077, 812 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$): δ 2.61 (s, 3H, SMe), 7.26 (dt, J = 8.92, 1.20, Hz, 1H, ArH), 7.53-7.59 (m, 1H, ArH), 7.69 (dt, J = 8.47, 1.20 Hz, 1H, ArH); ^{13}C NMR (100 MHz, $CDCl_3$): δ 13.9, 113 (d, J_{C-F} = 18.1 Hz), 123.1 (d, J_{C-F} = 4.9 Hz), 129.2, 129.8 (d, J_{C-F} = 9.1 Hz), 142.2, 145.7, 155.2, 157.7 (d, J_{C-F} = 19.7 Hz); MS: (m/z , %): 228 (M^+ , 100), 230 (M^+ , 38). Anal. Calcd. for $C_9H_6ClFN_2S$ (228.67): C, 47.27; H, 2.64; N, 12.25 %. Found C, 47.39; H, 2.52; N, 12.47 %.

3-Chloro-2-(methylthio)benzo[h]quinoxaline (3k)

Yield 80 % (1.74 g); Yellow crystals; m.p. 141-142°C; R_f 0.80 (9.5:0.5 hexanes-EtOAc); IR (KBr): 1515, 1423, 1275, 1164, 936, 822 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$): δ 2.80 (s, 3H, SMe), 7.72 (dd, J = 6.1, 3.64 Hz, 2H, ArH), 7.80 (d, J = 9.04, Hz, 1H, ArH), 7.90-7.92 (m, 2H, ArH), 9.05-9.08 (m, 1H, ArH); ^{13}C NMR (100 MHz, $CDCl_3$): δ 14.0, 124.3, 125.4, 127.4, 128.0, 128.8, 129.7, 130.3, 133.4, 137.8, 139.2, 144.8, 154.4; MS: (m/z , %): 261 ($M+1$, 100), 263 ($M+1$, 37); Anal. Calcd. for $C_{13}H_9ClN_2S$ (260.74): C, 59.88; H, 3.48; N, 10.74 %. Found C, 59.65; H, 3.60; N, 10.48 %.

3-Chloro-2-(methylthio)pyrido[2,3-*b*]pyrazine

Yield 45 % (0.79 g); Yellow solid; m.p. 132-133 °C; R_f 0.5 (1:1 hexanes-EtOAc); IR (KBr): 2925, 1692, 1539, 1372, 1078, 788 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.68 (s, 3H, SMe), 7.68 (dd, $J = 8.32$, 4.14 Hz, 1H, ArH), 8.35 (d, $J = 7.32$ Hz, 1H, ArH), 8.98 (d, $J = 2.92$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.2, 125.4, 136.6, 136.7, 147.6, 148.5, 151.5, 158.4; MS: (m/z, %): 212 ($M+1$, 54), 214 ($M+1$, 22); Anal. Calcd. for $\text{C}_8\text{H}_6\text{N}_3\text{SCl}$ (211.67): C, 45.39; H, 2.86; N, 19.85 %. Found C, 45.65; H, 2.60; N, 19.52 %.

General Procedure for Cyclization of Bis(nitroketene-*N,S*-acetals) **9** and **11**: Synthesis of Pyrazinoquinoxalines **10** and **12**

The pyrazinoquinoxalines **10** and **12** were prepared following similar procedure for quinoxalines **3** by addition of 1.36 mL (14.6 mmol) of POCl_3 to bis(nitroketene-*N,S*-acetal) (1.0 g, 2.92 mmol) in CH_3CN (20 mL) at 0 °C. It was then warmed to 80 °C for 3-4 h and after standard workup and purification procedure **10** and **12** were obtained as crystalline compounds.

2,8-Bis(methylthio)-3,9-dichloropyrazino[2,3-*h*]quinoxaline (**10**)

Yield 55 % (0.55 g); Yellow solid; m.p. 244-245 °C; R_f 0.8 (10:1 hexanes-EtOAc); IR (KBr): 2921, 1513, 1276, 1177, 987 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.74 (s, 3H, SMe), 2.89 (s, 3H, SMe), 8.08 (d, $J = 9.28$ Hz, 1H, ArH), 8.12 (d, $J = 9.04$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.0, 14.1, 128.5, 130.2, 137.9, 138.7, 142.3, 146.5, 157.7, 158.6, 161.7, 165.4; MS: (m/z, %): 342 (M^+ , 58), 344 (M^+ , 41); Anal. Calcd. for $\text{C}_{12}\text{H}_8\text{N}_4\text{S}_2\text{Cl}_2$ (343.26): C, 41.99; H, 2.35; N, 16.32 %. Found C, 41.75; H, 2.60; N, 16.56 %.

2,7-Bis(methylthio)-3,8-dichloropyrazino[2,3-g]quinoxaline (12)

Yield 45 % (0.45 g); Yellow solid; m.p. 260-261°C; R_f 0.75 (9:1 hexanes-EtOAc); IR (KBr): 1512, 1273, 1176, 987, 833 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.71 (s, 6H, 2 $\times$ SMe), 8.16 (s, 2H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 14.0, 130.0, 135.0, 141.4, 146.6, 158.1; MS: (m/z , %): 343 (M^+ , 25), 347 (M^+ , 12); Anal. Calcd. for $\text{C}_{12}\text{H}_8\text{Cl}_2\text{N}_4\text{S}_2$ (343.26): C, 41.99; H, 2.35; N, 16.32 %. Found C, 41.81; H, 2.55; N, 16.13 %.

Procedure for the Preparation of 3-Chloro-7-methoxy (or 6-methoxy)-(2-methylsulfonyl)quinoxalines 13a and 13g

A solution of *m*CPBA (1.58 g, 9.15 mmol) in CH_2Cl_2 (25 mL) was added dropwise to a stirred solution of quinoxalines **3a** or **3g** (1.0 g, 4.15 mmol) in CH_2Cl_2 (25 mL) at 0 °C over a period of 30-minutes. The reaction mixture was further stirred at room temperature for 5-6 h (monitored by TLC). It was then poured into ice-cold water, washed with 10% NaHCO_3 solution (2 $\times$ 50 mL), water (50 mL) followed by brine (50 mL) and then dried over anhydrous Na_2SO_4 . The solvent was evaporated under vacuum to give crude products which were purified over silica gel using hexane:EtOAc as eluent (10:2) to give **13a** and **13g** as pale yellow crystals.

3-Chloro-7-methoxy-2-(methylsulfonyl)quinoxaline (13a)

Yield 95% (1.07 g); Pale yellow crystals; m.p. 173-174 °C; R_f 0.35 (10:1 hexanes-EtOAc); IR (KBr): 3022, 2935, 1311, 1223, 1149, 1127 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 3.53 (s, 3H, SO_2Me), 4.01 (s, 3H, OMe), 7.38 (d, $J = 2.68$ Hz, 1H, ArH), 7.58 (dd, $J = 9.28, 2.20$ Hz, 1H, ArH), 7.95 (d, $J = 9.28$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 40.1, 56.1, 106.5, 127.3, 129.0, 138.3, 138.8, 140.1, 149.7, 162.0; MS (m/z , %): 273 ($\text{M}+1$, 100), 275 ($\text{M}+1$, 43); Anal. Calcd. for $\text{C}_{10}\text{H}_9\text{N}_2\text{O}_3\text{SCl}$ (272.71): C, 44.04; H, 3.33; N, 10.27 %. Found C, 44.14; H, 3.56; N, 10.12 %.

3-Chloro-6-methoxy-2-(methylsulfonyl)quinoxaline (13g)

Yield 96% (1.08 g); Pale yellow crystals; m.p. 180-181 °C; R_f 0.23 (10:1 hexanes-EtOAc); IR (KBr): 1613, 1487, 1393, 1143, 1002, 759 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 3.49 (s, 3H, SO_2Me), 3.98 (s, 3H, OMe), 7.33 (d, $J = 2.68$ Hz, 1H, ArH), 7.49 (dd, $J = 9.28, 2.68$ Hz, 1H, ArH), 7.98 (d, $J = 9.28$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 40.2, 56.2, 105.7, 125.7, 130.5, 134.3, 141.9, 145.0, 147.1, 163.9; MS (m/z , %): 273 ($M+1$, 100), 275 ($M+1$, 36); Anal. Calcd. for $\text{C}_{10}\text{H}_9\text{N}_2\text{O}_3\text{SCl}$ (272.71): C, 44.04; H, 3.33; N, 10.27 %. Found C, 44.21; H, 3.12; N, 10.11 %.

General Procedure for the Displacement of 2-Methylsulfonyl/3-Chloro groups by Nitrogen Nucleophiles (primary and secondary aliphatic amines) in 13a/13g: Synthesis of 14, 16 and 18

A mixture of quinoxaline **13a** or **13g** (0.56 g, 2.05 mmol) and *n*-butyl amine or benzyl amine or *N*-benzylpiperazine (4.51 mmol) in 5 mL DMF was stirred at room temperature for 24 h (monitored by TLC). The reaction mixture was then diluted with CHCl_3 (1 $\times$ 50 mL), washed with water (50 mL), brine (50 mL) and the organic layer was dried over anhydrous Na_2SO_4 . The solvent was evaporated under vacuum to give crude products **14**, **16** and **18** which were purified by column chromatography using hexane:EtOAc (20:1) as eluent.

2-(*n*-Butylamino)-3-chloro-7-methoxyquinoxaline (14)

Yield 87% (0.47 g); Colorless viscous oil; R_f 0.81 (10:1 hexanes-EtOAc); IR (KBr): 3441, 3054, 2961, 1559, 1265, 739 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 0.86 (t, $J = 7.32$ Hz, 3H, CH_3), 1.29-1.38 (m, 2H, CH_2), 1.52-1.59 (m, 2H, CH_2), 3.42 (q, $J = 7.08$ Hz, 2H, CH_2), 3.76 (s, 3H, OMe), 6.86 (dd, $J = 9.02, 2.72$ Hz, 1H, ArH), 6.91 (d, $J = 2.68$ Hz, 1H, ArH), 7.51 (d, $J = 9.04$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.6, 20.0, 31.0, 41.0, 55.3, 104.8, 116.3, 128.4, 131.3, 134.6, 142.8, 148.1, 160.8; MS (m/z ,

%) : 266 (M+1, 100), 268 (M+1, 54); Anal. Calcd. for C₁₃H₁₆N₃OCl (265.74): C, 58.76; H, 6.07; N, 15.81 %. Found C, 58.53; H, 6.21; N, 15.71 %.

2-(Benzylamino)-3-chloro-7-methoxyquinoxaline (16)

Yield 94% (0.57 g); Pale yellow solid; R_f 0.75 (6:1 hexanes-EtOAc); IR (KBr): 3408, 3030, 2915, 1520, 1206 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 3.88 (s, 3H, OMe), 4.76 (d, *J* = 5.60 Hz, 2H, CH₂), 5.78 (t, *J* = 5.12 Hz, 1H, NH), 7.02 (dd, *J* = 8.92, 2.68 Hz, 1H, ArH), 7.06 (d, *J* = 2.68 Hz, 1H, ArH), 7.30 (t, *J* = 6.84 Hz, 1H, ArH), 7.35 (dd, *J* = 6.10, 1.72 Hz, 1H, ArH), 7.39 (t, *J* = 7.08 Hz, 3H, ArH), 7.66 (d, *J* = 9.00 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 45.4, 55.5, 105.0, 117.0, 127.5, 127.8, 128.7, 128.7, 131.8, 134.6, 138.0, 142.7, 148.0, 161.0; MS (*m/z*, %): 300 (M+1, 100), 302 (M+1, 34); Anal. Calcd. for C₁₆H₁₄N₃OCl (299.75): C, 64.11; H, 4.71; N, 14.02 %. Found C, 64.29; H, 4.63; N, 14.21 %.

3-(N-Benzylpiperazino)-6-methoxy-2-(methylsulfonyl)quinoxaline (18)

Yield 86% (0.72 g); Yellow crystals; R_f 0.24 (4:1 hexanes-EtOAc); IR (KBr): 2936, 2827, 1614, 1532, 1460, 1311, 1143, 1009, 830 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 2.68 (t, *J* = 4.16 Hz, 4H, CH₂NCH₂), 3.37 (s, 3H, SO₂Me), 3.59 (s, 2H, NCH₂), 3.73 (t, *J* = 4.40 Hz, 4H, CH₂NCH₂), 3.92 (s, 3H, OMe), 7.05 (d, *J* = 2.72 Hz, 1H, ArH), 7.15 (dd, *J* = 9.28, 2.44 Hz, 1H, ArH), 7.24 (t, *J* = 6.80 Hz, 1H, ArH), 7.31 (t, *J* = 7.56 Hz, 2H, ArH), 7.35 (d, *J* = 7.46 Hz, 2H, ArH), 7.74 (d, *J* = 9.28 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 41.6, 49.9, 52.7, 55.8, 62.8, 104.6, 120.5, 127.2, 128.2, 129.2, 129.9, 131.2, 137.4, 142.7, 144.1, 150.6, 163.1; MS (*m/z*, %): 413 (M+1, 100); Anal. Calcd. for C₂₁H₂₄N₄O₃S (412.51): C, 61.14; H, 5.86; N, 13.58 %. Found C, 61.02; H, 5.64; N, 13.72 %.

Procedure for Synthesis of 3-(Benzylamino)-2-(*n*-Butylamino)- / 2-(Benzylamino)-3-(*n*-butylamino)-7-methoxyquinoxalines (15** and **17**)**

A mixture of quinoxaline **14** or **16** (0.50 mmol) and benzylamine or *n*-butylamine (1.10 mmol) was heated at 100 °C for 22-24 h in a pressure tube with constant stirring (monitored by TLC). The reaction mixture was cooled to room temperature, diluted with CHCl₃ (1 × 15 mL), washed with water (2 × 10 mL), brine (1 × 10 mL) and the organic layer was dried over anhydrous Na₂SO₄. The solvent was evaporated under vacuum to give the products **15** and **17**, which were isolated by column chromatography over silica gel using hexane: EtOAc (20:1) as pale yellow viscous liquid.

3-(Benzylamino)-2-(*n*-butylamino)-7-methoxyquinoxaline (15**)**

Yield 75% (0.13 g); Pale yellow viscous liquid; *R_f* 0.46 (6:1 hexanes-EtOAc); ¹H NMR (400 MHz, CDCl₃): δ 0.84 (t, *J* = 7.32 Hz, 3H, CH₃), 1.29-1.39 (m, 2H, CH₂), 1.51-1.59 (m, 2H, CH₂), 3.44 (t, *J* = 6.84 Hz, 2H, CH₂), 3.78 (s, 3H, OMe), 4.53 (brs, 2H, NH), 4.58 (s, 2H, CH₂), 6.88 (dd, *J* = 8.92, 2.68 Hz, 1H, ArH), 7.02 (d, *J* = 2.92 Hz, 1H, ArH), 7.22 (t, *J* = 5.36 Hz, 1H, ArH), 7.26 (d, *J* = 6.84 Hz, 2H, ArH), 7.32 (d, *J* = 7.80 Hz, 2H, ArH), 7.49 (d, *J* = 9.00 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 13.8, 20.3, 31.3, 41.4, 46.1, 55.5, 106.0, 115.0, 126.5, 127.5, 128.5, 128.6, 131.3, 138.1, 138.8, 142.7, 144.8, 157.3; MS (*m/z*, %): 337 (M+1, 100); Anal. Calcd. for C₂₀H₂₄N₄O (336.43): C, 71.40; H, 7.19; N, 16.65 %. Found C, 71.59; H, 7.38; N, 16.43 %.

2-(Benzylamino)-3-(*n*-butylamino)-7-methoxyquinoxaline (17**)**

Yield 96% (0.16 g); Pale yellow viscous liquid; *R_f* 0.30 (10:1 hexanes-EtOAc); IR (KBr): 3415, 3052, 2935, 1465, 1260, 739 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 0.83 (t, *J* = 7.32 Hz, 3H, CH₃), 1.27-1.34 (m, 2H, CH₂), 1.49-1.56 (m, 2H, CH₂), 3.39 (t, *J* = 6.84

Hz, 2H, CH₂), 3.77 (s, 3H, OMe), 4.48 (brs, 1H, NH), 4.62 (s, 2H, CH₂), 4.96 (brs, 1H, NH), 6.88 (dd, *J* = 9.04, 2.68 Hz, 1H, ArH), 7.02 (d, *J* = 2.44 Hz, 1H, ArH), 7.16-7.25 (m, 3H, ArH), 7.29 (d, *J* = 6.60 Hz, 2H, ArH), 7.49 (d, *J* = 8.80 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 13.8, 20.3, 31.3, 41.6, 45.8, 55.5, 106.1, 115.1, 126.1, 127.5, 128.3, 128.6, 131.5, 137.5, 138.6, 143.0, 144.4, 157.1; MS (m/z, %): 337 (M+1, 89); Anal. Calcd. for C₂₀H₂₄N₄O (336.43): C, 71.40; H, 7.19; N, 16.65 %. Found C, 71.59; H, 7.02; N, 16.43 %.

Procedure for the Preparation of 3-(*N*-Benzylpiperazino)-2-cyano-6-methoxyquinoxaline **19**

A suspension of aminosulfonyl quinoxaline **18** (0.41 g, 1.0 mmol) and NaCN (0.10 g, 2.0 mmol) in 5 mL DMF was stirred at room temperature for 6 h. Upon the completion of the reaction, the reaction mixture was diluted with CHCl₃ (25 mL), washed with water (2 × 25 mL), brine (50 mL) and dried over anhydrous Na₂SO₄. The solvent was evaporated under vacuum to give crude product **19** which was purified over silica gel using hexane:EtOAc (19:1) as eluent.

3-(*N*-Benzylpiperazino)-2-cyano-6-methoxyquinoxaline (**19**)

Yield 97% (0.35 g); Yellow crystals; m.p. 147-148 °C; R_f 0.65 (4:1 hexanes-EtOAc); IR (KBr): 2812, 2225, 1611, 1535, 1384, 1225, 1009, 738 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 2.70 (t, *J* = 4.88 Hz, 4H, CH₂NCH₂), 3.62 (s, 2H, -NCH₂), 3.79 (t, *J* = 4.88 Hz, 4H, CH₂NCH₂), 3.93 (s, 3H, OMe), 7.00 (d, *J* = 2.68 Hz, 1H, ArH), 7.14 (dd, *J* = 9.16, 2.68 Hz, 1H, ArH), 7.26-7.37 (m, 5H, ArH), 7.76 (d, *J* = 9.28 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 48.0, 52.5, 55.9, 62.8, 104.7, 116.8, 118.6, 120.9, 127.4, 128.3, 129.3, 130.3, 133.7, 144.0, 154.8, 163.7; MS (m/z, %): 360 (M+1, 100); Anal. Calcd. for C₂₁H₂₁N₅O (359.42): C, 70.17; H, 5.89; N, 19.48 %. Found C, 70.11; H, 5.64; N, 19.22 %.

Procedure for Synthesis of 3-Cyano-6-methoxy-2-(methylsulfonyl)quinoxaline (**20**)

To a solution of **13g** (0.50 g, 1.83 mmol) in 5 mL DMF, NaCN (0.10 g, 2.01 mmol) was added and stirred at room temperature for 24 h. After the consumption of starting material the reaction mixture was diluted with CHCl₃ (25 mL), washed with water (2 × 25 mL), brine (50 mL) and dried over anhydrous Na₂SO₄. The solvent was evaporated under vacuum and upon purification by column chromatography over silica gel **20** was obtained as pale yellow crystals.

3-Cyano-6-methoxy-2-(methylsulfonyl)quinoxaline (**20**)

Yield 50 % (0.24 g); Pale yellow crystals; m.p. 169-170 °C; *R_f* 0.10 (6:1 hexanes-EtOAc); IR (KBr): 3022, 2927, 2366, 1613, 1318, 1222, 771 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 3.43 (s, 3H, SO₂Me), 4.00 (s, 3H, OMe), 7.41 (d, *J* = 2.72 Hz, 1H, ArH), 7.62 (dd, *J* = 9.28, 2.68 Hz, 1H, ArH), 8.05 (d, *J* = 9.28 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 40.2, 56.6, 106.53, 113.6, 121.2, 128.7, 130.7, 139.0, 142.3, 153.0, 164.9; MS (*m/z*, %): 264 (M+1, 100); Anal. Calcd. for C₁₁H₉N₃O₃S (263.27): C, 50.18; H, 3.45; N, 15.96 %. Found C, 50.04; H, 3.31; N, 15.78 %.

Procedure for Synthesis of 2-(*N*-Benzylpiperazino)-3-cyano-6-methoxyquinoxaline (**21**)

A homogenous mixture of **20** (0.20 g, 0.75 mmol) and *N*-benzylpiperazine (0.32 mL, 1.87 mmol) was heated at 100 °C for 2-3 h in a pressure tube with constant stirring (monitored by TLC). The reaction mixture was cooled to room temperature, diluted with CHCl₃ (1 × 15 mL), washed with water (2 × 10 mL), brine (1 × 10 mL) and the organic layer was dried over anhydrous Na₂SO₄. The solvent was evaporated under vacuum and the product **21** was isolated by column chromatography over silica gel using hexane: EtOAc (5:1) as yellow crystals.

2-(*N*-Benzylpiperazino)-3-cyano-6-methoxyquinoxaline (**21**)

Yield 87% (0.31 g); Yellow crystals; m.p. 82-83 °C; R_f 0.45 (5:1 hexanes-EtOAc); IR (KBr): 2814, 2227, 1621, 1536, 1449, 1390, 1213, 829 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.63 (t, J = 4.88 Hz, 4H, CH_2NCH_2), 3.55 (s, 2H, - NCH_2), 3.64 (t, J = 4.88 Hz, 4H, CH_2NCH_2), 3.84 (s, 3H, OMe), 7.14 (d, J = 2.68 Hz, 1H, ArH), 7.18-7.22 (m, 1H, ArH), 7.25-7.29 (m, 4H, ArH), 7.32 (dd, J = 9.40, 2.68 Hz, 1H, ArH), 7.60 (d, J = 9.24 Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 48.5, 52.6, 55.7, 62.8, 106.2, 116.5, 122.4, 126.7, 127.3, 128.2, 128.3, 129.2, 137.9, 138.7, 154.0, 159.0; MS (m/z , %): 360 ($M+1$, 100); Anal. Calcd. for $\text{C}_{21}\text{H}_{21}\text{N}_5\text{O}$ (359.42): C, 70.17; H, 5.89; N, 19.48 %. Found C, 70.32; H, 5.64; N, 19.32 %.

General Procedure for Iron Catalyzed Alkyl – 3-Chloroquinoxaline Cross-Coupling Reactions : Synthesis of **22a** and **23a**

A flame-dried 100 mL two-necked flask was charged with 3-chloro-7-methoxy-2-(methylthio)quinoxaline **3a** (1.0 g, 4.15 mmol), $\text{Fe}(\text{acac})_3$ (0.15 g, 0.41 mmol), THF (25 mL) and *N*-methylpyrrolidone (3.0 mL) and the mixture was cooled to 0 °C. A solution of methylmagnesium iodide or *n*-butylmagnesium bromide (2.0 M in Et_2O , 4.98 mmol) was added via syringe to the resulting red solution, causing a immediate color change to dark brown-black. The reaction mixture was further stirred at the same temperature for 5-10 min and quenched with brine solution. It was then extracted with CHCl_3 (3×25 mL), washed with water (2×25 mL), followed by brine (25 mL) and dried over anhydrous Na_2SO_4 . The solvent was removed under vacuum to give crude products **22a** and **23a** which were purified by column chromatography over silica gel using hexane:EtOAc (50:1) as eluent.

7-Methoxy-3-methyl-2-(methylthio)quinoxaline (22a)

Yield 82% (0.75 g); White crystals; m.p. 89-90 °C; R_f 0.78 (10:1 hexanes-EtOAc); IR (KBr): 2922, 1616, 1496, 1296, 1223, 817 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.64 (s, 3H, Me), 2.68 (s, 3H, SMe), 3.95 (s, 3H, OMe), 7.22-7.24 (m, 2H, ArH), 7.82 (d, $J = 10.0$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 12.7, 21.6, 55.5, 105.7, 119.9, 129.0, 134.6, 142.6, 149.0, 156.3, 159.9; MS (m/z , %): 221 (M+1, 100); Anal. Calcd. for $\text{C}_{11}\text{H}_{12}\text{N}_2\text{OS}$ (220.29): C, 59.97; H, 5.49; N, 12.72 %. Found C, 59.78; H, 5.32; N, 12.67 %.

3-(n-Butyl)-7-methoxy-2-(methylthio)quinoxaline (23a)

Yield 80% (0.87 g); Pale yellow crystals; m.p. 38-39 °C; R_f 0.70 (20:1 hexanes-EtOAc); IR (KBr): 2926, 1612, 1494, 1353, 1284, 1211, 824 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 0.87 (t, $J = 7.32$ Hz, 3H, CH_3), 1.32-1.42 (m, 2H, CH_2), 1.67-1.74 (m, 2H, CH_2), 2.54 (s, 3H, SMe), 2.80 (t, $J = 8.04$ Hz, 2H, CH_2), 3.80 (s, 3H, OMe), 7.07-7.10 (m, 2H, ArH), 7.71 (d, $J = 5.96$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 12.6, 13.8, 22.6, 29.6, 34.4, 55.4, 105.7, 119.8, 129.2, 134.7, 142.5, 152.5, 156.0, 159.8; MS (m/z , %): 263 (M+1, 100); Anal. Calcd. for $\text{C}_{14}\text{H}_{18}\text{N}_2\text{OS}$ (262.37): C, 64.09; H, 6.91; N, 10.68 %. Found C, 63.82; H, 6.78; N, 10.82 %.

General Procedure for Iron Catalyzed Aryl-3-Chloroquinoxaline Cross-Coupling Reactions : Synthesis of 22b and 23b

A flame-dried 100 mL two-necked flask was charged with 3-chloro-7-methoxy-2-(methylthio)quinoxaline **3a** (1.0 g, 4.15 mmol), $\text{Fe}(\text{acac})_3$ (0.15 g, 0.41 mmol), THF (25 mL) and the mixture was cooled to -30 °C. A solution of phenylmagnesium bromide or 4-methoxyphenylmagnesium bromide (1.0 M in THF, 9.13 mmol) was added via syringe to the resulting red solution, causing a immediate color change to dark brown-black. The

reaction mixture was further stirred for 10-15 min and quenched with brine solution. It was then extracted with CHCl_3 (3×25 mL), washed with water (2×25 mL), followed by brine (25 mL) and dried over anhydrous Na_2SO_4 . Standard purification by column chromatography using hexane:EtOAc as eluent provides the products **22b** and **23b**.

7-Methoxy-2-(methylthio)-3-phenylquinoxaline (**22b**)

Yield 68% (0.79 g); Colorless crystals; m.p. 142-143 °C; R_f 0.65 (10:1 hexanes-EtOAc); IR (KBr): 3051, 2920, 1607, 1484, 1214, 1091, 826 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.61 (s, 3H, SCH_3), 3.95 (s, 3H, OMe), 7.25 (dd, $J = 9.02, 2.44$ Hz, 1H, ArH), 7.28 (d, $J = 2.68$ Hz, 1H, ArH), 7.47-7.53 (m, 3H, ArH), 7.74-7.77 (m, 2H, ArH), 7.92 (d, $J = 9.04$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.6, 55.6, 105.6, 120.6, 128.3, 128.9, 129.3, 130.0, 134.9, 137.4, 143.0, 150.8, 155.7, 160.7; MS (m/z , %): 283 ($M+1$, 100), 235 ($M-47$, 63); Anal. Calcd. for $\text{C}_{16}\text{H}_{14}\text{N}_2\text{SO}$ (282.36): C, 68.06; H, 5.00; N, 9.92 %. Found C, 68.23; H, 5.19; N, 9.73 %.

7-Methoxy-3-(4'-methoxyphenyl)-2-(methylthio)quinoxaline (**23b**)

Yield 64% (0.83 g); Colorless solid; m.p. 115-116 °C; R_f 0.80 (10:1 hexanes-EtOAc); IR (KBr): 3002, 2920, 1613, 1513, 1253, 1097, 1032, 825 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.62 (s, 3H, SCH_3), 3.86 (s, 3H, OMe), 3.96 (s, 3H, OMe), 7.02 (d, $J = 8.80$ Hz, 2H, ArH), 7.24 (dd, $J = 8.90, 2.68$ Hz, 1H, ArH), 7.28 (d, $J = 2.68$ Hz, 1H, ArH), 7.72 (d, $J = 8.86$ Hz, 2H, ArH), 7.91 (d, $J = 9.04$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.7, 55.2, 55.6, 105.6, 113.7, 120.5, 129.8, 129.9, 130.4, 134.9, 142.7, 150.5, 155.8, 160.5, 160.5; MS (m/z , %): 313 ($M+1$, 100); Anal. Calcd. for $\text{C}_{17}\text{H}_{16}\text{N}_2\text{SO}_2$ (312.39): C, 65.36; H, 5.16; N, 8.97 %. Found C, 65.21; H, 5.32; N, 8.81 %.

General Procedure for Nickel-Catalyzed Cross-Coupling Reactions of **22** and **23** with Grignard Reagents: Synthesis of **24a-c** and **25a-c**

A solution of respective Grignard reagent (0.10 mmol) in Et₂O was added dropwise to a stirring suspension of (PPh₃)₂NiCl₂ (30 mol %) in 25 mL dry benzene under argon atmosphere and the mixture was refluxed for 15 minutes. After the catalyst reduction, 1.90 mmol of the required Grignard reagent (MeMgI/PhMgBr/4-MeOC₆H₄Br) and a solution of 2-methylthio-3-alkyl/arylquinoxalines **22** or **23** (1.0 mmol) in dry benzene (20 mL) were added to the reaction mixture and refluxed for 8-12 h. It was then cooled, poured into 50 mL of saturated NH₄Cl solution and extracted with CHCl₃ (3 × 50 mL). The organic layer was dried (anhydrous Na₂SO₄) and evaporated to give crude products **24a-c** or **25a-c**, which were purified by column chromatography using hexane: EtOAc as eluent. In the case of *n*-butylmagnesium bromide 30 mol% of NiCl₂(dppp) was used as catalyst to avoid any reduction of methylthio group.

2-(*n*-Butyl)-7-methoxy-3-methylquinoxaline (**24a**)

Yield 80% (0.18 g); Viscous liquid; R_f 0.75 (10:1 hexanes-EtOAc); IR (KBr): 2364, 1598, 1353, 768 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 0.90 (t, *J* = 7.56 Hz, 3H, CH₃), 1.38-1.43 (m, 2H, CH₂), 1.66-1.73 (m, 2H, CH₂), 2.62 (s, 3H, CH₃), 2.87 (t, *J* = 8.04 Hz, 2H, CH₂), 3.84 (s, 3H, OMe), 7.19-7.22 (m, 2H, ArH), 7.76 (dd, *J* = 8.30, 1.20 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 13.8, 22.3, 22.7, 30.3, 35.5, 55.5, 106.0, 121.6, 129.0, 136.6, 142.35, 150.14, 156.7, 159.83; MS (*m/z*, %): 231 (M+1, 100); Anal. Calcd. for C₁₄H₁₈N₂O (230.31): C, 73.01; H, 7.88; N, 12.16 %. Found C, 73.10; H, 7.72; N, 12.31 %.

2-(*n*-Butyl)-7-methoxy-3-phenylquinoxaline (24b)

Yield 70% (0.20 g); Colorless viscous liquid; R_f 0.58 (10:1 hexanes-EtOAc); IR (film): 2958, 1618, 1489, 1348, 1223, 1026, 831 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 0.76 (t, $J = 7.32$ Hz, 3H, CH_3), 1.19-1.28 (m, 2H, CH_2), 1.59-1.66 (m, 2H, CH_2), 2.93 (t, $J = 7.80$ Hz, 2H, CH_2), 3.90 (s, 3H, OMe), 7.27 (d, $J = 2.44$ Hz, 1H, ArH), 7.30-7.31 (m, 1H, ArH), 7.38-7.44 (m, 3H, ArH), 7.51 (d, $J = 7.08$ Hz, 2H, ArH), 7.90 (d, $J = 8.76$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.7, 22.5, 31.3, 35.6, 55.7, 105.8, 122.4, 128.4, 128.5, 128.8, 130.1, 136.7, 139.2, 142.7, 152.4, 156.1, 160.6; MS (m/z , %): 293 ($M+1$, 100); Anal. Calcd. for $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}$ (292.37): C, 78.05; H, 6.89; N, 9.58 %. Found C, 78.20; H, 6.68; N, 9.69 %.

7-Methoxy-2-(4'-methoxyphenyl)-3-phenylquinoxaline (24c)

Yield 72% (0.25 g); Pale yellow crystals; m.p. 154-155 $^{\circ}\text{C}$; R_f 0.35 (10:1 hexanes-EtOAc); IR (KBr): 2920, 1517, 1277, 1165, 937 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 3.79 (s, 3H, OMe), 3.96 (s, 3H, OMe), 6.83 (d, $J = 8.80$ Hz, 2H, ArH), 7.31-7.33 (m, 2H, ArH), 7.37 (dd, $J = 9.28, 2.68$ Hz, 1H, ArH), 7.44 (d, $J = 8.52$ Hz, 2H, ArH), 7.47-7.50 (m, 4H, ArH), 8.01 (d, $J = 9.28$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 55.2, 55.8, 106.0, 113.6, 123.0, 128.2, 128.4, 129.6, 130.0, 131.0, 131.3, 137.1, 139.3, 142.3, 150.8, 152.5, 160.1, 160.9; MS (m/z , %): 343 ($M+1$, 100); Anal. Calcd. for $\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}_2$ (342.39): C, 77.17; H, 5.30; N, 8.18 %. Found C, 77.41; H, 5.12; N, 8.41 %.

3-(*n*-Butyl)-7-methoxy-2-methylquinoxaline (25a)

Yield 80% (0.18 g); Pale yellow crystals; m.p. 55-56 °C; R_f 0.40 (10:1 hexanes-EtOAc); IR (KBr): 2939, 1615, 1355, 1216, 1023, 835 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 0.94 (t, $J = 7.32$ Hz, 3H, CH_3), 1.41-1.47 (m, 2H, CH_2), 1.71-1.75 (m, 2H, CH_2), 2.68 (s, 3H, CH_3), 2.90 (t, $J = 8.04$ Hz, 2H, CH_2), 3.88 (s, 3H, OMe), 7.23-7.27 (m, 2H, ArH), 7.82 (d, $J = 8.76$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.9, 22.6, 22.7, 30.4, 35.3, 55.6, 105.9, 121.5, 129.3, 137.0, 142.1, 152.9, 154.1, 159.8; MS (m/z , %): 231 ($M+1$, 100); Anal. Calcd. for $\text{C}_{14}\text{H}_{18}\text{N}_2\text{O}$ (230.31): C, 73.01; H, 7.88; N, 12.16 %. Found C, 72.91; H, 7.68; N, 12.01 %.

3-(*n*-Butyl)-7-methoxy-2-phenylquinoxaline (25b)

Yield 89% (0.26 g); Colorless viscous liquid; R_f 0.58 (20:1 hexanes-EtOAc); IR (film): 2955, 1594, 1360, 1220, 834 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 0.75 (t, $J = 7.32$ Hz, 3H, CH_3), 1.17-1.26 (m, 2H, CH_2), 1.57-1.64 (m, 2H, CH_2), 2.91 (t, $J = 8.04$ Hz, 2H, CH_2), 3.85 (s, 3H, OMe), 7.28-7.31 (m, 2H, ArH), 7.38-7.44 (m, 3H, ArH), 7.51 (d, $J = 6.32$ Hz, 2H, ArH), 7.87 (d, $J = 9.04$ Hz, 1H, ArH); ^{13}C NMR (100 MHz, CDCl_3): δ 13.7, 22.5, 31.3, 35.3, 55.6, 106.5, 122.8, 128.4, 128.6, 128.7, 129.3, 137.4, 139.2, 142.0, 153.4, 154.8, 160.1; MS (m/z , %): 293 ($M+1$, 100); Anal. Calcd. for $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}$ (292.37): C, 78.05; H, 6.89; N, 9.58 %. Found C, 78.21; H, 6.67; N, 9.40 %.

7-Methoxy-3-(4'-methoxyphenyl)-2-phenylquinoxaline (25c)

Yield 68% (0.23 g); Pale yellow crystals; m.p. 184-185 °C; R_f 0.43 (10:1 hexanes-EtOAc); IR (KBr): 3062, 2962, 1609, 1476, 1344, 1239, 1017, 830 cm^{-1} ; ^1H NMR (400

MHz, CDCl₃): δ 3.79 (s, 3H, OMe), 3.96 (s, 3H, OMe), 6.84 (d, J = 9.14 Hz, 2H, ArH), 7.32-7.35 (m, 3H, ArH), 7.39 (dd, J = 9.04, 2.68 Hz, 1H, ArH), 7.42-7.45 (m, 3H, ArH), 7.52 (dd, J = 7.56, 1.96 Hz, 2H, ArH), 8.02 (d, J = 9.00 Hz, 1H, ArH); ¹³C NMR (100 MHz, CDCl₃): δ 55.1, 55.7, 106.3, 113.5, 123.1, 128.2, 128.5, 129.6, 129.9, 131.1, 131.4, 137.3, 139.4, 142.3, 150.4, 153.1, 159.8, 160.5; MS (m/z, %): 343 (M+1, 100); Anal. Calcd. for C₂₂H₁₈N₂O₂ (342.39): C, 77.17; H, 5.30; N, 8.18 %. Found C, 77.38; H, 5.16; N, 8.37 %.