SUPPORTING INFORMATION

Highly Convergent Synthesis of C₃- or C₂-Symmetric Carbohydrate Macrocycles

Kyle D. Bodine, David Y. Gin and Mary S. Gin*

*Department of Chemistry, University of Illinois at Urbana-Champaign, 600 S. Mathews Ave,
Urbana, Illinois 61801*

mgin@scs.uiuc.edu

CONTENTS

General Information	S1
Experimental Procedures	S3
Modeling Images	S19
¹H and ¹³C NMR Plots	S22

General Procedures. All reactions were performed in flame-dried modified Schlenk (Kjeldahl shape) flasks fitted with a stopper or sealed tubes under a positive pressure of argon, unless otherwise noted. Air- and moisture-sensitive liquids and solutions were transferred via syringe or stainless steel cannula. Organic solutions were concentrated by rotary evaporation below 35 °C at ca. 25 Torr or by evaporation via a stream of dry nitrogen gas. Flash column chromatography was performed employing 230-400 mesh silica gel. Thin-layer chromatography was performed using glass plates pre-coated to a depth of 0.25 mm with 230-400 mesh silica gel impregnated with a fluorescent indicator (254 nm). TLC plates were visualized with UV light and/or ceric ammonium molybdate (CAM). All saccharide intermediates and diphenylsulfoxide were dried by removal of water as an azeotrope with toluene.

Materials. Dichloromethane (CH₂Cl₂), triethylamine (Et₃N), trimethylsilylazide (Me₃SiN₃), and 1,8-diaza[5.4.0]bicycloundec-7-ene (DBU) were distilled from CaH₂ and

toluene was distilled from Na metal. Trifluoromethanesulfonic anhydride (Tf₂O) was triply distilled from P₂O₅ and TfOH. H₂O was purified by a Millipore Milli-Q filtration unit. 2,4-Hexadiyne-1,6-diol was twice recrystallized from CHCl₃/MeOH (20:1).

Instrumentation. Infrared (IR) spectra were obtained using a Perkin Elmer Spectrum BX spectrophotometer referenced to a polystyrene standard. Data are presented as the frequency of absorption (cm⁻¹), and intensity of absorption (s = strong, m = medium, w = weak). Proton and carbon-13 nuclear magnetic resonance (¹H NMR or ¹³C NMR) spectra were recorded on a Varian Unity 400/500 or a Varian Unity Inova 500 NMR spectrometer; chemical shifts are expressed in parts per million (δ scale) downfield from tetramethylsilane and are referenced to the residual protium in the NMR solvent (CHCl₃: δ 7.26 for ¹H NMR, δ 77.23 for ¹³C NMR; HOD: δ 4.79 for ¹H NMR; Methanol: δ 3.31 for ¹H NMR, δ 49.00 for ¹³C NMR). Data are presented as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, m = multiplet and/or multiple resonances), integration, coupling constant in Hertz (Hz). GCOSY, GHMQC, TOCSY 1D, and NOESY 1D spectra were recorded on a Varian Unity Inova 500 or a Varian Unity Inova 600 NMR spectrometer. Low resolution fast atom bombardment (FAB) mass spectra were obtained on a Micromass ZAB-SE spectrometer. High resolution FAB mass spectra were obtained on a Micromass 70-SE-4F spectrometer. High resolution electrospray ionization (ESI) mass spectra were obtained on a Micromass Q-Tof Ultima spectrometer. Matrix assisted laser desorption ionization (MALDI) mass spectra were obtained using an Applied Biosystems Voyager-DE STR. Optical rotations were determined on a Jasco P-1020 polarimeter using a sodium lamp at 25 °C (concentrations provided in units of g/100mL).

Experimental Procedures.

Azido-alkynyl monosaccharides (6 α and 6 β). Glycosyl acceptor **3**¹ and diphenylsulfoxide were dried by azeotropic removal of water with toluene immediately prior to reaction. To a solution of **3** (465 mg, 0.952 mmol, 1.00 equiv) and diphenylsulfoxide (540 mg, 2.67 mmol, 2.80 equiv) in 35.3 mL of toluene and 11.7 mL of CH₂Cl₂ at -78 °C was added Tf₂O (0.224 mL, 1.33 mmol, 1.40 equiv). After stirring for 15 min at -78 °C, the reaction was warmed to -45 °C and stirred at that temperature for 30 min. Azidotrimethylsilane (0.632 mL, 4.76 mmol, 5.00 equiv) was injected to the reaction mixture and the reaction was stirred at -45 °C for an additional 30 min. The solution was warmed to -20 °C and stirred at that temperature for 12 h, then Et₃N (1.33 mL, 9.52 mmol, 10.0 equiv) was added and the reaction warmed to 23 °C. The reaction mixture was diluted with 200 mL EtOAc, and washed with saturated aqueous NaHCO₃ (2 x 100 mL) and saturated aqueous NaCl (1 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The crude mixture was purified by silica gel flash column chromatography (hexanes/benzene/EtOAc 5:2:1) to afford 290 mg **6 α** (60% yield) as a colorless oil and 172 mg (35% yield) **6 β** as a colorless oil.

6 α : R_f = 0.61 (benzene/EtOAc 5:1); $[\alpha]_D$: +84.94° (c = 1.6, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.38-7.25 (m, 15H), 5.38 (d, 1H, J = 1.9), 4.68 (m, 3H), 4.58 (m, 3H), 4.44 (dd, 1H, J = 15.4, 2.5), 4.28 (dd, 1H, J = 15.4, 2.5), 3.92 (t, 1H, J = 9.1), 3.87-3.82 (m, 3H), 3.76 (dd, 1H, J = 9.1, 2.9), 3.61 (m, 1H), 2.39 (t, 1H, J = 2.5);

$^{13}\text{C}_1\text{-}^1\text{H}_1$ $J = 169$ Hz; ^{13}C NMR (125 MHz, CDCl_3) δ 138.5, 138.2, 137.9, 128.67, 128.66, 128.5, 128.06, 128.04, 128.00, 127.94, 127.91, 127.7, 88.2, 80.2, 79.1, 74.6, 74.5, 74.1, 73.7, 73.6, 72.9, 72.5, 69.3, 60.2; IR (neat film) 3284 (w), 3025 (m), 2912 (m), 2857 (m), 2112 (s), 1495 (m), 1454 (m), 1358 (m), 1123 (s), 1047 (s), 826 (m), 739 (s); HRMS (ESI) m/z : Calcd for $\text{C}_{30}\text{H}_{31}\text{N}_3\text{O}_5\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 536.2161, found 536.2158.

6 β : $R_f = 0.53$ (benzene/EtOAc 5:1); $[\alpha]_D: -57.73^\circ$ ($c = 3.3$, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ 7.45-7.26 (m, 15H), 4.92 (d, 1H, $J = 11.8$), 4.85 (d, 1H, $J = 11.8$), 4.71 (d, 1H, $J = 12.0$), 4.63 (m, 3H), 4.47 (dd, 1H, $J = 15.3, 2.4$), 4.39 (d, 1H, $J = 1.0$), 4.36 (dd, 1H, $J = 15.3, 2.4$), 3.93-3.87 (m, 3H), 3.83 (dd, 1H, $J = 11.5, 5.8$), 3.55-3.51 (m, 2H), 2.43 (t, 1H, $J = 2.4$); $^{13}\text{C}_1\text{-}^1\text{H}_1$ $J = 156$ Hz; ^{13}C NMR (125 MHz, CDCl_3) δ 138.5, 138.1, 137.9, 128.7, 128.5, 128.4, 128.3, 128.1, 127.94, 127.87, 127.78, 127.68, 86.6, 82.7, 80.1, 78.0, 75.8, 74.8, 74.7, 74.0, 73.7, 72.3, 69.4, 60.3; IR (neat film) 3284 (w), 3025 (m), 2912 (m), 2857 (m), 2112 (s), 1495 (m), 1454 (m), 1358 (m), 1123 (s), 1047 (s), 826 (m), 739 (s); HRMS (ESI) m/z : Calcd for $\text{C}_{30}\text{H}_{31}\text{N}_3\text{O}_5\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 536.2161, found 536.2159.

Azido-alkynyl disaccharides (7 α and 7 β). Glycosyl acceptor **5**¹ and diphenylsulfoxide were dried by azeotropic removal of water with toluene immediately prior to reaction.

To a solution of acceptor **5** (700 mg, 0.760 mmol, 1.00 equiv) and diphenylsulfoxide (430 mg, 2.13 mmol, 2.80 equiv) in 37.5 mL of toluene and 12.5 mL of CH₂Cl₂ at -78 °C was added Tf₂O (180 μL, 1.06 mmol, 1.40 equiv). After stirring for 20 min, the reaction was warmed to -45 °C and stirred at that temperature for 30 min. Trimethylsilylazide (0.505 mL, 3.80 mmol, 5.00 equiv) was added and the reaction was stirred at -45 °C for an additional 30 min. The solution was then warmed to -20 °C and stirred at that temperature for 16 hours, then Et₃N (1.06 mL, 7.60 mmol, 10.0 equiv) was added and the reaction was warmed to 23 °C. The reaction mixture was diluted with 200 mL of EtOAc and washed with saturated aqueous NaHCO₃ (2 x 50 mL) and saturated aqueous NaCl (1 x 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The crude mixture was purified by silica gel flash column chromatography (benzene/hexanes/EtOAc 7:6:2) to afford 528 mg **7α** (73% yield) as a colorless oil and 169 mg **7β** (24% yield) as a white foam.

7α: R_f = 0.61 (benzene/EtOAc 5:1); $[\alpha]_D$: +56.82° (c = 1.7, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.40-7.14 (m, 30H), 5.41 (d, 1H, J = 1.8), 5.28 (d, 1H, J = 1.8), 4.61 (m, 3H), 4.58 (d, 1H, J = 11.8), 4.53 (m, 2H), 4.48 (m, 3H), 4.43 (dd, 1H, J = 15.4, 2.4), 4.35 (d, 1H, J = 11.8), 4.29 (app s, 2H), 4.24 (dd, 1H, J = 15.4, 2.4), 4.11 (t, 1H, J = 9.5), 3.92-3.85 (m, 2H), 3.84-3.78 (m, 3H), 3.76-3.64 (m, 5H), 3.61 (t, 1H, J = 2.6), 2.37 (t, 1H, J = 2.4); ¹³C-¹H J = 169 Hz, and 171 Hz; ¹³C NMR (125 MHz, CDCl₃) δ 138.74, 138.71, 138.70, 138.6, 137.9, 137.8, 128.7, 128.6, 128.54, 128.46, 128.42, 128.26, 128.10, 128.06, 127.9, 127.83, 127.79, 127.76, 127.74, 127.72, 127.58, 127.43, 127.38, 100.0, 87.9, 80.5, 80.0, 75.6, 74.6, 74.2, 73.93, 73.87, 73.78, 73.6, 73.4, 72.8, 72.7, 72.12, 72.08, 71.6, 69.9, 69.6, 60.2; IR (neat film) 3280 (w), 3032 (m), 2910 (m), 2863 (m),

2110 (s), 1495 (m), 1454 (m), 1358 (m), 1123 (s), 1047 (s), 826 (m), 742 (s); HRMS (ESI) m/z : Calcd for $C_{57}H_{59}N_3O_{10}Na$ ($M+Na$)⁺ 968.4001, found 968.4004.

7β: R_f = 0.57 (benzene/EtOAc 5:1); $[\alpha]_D$: -20.87° (c = 2.0, $CHCl_3$); 1H NMR (500 MHz, $CDCl_3$) δ 7.43-7.14 (m, 30H), 5.36 (d, 1H, J = 1.7), 4.83 (app s, 2H), 4.64 (d, 1H, J = 12.1), 4.60 (d, 1H, J = 11.7), 4.56 (m, 3H), 4.53 (d, 1H, J = 11.7), 4.50 (d, 1H, J = 12.1), 4.45 (dd, 1H, J = 15.4, 2.3), 4.43 (app s, 1H), 4.35 (d, 1H, J = 11.5), 4.28 (app s, 2H), 4.25 (dd, 1H, J = 15.4, 2.3), 4.10 (t, 1H, J = 9.6), 3.94-3.86 (m, 3H), 3.81-3.74 (m, 2H), 3.74-3.71 (m, 1H), 3.70-3.67 (m, 2H), 3.64 (dd, 1H, J = 10.5, 1.2), 3.55 (m, 1H), 3.45 (dd, 1H, J = 9.5, 2.4), 2.39 (t, 1H, J = 2.3); ^{13}C - 1H J = 172 Hz, and 158 Hz; ^{13}C NMR (125 MHz, $CDCl_3$) δ 138.70, 138.69, 138.67, 138.59, 138.04, 137.5, 128.8, 128.51, 128.50, 128.49, 128.42, 128.24, 128.21, 128.15, 127.93, 127.91, 127.82, 127.73, 127.71, 127.63, 127.56, 127.53, 127.39, 127.37, 109.9, 100.1, 86.4, 83.3, 80.4, 79.9, 77.9, 75.6, 75.1, 74.8, 74.6, 74.2, 73.7, 73.63, 73.57, 72.7, 72.1, 71.7, 69.9, 69.4, 60.1; IR (neat film) 3271 (w), 3028 (m), 2914 (m), 2874 (m), 2112 (s), 1497 (m), 1450 (m), 1361 (m), 1125 (s), 1050 (s), 831 (m), 732 (s); HRMS (ESI) m/z : Calcd for $C_{57}H_{59}N_3O_{10}Na$ ($M+Na$)⁺ 968.4001, found 968.3998.

Cyclotrimer (8). To a solution of monosaccharide **6** (320 mg, 0.623 mmol, 1.00 equiv) and DBU (1.86 mL, 12.5 mmol, 20.0 equiv) in 100.0 mL of toluene was

added CuI (119 mg, 0.623 mmol, 1.00 equiv) and the reaction mixture was stirred for 20 hours. The crude mixture was filtered through a short column of silica gel with 300 mL (1:1) EtOAc/CH₂Cl₂ to remove the insoluble material, and the filtrate was concentrated in vacuo. The crude organic material was purified by silica gel flash column chromatography (toluene/CH₂Cl₂/EtOAc/MeCN 12:4:2:1) to afford 198 mg **8** (62% yield) as a waxy white solid.

8: R_f = 0.48 (benzene/EtOAc 2:1); $[\alpha]_D$: +40.56° (c = 0.4, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.42-7.20 (m, 45H), 6.00 (br s, 3H), 4.98 (br s, 3H), 4.66 (br m, 19H), 4.46 (br m, 8H), 3.99 (br s, 7H), 3.74 (br m, 4H), 3.65 (br s, 4H); ¹³C NMR (125 MHz, CDCl₃) δ 138.08, 138.04, 137.67, 128.75, 128.69, 128.66, 128.16, 128.13-128.03 (broad signal), 121.4 (broad), 85.0 (broad), 75.5-74.4 (broad), 73.9-73.1 (broad), 68.5, 66.4-65.3 (broad); IR (thin film) 3062 (w), 3024 (w), 2922 (m), 2863 (m), 1726 (w), 1495 (m), 1455 (m), 1366 (m), 1323 (m), 1255 (m), 1207 (m), 1108 (s), 1038 (s), 906 (m), 820 (m), 799 (m), 737 (s); HRMS (ESI) m/z : Calcd for C₉₀H₉₃N₉O₁₅ (M^{•+}) = 1539.6791, found 1539.6793.

Cyclodimer (10). To a solution of disaccharide **7** (425 mg, 0.449 mmol, 1.00 equiv) and DBU (1.34 mL, 8.98 mmol, 20.0 equiv) in 75 mL of toluene was added CuI (86.0 mg, 0.449 mmol, 1.00 equiv) and the reaction mixture was stirred for 18 hours. The

crude mixture was filtered through a short column of silica gel with 600 mL (1:1) EtOAc/CH₂Cl₂ to remove the insoluble material, and the filtrate was concentrated in vacuo. The crude organic material was purified by silica gel flash column chromatography (hexanes/benzene/EtOAc 3:3:2) to afford 378 mg **10** (89% yield) as a waxy white solid.

10: R_f = 0.32 (benzene/EtOAc 5:1); $[\alpha]_D$: +27.77° (c = 1.9, CHCl₃); ¹H NMR (500 MHz, CDCl₃, 45 °C) δ 7.56 (s, 1H), 7.36-7.18 (m, 61H), 6.03 (s, 1H), 5.15 (d, 1H, J = 10.1), 4.97 (s, 1H), 4.70 (d, 1H, J = 11.6), 4.65-4.48 (m, 12H), 4.41 (d, 1H, J = 11.9), 4.36 (d, 1H, J = 11.9), 4.03-3.93 (m, 6H), 3.87-3.78 (m, 16H), 3.74-3.62 (m, 13H), 3.47 (m, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 146.2 (broad), 138.58, 138.55, 138.46, 138.3, 137.8, 137.5, 128.71, 128.65, 128.61, 128.58, 128.54, 128.47, 128.24, 128.15, 128.14, 128.03, 127.99, 127.90, 127.81, 127.77, 127.74, 127.67, 127.64, 121.45 (broad), 101.7, 84.4 (broad), 79.3, 78.0 (broad), 75.2, 75.0, 73.6, 73.3, 73.1 (broad), 72.7 (broad), 72.5, 72.3, 69.0 (broad), 66.1 (broad), 53.6; IR (thin film) 3057 (w), 3030 (w), 2922 (m), 2866 (m), 1866 (w), 1957 (w), 1879 (w), 1602 (w), 1496 (m), 1454 (m), 1360 (m), 1315 (w), 1280 (w), 1204 (m), 1110 (s), 1063 (s), 909 (w), 818 (w), 794 (m), 736 (s); HRMS (FAB) m/z : Calcd for C₁₁₄H₁₁₉N₆O₂₀ (M+H)⁺ = 1891.8466, found 1891.8479.

Deprotected Cyclotrimer (9). To a solution of protected cyclotrimer **8** (67.0 mg, 0.0435 mmol, 1.00 equiv) in a mixture of 5.8 mL of EtOAc, 5.8 mL of EtOH, and 2.9 mL of H₂O was added Pd/C (10% w/w, 463 mg, 0.435 mmol Pd, 10 equiv Pd) and ammonium formate (480 mg, 7.61 mmol, 175 equiv). The reaction mixture was stirred for 16 h in a sealed tube. The crude mixture was filtered through a Millipore Millex-HN (0.45 μ m) HPLC sample preparation filter to remove insoluble materials. The filtrate was lyophilized to dryness and was then dissolved into 40 mL of Millipore grade H₂O. The crude solution was dialyzed in a SpectraPor CE membrane (MW cutoff: 100 g/mol) for 3 days to remove the ammonium salts. The solution of purified material was lyophilized to dryness to yield 31 mg **9** (>99% yield) as a white low density foam.

9: R_f = 0.56 (ⁱPrOH/H₂O/EtOAc/NH₄OH 5:4:3:1); $[\alpha]_D$: -31.91° (c = 1.2, H₂O); ¹H NMR (500 MHz, D₂O) δ 8.17 (s, 3H), 6.11 (d, 3H, J = 3.1), 4.98 (d, 3H, J = 11.0), 4.84 (t, 3H, J = 3.1), 4.73 (d, 3H, J = 11.0), 3.95 (dd, 3H, J = 8.3, 3.1), 3.83 (t, 3H, J = 8.6), 3.82 (dd, 3H, J = 12.5, 5.2), 3.79 (dd, 3H, J = 12.5, 2.5), 3.16 (broad m, 3H); ¹³C NMR (125 MHz, D₂O) δ 144.8, 123.6, 86.6, 75.4, 70.1, 68.3, 68.1, 64.9, 60.1; IR (KBr Pellet) 2915 (w), 2848 (w), 1696 (m), 1631 (m), 1571 (m), 1466 (m), 1432 (m), 1386 (w), 1300 (w), 1257 (w), 1215 (w), 1108 (s), 1050 (s), 737 (m); HRMS (ESI) m/z : Calcd for C₂₇H₃₉N₉O₁₅ (M^{+}) = 729.2566, found 729.2572.

Deprotected Cyclodimer (11). To a solution of protected cyclodimer **10** (170 mg, 0.0899 mmol, 1.00 equiv) in a mixture of 12 mL of EtOAc, 12 mL of EtOH, and 6 mL of H₂O was added Pd/C (10% w/w, 620 mg, 0.583 mmol Pd, 6.5 equiv Pd) and ammonium formate (992 mg, 15.7 mmol, 175 equiv). The reaction mixture was stirred for 21 h in a sealed tube. The crude mixture was filtered through a Millipore Millex-HN (0.45 μ m) HPLC sample preparation filter to remove insoluble materials. The filtrate was lyophilized to dryness and was then dissolved into 30 mL of Millipore grade H₂O. The crude solution was dialyzed in a SpectraPor CE membrane (MW cutoff: 100 g/mol) for 2 days to remove the ammonium salts. The solution of purified material was lyophilized to dryness to yield 72 mg **11** (>99% yield) as a white low density foam.

11: R_f = 0.59 (ⁱPrOH/H₂O/EtOAc/NH₄OH 5:4:3:1); $[\alpha]_D$: +31.89° (c = 2.6, H₂O); ¹H NMR (500 MHz, D₂O) δ 8.24 (s, 2H), 6.11 (d, 2H, J = 3.0), 4.96 (d, 2H, J = 10.7), 4.95 (d, 2H, J = 1.8), 4.89 (t, 2H, J = 3.0), 4.76 (d, 2H, J = 10.7), 3.99 (dd, 2H, J = 3.0, 2.2), 3.86-3.80 (m, 9H), 3.79-3.74 (m, 5H), 3.70 (t, 2H, J = 9.5), 3.55 (ddd, 2H, J = 9.5, 5.0, 2.2), 3.34 (broad m, 2H); ¹³C NMR (125 MHz, D₂O) δ 124.1, 110.0, 102.7, 86.6, 77.9, 75.5, 72.2, 71.1, 70.67, 70.61, 70.0, 68.0, 65.0, 60.75, 60.70; IR (KBr Pellet) 2916 (w), 2849 (w), 1696 (m), 1631 (m), 1571 (m), 1469 (m), 1432 (m), 1386 (w), 1301 (w), 1257 (w), 1215 (w), 1105 (s), 1056 (s), 730 (m); HRMS (ESI) m/z : Calcd for C₃₀H₄₇N₆O₂₀ (M+H)⁺ 811.2841, found 811.2845.

¹H NMR Binding Titrations.

Monosaccharide Cyclotrimer 9 (Host) with 2,4-Hexadiyne-1,6-diol 12 (Guest):

Preparation of Solutions.

- (A) 13.6 mg cyclotrimer **9** was dissolved into 3.00 mL D₂O to give a 6.2 mM clear solution.
- (B) 103 mg 2,4-Hexadiyne-1,6-diol was dissolved into 5.00 mL D₂O to give a 186 mM clear solution.
- (C) 264 mg 2,4-Hexadiyne-1,6-diol was dissolved into 5.00 mL D₂O to give a 480 mM clear solution.

Preparation of Guest and Host Mixtures.

The solutions were mixed according to the table below such that the total concentration of cyclotrimer **9** was always 3.1 mM and the total volume was 0.7 mL.

Guest : Host Ratio	Sol'n A	Sol'n B	Sol'n C	D₂O
0 : 1	350 µL	0 µL	0 µL	350 µL
1 : 1	350 µL	11.5 µL	0 µL	338.5 µL
2 : 1	350 µL	23 µL	0 µL	327 µL
5 : 1	350 µL	58 µL	0 µL	292 µL
10 : 1	350 µL	116 µL	0 µL	234 µL
20 : 1	350 µL	234 µL	0 µL	116 µL
30 : 1	350 µL	350 µL	0 µL	0 µL
50 : 1	350 µL	0 µL	219 µL	131 µL
80 : 1	350 µL	0 µL	350 µL	0 µL

^1H NMR of Mixtures.

Stacked plots of the mixed solutions are shown below.

Expansion of δ 4.0 – 3.1 is shown below.

Determination of K_{assoc}

A plot of $\delta - \delta_0$ versus guest concentration showed the rectangular hyperbolic shape expected for 1:1 complex formation (Figure 1, shown for those resonances with the most significant CIS). Non-linear regression analysis of this data using the method described by Gonzalez-Gaitano and Tardajos² gave an average K_{assoc} of $8.8 \pm 1.5 \text{ M}^{-1}$ over three trials. Non-linear regression was performed using the OriginLab 6.1 software package.³

Figure 1. Plot of $\delta - \delta_0$ versus guest concentration for H_f , H_g , and $H_{g'}$ (See Figure 3).

Disaccharide Cyclodimer 11 (Host) with 2,4-Hexadiyne-1,6-diol 12 (Guest):

Preparation of Solutions.

- (A) 24.0 mg cyclodimer **11** was dissolved into 3.00 mL D_2O to give a 9.9 mM clear solution.

- (B) 164 mg 2,4-Hexadiyne-1,6-diol was dissolved into 5.00 mL D₂O to give a 297 mM clear solution.
- (C) 262 mg 2,4-Hexadiyne-1,6-diol was dissolved into 3.00 mL D₂O to give a 792 mM clear solution.

Preparation of Guest and Host Mixtures.

The solutions were mixed according to the table below such that the total concentration of cyclodimer **11** was always 4.9 mM and the total volume was 0.7 mL.

Guest : Host Ratio	Sol'n A	Sol'n B	Sol'n C	D₂O
0 : 1	350 µL	0 µL	0 µL	350 µL
1 : 1	350 µL	11.5 µL	0 µL	338.5 µL
2 : 1	350 µL	23 µL	0 µL	327 µL
5 : 1	350 µL	58 µL	0 µL	292 µL
10 : 1	350 µL	116 µL	0 µL	234 µL
20 : 1	350 µL	234 µL	0 µL	116 µL
30 : 1	350 µL	350 µL	0 µL	0 µL
50 : 1	350 µL	0 µL	219 µL	131 µL
80 : 1	350 µL	0 µL	350 µL	0 µL

^1H NMR of Mixtures.

Stacked plots of the mixed solutions are shown below.

Expansion of δ 4.0 – 3.1 is shown below.

Determination of K_{assoc}

A plot of $\delta - \delta_0$ versus guest concentration showed the rectangular hyperbolic shape expected for 1:1 complex formation (Figure 2, shown for those resonances with the most significant CIS). Non-linear regression analysis of this data using the method described by Gonzalez-Gaitano and Tardajos² gave an average K_{assoc} of $10.4 \pm 1.1 \text{ M}^{-1}$ over three trials. Non-linear regression was performed using the OriginLab 6.1 software package.³

Figure 2. Plot of $\delta - \delta_0$ versus guest concentration for H_f , H_d , H_c , and H_l (See Figure 4).

NOESY1D Studies.

Monosaccharide Cyclotrimer 9 (Host) with 2,4-Hexadiyne-1,6-diol 12 (Guest):

A mixture of (10 : 1) **12** : **9** was degassed three times via freeze/pump/thaw in order to obtain the ^1H NMR and NOESY1D spectra shown in Figure 3. The singlet resulting from the methylene of **12** (highlighted in blue) was irradiated and NOE's were observed at various host resonances (highlighted in red).

Figure 3. Monosaccharide cyclotrimer **9** with excess 2,4-Hexadiyne-1,6-diol in D_2O .

bottom: ^1H NMR, top: NOESY1D

Disaccharide Cyclodimer 11 (Host) with 2,4-Hexadiyne-1,6-diol 12 (Guest):

A mixture of (10 : 1) **12** : **11** was degassed three times via freeze/pump/thaw in order to obtain the ^1H NMR and NOESY1D spectra shown in Figure 4. The singlet resulting from the methylene of **12** (highlighted in blue) was irradiated and NOE's were observed at various host resonances (highlighted in red).

Figure 4. Disaccharide cyclodimer **11** with excess 2,4-Hexadiyne-1,6-diol in D_2O .

bottom: ^1H NMR, top: NOESY1D

Modeling Images.

Each compound was entered into Molecular Operating Environment (MOE) and energy minimized using the MMFF94 forcefield to yield the structure images shown below.

α -Cyclodextrin: (Connolly surface included)

β -Cyclodextrin: (Connolly surface included)

Monosaccharide Cyclotrimer 11: (Connolly surface included)

Disaccharide Cyclodimer 9: (Connolly surface included)

Trisaccharide Cyclodimer 1: (Connolly surface included)

Scans of ^1H and ^{13}C NMR data for all new compounds follow below:

Applied Biosystems Voyager System 4066

Trimer 8

Voyager Spec #1=SM918P = 1553.4, 27269

+ Higher Oligomers

Mode of operation:	Linear
Extraction mode:	Delayed
Polarity:	Positive
Acquisition control:	Manual
Accelerating voltage:	20000 V
Grid voltage:	95%
Guide wire O:	0.1%
Extraction delay time:	125 nsec
Acquisition mass range:	100 - 10000 Da
Number of laser shots:	200/spectrum
Laser intensity:	1475
Laser Rep Rate:	3.0 Hz
Calibration type:	External - E:\Voyager02_2005\cal21405_AL_HABA_000
Calibration matrix:	HABA
Low mass gate:	500 Da
Digitizer start time:	9.98
Bin size:	2 nsec
Number of data points:	44309
Vertical scale:	1000 mV
Vertical offset:	0%
Input bandwidth:	200 MHz
Sample well:	55
Plate ID:	VOYAGER
Serial number:	4066
Instrument name:	Voyager-DE STR
Plate type filename:	C:\VOYAGER\100 well plate.plt
Lab name:	UI-UC MS Lab
Absolute x-position:	21417.9
Absolute y-position:	22240.6
Relative x-position:	-489.62
Relative y-position:	333.124
Shots in spectrum:	56
Source pressure:	1.246e-007
Mirror pressure:	1.345e-008
TCZ pressure:	0.02725
TIS gate width:	10
TIS flight length:	1131

Acquired: 10:30:00, February 14, 2005
KDB-III-0898, Bodine, HABA
E:\Voyager02_2005\17836_0004.dat

Printed: 10:31, February 14, 2005

10

References.

1. Bodine, K. D.; Gin, D. Y.; Gin, M. S. *J. Am. Chem. Soc.* **2004**, *126*, 1638.
2. Gonzalez-Gaitano, G.; Tardajos, G. *J. Chem. Ed.* **2004**, *81*, 270.
3. OriginLab Corporation *Origin* version 6.1; OriginLab Corporation: Northhampton, MA, **2000**.