**Highly Diastereoselective Thioglycosylation of Functionalized Peracetylated Glycosides
Catalyzed by MoO₂Cl₂**

Shiue-Shien Weng, Yow-Dzer Lin, and Chien-Tien Chen*

Department of Chemistry, National Taiwan Normal University, Taipei, Taiwan

#88, Sec. 4, Ding-jou Road, Taipei, Taiwan 11650

e-mail: chefv043@ntnu.edu.tw

Representative experimental procedures, spectral data (51 pages)

SUPPORTING INFORMATION

Table of contents

General experimental methods-----	S2
Materials and preparation of known starting materials by literature methods -----	S2-S3
Representative procedure for the thioglycosylation-----	S3
Effects of oxometallic species on catalytic thioglycosylation of penta- <i>O</i> -acetyl- β -D-glucose with thiophenol-----	S4
Analytical data for <i>S</i> -substituted-1-thio- β -D-glucopyranoside 2a-2n -----	S5-S10
Analytical data for phenyl-1-thio- β -glycosides 3-11 -----	S10-S13
Synthesis and analytical data for <i>p</i> -tolyl-1-thio- β -D-glucopyranoside- 12' -----	S14
Synthesis and analytical data for 4-methylphenyl-4,6- <i>O</i> -benzylidene-1-thio- β -D-glucopyranoside- 12 -----	S15
Preparation of thioglycosidic disaccharide- 15 -----	S15-18
References-----	S18-S19
¹ H and ¹³ C NMR spectra for <i>S</i> -substituted-1-thio- β -D-glucopyranosides 2a-2n -----	S20-S33
¹ H and ¹³ C NMR spectra for phenyl-1-thio- β -glycosides 3-14 -----	S34-S46
¹ H, ¹³ C, COSY, DEPT-90, DEPT-135, HMQC, and HMBC NMR spectra for thioglycosidic disaccharide- 15 -----	S47-S51

General experimental methods. ^1H -NMR and ^{13}C -NMR were recorded on Varian Gemini-2000 (400 MHz ^1H , 100 MHz ^{13}C) spectrometers in deuteriochloroform with tetramethylsilane (TMS) or chloroform as an internal reference unless otherwise stated. Chemical shifts are reported in ppm (δ), coupling constants, J , are reported in Hz. Mass spectra were recorded on a Finnigan TCQ-700 spectrometer with an ionization voltage of 70 eV unless otherwise stated. Electrospray ionization (ESI) mass spectra were recorded on a Finnigan MAT-95S spectrometer. Data are reported in the form m/e (intensity relative to base = 100%). Elemental analysis data were obtained on a Heraeus CHN-OS Rapid analyzer. Optical rotations were recorded on a JASCO DIP-1000 digital polarimeter and are reported as follows: (c = g/100ml, solvent). Analytical TLC was performed on Merck silica gel plates with QF-254 indicator. Visualization was accomplished with UV light or with phosphomolybdic acid (PMA) and KMnO_4 staining agents. Column (flash) chromatography was performed by using 32-63 μm silica gel. Solvents for extraction and chromatography were reagent grade. Optical rotations were recorded on a JASCO DIP-1000 digital polarimeter and are reported as follows: $[\alpha]_{\text{D}}^{\text{T}}$ (c = g/100mL, solvent). CH_2Cl_2 , $\text{Cl}(\text{CH}_2)_2\text{Cl}$, CHCl_3 , CCl_4 , and N,N -dimethylformamide (DMF) were dried over CaH_2 under nitrogen atmosphere and distilled before use. THF, Et_2O , 1,4-dioxane, and toluene were dried over benzophenone-ketyl intermediate under nitrogen atmosphere and distilled before use.. CH_3CN was dried over Mg turnings under argon atmosphere and distilled before use. All reactions were performed with oven-dried (120 $^\circ\text{C}$) or flame-dried glassware under an atmosphere of dry argon.

Materials and preparation of known starting materials by literature methods

D-Glucose, D-galactose, D-mannose, D-xylose, D-lactose, D-maltose, L-arabinose, and 2,3,5-tri-*O*-acetyl- β -D-ribofuranoside were purchased from ACROS. Methyltrioxorhenium(VII) (CH_3ReO_3), chromyl(VI) chloride (CrO_2Cl_2), bismuth(III) oxychloride (BiOCl), molybdenum(VI) dioxodichlororide (MoO_2Cl_2), molybdenum acetylacetonate ($\text{MoO}_2(\text{acac})_2$), oxomolybdenum (VI) tetrachloride (MoOCl_4), molybdenum(V) chloride (MoCl_5), hafnium(IV) oxychloride (HfOCl_2), and

vanadyl(IV) acetylacetonate ($\text{VO}(\text{acac})_2$) were purchased from Aldrich. The VOX_2 series of compounds (brand name as *Clip-all*® series, US patent # 6,541,659 B1, 2003) and TiOX_2 are now available directly from the institution (e-mail: chefv043@ntnu.edu.tw). Thiophenol, 4-methylthiophenol, 4-methoxythiophenol, 2-methoxythiophenol, 1,3-propanedithiol, ethanethiol, cyclohexyl mercaptan, *tert*-butylmercaptan, benzylmercaptan, and 2-mercaptoethanol were purchased from ACROS. They are freshly distilled from CaSO_4 before use. 11-mercaptoundecanoic acid (re-crystallized from ether), 4-methyl-thiophenol, 2-naphthalenthio, and 4-chlorothiophenol (re-crystallized from benzene) were purchased from ACROS. Penta-*O*-acetyl- β -D-glucose,¹ penta-*O*-acetyl- β -D-galactose,² penta-*O*-acetyl- β -D-mannose,³ tetra-*O*-acetyl- β -D-xylose,⁴ tetra-*O*-acetyl- α -L-arabinose,⁵ hepta-*O*-acetyl- β -D-lactose, hepta-*O*-acetyl- β -D-maltose,⁶ 2-(*tert*-butyldimethylsiloxy)ethylmercaptan,⁷ and methyl 11-mercaptoundecanoate⁸ were prepared by following literature methods.

Representative Procedure for the catalytic thioglycosylation

In a 25-mL, two-necked, round-bottomed flask equipped with an addition funnel was placed MoO_2Cl_2 (5.9 mg, 0.03 mmol, 3 mol%) followed by addition of anhydrous CH_2Cl_2 (1 mL). To this solution was slowly added a solution of a per-acetylated sugar (1 mmol) in CH_2Cl_2 (3 mL) at ambient temperature via the addition funnel. After 10 min, a solution of thiol (1.3-1.5 equiv or 3.0 equiv for *t*-butanethiol) in CH_2Cl_2 (1 mL) was slowly added to the reaction mixture. The reaction solution immediately turned deep blue or green and gradually changed to yellowish brown in 10 min. After completion of the reaction as evidenced by the solution color (deep brown) and by TLC analysis, the reaction mixture was quenched with cold saturated aqueous NaHCO_3 (5 mL). The organic layer was separated and washed with brine, dried (MgSO_4), filtered, and evaporated. The crude residue was purified by column chromatography on silica gel. The product was characterized by routine spectroscopic methods.

Effect of oxometallic species on catalytic thioglycosylation of penta-*O*-acetyl- β -D-glucose with thiophenol.

entry ^a	MO _m X _n	time (h)	yield, ^a %
1	VO(OTf) ₂	24	32 (67 ^c)
2	VO(OTs) ₂ ^b	24	16
3	VOCl ₂	24	6
4	VO(acac) ₂	24	7
5	BiOCl	24	none
6	CrO ₂ Cl ₂	24	none
7	WO ₂ Cl ₂	24	none
8	CH ₃ ReO ₃	24	7
9	MoO ₂ (acac) ₂	24	trace
10	MoO ₂ Cl ₂	14	88 (93 ^c)
11	MoOCl ₄	14	58%
12	TiO(OTf) ₂	24	none
13	ZrO(OTf) ₂	24	none
14	HfOCl ₂	24	none
16	MoCl ₅	24	none
17	Cu(OTf) ₂	24	none
18	(n-Bu) ₂ SnO	24	none

^a Isolated yields and no α -anomer was observed. ^bVanadyl *p*-toluenesulfonate. ^c

Three mol % of catalyst was used.

Analytical data for *S*-substituted -1-thio- β -D-glucopyranosides

Phenyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside **2a**⁹

¹H NMR (400 MHz, CDCl₃) δ 7.51-7.50 (m, 2 H), 7.32-7.26 (m, 3 H), 5.23 (t, J = 9.2, 1H), 5.04 (t, J = 9.6, 1H), 4.97 (t, J = 9.6, 1H), 4.70 (d, J = 10.4, 1H), 4.21-4.19 (m, 2H), 3.72 (m, 1H), 2.09 (s, 3H, COCH₃), 2.07 (s, 3H, COCH₃), 2.01 (s, 3H, COCH₃), 1.98 (s, 3H, COCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.51, 170.12, 169.34, 169.2, 133.10, 131.61, 128.9, 128.4, 85.7, 75.8, 73.9, 69.9, 68.2, 62.1, 20.70, 20.7, 20.6, 20.5; M.W. (C₂₀H₂₄O₉S) 440.46; ESI-MS m/z 463 (M+Na⁺, 71); mp 118-120 °C; $[\alpha]_D^{23}$ -22.7 (c 1.0, CHCl₃); TLC R_f 0.35 (Ethyl acetate/hexane, 1/ 2).

p-Tolyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside **2b**¹⁰

¹H NMR (400 MHz, CDCl₃) δ 7.39 (d, J = 8.1, 2 H), 7.39 (d, J = 7.9, 2 H), 5.21 (t, J = 9.4, 1H), 5.02 (t, J = 9.7, 1H), 4.94 (t, J = 10.1, 1H), 4.63 (d, J = 10.0, 1H), 4.21-4.18 (m, 2H), 3.70 (ddd, J = 10.1, 4.7, 2.7, 1H), 2.35 (s, 3H), 2.09 (s, 3H, COCH₃), 2.08 (s, 3H, COCH₃), 2.01 (s, 3H, COCH₃), 1.98 (s, 3H, COCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.6, 170.2, 169.4, 169.2, 138.8, 129.7, 127.6, 85.8, 75.8, 74.0, 69.9, 68.2, 62.1, 21.2, 20.7, 20.7, 20.6, 20.6; M.W. (C₂₁H₂₆O₉S) 454.49; ESI-MS m/z 477 (M+Na⁺, 27), 478 (12); mp 114-116 °C; $[\alpha]_D^{24}$ -18.2 (c 1.2, CHCl₃); TLC R_f 0.32 (Ethyl acetate/hexane, 1/ 2).

4-Chlorophenyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside **2c**¹¹

¹H NMR (400 MHz, CDCl₃) δ 7.44 (d, J = 6.4, 2 H), 7.27 (d, J = 6.4, 2 H), 5.21 (t, J = 9.2, 1H), 5.01 (t, J = 9.6, 1H), 4.93 (t, J = 10.0, 1H), 4.64 (d, J = 10.0, 1H), 4.25-4.11 (m, 2H), 3.71 (ddd, J = 10.0, 4.8, 2.8, 1H), 2.07 (s, 3H, COCH₃), 2.06 (s, 3H, COCH₃), 2.00 (s, 3H, COCH₃), 1.97 (s, 3H, COCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.4, 170.1, 169.3, 169.1, 134.9, 129.4, 85.2, 75.8, 73.8, 69.8, 68.1,

62.0, 20.7, 20.6; M.W. ($C_{20}H_{23}ClO_9S$) 474.19; ESI-MS m/z 497 ($M+Na^+$, 100), 498 (67); mp 112-113 °C; $[\alpha]_D^{26}$ -26.4 (c 2.0, $CHCl_3$); TLC R_f 0.36 (Ethyl acetate/hexane, 1/ 2).

4-Methoxyphenyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside 2d¹²

¹H NMR (400 MHz, $CDCl_3$) δ 7.49 (d, J = 10.8, 2 H), 6.84 (d, J = 10.8, 2 H), 5.17 (t, J = 9.2, 1H), 4.97 (t, J = 10.0, 1H), 4.87 (t, J = 9.6, 1H), 4.53 (d, J = 10.0, 1H), 4.19 (dd, J = 12.0, 4.4, 1H), 4.15 (dd, J = 12.3, 2.2, 1H), 3.80 (s, 3H, OCH_3), 3.66 (ddd, J = 7.2, 4.0, 3.2, 1H), 2.08 (s, 3H, $COCH_3$), 2.05 (s, 3H, $COCH_3$), 1.99 (s, 3H, $COCH_3$), 1.96 (s, 3H, $COCH_3$); ¹³C NMR (100 MHz, $CDCl_3$) δ 170.5, 170.1, 169.3, 169.2, 160.4, 136.5, 120.8, 114.4, 85.6, 75.7, 74.0, 69.9, 68.2, 62.0, 55.3, 20.7, 20.7, 20.5, 20.5; M.W. ($C_{21}H_{26}O_{10}S$) 470.49; ESI-MS m/z 493 ($M+Na^+$, 30), 494 (18); mp 92-93 °C; $[\alpha]_D^{26}$ -24.3 (c 2.0, $CHCl_3$); TLC R_f 0.31 (Ethyl acetate/hexane, 1/ 2).

2-Naphthyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside 2e¹³

¹H NMR (400 MHz, $CDCl_3$) δ 7.99 (s, 1H), 7.83-7.77 (m, 3H), 7.57-7.46 (m, 3H), 5.24 (t, J = 9.7, 1 H), 5.07-4.99 (m, 2H), 4.78 (d, J = 10.0, 1H), 4.23 (dd, J = 12.3, 5.1, 1H), 4.18 (dd, J = 12.3, 2.4, 1H), 3.72 (ddd, J = 7.4, 4.9, 2.4, 1H), 2.11 (s, 3H, $COCH_3$), 2.02 (s, 3H, $COCH_3$), 2.00 (s, 3H, $COCH_3$), 1.98 (s, 3H, $COCH_3$); ¹³C NMR (100 MHz, $CDCl_3$) δ 170.5, 170.1, 169.3, 169.2, 133.4, 132.9, 132.9, 132.7, 130.2, 128.7, 128.4, 127.7, 126.7, 126.6, 85.8, 75.9, 73.9, 70.0, 68.2, 62.1, 20.7, 20.6, 20.52, 20.50; M.W. ($C_{24}H_{26}O_9S$) 490.52; ESI-MS m/z 513 ($M+Na^+$, 53); mp 108-109 °C; $[\alpha]_D^{24}$ -16.2 (c 1.0, $CHCl_3$); TLC R_f 0.36 (Ethyl acetate/hexane, 1/ 2).

2-Methoxyphenyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside 2f

¹H NMR (400 MHz, $CDCl_3$) δ 7.47 (d, J = 8.8, 1 H), 7.29 (m, 1 H), 6.91-6.86 (m, 2 H), 5.22 (t, J = 9.4, 1 H), 5.05 (t, J = 9.8, 1H), 4.99

(t, $J = 9.2$, 1H), 4.79 (d, $J = 10.0$, 1H), 4.19 (dd, $J = 12.2$, 5.2, 1H), 4.15 (dd, $J = 12.2$, 2.4, 1H), 3.84 (s, 3H, OCH₃), 3.75 (ddd, $J = 7.0$, 3.8, 2.8, 1H), 2.05 (s, 3H, COCH₃), 2.04 (s, 3H, COCH₃), 2.01 (s, 3H, COCH₃), 1.98 (s, 3H, COCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.5, 170.2, 169.4, 169.3, 158.5, 133.6, 129.6, 121.0, 120.0, 111.0, 84.3, 75.7, 74.1, 70.1, 68.3, 62.2, 55.7, 20.7, 20.6, 20.5; M.W. (C₂₁H₂₆O₁₀S) 470.49; ESI-MS m/z 493 (M+Na⁺, 15); mp 88-90 °C; $[\alpha]_D^{26}$ -19.6 (c 2.0, CHCl₃); TLC R_f 0.33 (Ethyl acetate/hexane, 1/ 2); Anal. calcd for C₂₁H₂₆O₁₀S: C, 53.61; H, 5.57. Found: C, 53.67; H, 5.34.

***o*-Tolyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside 2g¹⁴**

¹H NMR (400 MHz, CDCl₃) δ 7.49 (d, $J = 7.7$, 1 H), 7.21-7.13 (m, 3 H), 5.20 (t, $J = 12.0$, 1H), 5.08-5.01 (m, 2H), 4.66 (d, $J = 10.2$, 1H), 4.21 (dd, $J = 12.3$, 5.6, 1H), 4.12 (dd, $J = 12.3$, 2.2, 1H), 3.67 (m, 2H), 2.38 (s, 3H, CH₃), 2.07 (s, 3H, COCH₃), 2.05 (s, 3H, COCH₃), 2.00 (s, 3H, COCH₃), 1.98 (s, 3H, COCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.4, 170.1, 169.3, 169.2, 140.4, 131.8, 130.3, 128.4, 126.5, 86.3, 75.7, 73.9, 70.1, 68.3, 62.2, 20.8, 20.61, 20.59, 20.49, 20.47; M.W. (C₂₁H₂₆O₉S) 454.49; ESI MS m/z 477 (M+Na⁺, 73); mp 114-116 °C; $[\alpha]_D^{26}$ -10.2 (c 1.0, CHCl₃); TLC R_f 0.33 (Ethyl acetate/hexane, 1/ 2).

Ethyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside 2h¹⁵

¹H NMR (400 MHz, CDCl₃) δ 5.20 (t, $J = 9.2$, 1H), 5.06 (t, $J = 10.0$, 1H), 5.01 (t, $J = 10.0$, 1H), 4.48 (d, $J = 10.0$, 1H), 4.24 (dd, $J = 12.0$, 3.2, 1H), 4.11 (dd, $J = 12.4$, 2.0, 1H), 3.69 (ddd, $J = 10.0$, 4.8, 2.4, 1H), 2.74-2.63 (m, 2H), 2.05 (s, 3H, COCH₃), 2.03 (s, 3H, COCH₃), 1.99 (s, 3H, COCH₃), 1.97 (s, 3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.6, 170.1, 169.3, 169.3, 83.5, 75.8, 73.9, 69.8, 68.3, 62.1, 24.1, 20.7, 20.53, 20.51, 14.8; M.W. (C₁₆H₂₄O₉S) 392.42; ESI-MS m/z 415 (M+Na⁺, 93); mp 82-83 °C; $[\alpha]_D^{26}$ -25.6 (c 1.0, CHCl₃); TLC R_f 0.36 (Ethyl acetate/hexane, 1/ 2).

Benzyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside 2i¹⁶

¹H NMR (400 MHz, CDCl₃) δ 7.33-7.27 (m, 5 H), 5.15-5.03 (m, 3H), 4.28 (d, J = 9.7, 1H), 4.25 (dd, J = 12.4, 5.12, 1H), 4.13 (dd, J = 12.4, 2.28, 1H), 3.94 (d, J = 12.9, 1H), 3.83 (d, J = 12.9, 1H), 3.59 (ddd, J = 9.7, 5.0, 2.3, 1H), 2.11 (s, 3H, COCH₃), 2.06 (s, 6H, 2 $\times$ COCH₃), 1.99 (s, 3H, COCH₃); ¹³C NMR (100 MHz, CDCl₃) δ 170.6, 170.1, 169.4, 169.4, 136.8, 129.0, 128.6, 127.4, 81.9, 75.8, 73.8, 69.6, 62.2, 33.7, 20.7, 20.6, 20.54, 20.52; M.W. (C₂₁H₂₆O₉S) 454.49; ESI-MS m/z 477 (M+Na⁺, 32), 478 (18); mp 98-99 °C; $[\alpha]_D^{26}$ -88.2 (c 1.0, CHCl₃); TLC R_f 0.38 (Ethyl acetate/hexane, 1/ 2).

Cyclohexyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside 2j¹⁷

¹H NMR (400 MHz, CDCl₃) δ 5.21 (t, J = 9.4, 1H), 5.06 (t, J = 9.9, 1H), 4.98 (t, J = 10.1, 1H), 4.59 (d, J = 10.1, 1H), 4.21 (dd, J = 12.3, 5.3, 1H), 4.12 (dd, J = 12.4, 2.5, 1H), 3.69 (ddd, J = 10.0, 5.4, 2.5, 1H), 2.89 (m, 2H), 2.06 (s, 3H, COCH₃), 2.03 (s, 3H, COCH₃), 2.02 (s, 3H, COCH₃), 1.99 (s, 3H, COCH₃), 1.98 (m, 2H), 1.75 (m, 2H), 1.43-1.21 (m, 6H); ¹³C NMR (CDCl₃, 100 MHz) δ 170.6, 170.2, 169.4, 169.3, 83.2, 75.7, 74.0, 70.3, 68.5, 62.3, 43.9, 34.2, 33.9, 26.0, 25.9, 25.6, 20.72, 20.67, 20.58, 20.56; M.W. (C₂₀H₃₀O₉S) 446.51; ESI-MS m/z 469 (M+Na⁺, 27); mp 120-122 °C; $[\alpha]_D^{27}$ -38.2 (c 1.0, CHCl₃); TLC R_f 0.40 (Ethyl acetate/hexane, 1/ 2).

***tert*-Butyl-2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside 2k¹⁸**

¹H NMR (CDCl₃, 400 MHz) δ 5.25 (t, J = 9.3, 1H), 5.03 (t, J = 9.7, 1H), 4.94 (t, J = 9.8, 1H), 4.63 (d, J = 10.0, 1H), 4.19 (dd, J = 12.2, 6.0, 1H), 4.10 (dd, J = 12.2, 2.2, 1H), 3.71 (ddd, J = 6.4, 6.0, 2.3, 1H), 2.04 (s, 3H, COCH₃), 2.03 (s, 3H, COCH₃), 2.02 (s, 3H, COCH₃), 2.00 (s, 3H, COCH₃), 1.36 (s, 9H, C(CH₃)₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.6, 170.2, 169.4, 169.3, 82.3, 75.6, 74.0, 70.2, 68.6, 62.6, 44.3, 31.4, 20.7, 20.66, 20.58, 20.56; M.W. (C₁₈H₂₈O₉S) 420.47; ESI-MS m/z 443 (M+Na⁺,

13); mp 140-142 °C; $[\alpha]_D^{27}$ -6.8 (*c* 0.65, CHCl₃); TLC *R_f* 0.23 (Ethyl acetate/hexane, 1/ 2).

3'-Mercapto-propyl 2, 3, 4, 6-tetra-*O*-acetyl-1-thio-β-D-glucopyranoside 2l¹⁹

¹H NMR (CDCl₃, 400 MHz) δ 5.20 (t, *J* = 9.2, 1H), 5.08 (t, *J* = 11.6, 1H), 4.99 (t, *J* = 10.0, 1H), 4.48 (d, *J* = 10.0, 1H), 4.23 (dd, *J* = 12.4, 4.8, 1H), 4.13 (dd, *J* = 12.0, 2.0, 1H), 3.76- 3.72 (m, 1H), 2.81 (dt, *J* = 13.0, 6.8, 1H), 2.71 (dt, *J* = 12.8, 6.8 1H), 2.61 (td, *J* = 14.8, 7.6, 2H), 2.07 (s, 3H, COCH₃), 2.05 (s, 3H, COCH₃), 2.01 (s, 3H, COCH₃), 1.97 (s, 3H, COCH₃), 2.24-1.83 (m, 4H), 1.89 (t, *J* = 7.2, 2H), 1.34 (t, *J* = 7.8, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ 170.5, 170.1, 169.33, 169.32, 83.6, 75.9, 73.8, 69.8, 68.3, 62.1, 33.4, 28.2, 23.1, 20.69, 20.64, 20.53, 20.51; M.W. (C₁₇H₂₆O₉S₂) 438.51; MS (EI, 70 eV) 438 (M⁺, 39), 422 (18), 421 (100), 409 (18); $[\alpha]_D^{28}$ -18.7 (*c* 1.0, CHCl₃); TLC *R_f* 0.42 (Ethyl acetate/hexane, 1/ 2).

11-Methoxycarbonyl undecanyl- 2, 3, 4, 6-tetra-*O*-acetyl 1-1-thio-β-D-glucopyranoside 2m²⁰

¹H NMR (CDCl₃, 400 MHz) δ .21 (t, *J* = 9.2, 1H), 5.07 (t, *J* = 10.0, 1H), 5.02 (dd, *J* = 11.6, 11.2, 1H), 4.47 (d, *J* = 10.4, 1H), 4.23 (dd, *J* = 12.4, 4.8, 1H), 4.12 (dd, *J* = 12.4, 2.4, 1H), 3.70 (ddd, *J* = 9.2, 4.8, 2.4, 1H), 3.65 (s, 3H, OCH₃), 2.68-2.61 (m, 2H), 2.29 (t, *J* = 7.6, 1H), 2.07 (s, 3H, COCH₃), 2.05 (s, 3H, COCH₃), 2.01 (s, 3H, COCH₃), 1.99 (s, 3H, COCH₃), 1.65-1.52 (m, 5H), 1.36-1.26 (m, 13H); ¹³C NMR (CDCl₃, 100 MHz) δ 174.3, 170.6, 170.2, 169.4, 169.3, 83.6, 75.8, 73.9, 69.9, 68.3, 62.2, 51.4, 34.1, 30.0, 29.6, 29.4, 29.3, 29.2, 29.1, 28.7, 24.9, 20.7, 20.6, 20.5; M.W. (C₂₆H₄₂O₁₁S) 562.67; ESI-MS *m/z* 585 (M+Na⁺, 14); mp 54-56 °C; $[\alpha]_D^{27}$ -42.2 (*c* 1.0, CHCl₃); TLC *R_f* 0.38 (Ethyl acetate/hexane, 1/ 3).

2'-*tert*-Butyl dimethyl silyl 2, 3, 4, 6-tetra-*O*-acetyl-1-thio-β-D-glucopyranoside 2n

¹H NMR (CDCl₃, 400 MHz) δ 5.20 (t, *J* = 9.6, 1H), 5.08 (t,

$J = 9.6$, 1H), 5.01 (t, $J = 10.0$, 1H), 4.58 (d, $J = 10.0$, 1H), 4.24 (dd, $J = 12.4$, 4.8, 1H), 4.13 (dd, $J = 12.4$, 2.4, 1H), 3.85-3.73 (m, 2H), 3.69 (ddd, $J = 7.2$, 4.8, 2.4, 1H), 2.88 (dt, $J = 14.0$, 7.6, 1H), 2.71 (dt, $J = 13.6$, 6.0, 1H), 2.08 (s, 3H, COCH₃), 2.05 (s, 3H, COCH₃), 2.02 (s, 3H, COCH₃), 1.99 (s, 3H, COCH₃), 0.90 (s, 9H, C(CH₃)₃), 0.07 (s, 6H, 2×CH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.6, 170.1, 169.4, 169.3, 83.6, 75.9, 73.8, 70.0, 68.3, 63.3, 62.1, 32.3, 25.8, 20.68, 20.65, 20.55, 20.53, 18.2, -5.31, -5.32; M.W. (C₂₂H₃₈O₁₀SSi) 522.68; ESI-MS m/z 545 (M+Na⁺, 45); colorless oil; $[\alpha]_D^{23}$ -18.2 (c 1.0, CHCl₃); TLC R_f 0.34 (Ethyl acetate/hexane, 1/ 3); Anal. calcd for C₂₂H₃₈O₁₀SSi: C, 50.55; H, 7.33. Found: C, 50.73; H, 7.08.

Analytic data for *S*-Phenyl-1-thio- β -glycosides

Phenyl-3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside **3**²¹

¹H NMR (CDCl₃, 400 MHz) δ 7.87 (dd, $J = 5.6$, 3.2, 2 H), 7.75 (dd, $J = 5.6$, 2.8, 2H), 7.41 (dd, $J = 7.2$, 1.6, 2H), 7.30-7.26 (m, 3H), 5.79 (dd, $J = 10.2$, 9.2, 1H), 5.71 (d, $J = 10.0$, 1H), 5.13 (dd, $J = 10.1$, 9.2, 1H), 4.35 (t, $J = 10.4$, 1H), 4.29 (ddd, $J = 12.3$, 5.0, 2.4, 1H), 4.21 (ddd, $J = 12.3$, 5.0, 2.2, 1H), 3.91 (ddd, $J = 10.1$, 5.2, 2.6, 1H), 2.10 (s, 3H, COCH₃), 2.02 (s, 3H, COCH₃), 1.84 (s, 3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.6, 170.1, 169.4, 167.8, 166.9, 134.4, 134.3, 133.3, 131.6, 131.2, 131.0, 128.9, 128.4, 123.7, 83.1, 75.9, 71.6, 68.8, 62.2, 53.6, 20.7, 20.6, 20.4; M.W. (C₂₆H₂₅NO₉S) 527.54; ESI-MS m/z 550 (17), 551 (M+Na⁺, 21); mp 142-143 °C; $[\alpha]_D^{26}$ +56.6 (c 1.0, CHCl₃); TLC R_f 0.23 (Ethyl acetate/hexane, 1/ 1).

Phenyl 2, 3, 4, 6-tetra-*O*-acetyl-1-thio- β -D-galactopyranoside **4**²²

¹H NMR (CDCl₃, 400 MHz) δ 7.47-7.45 (m, 2 H), 7.26-7.24 (m, 3 H), 5.36 (d, $J = 3.2$, 1H), 5.18 (t, $J = 10.0$, 1H), 5.01 (dd, $J = 10.0$, 3.2, 1H), 4.69 (d, $J = 10.0$, 1H), 4.13 (dd, $J = 11.6$, 7.2, 1H), 4.06 (dd, $J = 11.6$, 6.0, 1H), 3.91 (t, $J = 6.6$, 1H), 2.05 (s, 3H, COCH₃), 2.03 (s, 3H, COCH₃), 1.97 (s, 3H, COCH₃), 1.91 (s,

3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.0, 169.8, 169.6, 169.1, 132.2, 128.6, 127.8, 86.1, 74.1, 71.7, 67.0, 61.4, 20.5, 20.32, 20.27, 20.1; M.W. (C₂₀H₂₄O₉S) 440.46; ESI-MS *m/z* 463 (M+Na⁺, 100), 464 (73); mp 65-67 °C; [α]_D²⁶ +4.2 (*c* 1.0, CHCl₃); TLC R_f 0.33 (Ethyl acetate/hexane, 1/ 2).

Phenyl 2, 3, 4, 6-tetra-*O*-acetyl-1-thio-α-D-mannopyranoside 5²¹

¹H NMR (CDCl₃, 400 MHz) δ 7.50-7.48 (m, 2 H), 7.33-7.29 (m, 3 H), 5.53-5.49 (m, 2H), 5.36-5.29 (m, 2H), 4.56-4.53 (m, 1H), 4.30 (dd, *J* = 12.4, 6.0, 1H), 4.10 (dd, *J* = 12.4, 2.4, 1H), 2.15 (s, 3H, COCH₃), 2.08 (s, 3H, COCH₃), 2.05 (s, 3H, COCH₃), 2.02 (s, 3H, COCH₃); ¹³C NMR

(CDCl₃, 100 MHz) δ 170.5, 169.9, 169.8, 169.1, 132.6, 132.0, 129.2, 128.1, 85.7, 70.9, 69.5, 69.4, 66.4, 62.4, 20.9, 20.69, 20.67, 20.6; M.W. (C₂₀H₂₄O₉S) 440.46; ESI-MS *m/z* 463 (M+Na⁺, 79), 464 (57); [α]_D²⁴ +85.6 (*c* 2.0, CH₂Cl₂); TLC R_f 0.27 (Ethyl acetate/hexane, 1/ 3).

Phenyl 2, 3, 5-tetra-*O*-acetyl-β-D-ribofuranoside 6²³

¹H NMR (CDCl₃, 400 MHz) δ 7.53-7.51 (m, 2 H), 7.34-7.32 (m, 3 H), 5.31 (d, *J* = 4.8, 1H), 5.26-5.19 (m, 2H), 4.27-4.25 (m, 2H), 4.09 (dd, *J* = 12.8, 5.2, 1H), 2.09 (s, 3H, COCH₃), 2.07 (s, 3H, COCH₃), 2.06 (s,

3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.5, 169.6, 169.4, 133.4, 131.7, 129.0, 128.4, 87.9, 80.1, 73.9, 71.4, 63.4, 20.8, 20.5; M.W. (C₁₇H₂₀O₇S) 368.40; ESI-MS *m/z* 391 (M+Na⁺, 53), 392 (42); [α]_D²⁷ +12.3 (*c* 1.0, CHCl₃); TLC R_f 0.31 (Ethyl acetate/hexane, 1/ 3).

phenyl 2, 3, 4-tri-*O*-acetyl-1-thio-β-D-xylopyranoside 7²⁴

¹H NMR (CDCl₃, 400 MHz) δ 7.50-7.47 (m, 2 H), 7.33-7.31 (m, 3 H), 5.19 (t, *J* = 8.2, 1H), 4.90 (t, *J* = 8.2, 1H), 4.92-4.89 (m, 1H). 4.80 (dd,

$J = 8.3$, 1H), 4.28 (dd, $J = 11.8$, 4.9, 1H), 3.42 (dd, $J = 12.0$, 8.8, 1H), 2.11 (s, 3H, COCH₃), 2.04 (s, 6H, 2 × COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 169.9, 169.8, 169.3, 132.7, 132.2, 129.0, 128.2, 86.2, 72.0, 69.8, 68.4, 65.2, 20.8, 20.7; M.W. (C₁₇H₂₀O₇S) 368.40; ESI-MS m/z 391 (M+Na⁺, 50), 392 (41); mp 76-77 °C; $[\alpha]_D^{26}$ -56.6 (*c* 1.0, CHCl₃); TLC R_f 0.26 (Ethyl acetate/hexane, 1/ 3).

Phenyl 2, 3, 4-tri-*O*-acetyl-1-thio-β-L-arabinopyranoside 8²⁵

¹H NMR (CDCl₃, 400 MHz) δ 7.49-7.46 (m, 2 H) 7.28-7.27 (m, 3 H), 5.26-5.20 (m, 1H), 5.23 (t, $J = 8.2$, 1H), 5.09 (dd, $J = 8.4$, 3.4, 1H), 4.80 (d, $J = 7.8$, 1H), 4.12 (dd, $J = 12.7$, 4.2, 1H), 3.65 (dd, $J = 12.7$, 2.0, 1H), 2.06 (s, 6H, 2 × COCH₃), 2.02 (s, 3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 169.9, 169.7, 169.2, 133.1, 132.1, 128.8, 127.7, 86.5, 70.3, 68.3, 67.4, 65.1, 20.63, 20.61, 20.5; M.W. (C₁₇H₂₀O₇S) 368.40; ESI-MS m/z 391 (M+Na⁺, 50), 392 (41); $[\alpha]_D^{26}$ -58.6 (*c* 1.0, CH₂Cl₂); TLC R_f 0.28 (Ethyl acetate/hexane, 1/ 3).

Phenyl 4, 6-di-*O*-acetyl-2, 3-dideoxy-1-thio-D-erythro-hex-2-enopyranoside 9²⁶

¹H NMR (CDCl₃, 400 MHz) δ 7.55-7.53 (m, 2 H), 7.33-7.27 (m, 3 H), 6.06 (dd, $J = 10.0$, 2.8, 1H), 5.23 (d, $J = 10.0$, 1H), 5.76 (bs, 1H), 5.37 (dd, $J = 9.2$, 1.2, 1H), 4.47 (dt, $J = 6.0$, 2.4, 1H), 4.12 (dd, $J = 12.2$, 6.0, 1H), 4.20 (dd, $J = 12.0$, 2.0, 1H), 2.06 (major, s, 3H, COCH₃), 2.07 (major, s, 3H, COCH₃), 2.00 (minor, s, 3H, COCH₃), 1.94 (minor, s, 3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.7, 170.2, 134.7, 131.7, 128.9, 128.5, 127.8, 127.6, 83.0, 67.2, 65.1, 63.0, 20.9, 20.7; M.W. (C₁₆H₁₈O₃S) 322.37; ESI-MS m/z 345 (M+Na⁺, 100), 346 (45); mp 64-65 °C; $[\alpha]_D^{26}$ +24.4 (*c* 5.0, CHCl₃); TLC R_f 0.34 (Ethyl acetate/hexane, 1/ 3).

Phenyl 2, 3, 6-tri-*O*-acetyl-4-*O*-(2', 3', 4', 6'-tetra-*O*-acetyl-α-D-glucopyranosyl)-1-thio-β-D-glucopyranoside 10²⁷

¹H NMR (CDCl₃, 400 MHz) δ 7.48-7.46 (m, 2H), 7.34-7.27 (m, 3H), 5.37 (d, $J = 4.0$, 1H), 5.33 (dd,

$J = 10.3, 9.7, 1\text{H}$), 5.27 (t, $J = 8.9, 1\text{H}$), 5.02 (dd, $J = 10.0, 9.7, 1\text{H}$), 4.89-4.79 (m, 2H), 4.52 (d, $J = 10.0, 1\text{H}$), 4.27-4.21 (m, 2H), 4.16 (dd, $J = 12.4, 7.7, 1\text{H}$), 4.03 (dd, $J = 12.4, 2.1, 1\text{H}$), 3.95-3.87 (m, 2H), 3.70 (ddd, $J = 7.5, 4.8, 2.6, 1\text{H}$), 2.11 (s, 3H, COCH₃), 2.09 (s, 3H, COCH₃), 2.05 (s, 3H, COCH₃), 2.02 (s, 3H, COCH₃), 2.01 (s, 3H, COCH₃), 1.98 (s, 3H, COCH₃), 1.97 (s, 3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.49, 170.47, 170.3, 170.1, 169.9, 169.5, 169.4, 133.4, 131.3, 128.9, 128.4, 95.6, 85.1, 76.5, 76.1, 72.6, 70.7, 70.0, 69.3, 68.6, 68.1, 62.9, 61.6, 20.9, 20.8, 20.7, 20.6, 20.5; M.W. (C₃₂H₄₀O₁₇S) 728.72; ESI-MS m/z 751 (M+Na⁺, 13); mp 89-90 °C; $[\alpha]_{\text{D}}^{25} +44.2$ (c 1.0, CHCl₃); TLC R_f 0.12 (Ethyl acetate/hexane, 1/ 1).

Phenyl 2, 3, 6-tri-*O*-acetyl-4-*O*-(2', 3', 4', 6'-tetra-*O*-acetyl- β -D-galactopyranosyl)-1-thio- β -D-glucopyranoside 11²⁸

¹H NMR (CDCl₃, 400 MHz) δ 7.49-7.46 (m, 2H), 7.33-7.27 (m, 3H), 5.34 (dd, $J = 3.4, 1.0, 1\text{H}$), 5.23 (t, $J = 9.2, 1\text{H}$), 5.09 (dd, $J = 10.4, 7.8, 1\text{H}$), 4.94 (dd, $J = 10.4, 3.4, 1\text{H}$), 4.88 (dd, $J = 10.0, 9.2, 1\text{H}$), 4.66 (d, $J = 10.1, 1\text{H}$), 4.52 (dd, $J = 12.0, 2.0, 1\text{H}$), 4.46 (d, $J = 8.0, 1\text{H}$), 4.14-4.04 (m, 3H), 3.85 (bt, $J = 7.2, 1\text{H}$), 3.63-3.62 (m, 2H), 2.15 (s, 3H, COCH₃), 2.10 (s, 3H, COCH₃), 2.09 (s, 3H, COCH₃), 2.04 (s, 6H, 2 $\times$ COCH₃), 2.03 (s, 3H, COCH₃), 1.95 (s, 3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.21, 170.16, 170.0, 169.9, 169.6, 169.5, 169.0, 133.0, 131.7, 128.8, 128.2, 100.9, 85.4, 76.1, 73.8, 70.9, 70.7, 70.3, 69.1, 66.6, 62.1, 60.8, 20.72, 20.67, 20.5, 20.4; M.W. (C₃₂H₄₀O₁₇S) 728.72; ESI-MS m/z 752 (M+Na⁺, 48), 753 (43); mp 170-171 °C; $[\alpha]_{\text{D}}^{25} -19.6$ (c 1.0, CHCl₃); TLC R_f 0.13 (Ethyl acetate/hexane, 1/ 1).

***p*-Tolyl-1-thio- β -D-glucopyranoside-**12'**²⁹**

In a flame-dried, 25-mL, two-necked, round-bottomed flask equipped with an addition funnel, under argon atmosphere, was placed MoO_2Cl_2 (11.8 mg, 0.06 mmol, 3 mol%) in anhydrous CH_2Cl_2 (3 mL). To the catalyst solution, a solution of pentaacetyl- β -D-glucose (880 mg, 2 mmol) in CH_2Cl_2 (5 mL) was slowly added at ambient temperature via the addition funnel. After 10 min, a solution of 4-methyl-thiophenol (323 mg, 2.6 mmol, 2.6 equiv) in CH_2Cl_2 (2 mL) was slowly added to the reaction mixture. After completion of the reaction (about 14 h), the reaction mixture was charged with a solution of MoO_2Cl_2 (1.9 mg, 0.02 mmol, 1mol%) in anhydrous ethanol (5 mL) and then heated to reflux for 12 h. After having been cooled to ambient temperature, the mixture was evaporated to dryness under reduced pressure, and the residue was solidified by dilution with 20 mL of warm diethyl ether. The resulting solid was filtered and washed with diethyl ether (2×20 mL). The resultant white solid was dissolved in ethyl alcohol (20 mL) and the solution was filtered through a short plug of silica gel. The filtrate was concentrated to give 556 mg (94 %) of **12'** as white crystals: ^1H NMR (CDCl_3 , 400 MHz) δ 7.46 (d, $J = 8.0$, 2 H), 7.13 (d, $J = 8.0$, 2 H), 4.58 (bs, 2H, OH), 4.52 (d, $J = 7.4$, 1H), 3.86 (dd, $J = 11.0$, 1.4, 1H), 3.67 (dd, $J = 11.1$, 4.9, 1H), 3.41 (t, $J = 1.9$, 1H), 3.36-3.26 (m, 2H), 3.20 (t, $J = 9.6$, 1H); ^{13}C NMR (DMSO-d_6 , 100 MHz) δ 138.9, 133.6, 131.2, 130.7, 89.7, 82.1, 79.7, 73.8, 71.5, 63.0, 21.2; M.W. ($\text{C}_{13}\text{H}_{18}\text{O}_5\text{S}$) 286.34; ESI-MS m/z 309 ($\text{M}+\text{Na}^+$, 48), 310 (20); mp 147-148 $^\circ\text{C}$; $[\alpha]_{\text{D}}^{22} +32.0$ (c 1.0, acetone); TLC R_f 0.14 (EtOAc/hexane, 2/ 1).

4-Methylphenyl 4, 6-*O*-benzylidene-1-thio- β -D-glucopyranoside-**12**³⁰

In a dry 50-mL, two-necked, round-bottomed flask was placed MoO_2Cl_2 (17.3 mg, 0.087 mmol, 5 mol%) in anhydrous 1,4-dioxane (5 mL). To the above solution, benzaldehyde (267.5 mg, 273 μL , 2.62 mmol, 1.5 equiv) was slowly added via a micro-syringe at ambient temperature. After 20 min, *p*-tolyl-1-thio- β -D-glucopyranoside (500 mg, 1.75 mmol) was added to the above solution in one portion. The resulting reaction mixture was stirred the reaction for 20-24 hours. After completion of the reaction as evidenced by TLC analysis, the reaction mixture was sequentially treated with Et_3N (0.2 mL) and cold saturated aqueous NaHCO_3 (5 mL). The organic layer was separated and the aqueous layer was further extracted ethyl acetate (50 mL). The combined organic layer was washed with brine, dried (MgSO_4), filtered, and evaporated. The resultant crude product was purified by column chromatography (ethyl acetate/hexane, 1/2; with 1% Et_3N) to give 524 mg of acetal-**12** (80 %) as a white solid: ^1H NMR (400 MHz, CDCl_3) δ 7.48-7.43 (m, 4H), 7.38-7.35 (m, 3H), 7.15 (d, $J = 8.0$, 2H), 5.53 (s, 1H), 4.58 (d, $J = 9.6$, 1H), 4.37 (dd, $J = 10.7$, 4.4, 1H), 3.85 (dd, $J = 7.5$, 2.2, 1H), 3.77 (dd, $J = 10.7$, 4.4, 1H), 3.52-3.48 (m, 2H), 3.43 (dd, $J = 4.4$, 2.2, 1H), 2.70 (br, 1H, OH), 2.60 (br, 1H, OH), 2.36 (s, 3H, CH_3); ^{13}C NMR (100 MHz, CDCl_3) δ 138.9, 136.9, 133.7, 129.9, 129.3, 128.3, 127.2, 126.3, 101.9, 88.7, 80.3, 76.7, 74.6, 72.5, 70.6, 68.6, 21.2; M.W. ($\text{C}_{20}\text{H}_{22}\text{O}_5\text{S}$) 374; MS (EI, 70 eV) 374 (M^+ , 34), 250 (100), 145 (10), 127 (19), 124 (77), 107 (71), 91 (43), 79 (25); $[\alpha]_D^{22}$ -34.40 (c 1.0, CHCl_3); TLC R_f 0.21 (Ethyl acetate/hexane, 1/2).

Preparation of thioglycosidic disaccharide-15

Methyl 6-*S*-Acetyl-2,3,4-tri-*O*-benzyl-6-thio- α -D-glucopyranoside³¹

Diisopropyl azodicarboxylate (3.24 mL, 16.4 mmol) was added dropwise to a solution of PPh_3 (4.32 g, 16.48 mmol) in THF (100 mL) at 0 $^\circ\text{C}$. The mixture was stirred at this temperature for 30 min, after which time a white precipitate

was formed. A solution of methyl-2,3,4-tri-*O*-benzyl- α -D-glucopyranoside³² (3.808 g, 8.2 mmol) and thioacetic acid (1.2 mL, 16.4 mmol, 2 equiv) in THF (100 mL) was then added. The reaction mixture was stirred at 0 °C for 1 h and then allowed to warm to ambient temperature overnight. The reaction mixture was evaporated and the resulting red residue was purified by column chromatography. The product was further purified by re-crystallization from hexane to afford methyl 6-*S*-acetyl-2,3,4-tri-*O*-benzyl-6-thio- α -D-glucopyranoside as a crystalline solid (2.657 g, 62%): ¹H NMR (CDCl₃, 400 MHz) δ 7.35-7.24 (m, 15 H), 4.98 (d, J = 10.8, 1H, OCH_aH_bPh), 4.89 (d, J = 10.8, OCH_aH_bPh), 4.81 (d, J = 11.0, 1H, OCH_aH_bPh), 4.78 (d, J = 11.0, 1H, OCH_aH_bPh), 4.65 (d, J = 12.0, 1H, OCH_aH_bPh), 4.62 (d, J = 12.0, 1H, OCH_aH_bPh), 4.54 (d, J = 3.2, 1H, HC(1)), 3.97 (d, J = 9.2, 1H), 3.75 (td, J = 7.5, 2.2, 1H), 3.51 (dd, J = 9.6, 3.3, 1H), 3.43 (dd, J = 13.6, 2.3, 1H), 3.36 (s, 3H, OCH₃), 3.30 (t, J = 9.3, 1H), 3.03 (dd, J = 13.7, 7.8, 1H), 2.34 (s, 3H, CH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 194.8, 138.6, 138.0, 137.9, 128.41, 128.36, 128.1, 128.0, 127.92, 127.89, 127.8, 127.6, 97.8, 81.8, 80.5, 79.9, 75.7, 75.1, 73.3, 69.3, 55.1, 30.8, 30.5; M.W.(C₃₀H₃₄O₆S) 522.65; ESI-MS m/z 545 (M+Na⁺, 29); mp 82-83 °C; [α]_D²² +24.4 (c 1.0, CHCl₃); TLC R_f 0.51 (Ethyl acetate/hexane, 1/ 3).

Methyl 2,3,4-tri-*O*-benzyl-6-thio- α -D-glucopyranoside-14^{31a}

A solution of anhydrous NaOMe (103 mg, 1.9 mmol) in MeOH (3 mL) was added to a solution of methyl 6-*S*-acetyl-2,3,4-tri-*O*-benzyl-6-thio- α -D-glucopyranoside (1.0 g, 1.9 mmol) in degassed, anhydrous MeOH (40 mL).

The reaction mixture was stirred for 2 h. Amberlyst-15 ion-exchange resin was then added until the pH of the solution became neutral. The resin in the reaction mixture was removed by filtration and the collected filtrate was evaporated. The resulting residue was purified by column chromatography under nitrogen atmosphere to afford 483 mg (53 %) of **14** as a colorless oil: ¹H NMR (CDCl₃, 400 MHz) δ 7.34-7.29 (m, 15 H), 4.99 (d, J = 10.8, 1 H, OCH_aH_bPh), 4.89 (d, J = 10.8, 1 H, OCH_aH_bPh), 4.80 (d, J = 12.6, 1 H, OCH_aH_bPh), 4.77 (d, J = 12.6, 1 H, OCH_aH_bPh), 4.63 (d, J =

13.6, 1 H, $\text{OCH}_a\text{H}_b\text{Ph}$), 4.63 (dd, $J = 13.6$, 1 H, $\text{OCH}_a\text{H}_b\text{Ph}$), 4.56 (d, $J = 3.6$, HC(1) 1H), 3.98 (d, $J = 8.8$, 1H), 3.87 (t, $J = 8.4$, 1H), 3.51 (dd, $J = 9.2$, 3.2, 1H), 3.38 (s, 3H, OCH_3), 3.31 (t, $J = 9.2$, 1H), 3.16 (dd, $J = 13.6$, 1.6, 1H), 2.76 (dd, $J = 13.2$, 8.7, 1H), 1.28 (t, $J = 8.8$, 1H, SH); ^{13}C NMR (CDCl_3 , 100 MHz) δ 138.6, 138.0, 128.43, 128.40, 128.37, 128.1, 127.95, 127.90, 127.8, 127.6, 97.8, 81.9, 80.6, 80.0, 75.7, 75.0, 73.3, 69.0, 55.3, 42.1; M.W. ($\text{C}_{28}\text{H}_{32}\text{O}_5\text{S}$) 480.6; ESI-MS m/z 503 ($\text{M}+\text{Na}^+$, 53); $[\alpha]_{\text{D}}^{22} +58.6$ (c 0.9, CHCl_3); TLC R_f 0.41 (Ethyl acetate/hexane, 1/ 3).

Methyl 2,3,4- tri-*O*-benzyl -6-S-(2',3',4',6'-tetra-*O*-acetyl - β -D- glucopyranosyl-6-thio- α -D- glucopyranoside-15

In a flame-dried, 25-mL, two-necked, round-bottomed flask equipped with an addition funnel was placed MoO_2Cl_2 (12.9 mg, 0.065 mmol, 10 mol%) in anhydrous CH_2Cl_2 (1 mL). To this solution was slowly added a solution of per-acetylated-D-glucose -**2a** (329 mg, 0.837 mmol, 1.3 equiv) in CH_2Cl_2 (2 mL) at ambient temperature via the addition funnel. After 10 min, the reaction was slowly treated with a solution of methyl 2,3,4-tri-*O*-benzyl-6-thio- α -D-glucopyranoside-**15** (312 mg, 0.649 mmol) in CH_2Cl_2 (2 mL). After completion of the reaction as evidenced by TLC analysis (18 h), the reaction mixture was treated with cold saturated aqueous NaHCO_3 (5 mL). The organic layer was separated and washed with brine, dried (MgSO_4), filtered, and evaporated. The resultant crude residue was purified by column chromatography to afford 389 mg of thioglycosidic disaccharide-**15** in 74 % yield. ^1H NMR (CDCl_3 , 400 MHz) δ 7.37-7.27 (m, 15 H), 5.19 (t, $J = 9.6$, 1 H, HC(4')), 5.05 (t, $J = 9.6$, 1H, HC(3')), 4.95 (t, $J = 7.2$, 1 H, HC(2')), 4.98 (d, $J = 11.0$, 1H, $\text{OCH}_a\text{H}_b\text{Ph}$), 4.92 (d, 1 H, $J = 11.0$, $\text{OCH}_a\text{H}_b\text{Ph}$), 4.81-4.77 (m, 2 H, OCH_2Ph), 4.66 (d, $J = 13.2$, 1 H, $\text{OCH}_a\text{H}_b\text{Ph}$), 4.64 (d, $J = 13.2$, 1 H, $\text{OCH}_a\text{H}_b\text{Ph}$), 4.60 (d, $J = 7.2$, 1 H, HC(1')), 4.57 (d, $J = 3.6$, 1 H, HC(1)), 4.18 (dd, $J = 12.4$, 4.8, 1 H, $\text{H}_a\text{H}_b\text{C}(6')$), 4.05 (dd, $J = 12.4$, 2.0, 1 H, $\text{H}_a\text{H}_b\text{C}(6')$), 3.96 (t, $J = 9.2$, 1 H, HC(3)), 3.83 (td, $J = 8.0$, 2.0, 1 H, HC(5)), 3.59 (ddd, $J = 10.0$, 4.4, 2.0, 1 H, HC(5')), 3.49 (dd, $J = 9.6$, 3.2, 1 H, HC(2)), 3.40 (s, 3H, OCH_3), 3.31

(t, $J = 9.2$, 1 H, HC(4)), 3.00 (dd, $J = 13.6$, 2.0, 1 H, $H_aH_bC(6)$), 2.69 (dd, $J = 13.6$, 8.8, 1 H, $H_aH_bC(6)$), 2.04 (brs, 3H, COCH₃), 2.02 (brs, 3H, COCH₃), 2.01 (brs, 3H, COCH₃), 2.00 (brs, 3H, COCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 170.6 (C=O), 170.2 (C=O), 169.3 (C=O), 169.1 (C=O), 138.6, 138.2, 138.1, 128.47, 128.45, 128.37, 128.0, 127.94, 127.86, 127.83, 127.6, 97.8 (C(1)), 83.9 (C(1')), 81.9 (C(3)), 80.7 (C(4)), 79.9 (C(2)), 75.71(OCH₂Ph), 75.70 (C(5')), 75.0 (OCH₂Ph), 73.9 (C(4)), 73.3 (OCH₂Ph), 71.0 (C(5)), 70.2 (C(2')), 68.3 (C(3')), 62.0 (C(6')), 55.1 (OCH₃), 31.6 (C(6)), 20.61 (COCH₃), 20.58 (COCH₃), 20.55 (COCH₃), 20.51 (COCH₃); M.W. (C₄₂H₅₀O₁₄S) 810.9; ESI-MS m/z 833 (M+Na⁺, 100); $[\alpha]_D^{22} +14.6$ (c 1.0, CHCl₃); TLC R_f 0.38 (Ethyl acetate/hexane, 1/2); Anal. calcd for C₄₂H₅₀O₁₄S: C, 62.21; H, 6.21; S, 3.95. Found: C, 62.47; H, 5.98; S, 3.62.

References:

- (1) Bretting, H.; Jacobs, G.; Benecke, I.; Koenig, W. A.; Thiem, J. *Carbohydr. Res.* **1985**, *139*, 225.
- (2) (a) Hodosi, G.; Kovac, P. *Carbohydr. Res.* **1997**, *303*, 239. (b) Utille, J.-P.; Gagnaire, D. *Carbohydr. Res.* **1982**, *106*, 43.
- (3) Ishii, H.; Kitagawa, I.; Matsushita, K.; Shirakawa, K.; Tori, K. *Tetrahedron Lett.* **1981**, *22*, 1529.
- (4) Xue, J.; Pan, Y.; Guo, Z. *Tetrahedron Lett.* **2002**, *43*, 1599.
- (5) Lee, E. E.; Wood, J. O. *Carbohydr. Res.* **1981**, *89*, 329-333.
- (6) Yoshida, A.; Hayashi, T.; Takeda, N.; Oida, S.; Ohki, E. *Chem. Pharm. Bull.* **1981**, *29*, 1854.
- (7) Magnusson, G.; Ahlfors, S.; Dahmen, J.; Jansson, K.; Nilsson, U. *J. Org. Chem.* **1990**, *55*, 3932.
- (8) Cheng, M. S.; Wang, Q. L.; Tian, Q.; Song, H. Y.; Liu, Y. X.; Li, Q.; Xu, X.; Miao, H. D.; Yao, X. S.; Yang, Z. *J. Org. Chem.*, **2003**, *68*, 3658.
- (9) Kondo, H.; Aoki, S.; Ichikawa, Y.; Halcomb, R. L.; Ritzen, H.; Wong, C.-H. *J. Org. Chem.* **1994**, *59*, 864.
- (10) Ding, Y.; Liu, Y. *Carbohydr. Res.* **1991**, *209*, 306.
- (11) Pakulski, Z.; Pierozynski, D.; Zamojski, A. *Tetrahedron*, **1996**, *50*, 2975.
- (12) Shimadate, T.; Chiba, S.; Inouye, K.; Iino, T.; Hosoyama, Y. *Bull. Chem. Soc. Jpn.* **1982**, *55*, 3552.
- (13) Caddick, S.; Gazzard, L.; Motherwell, W. B.; Wilkinson, J. A. *Tetrahedron* **1996**, *52*, 149.
- (14) Griffin, G. W.; Bandara, N. C.; Clarke, M. A.; Tsang, W.-S.; Garegg, P. J. *Heterocycles* **1990**, *30*, 939.
- (15) Ogawa, M. *Carbohydr. Res.* **1977**, *54*, C17-C19.
- (16) Vic, G.; Hastings, J. J.; Howarth, O. W.; Crout, D. H. G. *Tetrahedron: Asymmetry*, **1996**, *7*, 709.

- (17) Yanase, M.; Funabashi, M. *J. Carbohydr. Chem.* **2000**, *19*, 53.
- (18) Chervin, S. M.; Abada, P.; Koreeda, M. *Org. Lett.* **2000**, *2*, 369.
- (19) Suzuki, K.; Susaki, H.; Okuno, S.; Sugiyama, Y. *J. Pharmacol. Exp. Ther.* **1999**, *288*, 57.
- (20) Xia, J.; Alderfer, J. L.; Piskorz, C. F.; Locke, R. D.; Matta, K. L. *Carbohydr. Res.* **2000**, *328*, 147.
- (21) Fuente, J. M. de la; Penades, S. *Tetrahedron: Asymmetry* **2002**, *13*, 1879.
- (22) Kametani, T.; Kawamura, K.; Honda, T. *J. Am. Chem. Soc.*, **1987**, *109*, 3010.
- (23) (a) Kim, J. M.; Roy, R. *Carbohydr. Res.* **1997**, *298*, 173. (b) Ref-22.
- (24) Kim, J. M.; Roy, R. *Carbohydr. Res.* **1997**, *298*, 173.
- (25) Yu, W.; Jin, Z. *J. Am. Chem. Soc.* **2001**, *123*, 3369.
- (26) Raadt, A. de; Ferrier, R. J., *Carbohydr. Res.* **1991**, *216*, 93.
- (27) Tropper, F. D.; Andersson, F. O.; Grand-Maitre, C.; Roy, R. *Synthesis* **1991**, *9*, 734.
- (28) Zhang, Z.; Magnusson, G. *Carbohydr. Res.* **1996**, *295*, 41.
- (29) Lin, C.-C.; Hsu, T.-S.; Lu, K.-C.; Huang, I.-T. *J. Chin. Chem. Soc.* **2000**, *47*, 921.
- (30) Wong, C.-H.; Ye, X.-S.; Zhang, Z. *J. Am. Chem. Soc.* **1998**, *120*, 7137.
- (31) (a) Paterson, D. E.; Griffin, F. K.; Alcaraz, M.-L.; Taylor, R. J. K. *Eur. J. Org. Chem.* **2002**, *7*, 1323. (b) Kvaerno, L.; Werder, M.; Hauser, H.; Carreira, E. M. *Org. Lett.* **2005**, *6*, 1145.

Current Data Parameters
NAME PhSglucosepreacetate
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031222
Time 0.19
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 1
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 228.1
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.5000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SF01 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300991 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 9.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -20.07 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME PhSglucosepreacetate
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031222
Time 0.31
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 13191
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 4096
DW 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.2000000 sec
d11 0.0360000 sec
d12 0.0000200 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SF01 100.6242395 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SF02 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127721 MHz
WDW EN
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 10.00 cm
F1P 220.000 ppm
F1 22134.61 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 11.50000 ppm/cm
HZCM 1157.04588 Hz/cm

Current Data Parameters
NAME 02
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031023
Time 16.47
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 1
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 322.5
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300095 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 8.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.06 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07150 Hz/cm

Current Data Parameters
NAME Stoglucosperacetate
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031221
Time 3.22
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 7721
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 4096
DW 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000005 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SFO1 100.6242995 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SFO2 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127706 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 7.00 cm
F1P 210.000 ppm
F1 21108.148 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 11.00000 ppm/cm
HZCM 1106.74048 Hz/cm

Current Data Parameters
NAME 4C1PhSglucosperacetate
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031227
Time 23.45
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 16384
SOLVENT CDCl3
NS 1
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 228.1
DM 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300095 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 8.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME 4C1PhSglucosperacetate
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031227
Time 22.40
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1844
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 4096
DM 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000005 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SFO1 100.6242995 MHz

===== CHANNEL f2 =====
CPOPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SFO2 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127735 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 9.00 cm
F1P 240.000 ppm
F1 24147.07 Hz
F2P -10.000 ppm
F2 -1008.13 Hz
PPMCM 12.50000 ppm/cm
HZCM 1257.65967 Hz/cm

Current Data Parameters
NAME 4MeOPhSglucoseperacetate
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040421
Time 22.56
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 16
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664766 sec
RG 71.8
W 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.5000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300095 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PRNCH 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME 4MeOPhSglucoseperacetate
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040421
Time 23.07
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 748
DS 0
SWH 25125.629 Hz
FIDRES 0.38387 Hz
AQ 1.3042164 sec
RG 4056
W 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.2000000 sec
d11 0.0300000 sec
d12 0.0000000 sec

***** CHANNEL f1 *****
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SFO1 100.6242995 MHz

***** CHANNEL f2 *****
CPDPRG2 waltz16
NUC2 1H
P2P2 50.00 usec
PL2 -3.00 dB
PL12 11.20 dB
PL13 14.20 dB
SFO2 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127736 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 7.00 cm
F1P 220.000 ppm
F1 22134.81 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PRNCH 11.50000 ppm/cm
HZCM 1157.04590 Hz/cm

Current Data Parameters
NAME 2Naphthylthioglycosperac
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031226
Time 21:57
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 16384
SOLVENT CDCl3
NS 0
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.386456 sec
RG 50.5
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.5000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300088 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PCHN 0.55000 ppm/cm
HCN 220.07152 Hz/cm

Current Data Parameters
NAME 2Naphthylthioglycosperac
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031226
Time 22:56
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1439
DS 0
SWH 25125.620 Hz
FIDRES 0.303287 Hz
AQ 1.3042154 sec
RG 4096
DW 15.900 usec
DE 5.50 usec
TE 300.0 K
D1 1.2000000 sec
D11 0.0300000 sec
D12 0.0000000 sec

***** CHANNEL f1 *****
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SFO1 100.624996 MHz

***** CHANNEL f2 *****
CPDPRG2 waltz16
NUC2 1H
PCPD2 30.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SFO2 400.1315000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127736 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 30.00 cm
CY 10.00 cm
F1P 230.000 ppm
F1 28134.81 Hz
F2P -10.000 ppm
F2 -100.613 Hz
PCHN 11.50000 ppm/cm
HCN 1157.84608 Hz/cm

Current Data Parameters
NAME 4MePhSglucoseperacetate
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040131
Time 20.09
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 16384
SOLVENT CDCl3
NS 1
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3564756 sec
RG 181
DM 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300091 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 9.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PPHMC 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME 4MePhSglucoseperacetate
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040131
Time 20.16
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 2996
DS 0
SWH 25125.629 Hz
FIDRES 0.385387 Hz
AQ 1.3040164 sec
RG 4096
DM 19.900 usec
DE 5.50 usec
TE 300.0 K
D1 1.20000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SFO1 100.6242993 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
P2P2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SFO2 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127713 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 5.00 cm
F1P 210.000 ppm
F1 21328.68 Hz
F2P -1008.13 Hz
F2 -1008.13 Hz
PPHMC 11.00000 ppm/cm
HZCM 1108.74048 Hz/cm

Current Data Parameters
NAME BnSglucosepreacetate
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031215
Time 4.07
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 16384
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 32.0
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

----- CHANNEL f1 -----
NUC1 1H
P1 14.30 usec
PL1 5.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.130088 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.50 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME BnSglucosepreacetate
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031215
Time 3.03
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1395
DS 0
SWH 25125.629 Hz
FIDRES 0.38387 Hz
AQ 1.3042164 sec
RG 406.4
DW 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

----- CHANNEL f1 -----
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SFO1 100.6242995 MHz

----- CHANNEL f2 -----
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SFO2 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127744 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 5.00 cm
F1P 220.000 ppm
F1 22134.81 Hz
F2P -10.000 ppm
F2 -1005.13 Hz
PPMCM 11.50000 ppm/cm
HZCM 1157.04688 Hz/cm

Current Data Parameters
NAME tBuSglucoseperacetate
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031220
Time 23.37
INSTRUM spect
PROBHD 5 mm BBO
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 1
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 128
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.5000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300091 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.50 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PPHCH 0.35000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME tBuSglucoseperacetate
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031220
Time 23.52
INSTRUM spect
PROBHD 5 mm BBO
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1782
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 64
DW 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.2000000 sec
D11 0.0300000 sec
D12 0.0000200 sec

***** CHANNEL f1 *****
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SFO1 100.6242995 MHz

***** CHANNEL f2 *****
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SFO2 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127713 MHz
WDW EX
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 6.00 cm
F1P 220.000 ppm
F1 22134.81 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPHCH 11.50000 ppm/cm
HZCM 1157.04688 Hz/cm

Current Data Parameters
NAME 13Dithiopropane-glucose
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031225
Time 17.12
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 13384
SOLVENT CDCl3
NS 1
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 50.5
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1330000 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

ID NMR plot parameters
CX 20.00 cm
CY 9.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PRNCH 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME 13Dithiopropane-glucose
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031225
Time 17.14
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 3443
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 4896
DW 19.500 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000000 sec
d11 0.03000000 sec
d12 0.00000000 sec

***** CHANNEL f1 *****
NUC1 13C
P1 10.30 usec
PL1 0.00 dB
SFO1 100.6262995 MHz

***** CHANNEL f2 *****
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SFO2 400.1315000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127729 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

ID NMR plot parameters
CX 20.00 cm
CY 4.00 cm
F1P 220.000 ppm
F1 225134.61 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PRNCH 11.50000 ppm/cm
HZCM 1157.04696 Hz/cm

Current Data Parameters
NAME methylundecathiolegluco
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040507
Time 22:50
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 16384
SOLVENT CDCl3
NS 4
DS 0
SWH 5995.204 Hz
FIDRES 0.355018 Hz
AQ 1.3664756 sec
RG 154
DM 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.5000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SFO1 400.1300000 MHz

F2 - Processing parameters
SI 16384
SF 400.1300095 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 30.00 cm
CY 10.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PRCH 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME methylundecathiolegluco
EXPNO 4
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040507
Time 22:51
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 0
DS 0
SWH 25425.620 Hz
FIDRES 0.363367 Hz
AQ 1.3642164 sec
RG 4096
DM 19.500 usec
DE 6.50 usec
TE 300.0 K
D1 1.2000000 sec
d11 0.2000000 sec
d12 0.0000000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 19.00 usec
PL1 7.00 dB
SFO1 100.6242995 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 3.00 dB
PL12 21.83 dB
PL13 24.98 dB
SFO2 400.1315000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127728 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 4.00 cm
F1P 220.000 ppm
F1 20134.61 Hz
F2P -12.000 ppm
F2 -105.13 Hz
PRCH 11.50000 ppm/cm
HZCM 1157.04888 Hz/cm

Current Data Parameters
NAME NaphthPhSglucoseperacet
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040604
Time 1.28
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 16384
SOLVENT CDCl3
NS 8
DS 0
SMH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 228.1
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SFO1 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300095 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.50 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PWHCH 0.55000 ppm/cm
HZCH 220.07152 Hz/cm

Current Data Parameters
NAME NaphthPhSglucoseperacet
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040604
Time 1.38
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1432
DS 0
SMH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 4096
DW 19.500 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000000 sec
d11 0.03000000 sec
d12 0.0002000 sec

***** CHANNEL f1 *****
NUC1 13C
P1 10.80 usec
PL1 7.00 dB
SFO1 100.6242995 MHz

***** CHANNEL f2 *****
CROSS2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 3.00 dB
PL12 21.00 dB
PL13 24.80 dB
SFO2 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127713 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.46

1D NMR plot parameters
CX 20.00 cm
CY 5.00 cm
F1P 220.000 ppm
F1 22134.81 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PWHCH 11.50000 ppm/cm
HZCH 1157.04668 Hz/cm

Current Data Parameters
 NAME PhSmannoseAc
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20041122
 Time 14.12
 INSTRUM spect
 PROBHD 5 mm BBO BB-1H
 PULPROG zg30
 TD 16384
 SOLVENT CDCl3
 NS 16
 DS 0
 SWH 5995.204 Hz
 FIDRES 0.365918 Hz
 AQ 1.3664756 sec
 RG 406.4
 DW 83.400 usec
 DE 6.50 usec
 TE 300.0 K
 D1 1.5000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 10.10 usec
 PL1 3.00 dB
 SF01 400.1326008 MHz

F2 - Processing parameters
 SI 16384
 SF 400.1300091 MHz
 WDW EM
 SSB 0
 LB 0.10 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 CY 10.00 cm
 F1P 10.500 ppm
 F1 4201.37 Hz
 F2P -0.500 ppm
 F2 -200.06 Hz
 PPMCM 0.55000 ppm/cm
 HZCM 220.07150 Hz/cm

Current Data Parameters
 NAME PhSmannoseAc
 EXPNO 3
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20041122
 Time 12.47
 INSTRUM spect
 PROBHD 5 mm BBO BB-1H
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 1276
 DS 0
 SWH 25125.629 Hz
 FIDRES 0.383387 Hz
 AQ 1.3042164 sec
 RG 4096
 DW 19.900 usec
 DE 6.50 usec
 TE 300.0 K
 D1 1.20000000 sec
 d11 0.03000000 sec
 d12 0.00002000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 10.80 usec
 PL1 7.00 dB
 SF01 100.6242995 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 PCPRG2 90.00 usec
 PL2 3.00 dB
 PL12 21.83 dB
 PL13 24.80 dB
 SF02 400.1319000 MHz

F2 - Processing parameters
 SI 32768
 SF 100.6127706 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40
 1D NMR plot parameters
 CX 20.00 cm
 CY 9.00 cm
 F1P 240.000 ppm
 F1 24147.07 Hz
 F2P -10.000 ppm
 F2 -1006.13 Hz
 PPMCM 12.50000 ppm/cm
 HZCM 1257.65967 Hz/cm

Current Data Parameters
NAME PhSxylose
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050213
Time 23.48
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 8
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 228.1
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SF01 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.130088 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.50 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.05 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07150 Hz/cm

Current Data Parameters
NAME PhSxylose
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050213
Time 23.56
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 856
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042154 sec
RG 4096
DW 19.950 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000005 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.80 usec
PL1 7.00 dB
SF01 100.6242955 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 3.00 dB
PL12 21.83 dB
PL13 24.80 dB
SF02 400.1315000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127713 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 12.50 cm
F1P 220.000 ppm
F1 22134.61 Hz
F2P -1006.13 Hz
F2 -1006.13 Hz
PPMCM 11.50000 ppm/cm
HZCM 1157.04688 Hz/cm

Current Data Parameters
NAME PhSarabiose
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040427
Time 22.36
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 1
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 22.6
DM 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SF01 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300050 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME PhSarabiose
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040427
Time 22.38
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 81
DS 0
SWH 25125.529 Hz
FIDRES 0.361387 Hz
AQ 1.3042164 sec
RG 4095
DM 19.500 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000005 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.80 usec
PL1 7.00 dB
SF01 100.6242995 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 3.00 dB
PL12 21.83 dB
PL13 24.80 dB
SF02 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127890 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 11.00 cm
F1P 220.000 ppm
F1 22134.81 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 11.50000 ppm/cm
HZCM 1157.94700 Hz/cm

Current Data Parameters
NAME 134-2
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20041122
Time 12.06
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 16
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 71
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SFO1 400.132608 MHz

F2 - Processing parameters
SI 16384
SF 400.1300088 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.50 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.06 Hz
PPMCH 0.55000 ppm/cm
HZCM 220.07150 Hz/cm

Current Data Parameters
NAME 134-2
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20041122
Time 12.13
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 620
DS 0
SWH 25125.629 Hz
FIDRES 0.363387 Hz
AQ 1.3042154 sec
RG 4036
DW 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.80 usec
PL1 7.00 dB
SFO1 100.6242995 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 3.00 dB
PL12 21.83 dB
PL13 24.80 dB
SFO2 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127767 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 10.00 cm
F1P 240.000 ppm
F1 24147.07 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCH 12.50000 ppm/cm

Current Data Parameters
NAME PhSnitosperacetate
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040420
Time 23.56
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 16384
SOLVENT CDCl3
NS 1
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 143.7
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 14.30 usec
PL1 -5.00 dB
SF01 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300095 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.50 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.97 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME PhSnitosperacetate
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040420
Time 23.59
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 2781
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 4096
DW 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.20000005 sec
d11 0.03000000 sec
d12 0.00020000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SF01 100.6242995 MHz

===== CHANNEL f2 =====
CROSSG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SF02 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127705 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 5.00 cm
F1P 220.000 ppm
F1 22134.81 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 11.50000 ppm/cm
HZCM 1157.04688 Hz/cm

Current Data Parameters
NAME Phlactoseperacetate
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040429
Time 21.24
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 4
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 256
DM 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SF01 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.130099 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME Phlactoseperacetate
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040422
Time 19.46
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 1251
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 4096
DM 19.900 usec
DE 5.50 usec
TE 300.0 K
D1 1.20000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.30 usec
PL1 8.00 dB
SF01 100.6242395 MHz

===== CHANNEL f2 =====
CPOPRG2 waltz16
NUC2 1H
P2P2 90.00 usec
PL2 -5.00 dB
PL12 11.20 dB
PL13 14.20 dB
SF02 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127752 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 7.00 cm
F1P 220.000 ppm
F1 22134.81 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 11.50000 ppm/cm
HZCM 1157.04688 Hz/cm

Current Data Parameters
 NAME Stoglucose
 EXPNO 2
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20040701
 Time 1.49
 INSTRUM spect
 PROBHD 5 mm BBO BB-1H
 PULPROG zg30
 TD 16384
 SOLVENT MeOD
 NS 8
 DS 0
 SMH 5995.204 Hz
 FIDRES 0.365918 Hz
 AQ 1.3664756 sec
 RG 45.3
 DW 83.400 usec
 DE 6.50 usec
 TE 300.0 K
 D1 1.5000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 10.10 usec
 PL1 3.00 dB
 SF01 400.1326008 MHz

F2 - Processing parameters
 SI 16384
 SF 400.1300034 MHz
 WDW EM
 SSB 0
 LB 0.10 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 CY 15.00 cm
 F1P 10.500 ppm
 F1 4201.37 Hz
 F2P -0.500 ppm
 F2 -200.06 Hz
 PPMCM 0.55000 ppm/cm
 HZCM 220.07150 Hz/cm

Current Data Parameters
 NAME Stoglucose
 EXPNO 3
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040701
 Time 2.00
 INSTRUM spect
 PROBHD 5 mm BBO BB-1H
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 389
 DS 0
 SMH 25125.629 Hz
 FIDRES 0.383387 Hz
 AQ 1.3042154 sec
 RG 4036
 DW 19.900 usec
 DE 5.50 usec
 TE 300.0 K
 D1 1.2000000 sec
 d11 0.0300000 sec
 d12 0.0000200 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 10.80 usec
 PL1 7.00 dB
 SF01 100.6242995 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 90.00 usec
 PL2 3.00 dB
 PL12 21.83 dB
 PL13 24.80 dB
 SF02 400.1319000 MHz

F2 - Processing parameters
 SI 32768
 SF 100.5126190 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR plot parameters
 CX 20.00 cm
 CY 9.00 cm
 F1P 240.000 ppm
 F1 24147.03 Hz
 F2P -10.000 ppm
 F2 -1006.13 Hz
 PPMCM 12.50000 ppm/cm
 HZCM 1257.69771 Hz/cm

Current Data Parameters
NAME 240-1
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060316
Time 16.26
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDCl3
NS 12
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 256
OW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.5000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SF01 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300099 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 10.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.06 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07150 Hz/cm

Current Data Parameters
NAME 240-1
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060316
Time 14.52
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 240
DS 0
SWH 25125.629 Hz
FIDRES 0.383387 Hz
AQ 1.3042164 sec
RG 4096
OW 19.900 usec
DE 6.50 usec
TE 300.0 K
D1 1.2000000 sec
d11 0.0300000 sec
d12 0.0000200 sec

===== CHANNEL f1 =====
NUC1 13C
P1 10.80 usec
PL1 7.00 dB
SF01 100.6242995 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 90.00 usec
PL2 3.00 dB
PL12 21.83 dB
PL13 24.80 dB
SF02 400.1319000 MHz

F2 - Processing parameters
SI 32768
SF 100.6127782 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 10.00 cm
F1P 220.000 ppm
F1 22134.81 Hz
F2P -10.000 ppm
F2 -1006.13 Hz
PPMCM 0.55000 ppm/cm
HZCM 1157.04688 Hz/cm

Current Data Parameters
NAME 240-final
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060329
Time 10.24
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDC13
NS 32
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 128
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SF01 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300084 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 9.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

Current Data Parameters
NAME 240-final
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060329
Time 10.24
INSTRUM spect
PROBHD 5 mm BBO BB-1H
PULPROG zg30
TD 16384
SOLVENT CDC13
NS 32
DS 0
SWH 5995.204 Hz
FIDRES 0.365918 Hz
AQ 1.3664756 sec
RG 128
DW 83.400 usec
DE 6.50 usec
TE 300.0 K
D1 1.50000000 sec

***** CHANNEL f1 *****
NUC1 1H
P1 10.10 usec
PL1 3.00 dB
SF01 400.1326008 MHz

F2 - Processing parameters
SI 16384
SF 400.1300084 MHz
WDW EM
SSB 0
LB 0.10 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
CY 9.00 cm
F1P 10.500 ppm
F1 4201.37 Hz
F2P -0.500 ppm
F2 -200.07 Hz
PPMCM 0.55000 ppm/cm
HZCM 220.07152 Hz/cm

spectrum of 240

PT-135 spectrum of 240

PT-90 spectrum of 240

Current Data Parameters
NAME CTC-030052
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060330
Time 19.01
INSTRUM spect
PROBHD 5 mm Multinucl
PULPROG cosygpgqf
TD 2048
SOLVENT CDCl3
NS 4
DS 16
SWH 6009.615 Hz
FIDRES 2.934382 Hz
AQ 0.1705268 sec
RG 24
DW 83.200 usec
DE 6.50 usec
TE 300.0 K
d0 0.00000100 sec
D1 2.00000000 sec
d13 0.00000400 sec
D16 0.00020000 sec
IN0 0.00016640 sec
MCREST 0.00000000 sec
MCWRK 2.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P0 9.65 usec
P1 9.65 usec
PL1 2.00 dB
SFO1 500.1325087 MHz

===== GRADIENT CHANNEL =====
GPRAM1 SINE.100
GPRAM2 SINE.100
GPX1 0.00 %
GPX2 0.00 %
GPY1 0.00 %
GPY2 0.00 %
GPZ1 10.00 %
GPZ2 10.00 %
P16 1000.00 usec

F1 - Acquisition parameters
ND0 1
TD 400
SFO1 500.1325 MHz
FIDRES 15.024038 Hz
SW 12.016 ppm
FMODE QP

F2 - Processing parameters
SI 1024
SF 500.1300134 MHz
WDW SINE
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

F1 - Processing parameters
SI 1024
MC2 QP
SF 500.1300134 MHz
WDW SINE

Current Data Parameters
NAME CTC-030052
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20060330
Time 20.01
INSTRUM spect
PROBHD 5 mm Multinucl
PULPROG hmqcetgpa1.2
TD 2048
SOLVENT CDCl3
NS 8
DS 16
SWH 6009.615 Hz
FIDRES 2.934382 Hz
AQ 0.1705268 sec
RG 5120
DW 83.200 usec
DE 6.50 usec
TE 300.0 K
CNST2 145.0000000
d0 0.00000300 sec
D1 2.00000000 sec
d4 0.00172414 sec
D11 0.03000000 sec
D16 0.00020000 sec
DELTA 0.00002530 sec
DELTA1 0.00120800 sec
DELTA2 0.00097045 sec
DELTA3 0.00148594 sec
LNO 0.00001500 sec
MCREST 0.00000000 sec
MCWRK 0.33333400 sec
STCNT 128

===== CHANNEL f1 =====
NUC1 1H
P1 9.65 usec
P2 19.30 usec
PL1 2.00 dB
SFO1 500.1325087 MHz

===== CHANNEL f2 =====
CPDPRG2 gart
NUC2 13C
P3 11.80 usec
P4 23.60 usec
P14 500.00 usec
PCPD2 55.00 usec
PL0 120.00 dB
PL2 0.00 dB
PL12 12.25 dB
SFO2 125.7725155 MHz
SP1 6.20 dB
SFRAM3 Crp60.0.5.20.1
SPOFF3 0.00 Hz

===== GRADIENT CHANNEL =====
GPRAM1 SINE.100
GPRAM2 SINE.100
GPX1 0.00 %
GPX2 0.00 %
GPY1 0.00 %
GPY2 0.00 %
GPZ1 80.00 %
GPZ2 40.20 %
P16 1000.00 usec