Supporting Information

Highly efficient and diastereoselective synthesis of (+)-lineatin

Ramon Alibés,^{*} Pedro de March, Marta Figueredo, Josep Font,^{*} Marta Racamonde, and Teodor Parella

Departament de Química, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

Experimental Section S2-S10

¹H and ¹³C NMR spectra for compounds **8** and **9** S11-S14

Experimental Section

General Methods.

Solutions were concentrated using an evaporator at 15-20 Torr. Flash column chromatographies were carried out on silica gel (230-400 mesh). Melting points were determined on hot stage and are uncorrected. ^1H NMR and ^{13}C NMR spectra were recorded at the Servei de Resonància Magnètica Nuclear de la Universitat Autònoma de Barcelona at 250 and 62.5 MHz or 400 and 100 or 500 MHz. HRMS spectrum were performed at the SIDI in the Universidad Autónoma de Madrid. Microanalyses were performed at the Servei d'Anàlisi Elemental de la Universitat Autònoma de Barcelona. Microwave irradiations were conducted in a focused microwave reactor (Prolabo MX350) with measurement and control of power and temperature by infrared detection at the Universidad de Castilla-La Mancha. Enantiomeric excesses (*ee*) were determined by Chiral GC analyses using a FS-LIPODEX[®] B column (30 m x 0.25 mm). Optical rotations were measured at 22 ± 2 °C.

(1*R*,4*S*,5*S*)-5-Methyl-4-pivaloyloxymethyl-3-oxabicyclo[3.2.0]hept-6-en-2-one (**3**)

A solution of 2(5*H*)-furanone **4** (600 mg, 2.8 mmol) and (*Z*)-1,2-dichloroethylene (1.1 ml, 14.1 mmol) in acetonitrile (280 mL) was placed in a photochemical reactor (two-necked 300 mL vessel fitted with a Quartz immersion type cooling jacket). The reactor was immersed in a cooling bath at -20 °C and a stream of MeOH at -15 °C was circulated throughout the refrigeration jacket. The reaction mixture was irradiated using a high pressure 125W mercury lamp (Cathodeon HPK-125) for 7.5 h. The progress of the reaction was monitored by GC. Evaporation of the solvent and chromatography (from 8:1 hexane-EtOAc to 3:1 hexane-EtOAc) afforded a mixture of seven dichlorocyclobutane stereoisomers **5** (774 mg, 2.50 mmol, 89% yield) in a ratio (34:5:6:11:29:14:1). A mixture of dichlorides **5** (200 mg, 0.65 mmol) and activated Zn dust (1.5 g, 24.0 mmol) in (80%) aqueous EtOH (3.6 mL) was irradiated under pressure in a focused microwave reactor at 105 °C for 20 min. After cooling, the reaction mixture was filtered through Celite. The solid was washed several times with EtOH and EtOAc. Evaporation of the solvent gave a residue, which was subjected to column

chromatography (12:1 hexane-EtOAc) to afford the *syn* cycloadduct **6**¹ (15 mg, 0.06 mmol, 10 % yield) and the *anti* cycloadduct **3**¹ (107 mg, 0.45 mmol, 69 % yield).

(1S)-1-[(1S,4R)-4-(1-Hydroxy-1-methylethyl)-1-methyl-2-cyclobutenyl]ethane-1,2-diol (7)

7

To a solution of **3** (880 mg, 3.70 mmol) in anhydrous THF (80 mL) at -78 °C was added MeLi 1.6 M in ether (7.2 ml, 25.6 mmol) dropwise and the mixture was stirred at -78 °C for 1 h and at room temperature for 2 h. Then, saturated solution of NH₄Cl (40 mL) was slowly added, the organic layer was separated, and the aqueous phase was extracted with CH₂Cl₂ and EtOAc, successively. The organic extracts were washed with brine, dried, and the solvents removed. Chromatography of the residue (1:1 hexane-EtOAc) gave **7** (591 mg, 3.18 mmol, 86% yield) as a white solid: mp 88-90 °C (from EtOAc-pentane); [α]_D -51.7 (c 1.04, CHCl₃); **IR** (ATR) 3224 (br), 2958, 2887, 1365, 1187 cm⁻¹; ¹H NMR (250 MHz, CDCl₃) δ 6.10 (dd, *J*_{2',3'}=3.0 Hz, *J*_{2',4'}=1.1 Hz, 1H, H-2'), 6.00 (dd, *J*_{3',2'}=3.0 Hz, *J*_{3',4'}=1.2 Hz, 1H, H-3'), 4.16 (t, *J*_{1,2}=5.8 Hz, *J*_{1,2}=5.8 Hz, 1H, H-1), 3.63 (d, *J*_{2,1}=5.8 Hz, 2H, 2H-2), 2.74 (br s, 3H, OH), 2.59 (s, 1H, H-4'), 1.31 (s, 3H, CH₃), 1.29 (s, 3H, CH₃), 1.17 (s, 3H, CH₃); ¹³C NMR (62.5 MHz, CDCl₃) δ 142.0 (CH, C-3'), 135.9 (CH, C-2'), 74.5 (CH, C-1), 72.5 (C, C-1'), 66.2 (CH, C-4'), 64.1 (CH₂, C-2), 54.4 (C, C-1'), 31.7 (CH₃), 26.9 (CH₃), 19.4 (CH₃); **MS** *m/z* (CI⁺, NH₃) 204 ([M+NH₄]⁺, 18), 187 ([M+H]⁺, 44), 169 ([M-OH]⁺, 100). Anal. Calcd for (C₁₀H₁₈O₃): C, 64.49; H, 9.74. Found: C, 64.34; H, 9.86.

2-[(1R,4S)-4-[(4S)-2,2-Dimethyl-1,3-dioxolan-4-yl]-4-methyl-2-cyclobutenyl]-2-propanol (8)

8

¹ Alibés, R.; de March, P.; Figueredo, M.; Font, J.; Fu, X.; Racamonde, M.; Álvarez-Larena, A.; Piniella, J. F.; Parella, T. *J. Org. Chem.* **2003**, 68, 1283-1289.

To a stirred solution of triol **7** (400 mg, 2.15 mmol) in acetone (6 mL) was added *p*-TsOH 0.05M in acetone (0.35 mL, 0.02 mmol). After being stirred for 12 h, the reaction mixture was diluted with CH₂Cl₂ and successively washed with a saturated aqueous solution of NaHCO₃ and brine before being dried and concentrated. The product was purified by chromatography (4:1 hexane-EtOAc) to give **8** (440 mg, 1.95 mmol, 91% yield) as a colorless oil: $[\alpha]_D$ -59.6 (*c* 1.25, CHCl₃); **IR** (ATR) 3481 (br), 2969, 2878, 1456, 1377, 1211 cm⁻¹; **¹H NMR** (250 MHz, CDCl₃) δ 6.09 (dd, $J_{2',3'}=3.0$ Hz, $J_{2'/3',1'}=1.2$ Hz, 1H, H-2'/H-3'), 5.95 (dd, $J_{2',3'}=3.0$ Hz, $J_{2'/3',1'}=1.5$ Hz, 1H, H-2'/H-3'), 4.59 (t, $J_{4'',5''}=6.8$ Hz, $J_{4'',5''}=6.8$ Hz, 1H, H-4''), 3.98 (dd, $J_{\text{gem}}=8.2$ Hz, $J_{5'',4''}=6.8$ Hz, 1H, H-5''), 3.78 (dd, $J_{\text{gem}}=8.2$ Hz, $J_{5'',4''}=6.8$ Hz, 1H, H-5''), 3.56 (br s, 1H, OH), 2.66 (s, 1H, H-1'), 1.43 (s, 3H, CH₃), 1.35 (s, 3H, CH₃), 1.27 (s, 3H, CH₃), 1.24 (s, 3H, CH₃), 1.20 (s, 3H, CH₃); **¹³C NMR** (62.5 MHz, CDCl₃) δ 140.9 (CH, C-2'/C-3'), 136.9 (CH, C-2'/C-3'), 109.3 (C, C-2''), 77.9 (CH, C-4''), 71.1 (C, C-2), 65.9 (CH₂, C-5''), 65.9 (CH, C-1'), 52.8 (C, C-4'), 30.0 (CH₃), 27.7 (CH₃), 26.1 (CH₃), 24.9 (CH₃), 18.2 (CH₃); **MS** *m/z* (CI⁺, CH₄) 227 ([M+H]⁺, 10), 209 ([M-OH]⁺, 100); **HRMS** *m/z* calcd for [C₁₃H₂₂O₃+H]⁺ 227.1647, found 227.1636.

(1*S*,2*S*,3*S*)-3-[(4*S*)-2,2-Dimethyl-1,3-dioxolan-4-yl]-2-(1-hydroxy-1-methylethyl)-3-methylcyclobutan-1-ol (9)

To a vigorously stirred yellow suspension of mercuric acetate Hg(OAc)₂ (538 mg, 1.69 mmol) in a mixture of H₂O (2 mL) and THF (1 mL), a solution of cyclobutene **8** (381 mg, 1.69 mmol) in THF (2 mL) was added dropwise. The reaction mixture was stirred for 2.5 h at rt. After 40 min the yellow color had disappeared. Then, a solution of NaOH 3 M (2 mL) followed by a solution of NaBH₄ 0.5 M (2 mL) in NaOH 3 M were added all at once with vigorous stirring. After stirring for 30 min, most of the mercury had coagulated; K₂CO₃ (30 mg) was added, the organic layer was separated and extracted with ether. The organic extracts were washed with brine, dried, and the solvents removed. Chromatography of the residue (2:1 hexane-EtOAc) gave diol **9** (377 mg, 1.55 mmol, 92% yield) as a colorless oil: $[\alpha]_D$ -3.7 (*c* 1.21, CHCl₃); **IR** (ATR) ν 3419 (br),

2969, 2932, 1455, 1371, 1157, 1050 cm^{-1} ; $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 4.41 (t, $J_{4',5'}=6.9$ Hz, $J_{4',5''}=6.9$ Hz, 1H, H-4'), 4.25 (m, 1H, H-1), 3.97 (dd, $J_{\text{gem}}=8.2$ Hz, $J_{5',4'}=6.9$ Hz, 1H, H-5'), 3.70 (dd, $J_{\text{gem}}=8.2$ Hz, $J_{5'',4''}=6.9$ Hz, 1H, H-5''), 3.65 (s, 1H, OH), 2.29 (s, 1H, OH), 1.88 (dd, $J_{\text{gem}}=11.2$ Hz, $J_{4\text{endo},1}=7.5$ Hz, 1H, H-4endo), 1.86 (d, $J_{2,1}=8.6$ Hz, 1H, H-2), 1.52 (dd, $J_{\text{gem}}=11.2$ Hz, $J_{4\text{exo},1}=8.1$ Hz, 1H, H-4exo), 1.40 (s, 3H, CH_3), 1.32 (s, 3H, CH_3), 1.28 (s, 3H, CH_3), 1.20 (s, 6H, CH_3); $^{13}\text{C NMR}$ (62.5 MHz, CDCl_3) δ 109.6 (C, C-2'), 78.4 (CH, C-4'), 70.9 (C, C-1''), 66.0 (CH, C-2), 65.6 (CH_2 , C-5'), 64.0 (CH, C-1), 39.7 (CH_2 , C-4), 36.4 (C, C-3), 29.5 (CH_3), 27.5 (CH_3), 26.0 (CH_3), 24.7 (CH_3), 22.6 (CH_3); **MS** m/z (FAB+) 245 ($[\text{M}+\text{H}]^+$, 39), 227 ($[\text{M}-\text{OH}]^+$, 90) 169 (100); **HRMS** m/z calcd for $[\text{C}_{13}\text{H}_{24}\text{O}_4+\text{H}]^+$ 245.1753 and $[\text{C}_{13}\text{H}_{24}\text{O}_4-\text{OH}]^+$: 227.1647, found 245.1749 and 227.1640.

2-[(1S,2S,4S)-4-(Benzyloxy)-2-[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]-2-methylcyclobutyl]-2-propanol (10)

To an ice-cooled suspension of NaH 60% (486 mg, 12.1 mmol) in dry THF (4 mL) under Ar atmosphere a solution of diol **9** (589 mg, 2.40 mmol) in THF (10 mL) was added dropwise. The solution was stirred for 2 h, at which point benzyl bromide (320 μL , 2.68 mmol) was added dropwise. The reaction mixture was stirred overnight as it warmed to room temperature. Excess sodium hydride was quenched with methanol and the solution was concentrated. The resulting residue was taken up in CH_2Cl_2 , washed with brine, dried and evaporated. Chromatography (12:1 hexane-EtOAc) gave the benzyl ether **10** (639 mg, 1.91 mmol, 79% yield) as a colorless oil: $[\alpha]_{\text{D}} +27.0$ (c 1.26, CHCl_3); **IR** (ATR) 3518 (br), 2973, 2931, 1728, 1454, 1370, 1210 cm^{-1} ; $^1\text{H NMR}$ (250 MHz, CDCl_3) δ 7.29 (m, 5H, Ph), 4.44 (t, $J_{4'',5''}=6.8$ Hz, $J_{4'',5'}=6.8$ Hz, 1H, H-4''), 4.42 (m, 2H, PhCH_2), 4.04 (ddd, $J_{4',1'}=8.9$ Hz, $J_{4',3'}=7.9$ Hz, $J_{4',3''}=6.9$ Hz, 1H, H-4'), 3.97 (dd, $J_{\text{gem}}=8.2$ Hz, $J_{5'',4''}=6.8$ Hz, 1H, H-5''), 3.70 (dd, $J_{\text{gem}}=8.2$ Hz, $J_{5',4'}=6.8$ Hz, 1H, H-5'), 3.64 (s, 1H, OH), 2.08 (d, $J_{1',4'}=8.9$ Hz, 1H, H-1'), 1.80 (dd, $J_{\text{gem}}=11.1$ Hz, $J_{3',4'}=6.9$ Hz, 1H, H-3'), 1.52 (dd, $J_{\text{gem}}=11.1$ Hz, $J_{3'',4''}=7.9$ Hz, 1H, H-3''), 1.42 (s, 3H, CH_3), 1.33

(s, 3H, $\underline{\text{CH}_3}$), 1.29 (s, 3H, $\underline{\text{CH}_3}$), 1.21 (s, 6H, 2 $\underline{\text{CH}_3}$); ^{13}C NMR (62.5 MHz, CDCl_3) δ 138.4 (C, Ph), 128.3/127.6 (3CH, Ph), 109.6 (C, C-2''), 78.5 (CH, C-4''), 70.8 (C, C-2), 70.7 (CH_2 , Ph $\underline{\text{CH}_2}$), 70.4 (CH, C-4'), 65.6 (CH_2 , C-5''), 63.1 (CH, C-1'), 37.5 (CH_2 , C-3'), 36.9 (C, C-2'), 29.6 ($\underline{\text{CH}_3}$), 27.7 ($\underline{\text{CH}_3}$), 26.0 ($\underline{\text{CH}_3}$), 24.7 ($\underline{\text{CH}_3}$), 22.6 ($\underline{\text{CH}_3}$); **MS** m/z (ESI+): 357 ($[\text{C}_{20}\text{H}_{30}\text{O}_4+\text{Na}]^+$, 100). Anal. Calcd for ($\text{C}_{20}\text{H}_{30}\text{O}_4$): C, 71.82; H, 9.04. Found: C, 71.75; H, 9.29.

(1S)-1-[(1S,2S,3S)-3-Benzoyloxy-2-(1-hydroxy-1-methylethyl)-1-methylcyclobutyl]ethane-1,2-diol (11)

To a stirred solution of alcohol **10** (614 mg, 1.84 mmol) in a mixture of MeOH (5 mL) and H_2O (5 mL), trifluoroacetic acid (0.2 mL, 2.6 mmol) was added dropwise. The reaction mixture was stirred overnight at room temperature. Then it was diluted with EtOH and concentrated. The product was purified by chromatography (1:1 hexane-EtOAc) to give triol **11** (506 mg, 1.72 mmol, 94% yield) as a white solid: mp 100-102 °C (from pentane-EtOAc); $[\alpha]_{\text{D}}^{25} +65.7$ (c 1.05, CHCl_3); **IR** (ATR) 3247 (br), 2964, 2934, 2890, 1363, 1218, 1187 cm^{-1} ; ^1H NMR (250 MHz, CDCl_3) δ 7.29 (m, 5H, Ph), 4.46 (d, $J_{\text{gem}}=11.7$ Hz, 1H, Ph $\underline{\text{CH}_2}$), 4.37 (d, $J_{\text{gem}}=11.7$ Hz, 1H, Ph $\underline{\text{CH}_2}$), 4.00 (ddd, $J_{3',4'}=7.9$ Hz, $J_{3',4'}=7.9$ Hz, $J_{3',2'}=7.9$ Hz, 1H, H-3'), 3.79 (dd, $J_{1,2}=8.4$ Hz, $J_{1,2}=2.9$ Hz, 1H, H-1), 3.56 (m, 2H, 2H-2), 2.02 (m, 2H, H-2', H-4'), 1.47 (dd, $J_{\text{gem}}=11.1$ Hz, $J_{4',3'}=7.9$ Hz, 1H, H-4'), 1.34 (s, 3H, $\underline{\text{CH}_3}$), 1.27 (s, 3H, $\underline{\text{CH}_3}$), 1.16 (s, 3H, $\underline{\text{CH}_3}$); ^{13}C NMR (62.5 MHz, CDCl_3) δ 138.7 (C, Ph), 128.8/128.0 (3CH, Ph), 76.0 (CH, C-1), 72.1 (C, C-1'), 70.9 (CH_2 , Ph $\underline{\text{CH}_2}$), 70.5 (CH, C-3'), 64.1 (CH, C-2'), 63.3 (CH_2 , C-2), 39.2 (C, C-1'), 38.0 (CH_2 , C-4'), 29.8 ($\underline{\text{CH}_3}$), 29.4 ($\underline{\text{CH}_3}$), 24.2 ($\underline{\text{CH}_3}$); **MS** m/z (ESI+): 317 ($[\text{C}_{17}\text{H}_{26}\text{O}_4+\text{Na}]^+$, 100). Anal. Calcd for ($\text{C}_{17}\text{H}_{26}\text{O}_4$): C, 69.36; H, 8.90. Found: C, 69.32; H, 8.83.

(1S,5S,6S,8S)-8-Benzoyloxy-2,2,6-trimethyl-3-oxabicyclo[4.2.0]octan-5-ol (12)

(1*S*,5*S*,6*S*,8*S*)-8-Benzoyloxy-5-(imidazolylthiocarbonyl)oxy-2,2,6-trimethyl-3-oxabicyclo[4.2.0]octane (13)

S7

oil: $[\alpha]_D -20.2$ (c 0.8, CHCl_3); **IR** (ATR) 2971, 2864, 1460, 1382, 1365, 1323, 1278, 1226 cm^{-1} ; **^1H NMR** (250 MHz, CDCl_3) δ 8.36 (s, 1H, H-2'), 7.49 (s, 1H, H-5'), 7.32 (m, 5H, Ph), 7.05 (s, 1H, H-4'), 5.33 (t, $J_{5,4}=1.6$ Hz, $J_{5,4}=1.6$ Hz, 1H, H-5), 4.52 (d, $J_{\text{gem}}=12.0$ Hz, 1H, PhCH_2), 4.39 (d, $J_{\text{gem}}=12.0$ Hz, 1H, PhCH_2), 3.92 (ddd, $J_{8,1}=8.2$ Hz, $J_{8,7}=8.2$ Hz, $J_{8,7}=6.6$ Hz, 1H, H-8), 3.86 (dd, $J_{\text{gem}}=5.4$ Hz, $J_{4,5}=1.6$ Hz, 2H, 2H-4), 2.20 (dd, $J_{\text{gem}}=11.3$ Hz, $J_{7,8}=6.6$ Hz, 1H, H-7), 2.01 (d, $J_{1,8}=8.2$ Hz, 1H, H-1), 1.60 (dd, $J_{\text{gem}}=11.3$ Hz, $J_{7,8}=8.2$ Hz, 1H, H-7), 1.43 (s, 3H, CH_3), 1.28 (s, 3H, CH_3), 1.27 (s, 3H, CH_3); **^{13}C NMR** (62.5 MHz, CDCl_3) δ 183.7 (C=S), 137.8 (C, Ph), 137.2 (CH, C-2'), 131.0 (CH, C-4'), 128.5/127.8 (3CH, Ph), 117.6 (CH, C-5'), 81.0 (CH, C-5), 70.8 (C, C-2), 70.6 (CH_2 , PhCH_2), 69.5 (CH, C-8), 59.3 (CH_2 , C-4), 54.5 (CH, C-1), 37.7 (CH_2 , C-7), 31.6 (C, C-6), 28.1 (CH_3), 26.3 (CH_3), 24.4 (CH_3); **MS** m/z (ESI⁺): 409 ($[\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_3\text{S}+\text{Na}]^+$, 100). Anal. Calcd for ($\text{C}_{21}\text{H}_{26}\text{N}_2\text{O}_3\text{S}$): C, 65.26; N, 7.25; H, 6.78; S, 8.30. Found: C, 64.91; N, 7.02; H, 6.64; S, 8.08.

(1S,6S,8S)-8-Benzyloxy-2,2,6-trimethyl-3-oxabicyclo[4.2.0]octane (14)

14

To a heated (100 °C) solution of tributyltin hydride (1.07 mL, 3.97 mmol) in dry toluene (11 mL), a solution of imidazolylthiocarbonate **13** (383 mg, 0.99 mmol) and 2,2'-azobisisobutyronitrile AIBN (52 mg, 0.3 mmol) in dry toluene (11 mL) was added dropwise under Ar atmosphere. The mixture was heated for 20 min. After cooling, the solvent was evaporated and the residue was purified by chromatography (from 0 to 10 % of EtOAc in hexane) to afford **14** (240 mg, 0.92 mmol, 93% yield) as an oil: $[\alpha]_D +41.5$ (c 0.97, CHCl_3); **IR** (ATR) 2926, 2863, 1454, 1362, 1210, cm^{-1} ; **^1H NMR** (500 MHz, CDCl_3) δ 7.30 (m, 5H, Ph), 4.45 (d, $J_{\text{gem}}=11.7$ Hz, 1H, PhCH_2), 4.35 (d, $J_{\text{gem}}=11.7$ Hz, 1H, PhCH_2), 3.94 (ddd, $J_{8,1}=8.2$ Hz, $J_{8,7}=7.7$ Hz, $J_{8,7}=6.5$ Hz, 1H, H-8), 3.53 (d, $J_{\text{gem}}=8.1$ Hz, 2H, 2H-4), 1.96 (dd, $J_{\text{gem}}=10.1$ Hz, $J_{7,8}=6.5$ Hz, 1H, H-7), 1.76 (d, $J_{1,8}=8.2$ Hz, 1H, H-1), 1.55 (m, 1H, H-5), 1.44 (dd, $J_{\text{gem}}=10.1$ Hz, $J_{7,8}=7.7$ Hz, 1H, H-7), 1.34 (d, $J_{\text{gem}}=13.8$ Hz, 1H, H-5), 1.19 (s, 3H, CH_3), 1.18 (s, 3H, CH_3), 1.17 (s, 3H, CH_3); **^{13}C NMR** (62.5 MHz, CDCl_3) δ 138.5 (C, Ph), 128.3/127.8/127.5 (3CH, Ph), 70.4 (CH_2 , PhCH_2), 70.4 (CH, C-8), 70.4 (C, C-2), 57.5 (CH_2 , C-4), 54.9 (CH, C-1),

41.6 (CH₂, C-7), 35.5 (CH₂, C-5), 28.4 (CH₃), 27.3 (C, C-6), 26.8 (CH₃), 24.7 (CH₃).
MS *m/z* (ESI⁺): 283 ([C₁₇H₂₄O₂+Na]⁺, 100). Anal. Calcd for (C₁₇H₂₄O₂): C, 78.42; H, 9.29. Found: C, 78.42; H, 9.39.

(1*S*,6*S*,8*S*)-2,2,6-Trimethyl-3-oxabicyclo[4.2.0]octan-8-ol (15)

A stirred solution of bicyclic ether **14** (230 mg, 0.88 mmol) in a 9:1 mixture of EtOAc and HOAc (12 mL) was hydrogenated over 10% Pd/C (858 mg) under a flow of H₂ for 2 h. The catalyst was removed by filtration over Celite and the solvent was evaporated. Chromatography (4:1 hexane-EtOAc) afforded **15** (147 mg, 0.86 mmol, 98% yield) as a colorless oil: [α]_D -2.6 (*c* 0.59, CHCl₃); **IR** (ATR) 3249 (br), 2957, 2926, 2863, 1450, 1360, 1209, 1171 cm⁻¹; **¹H NMR** (250 MHz, CDCl₃) δ 4.18 (ddd, *J*_{8,1}=8.1 Hz, *J*_{8,7}=8.1 Hz, *J*_{8,7}=7.2 Hz, 1H, H-8), 3.53 (m, 2H, 2H-4), 2.05 (dd, *J*_{gem}=10.3 Hz, *J*_{7,8}=7.2 Hz, 1H, H-7), 1.58 (dd, *J*_{gem}=13.9 Hz, *J*_{5,4}=9.4 Hz, 1H, H-5), 1.54 (d, *J*_{1,8}=8.1 Hz, 1H, H-1), 1.44 (dd, *J*_{gem}=10.3 Hz, *J*_{7,8}=8.1 Hz, 1H, H-7), 1.33 (dt, *J*_{gem}=13.9 Hz, *J*_{5,4}=2.0 Hz, *J*_{5,4}=2.0 Hz, 1H, H-5), 1.18 (s, 9H, 3CH₃); **¹³C NMR** (62.5 MHz, CDCl₃) δ 70.5 (C, C-2), 64.4 (CH, C-8), 57.8 (CH, C-1), 57.5 (CH₂, C-4), 44.0 (CH₂, C-7), 35.3 (CH₂, C-5), 28.4 (CH₃), 26.7 (C, C-6), 26.6 (CH₃), 24.6 (CH₃); **MS** *m/z* (ESI⁺): 193 ([C₁₀H₁₈O₂+Na]⁺, 100). Anal. Calcd for (C₁₀H₁₈O₂): C, 70.55; H, 10.66. Found: C, 70.62; H, 10.80.

(1*R*,6*S*)-2,2,6-Trimethyl-3-oxabicyclo[4.2.0]octan-4,8-dione (2)

To a solution of cyclobutanol **15** (126 mg, 0.74 mmol) in anhydrous CH₂Cl₂ (11 mL), a solution of Dess-Martin periodinane 15% wt in CH₂Cl₂ (1.73 mL, 0.89 mmol) was added dropwise under Ar atmosphere. The reaction mixture was stirred at room temperature for 3 h and then diluted with diethyl ether. The organic layer was washed successively with a solution of Na₂S₂O₃ (2.6 g) in saturated aqueous NaHCO₃ (8 mL), a

saturated NaHCO₃ solution and brine. The organic layer was dried, filtered and concentrated. To a stirred solution of the resulting residue in carbon tetrachloride (1.1 mL) was added a solution of sodium periodate (542 mg, 2.10 mmol) in H₂O (3.6 mL). After 5 min of vigorous stirring, ruthenium (III) chloride hydrate (30 mg, 0.22 mmol) was added. Following 24 h of stirring, CH₂Cl₂ was added, the organic layer was separated, and the aqueous phase was extracted with CH₂Cl₂. The organic extracts were washed with saturated NaHCO₃ solution, dried, and the solvents removed. Chromatography of the residue (2:1 hexane-EtOAc) gave **2** (110 mg, 0.60 mmol, 81% yield) as colorless crystals; mp 135-8 °C (from EtOAc-pentane) [lit.² mp 132.5-135.0 °C]; [α]_D +209.1 (*c* 0.93, acetone) [lit.² [α]_D +206 (*c* 1.12, acetone)]. ¹H NMR (250 MHz, CDCl₃) δ 2.95 (br s, 1H, H-1), 2.89 (m, 2H, 2H-5), 2.83 (d, *J*_{gem}=16.8 Hz, 1H, H-7), 2.71 (d, *J*_{gem}=16.8 Hz, 1H, H-7), 1.53 (s, 3H, CH₃), 1.52 (s, 3H, CH₃), 1.40 (s, 3H, CH₃); ¹³C NMR (62.5 MHz, CDCl₃) δ 205.6 (C=O, C-8), 170.0 (C=O, C-4), 80.5 (C, C-2), 69.8 (CH, C-1), 58.0 (CH₂, C-7), 40.1 (CH₂, C-5), 29.5 (C, C-6), 28.1 (CH₃, CH₃), 26.9 (CH₃, CH₃), 26.7 (CH₃, CH₃). Anal. Calcd for (C₁₀H₁₄O₃): C, 65.91; H, 7.74. Found: C, 65.84; H, 7.85.

² Mori, K.; Nagano, E. *Liebigs Ann. Chem.* **1991**, 341-344.

S13