Supporting Information for
Highly Efficient and Reusable Dendritic Catalysts Derived from
***N*-Prolylsulfonamide for the Asymmetric Direct Aldol Reaction in**
Water

Yongyong Wu,^a Yazhu Zhang,^b Menglong Yu^b, Gang Zhao,^{b,*} Shaowu Wang^{a,*}

^a *Anhui Key Laboratory of Functional Molecular Solids, Institute of Organic Chemistry, School of Chemistry and Materials Science, Anhui Normal University, Wuhu, Anhui 241000, China*

^b *Shanghai Institute of Organic Chemistry, Chinese Academy of Science, Shanghai, 200032, China*

zhaog@mail.sioc.ac.cn swwang@mail.ahnu.edu.cn

Table of Contents

General information.....	S2
General procedure for the synthesis of catalysts 1a-f and data	S3-S6
General procedure for the Aldol reaction and data of 6a-v	S7-S11
List of the obvious difference between syn-6 and anti-6 on ¹ H NMR	S12
NMR spectra for catalysts 1a-f and 2	S13-S16
NMR spectra for compounds 6a-v	S17-S28
HPLC spectra for compounds 6a-v	S29-S50
References.....	S51

General Information:

Reactions were monitored by thin-layer chromatography (TLC) carried out on 0.25 mm silica gel plates visualized with UV light and/or by staining with ethanolic phosphomolybdic acid (PMA). Flash column chromatography was performed on silica gel H (10-40 μ). NMR spectra were recorded on 300 or 500 MHz instruments. Chemical shifts (δ) are given in ppm relative to TMS, coupling constants (J) in Hz. IR spectra were recorded on a spectrometer. Mass spectra (MS) were measured with a spectrometer. High resolution mass spectra (HRMS) were recorded on a mass spectrometer. Elemental analyses were performed. Melting point were measured on a digital melting- point apparatus. Optical rotations were measured on a JASCO P-1010 Polarimeter at $\lambda=589$ nm. Analytical high performance liquid chromatography (HPLC) was carried out on WATERS 510 instrument (2487 Dual λ Absorbance Detector and 515 HPLC Pump) using chiral column.

General procedure for the synthesis of catalysts 1a-f

Scheme 1

(a) DCC, DMAP, DCM, r.t. 24 h; (b) CF₃COOH, MeOH, r.t., 12 h; (c) Pd/C, H₂(1atm), MeOH r.t., 12 h.

N-Methanesulfonyl-L-proline Amide 1a^[1-4]

1a

Under a dry Ar atmosphere, to a stirred solution of Boc-L-proline (2.37 g, 11 mmol, 1.1 eq.) in dichloromethane (25 mL) were added methanesulfonamide (0.95 g, 10 mmol, 1.0 eq.), DMAP (366 mg, 3 mmol, 0.3 eq.) and DCC (2.47 g, 12 mmol, 1.2 eq.) respectively. The resulting mixture was stirred at room temperature for 2 days. Then the mixture was filtered, and washed with DCM (about 10 mL). The filtrate was added 1 M aqueous HCl (20 mL). The organic layer was washed with half-saturated brine (20 mL), dried (Na₂SO₄) and concentrated in vacuo. The crude product was purified by flash column chromatography (DCM - MeOH, 15:1) to give Boc-protected **1a** (1.75 g, 60%). Deprotection was performed by using 20% TFA in dichloromethane (50 mL) for one day at room temperature. After rotary evaporation, TFA salts were removed by triturating the

residue with 40 mL of methanol (saturated with ammonia). The crude product was purified by flash column chromatography (DCM–MeOH, 4:1) to give the **1a** as a white solid (832 mg, 74%). $[\alpha]_D^{26} = -37.6$ ($c = 0.58$, DMSO). ^1H NMR (DMSO- d_6) δ 8.54 (2H, br), 3.82 (1H, dd, $J = 8.1$, 6.9 Hz), 3.36–2.88 (2H, m), 2.79 (3H, s), 2.24–1.62 (4H, m). IR (film, cm^{-1}): 3102, 1574, 1244, 1133, 854.

N-Toluenesulfonyl-L-proline Amide 1b^[3-5]

1b

Under a dry Ar atmosphere, to a stirred solution of Boc-L-proline (2.37 g, 11 mmol, 1.1 eq.) in dichloromethane (25 mL) were added methanesulfonamide (1.72 g, 10 mmol, 1.0 eq.), DMAP (366 mg, 3 mmol, 0.3 eq.) and DCC (2.47 g, 12 mmol, 1.2 eq.) respectively. The resulting mixture was stirred at room temperature for 2 days. Then the mixture was filtered, and washed with DCM (about 10 mL). The filtrate was added 1 M aqueous HCl (20 mL). The organic layer was washed with half-saturated brine (20 mL), dried (Na_2SO_4) and concentrated in vacuo. The crude product was purified by flash column chromatography (Hexane-AcOEt, 8:1) to give Boc-protected **1b** (2.00 g, 54%). Deprotection was performed using 20% TFA in dichloromethane (50 mL) for one day at room temperature. After rotary evaporation, TFA salts were removed by triturating the residue with 40 mL of methanol (saturated with ammonia). The crude product was purified by flash column chromatography (Hexane-AcOEt, 2:1) to give the **1b** as a white solid (663 mg, 46%). $[\alpha]_D^{25} = 5.84$ ($c = 1.90$, MeOH). ^1H NMR (DCCl_3) δ 8.00 (1H, br), 7.81 (2H, d, $J = 8.4$ Hz), 7.26 (2H, d, $J = 7.8$ Hz), 3.93 (1H, t, $J = 12.0$ Hz), 3.72 (1H, br), 3.54–3.44 (1H, m), 3.40–3.30 (1H, m), 2.40 (3H, s), 2.20–1.68 (4H, m). IR (film, cm^{-1}): 2928, 1754, 1622, 1573, 1425, 1270, 1084, 963, 853, 818, 664, 569.

N-4-Aminobenzenesulfonyl-L-Boc-proline Amide 2^[5-6]

2

Under a dry Ar atmosphere, to a stirred solution of Boc-L-proline (9.46 g, 44 mmol, 1.1 eq.) in dichloromethane (100 mL) were added 4-nitrobenzenesulfonamide (7.68 g, 40 mmol, 1.0 eq.), DMAP (1.47 mg, 12 mmol, 0.3 eq.) and DCC (9.89 g, 48 mmol, 1.2 eq.) respectively. The resulting mixture was refluxed for 3 days. Then the mixture was filtered, and washed with DCM (about 50 mL). The filtrate was added 1 M aqueous HCl (100 mL). The organic layer was washed twice with half-saturated brine (50 mL), dried (Na_2SO_4) and concentrated in vacuo. It was the crude product of N-4-Nitrobenzenesulfonyl-L-Boc-proline Amide (**2'**). The pure **2'** was obtained by recrystallization from methanol (7.56 g, 68%). To a solution of **2** (3.99 g, 10 mmol, 1.0 eq.) in methanol was added 10% Pd-carbon (1% w/w) and subjected to hydrogenation under 1 atmospheric pressure at room temperature. The reaction was monitored by TLC, after completion of reaction the reaction mixture was filtered and concentrated in vacuum to afford crude product,

The pure **2** as obtained by recrystallization from methanol (2.58g, 86%). mp 142-143°C; 176-178°C. $[\alpha]_D^{25} = -7.64$ ($c = 0.62$, MeOH, mixture). ^1H NMR (300 MHz, DMSO- d_6): δ 11.77 (1H, s), 7.54 (2H, d, $J = 11.1$ Hz), 6.57 (2H, d, $J = 9.0$ Hz), 6.13 (2H, m), 4.14-3.96 (1H, m), 3.32-3.14 (2H, m), 2.16-1.92 (1H, m), 1.75-1.42 (3H, m), 1.34, 1.01 (9H, s). ^{13}C NMR (75MHz, DMSO- d_6): δ 176.2, 158.9, 158.1, 135.0, 128.9, 117.4, 83.9, 64.5, 51.5, 35.5, 32.8, 28.2. IR (film, cm^{-1}): 3429, 2252, 1689, 1596, 1400, 1027, 824, 762. MS (ESI) m/z $[\text{M} + \text{NH}_4]^+$ Calcd for $\text{C}_{16}\text{H}_{27}\text{N}_4\text{O}_5\text{S}$: 378.2, found: 378.2. Anal. Calcd for $\text{C}_{16}\text{H}_{23}\text{N}_3\text{O}_5\text{S}$: C, 52.02; H, 6.28; N, 11.37. Found: C, 52.01; H, 6.27; N, 11.47.

N*-4-Benzamidobenzenesulfonyl-L-Boc-proline Amide **1c*^[5]

1c

Under a dry Ar atmosphere, to a stirred solution of **2** (3.69 g, 10 mmol, 1.0 eq.) in DMF (50 mL) were added benzoic acid (1.34 g, 11 mmol, 1.1 eq.), DMAP (1.46 g, 12 mmol, 1.2 eq.) and DCC (2.47 g, 12 mmol, 1.2 eq.) respectively. The resulting mixture was stirred at room temperature for 1 day. Then the mixture was added EtOAc (50 mL) and 1 M aqueous HCl (100 mL). The organic layer was washed twice with saturated brine (50 mL), dried (Na_2SO_4) and concentrated in vacuo. The crude product was purified by flash column chromatography (Hexane-AcOEt, 20:1) to give Boc-protected **Boc-1c** (3.11 g, 66%). Deprotection was performed using 20% TFA in dichloromethane (50 mL) for 12 hours at room temperature. After rotary evaporation, TFA salts were removed by triturating the residue with 40 mL of methanol (saturated with ammonia). The crude product was purified by flash column chromatography (DCM - MeOH, 8:1) to give the **1c** as a white solid (2.07 g, 84%). mp 250-251°C. $[\alpha]_D^{25} = -42.5$ ($c=0.31$, MeOH). ^1H NMR (300 MHz, DMSO- d_6): δ 10.44 (1H, s), 7.94 (2H, d, $J = 7.2$ Hz), 7.82-7.70 (4H, m), 7.62-7.44 (3H, m), 3.82 (1H, t, $J = 7.2$ Hz), 3.18-2.96 (2H, m), 2.67 (1H, s), 2.19-2.04 (1H, m), 1.91-1.62 (3H, m). IR (film, cm^{-1}): 3372, 3188, 1669, 1400, 1086, 848. MS (MALDI) m/z $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{18}\text{H}_{20}\text{N}_3\text{O}_4\text{S}$: 374.1, found: 374.1. Anal. Calcd for $\text{C}_{18}\text{H}_{19}\text{N}_3\text{O}_4\text{S}$: C, 57.89; H, 5.13; N, 11.25. Found: C, 58.01; H, 5.23; N, 11.31.

Typical procedure for the synthesis of catalysts **1d-f**^[5,7]

1d $n=0$
1e $n=1$
1f $n=2$

Under a dry Ar atmosphere, to a stirred solution of **2** (1.0 eq.) in DCM were added dendritic wedges **3** with carboxyl groups (1.0 eq.), DMAP (1.5 eq.) and DCC (1.5 eq.) respectively. The resulting mixture was stirred at room temperature for 1 day. Then the mixture was added DCM and 1 M aqueous HCl. The organic layer was washed twice with saturated brine, dried (Na_2SO_4) and concentrated in vacuo. The crude product was purified by flash column chromatography

(DCM-Acetone) to give Boc-protected **Boc-1d**, **Boc-1e**, and **Boc-1f**. Deprotection was performed using 20% TFA in dichloromethane (50 mL) for 12 hours at room temperature. After rotary evaporation, TFA salts were removed by triturating the residue with methanol (saturated with ammonia). The crude product was purified by flash column chromatography (DCM–Acetone) to give the **1d-f** as a white solid.

Dendritic N-Aminobenzenesulfonyl-L-proline Amide (n=0) 1d

Total yield: 68%. mp 212-213 °C. $[\alpha]_D^{25} = -20.0$ ($c = 0.65$, CHCl_3). ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 10.33 (1H, s), 7.76(4H, s), 7.45-7.30 (10H, m), 7.20 (2H, d, $J = 1.5$ Hz), 6.90 (1H, s), 5.16 (1H, s), 3.82 (1H, t, $J = 7.2$ Hz), 3.20-3.02 (2H, m), 2.21-2.05 (1H, m), 1.85-1.70 (3H, m). IR (film, cm^{-1}): 3355, 3067, 1678, 1514, 1140, 834. MS (MALDI) m/z $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{32}\text{H}_{32}\text{N}_3\text{O}_6\text{S}$: 586.2, found: 585.9. Anal. Calcd for $\text{C}_{32}\text{H}_{31}\text{N}_3\text{O}_6\text{S}$: C, 65.62; H, 5.34; N, 7.17. Found: C, 65.72; H, 5.33; N, 7.18.

Dendritic N-Aminobenzenesulfonyl-L-proline Amide (n=1) 1e

Total yield: 63%. mp 188-189 °C. $[\alpha]_D^{25} = -29.5$ ($c = 0.44$, CHCl_3). ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 10.43 (1H, s), 7.88 (4H, s), 7.72-7.12 (22H, m), 7.02-6.60 (7H, m), 5.19 (12H, s), 3.93 (1H, t, $J = 7.2$ Hz), 3.24-3.10 (2H, m), 2.30-2.14 (1H, m), 2.00-1.72 (3H, m). IR (film, cm^{-1}): 3032, 1677, 1605, 1452, 1028, 836, 696. MS (MALDI) m/z $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{60}\text{H}_{56}\text{N}_3\text{O}_{10}\text{S}$: 1010.4, found: 1010.5. Anal. Calcd for $\text{C}_{60}\text{H}_{55}\text{N}_3\text{O}_{10}\text{S}$: C, 71.34; H, 5.49; N, 4.16. Found: C, 71.28; H, 5.56; N, 4.12.

Dendritic N-Aminobenzenesulfonyl-L-proline Amide (n=2) 1f

Total yield: 42%. mp 181-182 °C. $[\alpha]_D^{25} = -7.65$ ($c = 0.62$, CHCl_3). ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 10.32 (1H, br), 7.77 (4H, s), 7.42-6.90 (42H, m), 6.71-6.23 (19H, m), 5.13-4.44 (28H, m), 3.82 (1H, t, $J = 22.2$ Hz), 3.18-2.94 (2H, m), 2.19-2.04 (1H, m), 1.88-1.62 (3H, m). IR (film, cm^{-1}): 3306, 3064, 1596, 1448, 1160, 828, 699. MS (MALDI) m/z $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{116}\text{H}_{104}\text{N}_3\text{O}_{18}\text{S}$: 1858.7, found: 1858.7. Anal. Calcd for $\text{C}_{116}\text{H}_{103}\text{N}_3\text{O}_{18}\text{S}$: C, 74.94; H, 5.58; N, 2.26. Found: C, 75.04; H, 5.52; N, 2.21.

Typical procedure for the aldol reaction using the **1e catalyst in water:** The following procedure for the reaction of cyclohexanone with *p*-nitrobenzaldehyde (**2a**) in water using **1e** catalyst is representative. To a mixture of **1e** (51 mg, 0.05 mmol) in water (1.0 mL) *p*-nitrobenzaldehyde (**2a**, 76 mg, 0.5 mmol) was added at 20 °C under air in a closed system. The reaction mixture was stirred for 5 min, then cyclohexanone (156 μ L, 1.5 mmol) were added. The reaction mixture was stirred for 24 h, then it was added 1.5mL of hexane/AcOEt (1:1), and the organic phase was removed by centrifugal separation. This process was repeated 5 times. The combined organic extracts were dried over Na₂SO₄. Diastereoselectivity and conversion were determined by ¹H NMR analysis of the crude aldol product. Purification by flash column chromatography (silica gel, hexane/AcOEt) gave the aldol product **6a** as a colorless solid. The enantiomeric excess (ee) of **6a** was determined by chiral-phase HPLC analysis. The absolute configuration of aldol products **6** was extrapolated by comparison of the HPLC-data with those of **6a**, **6b**, **6c**, and **6e** whose absolute configuration were known. the catalyst **1e** could be recovered by filtering of the remaining aqueous phase, washing with MeOH (about 5mL $\times$ 2) and drying with P₂O₅ in vacuum.

(2S, 1'R)-2-(Hydroxy-(*p*-nitrophenyl)methyl)cyclohexan-1-one **6a**^[8,9]

$[\alpha]_D^{25} = +12.8$ ($c=1.85$, CHCl₃), >99% ee. ¹H NMR (300MHz, CDCl₃): δ 8.21 (2H, d, $J = 8.7$ Hz), 7.51 (2H, d, $J = 8.7$ Hz), 4.90 (1H, dd, $J = 8.4, 3.0$ Hz), 4.09 (1H, d, $J = 3.0$ Hz), 2.65-2.45 (2H, m), 2.36 (1H, td, $J = 13.2, 5.7$ Hz), 2.17-2.06 (1H, m), 1.87-1.78 (1H, m), 1.67-1.51 (3H, m), 1.45-1.31 (1H, m). IR (film): 3730, 2924, 1699, 1519, 1346, 854 cm⁻¹. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (4:1 hexane:2-propanol), 21 °C, 254 nm, 0.5 mL/min; major enantiomer tr = 52.7 min, minor enantiomer tr = 41.7 min.

(2S, 1'R)-2-(Hydroxy-(*m*-nitrophenyl)methyl)cyclohexan-1-one **6b**^[8]

$[\alpha]_D^{24} = +32.5$ ($c=1.35$, CHCl₃), >99% ee. ¹H NMR (300MHz, CDCl₃): δ 8.12 (1H, s), 8.16 (1H, d, $J = 7.8$ Hz), 7.68 (1H, d, $J = 7.5$ Hz), 7.53 (1H, t, $J = 7.8$ Hz), 4.90 (1H, dd, $J = 8.7, 3.0$ Hz), 4.15 (1H, d, $J = 3.0$ Hz), 2.68-2.58 (1H, m), 2.54-2.46 (1H, m), 2.38 (1H, td, $J = 12.3, 5.4$ Hz), 2.17-2.07 (1H, m), 1.87-1.78 (1H, m), 1.67-1.53 (3H, m), 1.46-1.31 (1H, m). IR (film, cm⁻¹): 3512, 2939, 1701, 1529, 1351, 736. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 20 °C, 254 nm, 0.5 mL/min; major enantiomer tr = 59.7 min, minor enantiomer tr = 75.1 min.

(2S, 1'R)-2-(Hydroxy-(*o*-nitrophenyl)methyl)cyclohexan-1-one **6c**^[8]

$[\alpha]_D^{24} = +19.8$ ($c=1.60$, CHCl₃), 99% ee. ¹H NMR (300MHz, CDCl₃): δ 7.84 (1H, d, $J = 8.1$ Hz), 7.77 (1H, d, $J = 7.8$ Hz), 7.63 (1H, t, $J = 7.5$ Hz), 7.43 (1H, t, $J = 7.8$ Hz), 5.45 (1H, d, $J = 6.6$ Hz), 3.90 (1H, br), 2.82-2.70 (1H, m), 2.50-2.40 (1H, m), 2.34 (1H, td, $J = 12.3, 5.7$ Hz), 2.15-2.06 (1H, m), 1.90-1.55 (4H, m). IR (film): 3512, 2940, 1699, 1526, 1351, 855 cm⁻¹. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 20 °C, 254 nm, 0.5 mL/min; major enantiomer tr = 59.4 min, minor enantiomer tr = 63.0 min.

(2S, 1'R)-2-(Hydroxy-(*p*-(trifluoromethyl)phenyl)methyl)cyclohexan-1-one **6d**^[10]

$[\alpha]_D^{25} = +2.3$ ($c = 1.00$, CHCl₃), 99% ee. ¹H NMR (300MHz, CDCl₃): δ 7.74-7.55 (3H, m), 7.40

(1H, t, $J = 7.2$ Hz), 5.30 (1H, d, $J = 9.3$ Hz), 4.03 (1H, t, $J = 3.0$ Hz), 2.81-2.69 (1H, m), 2.55-2.45 (1H, m), 2.37 (1H, td, $J = 12.9, 4.8$ Hz), 2.15-2.03 (1H, m), 1.81-1.49 (3H, m), 1.48-1.23 (1H, m). IR (film): 3512, 2943, 1701, 1453, 1312, 768 cm^{-1} . Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 21 °C, 254 nm, 0.5 mL/min; major enantiomer tr = 33.0 min, minor enantiomer tr = 35.3 min.

(2S, 1'R)-2-(Hydroxy-(*p*-cyanophenyl)methyl)cyclohexan-1-one 6e^[8]

$[\alpha]_{\text{D}}^{22} = +23.3$ ($c = 1.55$, CHCl_3), 98% ee. ^1H NMR (300MHz, CDCl_3): δ 7.65 (2H, d, $J = 8.1$ Hz), 7.45 (2H, d, $J = 8.1$ Hz), 4.85 (1H, dd, $J = 8.1, 3.0$ Hz), 4.11 (1H, d, $J = 3.0$ Hz), 2.65-2.44 (2H, m), 2.37 (1H, td, $J = 12.9, 6.0$ Hz), 2.17-2.06 (1H, m), 1.88-1.77 (1H, m), 1.72-1.47 (3H, m), 1.44-1.31 (1H, m). IR (film): 3509, 2932, 2227, 1701, 1609, 841 cm^{-1} . Enantiomeric excess was determined by HPLC with a Chiralpak AD column (4:1 hexane:2-propanol), 21 °C, 254 nm, 0.5 mL/min; major enantiomer tr = 46.6 min, minor enantiomer tr = 37.9 min.

(2S, 1'R)-2-(Hydroxy-(*p*-fluorophenyl)methyl)cyclohexan-1-one 6f^[11]

$[\alpha]_{\text{D}}^{24} = +27.5$ ($c = 0.35$, CHCl_3), 95% ee. ^1H NMR (300MHz, CDCl_3): δ 7.29 (2H, dd, $J = 8.7, 5.7$ Hz), 7.03 (2H, t, $J = 8.7$ Hz), 4.77 (1H, dd, $J = 8.7, 2.7$ Hz), 4.00 (1H, d, $J = 3.6$ Hz), 2.61-2.44 (2H, m), 2.35 (1H, td, $J = 12.9, 6.0$ Hz), 2.15-2.04 (1H, m), 1.85-1.75 (1H, m), 1.70-1.50 (3H, m), 1.37-1.20 (1H, m). IR (film, cm^{-1}): 3508, 2929, 1702, 1604, 1511, 1222, 839. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 21 °C, 220 nm, 0.5 mL/min; major enantiomer tr = 44.2 min, minor enantiomer tr = 39.6 min.

(2S, 1'R)-2-(Hydroxy-(*p*-chlorophenyl)methyl)cyclohexan-1-one 6g^[8]

$[\alpha]_{\text{D}}^{25} = +21.7$ ($c = 1.00$, CHCl_3), 99% ee. ^1H NMR (300MHz, CDCl_3): δ 7.29 (4H, dd, $J = 20.4, 8.4$ Hz), 4.76 (1H, dd, $J = 8.7, 2.7$ Hz), 3.99 (1H, d, $J = 3.0$ Hz), 2.61-2.44 (2H, m), 2.35 (1H, td, $J = 12.9, 5.4$ Hz), 2.15-2.05 (1H, m), 1.85-1.75 (1H, m), 1.70-1.50 (3H, m), 1.37-1.20 (1H, m). IR (film, cm^{-1}): 3491, 2934, 1698, 1489, 830. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 21 °C, 220 nm, 0.5 mL/min; major enantiomer tr = 31.8 min, minor enantiomer tr = 26.6 min.

(2S, 1'R)-2-(Hydroxy-(*p*-bromophenyl)methyl)cyclohexan-1-one 6h^[8]

$[\alpha]_{\text{D}}^{24} = +22.6$ ($c = 0.70$, CHCl_3), 99% ee. ^1H NMR (300MHz, CDCl_3): δ 7.47 (2H, d, $J = 8.1$ Hz), 7.20 (2H, d, $J = 8.7$ Hz), 4.75 (1H, dd, $J = 8.7, 2.7$ Hz), 3.99 (1H, d, $J = 3.0$ Hz), 2.61-2.44 (2H, m), 2.35 (1H, td, $J = 12.9, 6.3$ Hz), 2.15-2.04 (1H, m), 1.85-1.75 (1H, m), 1.70-1.50 (3H, m), 1.37-1.20 (1H, m). IR (film, cm^{-1}): 3361, 2928, 1692, 1460, 825 cm^{-1} . Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 21 °C, 220 nm, 0.5 mL/min; major enantiomer tr = 54.4 min, minor enantiomer tr = 45.4 min.

(2S, 1'R)-2-(Hydroxy-(2,4-dichlorophenyl)methyl)cyclohexan-1-one 6i^[12]

$[\alpha]_{\text{D}}^{22} = +23.9$ ($c = 1.55$, CHCl_3), 97% ee. ^1H NMR (300 MHz, CDCl_3): δ 7.50 (1H, d, $J = 8.4$ Hz), 7.35 (1H, s), 7.29 (1H, d, $J = 8.4$ Hz), 5.29 (1H, dd, $J = 7.5, 3.3$ Hz), 4.06 (1H, d, $J = 3.9$ Hz), 2.67-2.56 (1H, m), 2.52-2.42 (1H, m), 2.34 (1H, td, $J = 12.6, 5.7$ Hz), 2.16-2.04 (1H, m), 1.90-1.78 (1H, m), 1.73-1.50 (4H, m). IR (film, cm^{-1}): 3505, 2939, 1701, 1590, 1474, 821. Enantiomeric excess was determined by HPLC with a Chiralpak AS-H column (9:1

hexane:2-propanol), 20 °C, 220 nm, 0.5 mL/min; major enantiomer tr = 27.1 min, minor enantiomer tr = 23.2 min.

(2S, 1'R)-2-(Hydroxy(phenyl)methyl)cyclohexan-1-one 6j^[8]

$[\alpha]_D^{22} = +23.3$ ($c = 1.55$, CHCl₃), 98% ee. ¹H NMR (300MHz, CDCl₃): δ 7.50-7.24 (5H, m), 4.80 (1H, d, $J = 9.0$ Hz), 4.00 (1H, m), 2.70-2.56 (1H, m), 2.55-2.44 (1H, m), 2.34 (1H, td, $J = 12.3, 5.4$ Hz), 2.16-2.03 (1H, m), 1.87-1.73 (1H, m), 1.72-1.50 (3H, m), 1.40-1.22 (1H, m). IR (film, cm⁻¹): 3508, 2932, 1697, 1451, 702. Enantiomeric excess was determined by HPLC with a Chiralpak OD-H column (9:1 hexane:2-propanol), 20 °C, 220 nm, 0.5 mL/min; major enantiomer tr = 22.4 min, minor enantiomer tr = 32.0 min.

(2S, 1'R)-2-(Hydroxy-(p-tolyl)methyl)cyclohexan-1-one 6k^[13]

$[\alpha]_D^{24} = +12.9$ ($c = 0.17$, CHCl₃), 95% ee. ¹H NMR (300MHz, CDCl₃): δ 7.18 (4H, dd, $J = 17.1, 8.4$ Hz), 4.75 (1H, dd, $J = 9.0, 2.7$ Hz), 3.91 (1H, d, $J = 2.7$ Hz), 2.66-2.54 (1H, m), 2.51-2.43 (1H, m), 2.35 (1H, td, $J = 13.2, 6.0$ Hz), 2.34 (3H, s), 2.14-2.03 (1H, m), 1.82-1.72 (1H, m), 1.70-1.50 (3H, m), 1.38-1.18 (1H, m). IR (film, cm⁻¹): ν 3513, 2926, 1701, 1449, 820. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 21 °C, 220 nm, 0.5 mL/min; major enantiomer tr = 27.7 min, minor enantiomer tr = 34.7 min.

(2S, 1'R)-2-(Hydroxy-(p-methoxyphenyl)methyl)cyclohexan-1-one 6l^[9]

$[\alpha]_D^{24} = +28.9$ ($c = 1.50$, CHCl₃), 96% ee. ¹H NMR (300 MHz, CDCl₃): δ 7.24 (2H, d, $J = 8.4$ Hz), 6.88 (2H, d, $J = 8.7$ Hz), 4.74 (1H, dd, $J = 8.7, 2.4$ Hz), 3.93 (1H, d, $J = 2.7$ Hz), 3.80 (3H, s), 2.65-2.49 (1H, m), 2.52-2.43 (1H, m), 2.35 (1H, td, $J = 12.9, 5.4$ Hz), 2.15-2.04 (1H, m), 1.84-1.73 (1H, m), 1.70-1.45 (3H, m), 1.36-1.24 (1H, m). IR (film, cm⁻¹): 3514, 2925, 1699, 1514, 1248, 832. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 21 °C, 220 nm, 0.5 mL/min; major enantiomer tr = 64.2 min, minor enantiomer tr = 60.9 min.

(2S, 1'R)-2-(Hydroxy-(p-benzyloxyphenyl) methyl)cyclohexan-1-one 6m

mp 101-102 °C. $[\alpha]_D^{22} = +9.0$ ($c = 1.00$, CH₂Cl₂), >99% ee. ¹H NMR (300 MHz, CDCl₃): δ 7.45-7.28 (5H, m), 7.23 (2H, d, $J = 9.0$ Hz), 6.95 (2H, d, $J = 8.7$ Hz), 5.05 (2H, s), 4.74 (1H, dd, $J = 9.0, 2.7$ Hz), 3.93 (1H, d, $J = 2.4$ Hz), 2.64-2.53 (1H, m), 2.51-2.43 (1H, m), 2.34 (1H, m), 2.13-2.02 (1H, m), 1.82-1.72 (1H, m), 1.69-1.46 (3H, m), 1.34-1.21 (1H, m). ¹³C NMR (75 MHz, CDCl₃): 215.6, 158.5, 137.0, 133.5, 128.6, 128.2, 128.0, 127.5, 114.8, 74.3, 70.0, 57.5, 42.7, 30.9, 27.8, 24.7. IR (film, cm⁻¹): 3509, 2927, 1697, 1512, 1240, 832. MS (EI) m/z (%): 292 ([M-H₂O]⁺, 0.4), 91 (100), 65 (24.3), 55 (21.9). Anal. Calcd for C₂₀H₂₂O₃: C, 77.39; H, 7.14. Found: C, 77.24; H, 7.23. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 19 °C, 220 nm, 1.0 mL/min; major enantiomer tr = 78.3 min, minor enantiomer tr = 75.1 min.

(2S, 1'R)-2-(Hydroxy(naphthalen-1-yl)methyl)cyclohexan-1-one 6n^[14]

$[\alpha]_D^{24} = +8.8$ ($c = 1.10$, CHCl₃), 89% ee. ¹H NMR (300 MHz, CDCl₃): δ 8.18 (1H, d, $J = 9.0$ Hz), 7.79 (1H, d, $J = 9.0$ Hz), 7.72 (1H, d, $J = 8.4$ Hz), 7.52-7.35 (4H, m), 5.50 (1H, dd, $J = 8.7, 3.0$ Hz), 4.07 (1H, d, $J = 3.3$ Hz), 2.97-2.85 (1H, m), 2.48-2.39 (1H, br-d, $J = 12.9$ Hz), 2.31 (1H, td, J

= 12.9, 6.3 Hz), 2.05-1.94 (1H, m), 1.68-1.50 (2H, m), 1.45-1.23 (3H, m). IR (film, cm^{-1}): 3511, 2928, 1695, 1448, 803. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (4:1 hexane:2-propanol), 20 °C, 254 nm, 0.5 mL/min; major enantiomer t_r = 41.7 min, minor enantiomer t_r = 35.7 min.

(2S, 1'R)-2-(Hydroxy(naphthalen-2-yl)methyl)cyclohexan-1-one 6o^[9]

$[\alpha]_D^{24}$ = +6.7 (c = 1.50, CHCl_3), >99% ee. ^1H NMR (300 MHz, CDCl_3): δ 7.87-7.78 (3H, m), 7.75 (1H, s), 7.47 (3H, d, J = 7.2 Hz), 4.96 (1H, d, J = 8.7 Hz), 4.08 (1H, s), 2.77-2.65 (1H, m), 2.55-2.44 (1H, br-d, J = 13.8 Hz), 2.36 (1H, td, J = 12.9, 5.7 Hz), 2.14-2.00 (1H, m), 1.80-1.36 (5H, m). IR (film, cm^{-1}): 3496, 2931, 1698, 1594, 823. Enantiomeric excess was determined by HPLC with a Chiralpak AS-H column (9:1 hexane:2-propanol), 21 °C, 254 nm, 0.5 mL/min; major enantiomer t_r = 47.6 min, minor enantiomer t_r = 54.8 min.

(2S, 1'R)-2-(Hydroxy-(furan-2-yl)propyl)cyclohexan-1-one 6p^[9]

$[\alpha]_D^{22}$ = -18.3 (c = 0.58, CH_2Cl_2), 81% ee. ^1H NMR (300 MHz, CDCl_3): δ 7.39 (1H, d, J = 0.6 Hz), 6.33 (1H, dd, J = 3.0, 1.8 Hz), 6.28 (1H, d, J = 3.0 Hz), 4.83 (1H, dd, J = 8.4, 3.9 Hz), 3.88 (1H, d, J = 3.9 Hz), 2.97-2.86 (1H, m), 2.52-2.30 (2H, m), 2.17-2.04 (1H, m), 1.89-1.78 (1H, m), 1.75-1.56 (3H, m), 1.44-1.30 (1H, m). IR (film, cm^{-1}): ν 3511, 2926, 1701, 1129, 1007, 740. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 21 °C, 220 nm, 0.5 mL/min; major enantiomer t_r = 41.0 min, minor enantiomer t_r = 43.9 min.

(2S, 1'R)-2-(Hydroxy-(pyridin-4-yl)propyl)cyclohexan-1-one 6q^[9]

$[\alpha]_D^{22}$ = +11.4 (c = 0.40, CHCl_3), 94% ee. ^1H NMR (300 MHz, CDCl_3): δ 8.55 (2H, d, J = 6.0 Hz), 7.25 (2H, d, J = 6.0 Hz), 4.80 (1H, d, J = 8.4 Hz), 4.23 (1H, br-s), 2.65-2.55 (1H, m), 2.52-2.42 (1H, m), 2.35 (1H, td, J = 12.9, 6.9 Hz), 2.16-2.03 (1H, m), 1.81-1.78 (1H, m), 1.78-1.57 (3H, m), 1.56-1.30 (1H, m). IR (film, cm^{-1}): 3205, 2927, 1707, 1601, 1414, 1128. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 19 °C, 254 nm, 1.0 mL/min; major enantiomer t_r = 33.7 min, minor enantiomer t_r = 30.2 min.

(2S, 1'R)-2-(Hydroxy(cyclohexyl)methyl)cyclohexan-1-one 6r^[9]

$[\alpha]_D^{22}$ = +10.0 (c = 0.40, CHCl_3), 96% ee. ^1H NMR (300 MHz, CDCl_3): δ 3.49 (1H, dd, J = 13.2, 6.6 Hz), 3.25 (1H, d, J = 4.5 Hz), 2.57-2.25 (3H, m), 2.19-1.85 (3H, m), 1.81-1.58 (6H, m), 1.56-1.10 (7H, m), 1.01-0.75 (1H, m). IR (film, cm^{-1}): 3539, 2927, 1697, 1449, 1130, 533. Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 20 °C, 280 nm, 0.5 mL/min; major enantiomer t_r = 21.2 min, minor enantiomer t_r = 27.4 min.

(2S, 1'R)-2-(Hydroxy-(*p*-nitrophenyl)methyl)cyclopentan-1-one 6s^[8,9,15]

$[\alpha]_D^{22}$ = -30.6 (c = 0.56, CHCl_3), 98% ee. ^1H NMR (300 MHz, CDCl_3): δ 8.21 (2H, d, J = 8.7 Hz), 7.54 (2H, d, J = 9.0 Hz), 4.85 (1H, d, J = 9.0 Hz), 4.74 (1H, s), 2.54-2.18 (3H, m), 2.08-1.95 (1H, m), 1.81-1.48 (3H, m). IR (film, cm^{-1}): 3442, 2966, 1737, 1604, 1519, 1347, 857 cm^{-1} . Enantiomeric excess was determined by HPLC with a Chiralpak AD column (9:1 hexane:2-propanol), 20 °C, 254 nm, 1.0 mL/min; major enantiomer t_r = 44.3 min, minor

enantiomer tr = 42.4 min.

(2*S*, 1'*R*)-2-(Hydroxy-(*p*-nitrophenyl)methyl)cycloheptan-1-one 6t^[8,16]

$[\alpha]_{\text{D}}^{28} = -8.3$ ($c = 0.35$, CHCl_3), 99% ee. ^1H NMR (300 MHz, CDCl_3): δ 8.21 (2H, d, $J = 9.0$ Hz), 7.53 (2H, d, $J = 8.1$ Hz), 4.92 (1H, dd, $J = 6.9, 5.1$ Hz), 3.74 (1H, d, $J = 4.8$ Hz), 2.98 (1H, m), 2.58-2.40 (2H, m), 1.92-1.65 (4H, m), 1.44-1.20 (4H, m). IR (film, cm^{-1}): 3668, 2930, 1698, 1519, 1347, 972. Enantiomeric excess was determined by HPLC with a Chiralpak AS-H column (4:1 hexane:2-propanol), 19 °C, 254 nm, 0.75 mL/min; major enantiomer tr = 27.8 min, minor enantiomer tr = 33.4 min.

R-4-hydroxy-4-(4-nitrophenyl)butan-2-one 6u^[8,15]

$[\alpha]_{\text{D}}^{28} = +58.8$ ($c = 1.25$, CHCl_3), 87% ee. ^1H NMR (300 MHz, CDCl_3): δ 8.20 (2H, d, $J = 8.7$ Hz), 7.54 (2H, d, $J = 8.7$ Hz), 5.27 (1H, t, $J = 6.0$ Hz), 3.67 (1H, s), 2.86 (2H, d, $J = 6.6$ Hz), 2.22 (3H, s). IR (film, cm^{-1}): 3441, 2922, 1712, 1518, 1347, 857 cm^{-1} . Enantiomeric excess was determined by HPLC with a Chiralpak OJ column (4:1 hexane:2-propanol), 17 °C, 254 nm, 0.75 mL/min; major enantiomer tr = 29.8 min, minor enantiomer tr = 33.5 min.

(2*S*, 3*R*)-4-hydroxy-3-methyl-4-(4-nitrophenyl)butan-2-one 6v^[13,15]

$[\alpha]_{\text{D}}^{28} = +36.1$ ($c = 0.45$, CHCl_3), 89% ee. ^1H NMR (300 MHz, CDCl_3): δ 8.22 (2H, d, $J = 8.7$ Hz), 7.52 (2H, d, $J = 8.7$ Hz), 4.88 (1H, m), 3.29 (1H, s), 2.91 (1H, m), 2.22 (3H, s), 1.03 (3H, d, $J = 7.2$ Hz). IR (film, cm^{-1}): 3523, 2924, 1705, 1520, 1347, 856. Enantiomeric excess was determined by HPLC with a Chiralpak OJ column (4:1 hexane:2-propanol), 17 °C, 254 nm, 0.75 mL/min; major enantiomer tr = 30.1 min, minor enantiomer tr = 33.4 min.

List of the obvious difference between **syn-6** and **anti-6** on ^1H NMR ^[8]

entry	R ¹	R ²	R ³	product	^1H NMR(-CH $\underline{\text{O}}$ H)	
					syn	anti
1	4-O ₂ NC ₆ H ₄ -	—(CH ₂) ₄ —		6a	5.49 (d, J = 1.8 Hz)	4.90 (dd, J = 8.4, 3.0 Hz)
2	3-O ₂ NC ₆ H ₄ -	—(CH ₂) ₄ —		6b	5.48 (d, J = 2.1 Hz)	4.90 (dd, J = 8.7, 3.0 Hz)
3	2-O ₂ NC ₆ H ₄ -	—(CH ₂) ₄ —		6c	5.96 (d, J = 2.1 Hz)	5.45 (d, J = 6.6 Hz)
4	4-F ₃ CC ₆ H ₄ -	—(CH ₂) ₄ —		6d	5.81 (d, J = 1.8 Hz)	5.30 (d, J = 9.3 Hz)
5	4-NCC ₆ H ₄ -	—(CH ₂) ₄ —		6e	5.43 (brs)	4.85 (dd, J = 8.1, 3.0 Hz)
6	4-FC ₆ H ₄ -	—(CH ₂) ₄ —		6f	5.36 (brs)	4.00 (d, J = 3.6 Hz)
7	4-ClC ₆ H ₄ -	—(CH ₂) ₄ —		6g	5.35 (d, J = 1.8 Hz)	4.76 (dd, J = 8.7, 2.7 Hz)
8	4-BrC ₆ H ₄ -	—(CH ₂) ₄ —		6h	5.34 (d, J = 2.4 Hz)	4.75 (dd, J = 8.7, 2.7 Hz)
9	2,4-Cl ₂ C ₆ H ₃ -	—(CH ₂) ₄ —		6i	5.65 (d, J = 1.5 Hz)	5.29 (dd, J = 7.5, 3.3 Hz)
10	C ₆ H ₅ -	—(CH ₂) ₄ —		6j	5.39 (d, J = 2.4 Hz)	4.80 (d, J = 9.0 Hz)
11	4-MeC ₆ H ₄ -	—(CH ₂) ₄ —		6k	5.35 (d, J = 2.1 Hz)	4.75 (dd, J = 9.0, 2.7 Hz)
12	4-MeOC ₆ H ₄ -	—(CH ₂) ₄ —		6l	5.32 (d, J = 1.8 Hz)	4.74 (dd, J = 8.7, 2.4 Hz)
13	4-BnOC ₆ H ₄ -	—(CH ₂) ₄ —		6m	5.31 (d, J = 2.7 Hz)	4.74 (dd, J = 9.0, 2.7 Hz)
14	1-naphthyl-	—(CH ₂) ₄ —		6n	6.15 (brs)	5.50 (dd, J = 8.7, 3.0 Hz)
15	2-naphthyl-	—(CH ₂) ₄ —		6o	5.46 (d, J = 3.0 Hz)	4.96 (d, J = 8.7 Hz)
16	2-furanyl-	—(CH ₂) ₄ —		6p	5.26 (d, J = 3.0 Hz)	3.88 (d, J = 3.9 Hz)
17	4-pyridinyl-	—(CH ₂) ₄ —		6q	5.38 (brs)	4.80 (d, J = 8.4 Hz)
18	cyclohexyl-	—(CH ₂) ₄ —		6r	3.79 (d, J = 9.3 Hz)	3.49 (dd, J = 13.2, 6.6 Hz)
19	4-O ₂ NC ₆ H ₄ -	—(CH ₂) ₃ —		6s	5.42 (brs)	4.85 (d, J = 9.0 Hz)
20	4-O ₂ NC ₆ H ₄ -	—(CH ₂) ₅ —		6t	5.31 (brs)	4.92 (dd, J = 6.9, 5.1 Hz)
21	4-O ₂ NC ₆ H ₄ -	CH ₃	H	6u	-----	-----
22	4-O ₂ NC ₆ H ₄ -	CH ₃	CH ₃	6v	5.54 (m)	4.88 (m)

NMR spectra for catalysts **1a-f** and **2**

¹H NMR spectra of **1a**

¹H NMR spectra of **1b**

¹H NMR spectra of **1c**

¹H NMR spectra of **1d**

¹H NMR spectra of **1e**

¹H NMR spectra of **1f**

¹H NMR spectra of **2**

¹³C NMR spectra of **2**

NMR spectra for compounds **6a-v**

¹H NMR spectra of **6a**

¹H NMR spectra of **6b**

¹H NMR spectra of **6c**

¹H NMR spectra of **6d**

¹H NMR spectra of **6e**

¹H NMR spectra of **6f**

¹H NMR spectra of **6g**

¹H NMR spectra of **6h**

¹H NMR spectra of **6i**

¹H NMR spectra of **6j**

¹H NMR spectra of **6k**

¹H NMR spectra of **6l**

¹H NMR spectra of **6m**

¹³C NMR spectra of **6m**

¹H NMR spectra of **6n**

¹H NMR spectra of **6o**

¹H NMR spectra of **6p**

¹H NMR spectra of **6q**

¹H NMR spectra of **6r**

¹H NMR spectra of **6s**

¹H NMR spectra of **6t**

¹H NMR spectra of **6u**

¹H NMR spectra of **6v**

HPLC spectra for compounds **6a-v**

HPLC spectra of **6a**

Peak Results

	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		41.732	197801	0.19		908	0.07	
2		52.677	102131840	99.81		1286584	99.93	

HPLC spectra of **6b**

Peak Results

	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		59.696	249710699	99.79		2676666	99.72	
2		75.135	526330	0.21		7425	0.28	

HPLC spectra of **6c**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	59.404	160380096	99.70		2026769	99.56	
2	62.983	479420	0.30		8996	0.44	

HPLC spectra of **6d**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	32.952	10487716	99.30		242485	99.14	
2	35.251	73598	0.70		2103	0.86	

HPLC spectra of **6e**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	37.921	75654	1.03		1954	1.79	
2	46.571	7271856	98.97		107476	98.21	

HPLC spectra of **6f**

Peak Results

	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		39.595	2176250	3.00		53814	5.16	
2		44.244	70276113	97.00		988900	94.84	

HPLC spectra of **6g**

Peak Results

	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		26.579	124225	0.69		3287	1.05	
2		31.823	17967800	99.31		310153	98.95	

HPLC spectra of **6h**

Peak Results								
	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		45.365	243688	0.26		6738	0.53	
2		54.393	94546548	99.74		1260524	99.47	

HPLC spectra of **6i**

Peak Results

	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		23.229	1363466	1.51		58883	3.11	
2		27.064	68862644	98.49		1832992	96.89	

HPLC spectra of **6j**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	22.365	3506334	97.98		101399	98.10	
2	31.959	72464	2.02		1964	1.90	

HPLC spectra of **6k**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	27.670	160533082	97.51		2120232	97.28	
2	34.671	4106680	2.49		59218	2.72	

HPLC spectra of **6l**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	60.922	1140154	1.97		14286	2.29	
2	64.228	56751209	98.03		610719	97.71	

HPLC spectra of **6m**

Peak Results							
Name	RT	Area	% Area	Amount	Height	% Height	Units
1	49.816	271818	0.13		4827	0.28	
2	56.901	222389	0.11		4382	0.25	
3	75.117	139498	0.07		2013	0.12	
4	78.314	205341872	99.69		1718004	99.35	

HPLC spectra of **6n**

Peak Results

	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		35.723	1003876	5.38		14200	4.40	
2		41.654	17665444	94.62		308702	95.60	

HPLC spectra of **6o**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	47.553	70296028	99.85		839579	99.76	
2	54.757	105168	0.15		2054	0.24	

HPLC spectra of **6p**

Peak Results

	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		41.007	12181159	90.50		286707	92.08	
2		43.865	1279046	9.50		24665	7.92	

HPLC spectra of **6q**

Peak Results

	Name	RT	Area	% Area	Amount	Height	% Height	Units
1		30.229	223744	3.08		6760	5.63	
2		33.690	7029101	96.92		113221	94.37	

HPLC spectra of **6r**

Peak Results							
	Name	RT	Area	% Area	Amount	Height	% Height
1		21.243	582118	98.15		18553	98.11
2		27.384	10999	1.85		358	1.89

HPLC spectra of **6s**

Peak Results							
Name	RT	Area	% Area	Amount	Height	% Height	Units
1	24.153	7058717	25.27		186400	35.52	
2	31.865	7027911	25.16		133886	25.51	
3	42.428	6720595	24.06		107667	20.51	
4	44.156	7126910	25.51		96895	18.46

Peak Results							
Name	RT	Area	% Area	Amount	Height	% Height	Units
1	42.450	29965	1.12		703	1.95	
2	44.277	2633893	98.88		35404	98.05	

HPLC spectra of **6t**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	27.882	13936121	49.63		295428	65.03	
2	31.791	14146583	50.37		158887	34.97

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	27.846	27872242	99.36		590853	98.74	
2	32.382	178628	0.64		7542	1.26	

HPLC spectra of **6u**

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	28.319	67614737	49.79		959850	53.22	
2	31.436	68190809	50.21		843704	46.78

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	29.751	54579405	93.39		744821	90.75	
2	33.467	3860903	6.61		75928	9.25	

HPLC spectra of **6v**

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	28.319	67614737	49.79		959850	53.22	
2	31.436	68190809	50.21		843704	46.78

Peak Results

Name	RT	Area	% Area	Amount	Height	% Height	Units
1	30.051	53849262	94.57		743091	90.59	
2	33.417	3093136	5.43		77183	9.41	

References

- [1] Cobb, A. J. A.; Shaw, A. M.; Longbottom, D. A.; Gold, J. B.; Ley, S. V. *Org. Biomol. Chem.* **2005**, *3*, 84.
- [2] Sundén, H.; Ibrahem, I.; Eriksson, L.; Córdova, A. *Angew. Chem. Int. Ed.* **2005**, *44*, 4877.
- [3] Bellis, E.; Vasilatou, K.; Kokotos, G. *Synthesis* **2005**, *14*, 2407.
- [4] Sundén, H.; Dahlin, N.; Ibrahem, I.; Adolfsson, H.; Córdova, A. *Tetrahedron Letters* **2005**, *46*, 3385.
- [5] Berkessel, A.; Koch, B.; Lex, J. *Adv. Synth. Catal.* **2004**, *346*, 1141.
- [6] Tangallapally, R. P.; Yendapally, R.; Lee, R. E.; Lenaerts, A. J. M.; Lee, R. E. *J. Med. Chem.* **2005**, *48*, 8261.
- [7] Fan, Q. H.; Chen, Y. M.; Chen, X. M.; Jiang, D. Z.; Xi, F.; Chan, A. S. C. *Chem. Commun.* **2000**, 789.
- [8] Mase, N.; Nakai, Y.; Ohara, H.; Takabe, K.; Tanaka, F.; Barbas, C. F. *J. Am. Chem. Soc.* **2006**, *128*, 734.
- [9] Hayashi, Y.; Sumiya, T.; Takahashi, J.; Gotoh, H.; Urushima, T.; Shoji, M. *Angew. Chem. Int. Ed.* **2006**, *45*, 958.
- [10] Majewski, M.; Gleave, D. M. *Tetrahedron Letters* **1989**, *30*, 5681.
- [11] Flowers II, R. A.; Xu, X.; Timmons, C.; Li, G. *Eur. J. Org. Chem.* **2004**, 2988.
- [12] Ji, C.; Peng, Y.; Huang, C.; Wang, N.; Jiang, Y. *Synlett* **2005**, *6*, 986
- [13] Chen, J. R.; Lu, H. H.; Li, X. Y.; Cheng, L.; Wan, J.; and Xiao W. J. *Org. Lett.* **2005**, *7*, 4543
- [14] Yanagisawa, A.; Nakatsuka, Y.; Asakawa, K.; Wadamoao, M.; Kageyama, H.; Yamamoto, H. *Bnll. Chem. Soc. Jpn.* **2001**, *74*, 1477
- [15] Tang, Z.; Yang, Z.-H.; Chen, X.-H.; Cun, L.-F.; Mi, A.-Q.; Jiang, Y.-Z.; Gong, L.-Z. *J. Am. Chem. Soc.* **2005**, *127*, 9285.
- [16] Ranu, B. C.; Chakraborty, R. *Tetrahedron* **1993**, *49*, 5333.