Supporting information for:

**Highly Enantioselective Hydrovinylation of α -Alkyl Vinylarenes.
A New Approach to the Construction of All-Carbon Quarternary
Stereocenters**

Wen-Jian Shi, Qi Zhang, Jian-Hua Xie, Shou-Fei Zhu, Guo-Hua Hou,
and Qi-Lin Zhou*

*State Key Laboratory and Institute of Elemento-organic Chemistry, Nankai University,
Tianjin 300071, China.*

CONTENTS:

- (A) Preparation and physical data of new substrates α -alkyl vinylarenes.**
- (B) General procedure for the preparation of the precatalysts [Ni(allyl)Br(ligand)].**
- (C) General procedure for asymmetric hydrovinylation of α -alkyl vinylarenes.**
- (D) Representative examples, reaction conditions and analytical data of hydrovinylation products.**

General remarks. The reactions and manipulations were performed under an atmosphere of nitrogen by using standard Schlenk techniques and a Vacuum Atmospheres Drybox (VAC DRI-LAB HE 493). ^1H and ^{13}C NMR spectra were recorded on Bruker-300 and Varian-400 spectrometers. Chemical shifts were reported in ppm down field from internal Me_4Si . Optical rotations were determined using a Perkin Elmer Model 341 polarimeter. HRMS were recorded on Bruker APEX II

spectrometer. GC analyses were performed using Hewlett Packard Model HP 6890 Series. HPLC analyses were performed using a Hewlett Packard Model HP 1100 Series. A HP G1800A GCD System was used for GC-MS analysis. Methylene chloride and 1,2-dichloroethane were distilled from calcium hydride under nitrogen before use. Toluene was distilled under nitrogen from sodium/benzophenone ketyl. Ethylene (99.9%) was purchased from BAPB Gases Industry Company, Ltd., Beijing, China. The α -substituted vinylarenes were prepared from the corresponding aromatic ketones according to the literature methods, unless otherwise stated.¹ Ni(COD)₂, AgX (X = BF₄⁻, OTf, ClO₄⁻, PF₆⁻ and SbF₆⁻) and allylic bromide were purchased from Aldrich chemical company and used without further purification. The NaBARF was prepared according to the literature.² Ligands **4**, **7a** and **8** can be obtained from Strem Chemicals. Ligands **5**, **6**, **7b** and **7c** were prepared by previous methods.³ The chiral monophosphoramidites **3** was synthesized using the literature procedure.⁴

(A) Preparation and physical data of new substrates α -alkyl vinylarenes

α -Isopropyl *m*-methylstyrene (**9c**)

Typical procedure: Under a nitrogen atmosphere, to the mixture of PPh₃MeI (3.0 g, 0.75 mmol) and *m*-methylphenyl isopropyl ketone (811 mg, 0.5 mmol) in 10 mL of ether, KO^tBu (842 mg, 0.75 mmol) was added in one portion at room temperature. The suspension was stirred overnight, filtered. The filtrate was concentrated and purified by silica gel column chromatography with *n*-hexane to afford 760 mg (95% yield) of **9c** as a colorless liquid.

^1H NMR (CDCl_3 , 300 MHz): δ 7.25–7.07 (m, 4H, Ar-H), 5.12 (s, 1H, C=CH), 5.01 (s, 1H, C=CH), 2.89–2.75 (m, 1H, CH), 2.36 (s, 3H, Ar-CH₃), 1.09 (d, J = 6.9 Hz, 6H, CH₃); ^{13}C NMR (CDCl_3 , 75 MHz): δ 156.2, 143.1, 137.8, 128.2, 128.0, 127.6, 123.9, 110.0, 32.6, 22.3, 21.7; HRMS (EI) Calcd for $\text{C}_{12}\text{H}_{16}$: 160.1252. Found: 160.1251.

α -Isopropyl-*m*-methoxystyrene (**9d**)

α -Isopropyl *m*-methoxystyrene (**9d**) was prepared by the same procedure for the preparation of **9c** from *m*-methoxyphenyl isopropyl ketone. Colorless liquid, 96% yield.

^1H NMR (CDCl_3 , 300 MHz): δ 7.26–7.21 (m, 1H, Ar-H), 6.95–6.82 (m, 2H, Ar-H),

6.81–6.80 (m, 1H, Ar-H), 5.15 (s, 1H, C=CH), 5.03 (s, 1H, C=CH), 3.82 (s, 3H, OCH₃), 2.88–2.77 (m, 1H, CH), 1.09 (d, *J* = 6.9 Hz, 6H, CH₃); ¹³C NMR (CDCl₃, 75 MHz): δ 159.7, 155.9, 144.7, 129.3, 119.4, 112.8, 112.5, 110.3, 55.4, 32.6, 22.2; HRMS (EI) Calcd for C₁₂H₁₆O: 176.1201. Found: 176.1199.

α-Cyclohexyl-2'-vinyl-naphthalene (9k)

α-Cyclohexyl 2'-vinyl-naphthalene (**9k**) was prepared by the same procedure for the preparation of **9c** from cyclohexyl 2-naphthyl ketone. Colorless liquid, 96% yield.

¹H NMR (CDCl₃, 300 MHz): δ 7.80–7.76 (m, 4H, Ar-H), 7.51–7.25 (m, 3H, Ar-H), 5.26 (s, 1H, C=CH), 5.10 (s, 1H, C=CH), 2.59–2.52 (m, 1H, CH), 1.90–1.77 (m, 5H,

CH₂), 1.34–1.19 (m, 5H, CH₂); ¹³C NMR (CDCl₃, 75 MHz): δ 155.2, 140.6, 133.6, 132.9, 128.3, 127.8, 126.2, 125.8, 127, 125.2, 111.1, 42.8, 33.0, 27.1, 26.7; HRMS (EI) Calcd for C₁₈H₂₀: 236.1565. Found: 236.1562.

(B) General procedure for preparation of precatalysts [Ni(allyl)Br(ligand)] ⁵

Typical Procedure: Under a nitrogen atmosphere, Ni(COD)₂ (137.5 mg, 0.5 mmol) was suspended in 1, 5-cyclooctadiene (1 mL). Allylic bromide (63.5 mg, 0.525 mmol) was then added dropwise at 20 °C. The resulting slurry was stirred for 5–10 mins, forming a dark blood-red allylnickel(II) bromide dimer. Then, 2 mL of toluene was added to dissolve the solid materials. A solution of chiral monodentate phosphorous ligands **3–8** (0.5 mmol) in 4 mL toluene was added and stirred for another 2.0 hrs. The solution rapidly changed from maroon color to orange-red. The mixture was filtered through Celite under nitrogen atmosphere and the filtrate was concentrated *in vacuo*.

The red solid was dried *in vacuo* for 10 hrs to give precatalysts in 96–98% yield. The precatalysts were used directly for hydrovinylation without further purification.

(C) General procedure for asymmetric hydrovinylation.

Standard procedure: Under nitrogen atmosphere, to the suspension of NaBARF (21.0 mg, 0.021 mmol) in dichloromethane (2 mL), an orange-red solution of precatalyst [Ni(allyl)Br(*S_a,R,R*)-**8**] (13.8 mg, 0.02 mmol) in dichloromethane (2 mL) was added at room temperature, and stirred for 20 min. Then, ethylene was introduced into the resulting yellow solution, and α -alkyl vinylarene (0.2 mmol) was added dropwise with a microsyringe. The resulting reaction mixture was stirred for the desired time at 35 °C under an ethylene atmosphere (~1 atm). The reaction was quenched by addition of a saturated aqueous NH₄Cl solution, and the product was extracted with Et₂O, dried with MgSO₄ and purified by a short silica gel column chromatography. The conversion and chemoselectivity of product were determined by GC (column HP-5, 30 m) from the peak areas of **10** and all detectable products. This procedure was verified in selected reactions using tridecane as internal standard. The ee values of products were determined by chiral GC or HPLC with the conditions specified in the individual compounds.

(*S*)-3,4-Dimethyl-3-phenyl-1-pentene (10a) ⁶

¹H NMR (CDCl₃, 300 MHz): δ 7.35–7.14 (m, 5H, Ar-H), 6.16 (dd, J = 17.6, 11.0 Hz, C=CH), 5.17 (dd, J = 11.0, 1.5 Hz, 1H, C=CH), 5.05 (dd, J = 17.6, 1.5 Hz, 1H, C=CH), 2.26–2.12 (spt, 1H, CH), 1.32 (s, 3H, CH₃), 0.83 (d, J = 6.6 Hz, 3H, CH₃), 0.75 (d, J =

6.6 Hz, 3H, CH₃).

The ee of **10a** was determined by GC analysis using a Supelco β -DEXTM 120 capillary column (30 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 85 °C; rate, 0.5 °C/min; final temperature, 120 °C; *t_R* of (*R*)-isomer, 63.00 min; *t_R* of (*S*)-isomer, 63.66 min.

The *S* configuration of **10a** was determined by two different ways. One way is to compare the retention times of enantiomers of **10a** with those reported by Hoveyda using same type of chiral column (β -DEX) on GC, the enantiomer with a longer retention time is *S*.⁷ Another way is to oxidize **10a** to 2,3-dimethyl-2-phenyl-1-butyric acid and compare its optical rotation with that reported in literature.⁷

3,4-Dimethyl-3-(*p*-methylphenyl)-1-pentene (10b)

^1H NMR (CDCl_3 , 300 MHz) δ 7.22 (d, 2H, $J = 8.1$ Hz, Ar-H), 7.10 (d, 2H, $J = 8.1$ Hz, Ar-H), 6.14 (dd, 1H, $J = 17.7, 10.8$ Hz, C=CH), 5.14 (dd, 1H, $J = 10.8, 1.2$ Hz, C=CH),

5.03 (dd, 1H, $J = 17.7, 1.4$ Hz, C=CH), 2.31 (s, 3H, CH₃), 2.23–2.10 (spt, 1H, CH), 1.30 (s, 3H, CH₃), 0.84 (d, 3H, $J = 6.9$ Hz, CH₃), 0.75 (d, 3H, $J = 6.9$ Hz, CH₃); ¹³C NMR (CDCl₃, 75 MHz) δ 148.5, 145.0, 128.5, 126.5, 125.5, 113.2, 47.5, 36.3, 22.0, 20.7, 18.4, 18.0; HRMS (EI) Calcd for C₁₄H₂₀: 188.1565. Found: 188.1563.

The ee of **10b** was determined by GC analysis using Supelco β -DEXTM 120 capillary column (30 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 85 °C; rate, 0.5 °C/min; final temperature, 150 °C; t_R of minor isomer, 82.64 min; t_R of major isomer, 83.38 min.

3,4-Dimethyl-3-(*m*-methylphenyl)-1-pentene (10c)

^1H NMR (CDCl_3 , 300 MHz) δ 7.29–7.13 (m, 4H, Ar-H), 6.18 (dd, $J = 17.4$, 10.8 Hz, 1H, C=CH), 5.19 (dd, $J = 10.8$, 1.4 Hz, 1H, C=CH), 5.08 (dd, $J = 17.4$, 1.4 Hz, 1H, C=CH), 2.36 (s, 3H, CH_3), 2.28–2.14 (spt, 1H, CH), 1.34 (s, 3H, CH_3), 0.87 (d, $J = 6.9$

Hz, 3H, CH₃), 0.79 (d, *J* = 6.9 Hz, 3H, CH₃); ¹³C NMR (CDCl₃, 75 MHz) δ 145.5, 145.2, 135.1, 128.9, 126.7, 113.1, 47.3, 36.3, 21.1, 20.7, 18.4, 18.0; HRMS (EI) Calcd for C₁₄H₂₀: 188.1565. Found: 188.1564.

The ee of **10c** was determined by GC analysis using Varian CP-Chirasil-DexTM CB capillary column (25 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 85 °C; rate, 0.5 °C/min; final temperature, 160 °C; *t*_R of minor isomer, 71.56 min; *t*_R of major isomer, 71.99 min.

3,4-Dimethyl-3-(*m*-methoxyphenyl)-1-pentene (10d)

¹H NMR (CDCl₃, 400 MHz) δ 7.27–7.20 (m, 1H, Ar-H), 6.94–6.88 (m, 2H, Ar-H), 6.70 (m, 1H, Ar-H), 6.13 (dd, J = 17.6, 10.8 Hz, 1H, C=CH), 5.16 (d, J = 10.8 Hz, 1H, C=CH), 5.05 (d, J = 17.6 Hz, 1H, C=CH), 3.80 (s, 3H, OCH₃), 2.21–2.11 (spt, 1H, CH),

1.30 (s, 3H, CH₃), 0.83 (d, *J* = 6.8 Hz, 3H, CH₃), 0.75 (d, *J* = 6.8 Hz, 3H, CH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 159.6, 150.3, 144.7, 129.0, 119.4, 113.6, 113.4, 110.3, 55.3, 47.7, 36.4, 20.7, 18.4, 17.9; HRMS (EI) Calcd for C₁₄H₂₀O: 204.1514. Found: 204.1512.

The ee of **10d** was determined by GC analysis using Varian CP-Chirasil-DexTM CB capillary column (25 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 70 °C; rate, 0.5 °C/min; final temperature, 135 °C; *t*_R of minor isomer, 120.74 min; *t*_R of major isomer, 121.32 min.

3,4-Dimethyl-3-(*p*-methoxyphenyl)-1-pentene (10e)

^1H NMR (CDCl_3 , 300 MHz) δ 7.25 (d, 2H, $J = 8.7$ Hz, Ar-H), 6.83 (d, 2H, $J = 8.7$ Hz, Ar-H), 6.13 (dd, $J = 17.4$, 10.8 Hz, 1H, C=CH), 5.15 (dd, $J = 10.8$, 1.2 Hz, 1H, C=CH), 5.03 (dd, $J = 17.4$, 1.2 Hz, 1H, C=CH), 3.79 (s, 3H, OCH_3), 2.19–2.09 (spt, 1H, CH),

1.29 (s, 3H, CH₃), 0.83 (d, *J* = 6.6 Hz, 3H, CH₃), 0.75 (d, *J* = 6.6 Hz, 3H, CH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 157.6, 145.3, 140.5, 127.8, 113.5, 113.0, 55.4, 47.0, 36.4, 29.9, 18.3, 18.0; HRMS (EI) Calcd for C₁₄H₂₀O: 204.1514. Found: 204.1513.

The ee of **10e** was determined by GC analysis using Varian CP-Chirasil-DexTM CB capillary column (25 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 70 °C; rate, 0.5 °C/min; final temperature, 150 °C; *t*_R of minor isomer, 132.38 min; *t*_R of major isomer, 133.22 min.

3,4-Dimethyl-3-(*p*-chlorophenyl)-1-pentene (10f)

^1H NMR (CDCl_3 , 400 MHz) δ 7.26–7.25 (m, 4H, Ar-H), 6.10 (dd, $J = 17.6, 10.8$ Hz, 1H, C=CH), 5.18 (d, $J = 10.8$ Hz, 1H, C=CH), 5.04 (d, $J = 17.6$ Hz, 1H, C=CH), 2.18–2.08 (spt, 1H, CH), 1.29 (s, 3H, CH_3), 0.81 (d, $J = 6.8$ Hz, 3H, CH_3), 0.74 (d, $J =$

6.8 Hz, 3H, CH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 146.7, 144.2, 131.3, 128.1, 128.0, 113.6, 47.2, 36.2, 30.9, 18.0, 17.7; HRMS (EI) Calcd for C₁₃H₁₇Cl: 208.1019. Found: 208.1017.

The ee of **10f** was determined by GC analysis using Supelco β-DEXTM 120 capillary column (30 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 100 °C; rate, 1 °C/min; final temperature, 190 °C; *t*_R of minor isomer, 55.30 min; *t*_R of major isomer, 55.70 min.

(S)-3-Methyl-3-phenyl-1-pentene (10g) ⁷

¹H NMR (CDCl₃, 300 MHz) δ 7.33–7.15 (m, 5H, Ar-H), 6.02 (dd, J = 17.4, 10.8 Hz, 1H, C=CH), 5.10 (dd, J = 10.8, 1.2 Hz, 1H, C=CH), 5.03 (dd, J = 17.4, 1.2 Hz, 1H, C=CH), 1.88-1.72 (m, 2H, CH₂), 1.35 (s, 3H, CH₃), 0.77 (t, J = 7.4 Hz, 3H, CH₃).

The ee of **10g** was determined by GC analysis using Varian CP-Chirasil-DexTM CB

capillary column (25 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 75 °C; rate, 0.5 °C/min; final temperature, 120 °C; *t_R* of minor isomer, 65.26 min; *t_R* of major isomer, 66.31 min.

The absolute configuration of **10g** was determined by comparing the retention times of enantiomers with the data in literature using same chiral column (β-DEX 120).⁷

3-Methyl-3-phenyl-1-hexene (10h)

^1H NMR (CDCl_3 , 300 MHz) δ 7.33-7.14 (m, 5H, Ar-H), 6.03 (dd, J = 17.4, 10.8 Hz, 1H, C=CH), 5.08 (dd, J = 10.8, 1.4 Hz, 1H, C=CH), 5.03 (dd, J = 17.4, 1.4 Hz, 1H, C=CH), 1.81-1.64 (m, 2H, CH_2), 1.34 (s, 3H, CH_3), 1.28-1.03 (m, 2H, CH_2), 0.86 (t, J = 7.2 Hz, 3H, CH_3); ^{13}C NMR (CDCl_3 , 75 MHz) δ 147.8, 147.2, 128.0, 126.6, 125.7,

111.5, 44.4, 43.6, 25.0, 17.8, 14.8; HRMS (EI) Calcd for C₁₃H₁₈: 174.1409. Found: 174.1406.

The ee of **10h** was determined by GC analysis using Varian CP-Chirasil-DexTM CB capillary column (25 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 70 °C; rate, 0.25 °C/min; final temperature, 100 °C; *t*_R of minor isomer, 104.02 min; *t*_R of major isomer, 104.72 min.

3,5-Dimethyl-3-phenyl-1-hexene (10i)

^1H NMR (CDCl_3 , 300 MHz) δ 7.32–7.16 (m, 5H, Ar-H), 6.12–6.02 (dd, 1H, $J = 17.4$, 10.8 Hz), 5.05 (m, 2H, C=CH), 1.73–1.58 (m, 3H, CH_2 and CH), 1.40 (s, 3H, CH_3), 0.82 (d, $J = 6.6$ Hz, 3H, CH_3), 0.73 (d, $J = 6.6$ Hz, 3H, CH_3); ^{13}C NMR (CDCl_3 , 75 MHz) δ 148.0, 147.6, 127.9, 126.7, 125.7, 111.3, 50.2, 44.6, 25.3, 25.1, 24.9, 24.7; HRMS (EI)

Calcd for C₁₄H₂₀: 188.1565. Found: 188.1567.

The ee of **10i** was determined by GC analysis using Varian CP-Chirasil-DexTM CB capillary column (25 m). Conditions: carrier gas, N₂ (0.4 mL/min); inject temperature, 230 °C; initial temperature, 75 °C; rate, 0.5 °C/min; final temperature, 140 °C; *t_R* of minor isomer, 80.59 min; *t_R* of major isomer, 81.02 min.

3,4-Dimethyl-3-(2'-naphthalenyl)-1-pentene (10j)

^1H NMR (CDCl_3 , 300 MHz) δ 7.81–7.41 (m, 7H, Ar-H), 6.26 (dd, $J = 17.4$, 10.8 Hz, 1H, C=CH), 5.23 (d, $J = 10.8$ Hz, 1H, C=CH), 5.1 (d, $J = 17.4$ Hz, 1H, C=CH), 2.39–2.25 (spt, 1H, CH), 1.42 (s, 3H, CH_3), 0.87 (d, $J = 6.6$ Hz, 3H, CH_3), 0.78 (d, $J =$

6.6 Hz, 3H, CH₃); ¹³C NMR (CDCl₃, 75 MHz) δ 145.9, 144.8, 133.6, 132.0, 128.2, 127.7, 127.5, 125.9, 125.7, 125.5, 125.2, 113.7, 47.9, 36.1, 20.8, 18.4, 18.0; HRMS (EI) Calcd for C₁₇H₂₀: 224.1565. Found: 224.1562.

The ee of **10j** was determined by HPLC analysis using Daicel Chiralcel OJ column (25 cm × 0.46 cm ID) using *n*-hexane as an eluent. Conditions: UV detector, 254 nm; 0.3 mL/min; *t*_R of major isomer, 26.04 min; *t*_R of minor isomer, 36.26 min.

3-Cyclohexyl-3-(2'-naphthyl)-1-butene (10k)

^1H NMR (CDCl_3 , 300 MHz) δ 7.79-7.39 (m, 7H, Ar-H), 6.28 (dd, $J = 17.7, 11.1$ Hz,

1H, C=CH), 5.2 (d, $J = 11.1$ Hz, 1H, C=CH), 5.07 (d, $J = 17.7$ Hz, 1H, C=CH), 1.89–1.81 (m, 1H, CH), 1.73–0.87 (m, 10H, CH₂), 1.44 (s, 1H, CH₃); ¹³C NMR (CDCl₃, 75 MHz) δ 145.9, 145.4, 133.7, 131.9, 128.2, 127.7, 127.5, 126.0, 125.7, 125.5, 125.2, 113.4, 47.7, 47.3, 28.5, 28.2, 27.4, 27.3, 27.0, 21.1; HRMS (EI) Calcd for C₂₀H₂₄: 264.1878 Found: 264.1880.

The ee of **10k** was determined by HPLC analysis using Daicel Chiralcel OJ column (25 cm $\times$ 0.46 cm ID) using *n*-hexane as an eluent. Conditions: UV detector, 254 nm; 0.3 mL/min; t_R of major isomer, 22.49 min; t_R of minor isomer, 26.83 min.

Reference

- (1) Ohta, T.; Ikegami, H.; Miyake, T.; Takaya, H. *J. Organomet. Chem.* **1995**, 502, 169-176.
- (2) Brookhart, M.; Grant, B.; Volpe, A. *Organometallics*, **1992**, 11, 3920-3922.
- (3) (a) Zhou, H.; Wang, W.-H.; Fu, Y.; Xie, J.-H.; Shi, W.-J.; Wang, L.-X.; Zhou, Q.-L. *J. Org. Chem.* **2003**, 68, 1582-1584. (b) Fu, Y.; Hou, G.-H.; Xie, J.-H.; Xing, L.; Wang, L.-X.; Zhou, Q.-L. *J. Org. Chem.* **2004**, 69, 8157-8160. (c) Zhu, S.-F.; Yang, Y.; Wang, L.-X.; Liu, B.; Zhou, Q.-L. *Org. Lett.* **2005**, 7, 2333-2335.
- (4) Arnold, L. A.; Imbos, R.; Mandoli, I.; Vries, A. H. M. de; Naasz, R.; Feringa, B. L. *Tetrahedron* **2000**, 56, 2865-2878.
- (5) Dible, B. R.; Sigman, M. S. *J. Am. Chem. Soc.* **2003**, 125, 872-873.
- (6) van Veldhuizen, J. J.; Campbell, J.-E.; Giudici, R.-E.; Hoveyda, A. H. *J. Am. Soc. Chem.* **2005**, 127, 6877-6882.
- (7) Calas, M.; Calas, B.; Giral, L. *Bull. Soc. Chim. Fr.* **1976**, 857-861.