Supporting Information

Highly enantioselective organocatalytic direct aldol reaction in aqueous medium **

Vishnu Maya, Monika Raj and Vinod K. Singh*

Department of Chemistry, Indian Institute of Technology, Kanpur 208016, India

E-mail: vinodks@iitk.ac.in

General Methods	2
General procedure for enantioselective Aldol reaction	2
HPLC data for chiral Aldol products	3-33
¹ H NMR spectra of Aldol products	33-55
References	55

Experimental

General methods: ^1H NMR spectra were recorded on 400 MHz spectrometer. Chemical shifts are expressed in ppm downfield from TMS as internal standard, and coupling constants are reported in Hz. Routine monitoring of reaction was performed by TLC, using precoated silica gel TLC plates obtained by E-Merck. IUPAC names of ligands and aldol product were taken from Chem. ultra 8. All the column chromatography separations were done by using silica gel (Acme's, 100-200 mesh). Petroleum ether used was of boiling range 60-80 °C. The organic extracts were dried over *anhydrous* sodium sulfate. Evaporation of solvent was performed at reduced pressure. Brine refers to saturated solution of NaCl in water at 25 °C.

A representative procedure for the direct aldol reaction with the catalyst **1 in brine:**

An aldehyde (0.5 mmol) was added to a mixture of ketone (2 mmol) and an organocatalyst **1** (0.5 mol %) in brine (0.5 mL) at -5 °C. The reaction mixture was stirred and the progress of the reaction was monitored by TLC. After reaction is over, the reaction mixture was diluted with EtOAc (10 mL). The organic layer was separated and dried over *anhydrous* Na_2SO_4 . It was purified over silica gel by column chromatography. The enantiomeric excess (ee) of aldol product was determined by chiral HPLC analysis. Relative and absolute configurations of the products were determined by comparison with the known ^1H NMR, chiral HPLC analysis, and optical rotation values. All the compounds reported in **Table 1**, **Table 2**, **Table 3** and **Table 4** in main text are known in literature (corresponding references are mentioned over each compound) . The HPLC data and ^1H NMR spectra of all the compounds are given below.

(*R*)-4-Hydroxy-4-phenylbutan-2-one (3a)¹. It was obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1.0mL/min; t_R = 15.0 min (*R*), 16.4 min (*S*); $[\alpha]_D^{25} +60.3$ (c 1.0, CHCl_3), Anal. Calcd for $\text{C}_{10}\text{H}_{12}\text{O}_2$: C, 73.15; H, 7.37; Found: C, 73.25; H, 7.30.

3a (Racemic)

3a (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [μV·s]	Height [μV]	Area [%]	Area/Height [s]
1	15.012	9590718.00	302238.36	99.81	31.73
2	16.453	18020.00	913.70	0.19	19.72

(R)-4-(2-Chlorophenyl)-4-hydroxybutan-2-one (3b)¹. It was

obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; t_R = 10.4 min (*R*), 11.8 min (*S*); $[\alpha]_D^{25} +112.6$ (c 1.9, CHCl_3), Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{O}_2\text{Cl}$: C, 60.46; H, 5.58 Found: C, 60.42; H, 5.49.

3b (Racemic)

3b (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	10.394	10818905.00	418571.72	99.53	25.85
2	11.849	51241.00	5402.67	0.47	9.48

(R)-4-(3-Fluorophenyl)-4-hydroxybutan-2-one (3c)². It was

obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; t_R = 13.7 min (*R*), 16.2 min (*S*); $[\alpha]_D^{25} +44.8$ (c 2.1, CHCl_3), Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{O}_2\text{F}$: C, 65.92; H, 6.09; . Found: C, 65.99; H, 6.00.

3c (Racemic)

3c (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [μV·s]	Height [μV]	Area [%]	Area/Height [s]
1	13.727	1478786.00	50237.72	99.54	29.44
2	16.247	6776.00	386.58	0.46	17.53

(R)-4-(2,5-Difluorophenyl)-4-hydroxybutan-2-one (3d)². It was obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; t_R = 11.5 min (*R*), 13.7 min (*S*); $[\alpha]_D^{25}$ +80.9 (*c* 2.7, CHCl₃), Anal. Calcd for C₁₀H₁₀O₂F₂: C, 60.00; H, 5.04; . Found: C, 60.10; H, 5.09.

3d (Racemic)

3d (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	11.558	6969060.00	241257.92	99.55	28.89
2	13.742	31766.00	2047.77	0.45	15.51

(R)-4-(3-Chlorophenyl)-4-hydroxybutan-2-one (3e)². It was

obtained in a maximum of 99% ee. The optical purity was determined by HPLC on

chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; t_R = 15.4 min

(*R*), 17.9 min (*S*); $[\alpha]_D^{25} +33.2$ (*c* 2.1, CHCl_3), Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{O}_2\text{Cl}$: C, 60.46; H, 5.58; . Found: C, 60.40; H, 5.62.

3e (Racemic)

3e (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	15.479	774197.00	23067.84	99.36	33.56
2	17.900	4995.00	283.33	0.64	17.63

(*R*)-4-(2-Chloro-6-fluorophenyl)-4-hydroxybutan-2-one (**3f**)². It

was obtained in a maximum of >99% ee. The optical purity was determined by HPLC on

chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; $t_R = 16.5$ min(*R*), 18.3 min (*S*); $[\alpha]_D^{25} +13.7$ (*c* 1.9, CHCl₃), Anal. Calcd for C₁₀H₁₀O₂FCl: C, 55.44; H, 4.65; . Found: C, 55.35; H, 4.75.

3f (Racemic)

3f (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [μV·s]	Height [μV]	Area [%]	Area/Height [s]
1	16.122	2415217.00	66773.66	100.00	36.17

(R)-4-(4-Fluorophenyl)-4-hydroxybutan-2-one (3g)¹. It was obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 98:2]; flow rate 1.0 mL/min; t_R = 24.5min (*R*), 26.2 min (*S*); $[\alpha]_D^{25} +58.7$ (c 1.0, CHCl_3), Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{O}_2\text{F}$: C, 65.92; H, 6.09; . Found: C, 65.87; H, 6.00.

3g (Racemic)

3g (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	20.827	2315839.00	55494.16	100.00	41.73

(*R*)-4-Hydroxy-4-(3-methoxyphenyl)butan-2-one (3h)². It was obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 90:10]; flow rate 1.0 mL/min; $t_R = 23.9$ min (*R*), 27.2 min (*S*); $[\alpha]_D^{25} +30.5$ (c 1.3, CHCl_3), Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_3$: C, 68.02; H, 7.27; . Found: C, 68.10; H, 7.21.

3h (Racemic)

3h (Chiral)

DEMO REPORT

Peak #	Component Name	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]
0	peak1				
0	peak2				
1		23.910	2275499.00	42759.19	99.75
2		27.212	5692.00	185.95	0.25

(*R*)-4-Hydroxy-4-(4-nitrophenyl)butan-2-one (3i)¹. It was

obtained in a maximum of 85% ee. The optical purity was determined by HPLC on chiralpak AS-H column [hexane/2-propanol 70:30]; flow rate 1.0 mL/min; t_R = 9.9 min (*R*), 12.8min (*S*); $[\alpha]_D^{25} +14.0$ (c 1.0, CHCl_3), Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{O}_4\text{N}$: C, 57.41; H, 5.30; N, 6.70 . Found: C, 57.32; H, 5.39, N, 6.75.

3i (Racemic)

3i (Chiral)

DEFAULT REPORT

Peak #	Component Name	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]
1		9.939	9055624.00	222877.31	92.55
2		12.884	729062.50	16600.77	7.45

(R)-4-Hydroxy-4-(4-(trifluoromethyl)-phenyl)butan-2-one

(3j)⁷. It was obtained in a maximum of 91% ee. The optical purity was determined by HPLC on chiralpak AS-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; t_R = 12.9 min (*R*), 16.7 min (*S*); $[\alpha]_D^{25} +37.1$ (c 1.5, CHCl_3), Anal. Calcd for $\text{C}_{11}\text{H}_{11}\text{O}_2\text{F}_3$: C, 56.90; H, 4.77; . Found: C, 56.85; H, 4.75.

3j (Racemic)

3j (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	13.486	1523883.50	42657.48	95.57	35.72
2	18.227	70619.00	1742.39	4.43	40.53

(*R*)-4-(3-Bromophenyl)-4-hydroxybutan-2-one (3k)². It was

obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; t_R = 14.1 min (*R*), 15.7min (*S*); $[\alpha]_D^{25} +36.7$ (c 1.2, CHCl_3)., Anal. Calcd for $\text{C}_{10}\text{H}_{11}\text{O}_2\text{Br}$: C, 49.41; H, 4.56 . Found: C, 49.37; H, 4.50.

3k (Racemic)

3k (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	14.094	878028.90	25879.86	99.69	33.93
2	15.762	2713.50	158.26	0.31	17.15

(R)-4-Hydroxy-4-(4-methoxyphenyl)butan-2-one (3l)³. It

was obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; t_R = 22.3 min

(R), 26.5 min (S); $[\alpha]_D^{25} +27.2$ (c 1.3, CHCl_3), Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_3$: C, 68.02; H, 7.27; . Found: C, 68.09; H, 7.22.

3l (Racemic)

3l (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	22.382	1912344.50	35111.59	99.62	54.46
2	26.515	7257.00	228.83	0.38	31.71

(*R*)-4-Hydroxy-4-(3-methylphenyl)butan-2-one (3m)². It was

obtained in a maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 90:10]; flow rate 1.0 mL/min; t_R = 14.5min (*R*), 15.7 min (*S*); $[\alpha]_D^{25} +58.8$ (c 1.0, CHCl_3), Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_2$: C, 74.13; H, 7.92; . Found: C, 74.09; H, 7.99.

3m (Racemic)

3m (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	14.489	4475512.50	128191.38	100.00	34.91

(*R*)-4-Hydroxy-4-(naphthalene-2-yl)butan-2-one (3n)⁷. It

was obtained in a maximum of 84% ee. The optical purity was determined by HPLC on chiralpak AS-H column [hexane/2-propanol 95:5]; flow rate 1.0 mL/min; t_R = 25.3 min (*R*), 28.7 min (*S*); $[\alpha]_D^{25} +19.0$ (c 1.0, CHCl_3), Anal. Calcd for $\text{C}_{14}\text{H}_{14}\text{O}_2$: C, 78.48; H, 6.59; . Found: C, 78.54; H, 6.53.

3n (Racemic)

3n (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	25.290	15195803.50	191529.15	91.91	79.34
2	28.793	1338115.00	20994.61	8.09	63.74

(2S,1'R)-2-(hydroxy(phenyl) methyl) cyclohexan-1-one (4a)³. It

was obtained in a maximum of 85% Yield and >99% ee. The optical purity was determined by HPLC on chiralpak OD-H column [hexane/2-propanol 90:10]; flow rate 0.5mL/min; $t_{R(\text{major})} = 15.2$ min, $t_{R(\text{minor})} = 21.2$ min; $[\alpha]_D^{25} +19.0$ (c 1.0, CHCl_3), Anal. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_2$: C, 76.44; H, 7.90; . Found: C, 76.40; H, 7.95.

4a (Racemic)

4a (Chiral)

(2S,1'R)-2-(hydroxy(4-Nitrophenyl) methyl) cyclohexan-1-

one (4b)³. It was obtained in a maximum of 80% Yield and 91% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 80:20]; flow rate 0.5 mL/min; $t_{R(\text{minor})} = 26.2$ min, $t_{R(\text{major})} = 34.0$ min; $[\alpha]_{\text{D}}^{25} +11.0$ (c 1.5, CHCl_3), Anal. Calcd for $\text{C}_{13}\text{H}_{15}\text{O}_4\text{N}$: C, 62.64; H, 6.07; . Found: C, 62.69; H, 6.01.

4b (Racemic)

4b (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	26.247	1268103.00	27966.26	4.33	45.34
2	34.016	27986134.00	330880.62	95.67	84.58

(2S,1'R)-2-(hydroxy(4-methoxyphenyl)methyl) cyclohexan-

1-one (4c)³. It was obtained in a maximum of 71% Yield and 91% ee. The optical purity was determined by HPLC on chiralpak AS-H column [hexane/2-propanol 90:10]; flow rate 0.5 mL/min; $t_{R(\text{major})} = 29.5$ min, $t_{R(\text{minor})} = 42.4$ min; $[\alpha]_D^{25} +30.5$ (c 1.7, CHCl_3), Anal. Calcd for $\text{C}_{14}\text{H}_{18}\text{O}_3$: C, 71.77; H, 7.74; . Found: C, 71.69; H, 7.77.

4c (Racemic)

4c (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	29.540	1202185.00	14390.47	95.54	83.54
2	40.472	56103.00	1101.05	4.46	50.95

(2*S*,1'*R*)-4-(hydroxy(2-oxocyclohexyl) methyl)benzonitrile

(4d)³. It was obtained in a maximum of 80% Yield and 85% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 90:10]; flow rate 0.5 mL/min; $t_{R(\text{minor})}$ = 43.6 min, $t_{R(\text{major})}$ = 55.0 min; $[\alpha]_D^{25}$ +20.1 (c 1.4, CHCl_3), Anal. Calcd for $\text{C}_{14}\text{H}_{15}\text{NO}_2$: C, 73.34; H, 6.59, N, 6.11. Found: C, 73.39; H, 6.52; N, 6.09.

4d (Racemic)

4d (Chiral)

(2S,1'R)-2-(hydroxy(4trifluoromethyl)phenyl)methylcyclohexan-1-one (4e)⁴. It was obtained in a maximum of 77% Yield and >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 90:10]; flow rate 0.5 mL/min; $t_{R(\text{major})} = 21.5$ min, $t_{R(\text{minor})} = 28.1$ min; $[\alpha]_D^{25} +2.8$ (c 1.3, CHCl_3), Anal. Calcd for $\text{C}_{14}\text{H}_{15}\text{O}_2\text{F}_3$: C, 61.76; H, 5.55. Found: C, 61.71; H, 5.52.

4e (Racemic)

4e (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [μV·s]	Height [μV]	Area [%]	Area/Height [s]
1	21.465	20287848.50	268023.60	99.92	75.69
2	28.048	16511.00	268.77	0.08	61.43

(2S,1'R)- 2-(4-chlorophenyl)(hydroxy) methyl cyclohexan-1-

one (4f)³. It was obtained in a maximum of 83% Yield and 94% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 90:10]; flow

rate 0.5 mL/min; $t_{R(\text{minor})} = 26.6\text{min}$, $t_{R(\text{major})} = 30.5\text{ min}$; $[\alpha]_D^{25} +23.4$ (c 1.3, CHCl_3) , Anal.

Calcd for $\text{C}_{13}\text{H}_{15}\text{O}_2\text{Cl}$: C, 65.41; H, 6.33, Found: C, 65.35; H, 6.39.

4f (Racemic)

4f (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	26.575	98793.50	2105.09	2.85	46.93
2	30.492	3364836.50	45636.45	97.15	73.73

(2*S*,1'*R*)-2-furan-2-yl(hydroxy)methyl cyclohexan-1-one (4g)⁴. It

was obtained in a maximum of 70% Yield and 96% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 90:10]; flow rate 0.5 mL/min, 220 nm; $t_{R(\text{major})}$ = 26.8 min, $t_{R(\text{minor})}$ = 31.2 min; $[\alpha]_D^{25}$ +21 (c 1.0, CHCl₃), Anal. Calcd for C₁₁H₁₄O₃: C, 68.02; H, 7.27. Found: C, 68.09; H, 7.22.

4g (Racemic)

4g (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	26.812	35976095.00	529081.09	98.00	68.00
2	31.157	735679.00	15065.98	2.00	48.83

(2S,1'R)-2-(hydroxy(naphthalene-2-yl) methyl) cyclohexan-1-

one (4h)⁴. It was obtained in a maximum of 70% Yield and 98% ee. The optical purity was determined by HPLC on chiralpak AS-H column [hexane/2-propanol 90:10]; flow rate 0.5 mL/min; $t_{R(\text{major})} = 26.6$ min, $t_{R(\text{minor})} = 30.8$ min; $[\alpha]_{\text{D}}^{25} +5.8$ (c 1.3, CHCl_3), Anal. Calcd for $\text{C}_{17}\text{H}_{18}\text{O}_2$: C, 80.28; H, 7.13. Found: C, 80.18; H, 7.10.

4h (Racemic)

4h (Chiral)

(R)-4-hydroxy-5-methylhexan-2-one (5a)⁵. It was obtained in a

maximum of >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 97:3]; flow rate 0.5 mL/min; $t_{R(\text{major})}$ = 10.99 min, $t_{R(\text{minor})}$ = 12.83 min. $[\alpha]_D^{25} +3.0$ (c 0.4, CHCl_3), Anal. Calcd for $\text{C}_7\text{H}_{14}\text{O}_2$: C, 64.58; H, 10.84. Found: C, 64.65; H, 10.70.

5a (Racemic)

5a (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [μV·s]	Height [μV]	Area [%]	Area/Height [s]
1	18.108	43585.50	1153.79	99.85	37.78
2	19.533	63.69	-10.95	0.15	-5.82

(R)-4-cyclohexyl-4-hydroxybutan-2-one (5b)². It was obtained in a

maximum of 62% Yield and >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 1mL/min, 280 nm; $t_{R(\text{major})}$ = 9.7 min, $t_{R(\text{minor})}$ = 12.4 min; $[\alpha]_D^{25}$ +40.5 (c 1.2, CHCl_3), Anal. Calcd for $\text{C}_{10}\text{H}_{18}\text{O}_2$: C, 70.55; H, 10.66. Found: C, 70.59; H, 10.61.

5b (Racemic)

5b (Chiral)

DEFAULT REPORT

Peak #	Component Name	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]
1		9.750	1954040.49	71767.41	99.33
2		12.398	13226.51	468.33	0.67

(2S,1'R)-2-(cyclohexyl(hydroxy) methyl) cyclohexan-1-one (5c)⁴. It

was obtained in a maximum of 80% Yield and 99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 95:5]; flow rate 0.5 mL/min, 280 nm; $t_{\text{R(major)}}$ = 22.8 min, $t_{\text{R(minor)}}$ = 28.5 min; $[\alpha]_{\text{D}}^{25} +11.5$ (c 0.5, CHCl_3).

5c (Racemic)

5c (Chiral)

DEFAULT REPORT

Peak #	Component Name	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]
1		18.876	265625.00	4542.80	100.00

(2S,1'R) 3-(hydroxy(phenyl) methyl) tetrahydropyran-4-one (5d)⁶.

It was obtained in a maximum of 75% Yield and >99% ee. The optical purity was determined by HPLC on chiralpak AD-H column [hexane/2-propanol 85:15]; flow rate 1 mL/min, 220 nm; $t_{\text{R(major)}}$ = 28.3 min, $t_{\text{R(minor)}}$ = 30.9 min. $[\alpha]_{\text{D}}^{25} +20.8$ (c 1.8, CHCl_3),
 Anal. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_3$: C, 69.88; H, 6.84. Found: C, 69.81; H, 6.89.

5d (Racemic)

5d (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [μV·s]	Height [μV]	Area [%]	Area/Height [s]
1	28.376	40141000.00	574879.91	99.51	69.83
2	30.918	197019.41	5164.19	0.49	38.15

(2S,1'R) 3-(hydroxy(phenyl) methyl) tetrahydrothiopyran-4-one

(5e)⁶. It was obtained in a maximum of 73% Yield and >99% ee. The optical purity was determined by HPLC on chiralpak OD-H column [hexane/2-propanol 98:2]; flow rate 0.5 mL/min, $t_{R(\text{major})}$ = 60.6 min, $t_{R(\text{minor})}$ = 82.1 min. $[\alpha]_D^{25} +17.1$ (c 1.4, CHCl_3), Anal. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_2\text{S}$: C, 64.83; H, 6.35. Found: C, 64.89; H, 6.31.

5e (Racemic)

5e (Chiral)

DEFAULT REPORT

Peak #	Time [min]	Area [$\mu\text{V}\cdot\text{s}$]	Height [μV]	Area [%]	Area/Height [s]
1	60.636	9341909.00	36425.79	99.38	256.46
2	87.098	58023.00	503.75	0.62	115.18

^1H NMR spectra of enantiopure compounds

All the compounds mentioned in the manuscript (table 1-table 4) are literature known (corresponding references are mentioned over each compound). The ^1H NMR spectra of all the compounds are given below:

400 MHz ^1H NMR Spectrum of (**Compound 3a**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3b**) in CDCl_3

400 MHz ¹H NMR Spectrum of (**Compound 3c**) in CDCl₃

400 MHz ^1H NMR Spectrum of (**Compound 3d**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3e**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3f**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3g**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3h**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3i**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3j**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3k**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3l**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 3n**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 4a**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 4b**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 4c**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 4d**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 4e**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 4f**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 4g**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 4h**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 5b**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 5c**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 5d**) in CDCl_3

400 MHz ^1H NMR Spectrum of (**Compound 5e**) in CDCl_3

References:

- (1). Tang, Z.; Jiang, F.; Yu, L.-T.; Cui, X.; Gong, L.-Z.; Mi, A.-Q.; Jiang, Y.-Z.; Wu, Y.-
D. J. Am. Chem. Soc. **2005**, 127, 928.
- (2) Raj, M.; Maya, V.; Ginotra, S. K.; Singh, V. K.; *Org. Lett.* **2006**, 8, 4097.
- (3) Mase, N.; Nakai, Y.; Ohara, N.; Yoda, H.; Takabe, K.; Tanaka, F.; Barbas, C. F., III.
J. Am. Chem. Soc. **2006**, 128, 734.
- (4) Wu, Y.; Zhang, Y.; Yu, M.; Zhao, G.; Wang, S. *Org. Lett.* **2006**, 8, 4417.
- (5) Zhou, Y.; Shan, Z.; *J. Org. Chem.* **2006**, 71, 9510.
- (6) Rong, J.; Yong, L.-X.; Ning, X.-X.; Jing, X.-W. *J. Org. Chem.* **2006**, 71, 8198.
- 7) Guo, H.-M.; Cun, L.-F.; Gong, L.-Z.; Mi, A.-Q.; Jiang, Y.-Z. *Chem. Commun.* **2005**,
1450.