Highly Functionalized Organolithium Reagents for Enantiomerically Pure α -Amino Acid Synthesis

Martin N. Kenworthy, John Paul Kilburn and Richard J. K. Taylor*

Supplementary Material: Experimental Procedures and Characterisation

Synthesis of Lithiation Precursor 10d

(a) 2-tert-Butoxycarbonylamino-3-chloro-propionic acid methyl ester-

A solution of triphenylphosphine (19.2 g, 73.2 mmol) and hexachloroethane (17.3 g, 73.2 mmol) in dichloromethane (50 ml) was added in one portion to a solution of *N*-Boc-serine methyl ester **8** (14.6 g, 66.5 mmol) in dichloromethane (250 ml) under an atmosphere of argon. The reaction was stirred at rt for 2 h, then quenched with a saturated solution of sodium hydrogen carbonate (50 ml). The organic phase was separated and washed with brine (100 ml), dried (MgSO_4), evaporated then triturated with Et_2O (300 ml). After filtration and evaporation the subsequent residue was purified on silica gel, eluting with petrol(40-60):EtOAc 5:1, delivering 2-tert-butoxycarbonylamino-3-chloro-propionic acid methyl ester (13.2 g, 84%) as a white solid, m.p. 62-64°C. $\nu_{\text{max}}(\text{film})/\text{cm}^{-1}$ 3362 (NH), 2982 (CH), 1721 (CO); $[\alpha]_{\text{D}}^{21} +37.8$ (c. 1.5, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 5.37 (1H, brd, J 7.5), 4.65 (1H, m), 3.90 (1H, dd, J 3, 20), 3.78 (1H, dd, J 3.5, 20), 3.73 (3H, s), 1.40 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 169.6, 154.9, 80.5, 54.4, 52.9, 45.5, 28.2; (Found: MNH_4^+ , 255.1110. $\text{C}_8\text{H}_{16}\text{NO}_4^{35}\text{Cl}$ requires MNH_4 , 255.1112 (0.8 ppm)).

(b) (2-Chloro-1-hydroxymethyl-ethyl)-carbamic acid tert-butyl ester **9**-

Lithium borohydride (1.18 g, 54.1 mmol) was added in portions to a stirred solution of the above chloride (12.85 g, 54.1 mmol) in ethanol:THF 9:1 (700 ml) at 0°C under an atmosphere of argon. The reaction was stirred for 18 h at rt then quenched with a saturated solution of ammonium chloride (10 ml). The organics were then evaporated and the residue partitioned between water (300 ml) and Et_2O (300 ml). After

separation, the aqueous was further extracted with Et₂O (300 ml). The combined organics were dried (MgSO₄) and evaporated to give a gummy residue which was purified on silica gel, eluting with petrol(40-60):Et₂O 2:1, giving the *alcohol* **9** (8.1 g, 81%) as a gum which slowly solidified to a fluffy amorphous white solid. $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3353 (OH), 2978 (CH), 1689 (CO); $[\alpha]_{\text{D}}^{21}$ -4.6 (c. 0.7, CHCl₃); $\delta_{\text{H}}(400 \text{ MHz, CDCl}_3)$ 4.94 (1H, brs), 3.86 (1H, brs), 3.8-3.55 (4H, m), 1.42 (9H, s); $\delta_{\text{C}}(100 \text{ MHz; CDCl}_3)$ 155.6, 80.2, 62.0, 52.5, 44.2, 28.3; (Found: MH⁺, 210.0889. C₈H₁₆NO₃³⁵Cl requires *MH*, 210.0897 (4.0 ppm)).

[2-Chloro-1-(2-trimethylsilyl-ethoxymethoxymethyl)-ethyl]-carbamate tert-butyl ester **10d**-

SEMCl (0.99 ml, 5.5 mmol) was added dropwise to a stirred solution of alcohol **9** (1.16 g, 5.5 mmol) and Hunig's base (0.96 ml, 5.5 mmol) in dichloromethane (20ml) at 0°C under an atmosphere of argon. The reaction was allowed to warm to rt and stirred for 18 h. The reaction was evaporated and partitioned between Et₂O (30 ml) and water (20ml). After separation, further extraction with Et₂O (2 × 30 ml), the combined organics were dried (MgSO₄) and evaporated. The subsequent crude residue was purified on silica gel, eluting with petrol(40-60):Et₂O 4:1, giving the *ether* **10b** (1.49 g, 80%) as a colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3346 (NH), 2955, 2932, 2893 (CH), 1719 (CO); $[\alpha]_{\text{D}}^{21}$ +12.4 (c. 0.4, CHCl₃); $\delta_{\text{H}}(400 \text{ MHz, CDCl}_3)$ 5.03 (1H, brd, *J* 8.0), 4.67 (2H, s), 4.01 (1H, brs), 3.80 (1H, dd, *J* 4, 10), 3.71 (1H, dd, *J* 4, 11), 3.6-3.5 (4H, m), 1.44 (9H, s), 0.95 (2H, t, *J* 9), 0.02 (9H, s); $\delta_{\text{C}}(100 \text{ MHz; CDCl}_3)$ 155.4, 95.4, 80.2, 66.4, 65.8, 51.1, 44.3, 28.7, 18.4, -1.1; (Found: MH⁺, 340.1711. C₁₄H₃₀NO₄³⁵ClSi requires *MH*, 340.1711 (0.0 ppm)).

General Lithiation Reaction-for synthesis of adduct **29**

n-BuLi (0.21 ml, 0.42 mmol, 1.98*M* solution in hexanes) was added dropwise to a stirred solution of chloride **10d** (128 mg, 0.38 mmol) in THF (5 ml) at -78°C under an atmosphere of argon. Stirring was continued for 15 min, followed by the dropwise addition of LiNp¹ (2.2 ml, 1.1 mmol, 0.5*M* solution in THF) over 5 min. The dark solution was stirred at -78°C for 2 h. Subsequently CD₃OD (50 μl , 1.1 mmol) was added dropwise to the solution causing decolorisation. The reaction mixture was kept cold (*ca.* -78°C to -40°C) overnight then quenched with a saturated solution of ammonium chloride (1 ml) and allowed to warm to room temperature. The reaction was diluted with ether (20 ml) and saturated ammonium chloride (20 ml). The layers were separated and the aqueous layer was extracted with ether (2×10 ml). The combined organics were dried (MgSO₄), filtered and evaporated to give a crude product which was purified by flash column chromatography on silica gel, eluting with 1:3 diethyl ether-petrol ether (40-60) to give the *carbamate* **29** (101 mg, 87%) as a colorless oil; $\nu_{\text{max}}(\text{film})/\text{cm}^{-1}$ 3351 (NH), 2954 (CH), 2954 (CO); $[\alpha]_{\text{D}}^{21} -13.6$ (*c.* 1.0, CHCl₃); $\delta_{\text{H}}(400 \text{ MHz, CDCl}_3)$ 4.78 (1H, brs), 4.65 (2H, s), 3.91 (1H, brs), 3.60 (2H, t, *J* 8.5), 3.48 (2H, m), 1.42 (9H, s), 1.15 (2H, m), 0.93 (2H, t, *J* 8.5), 0.00 (9H, s); $\delta_{\text{C}}(100 \text{ MHz; CDCl}_3)$ 155.5, 92.2, 79.2, 71.4, 65.3, 46.3, 28.5, 18.2, 17.8 (t, *J* 19.8, CH₂D), -1.3; (Found: MH⁺, 307.2162. C₁₄H₃₁NDO₄Si requires *MH*, 307.2163 (0.5 ppm)).

¹ Prepared by literature method see Foubelo, F.; Yus, M. *Tetrahedron: Asymmetry* **1996**, 7, 2911-2922, and the references therein.

Similar procedures were employed to prepare the following compounds:

Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3448 (OH), 2932, 2867 (CH), 1691 (CO); $[\alpha]_{\text{D}}^{21} -5.0$ (c. 0.6, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 5.13 (1H, brs), 4.65 (2H, s), 3.95 (1H, brs), 3.6-3.5 (4H, m), 2.98 (1H, brs), 1.65-1.2 (21H, m), 0.94 (2H, t, J 8.5), 0.01 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 156.3, 95.3, 79.8, 71.9, 70.4, 65.5, 46.8, 44.6, 38.5, 37.9, 29.5, 26.0, 22.4, 22.3, 18.2, -1.3 ; (Found: MH^+ , 404.2831. $\text{C}_{20}\text{H}_{41}\text{NO}_5\text{Si}$ requires MH , 404.2832 (0.4 ppm)).

Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3397 (OH), 2977, 2953, 2934 (CH), 1686 (CO); $[\alpha]_{\text{D}}^{21} -1.6$ (c. 0.2, CHCl_3); $\delta_{\text{H}}(270 \text{ MHz}, \text{CDCl}_3)$ 5.19 (1H, brd, J 8), 4.68 (2H, s), 4.41 (1H, brs), 3.97 (1H, brm), 3.7-3.6 (4H, m), 2.2-1.8 (6H, m), 1.7-1.5 (2H, m), 1.44 (9H, s), 0.96 (2H, t, J 8.5), 0.03 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 156.3, 95.3, 80.1, 73.3, 71.6, 65.4, 47.3, 42.5, 35.9, 35.6, 28.4, 18.1, 13.0, -1.5 ; (Found: MH^+ , 376.2521. $\text{C}_{18}\text{H}_{37}\text{NO}_5\text{Si}$ requires MH , 376.2519 (-0.4 ppm)).

Diastereomer 1- Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3422 (OH), 2954, 2894 (CH), 1689 (CO); $[\alpha]_{\text{D}}^{21} -0.3$ (c. 0.6, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 7.5-7.2 (5H, m), 5.29 (1H, brd, J 8), 4.8-4.6 (3H, m), 4.46 (1H, d, J 3.5), 4.10 (1H, brs), 3.7-3.5 (4H, m), 1.93 (1H, m), 1.75 (1H, m), 1.48 (9H, s), 0.93 (2H, t, J 8.5), 0.00 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 156.3, 144.1, 128.2, 127.0, 125.5, 95.2, 80.1, 70.8, 69.9, 65.4, 47.5, 43.3, 28.3, 18.0, -1.5 ; (Found: MH^+ , 412.2519. $\text{C}_{21}\text{H}_{37}\text{NO}_5\text{Si}$ requires MH , 412.2519 (0.0 ppm)).

Diastereomer 2- Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3373 (OH), 2953, 2893 (CH), 1696 (CO); $[\alpha]_{\text{D}}^{21} +17.0$ (c. 0.3, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 7.4-7.2 (5H, m), 5.10 (1H, brd), 4.78 (1H, t, J 6.5), 4.65 (2H, s), 3.89 (1H, brs), 3.7-3.5 (4H, m), 3.26 (1H, brs), 1.97 (2H, t, J 6.5) 1.43 (9H, s), 0.93 (2H, t, J 8.5), 0.00 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 155.8, 144.6, 128.4, 127.4, 125.7, 95.1, 79.5, 72.0, 70.2, 65.4, 48.6, 42.0, 28.4, 18.0, -1.5.

Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3348 (NH), 2954 (CH), 1715 (CO); $[\alpha]_{\text{D}}^{21} -2.3$ (c. 1.2, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 5.15 (1H, brd, J 7), 4.63 (2H, s), 4.11 (1H, brm), 3.7-3.5 (4H, m), 2.8-2.65 (2H, m), 2.15 (3H, s), 1.42 (9H, s), 0.93 (2H, t, J 8.5), 0.02 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 207.5, 155.4, 95.3, 79.6, 69.2, 65.5, 47.1, 44.9, 30.5, 28.5, 18.2, -1.3; (Found: MH^+ , 348.2201. $\text{C}_{16}\text{H}_{33}\text{NO}_5\text{Si}$ requires MH , 348.2206 (1.4 ppm)).

Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3356 (NH), 2954 (CH), 1715, 1688 ($2 \times \text{CO}$); $[\alpha]_{\text{D}}^{21} +6.6$ (c. 1.0, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 7.96 (2H, d, J 7.5), 7.56 (1H, t, J 7.5), 7.45 (2H, t, J 7.5), 5.32 (1H, brd, J 7.5), 4.63 (2H, s), 4.28 (1H, brm), 3.75 (1H, dd, J 4, 9.5), 3.64 (1H, dd, J 5, 9.5), 3.54 (2H, t, J 8.5), 3.36 (1H, brm), 3.23 (1H, dd, J 7, 16.5), 1.42 (9H, s) 0.89 (2H, t, J 8.5), -0.03 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 199.0, 155.6, 137.2, 133.6, 129.0, 128.4, 95.5, 79.7, 69.3, 65.7, 47.7, 39.9, 28.7, 18.4, -1.1; (Found: MH^+ , 410.2362. $\text{C}_{21}\text{H}_{35}\text{NO}_5\text{Si}$ requires MH , 410.2363 (0.3 ppm)).

Colorless oil;² $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3356 (NH), 2954 (CH), 1741, 1718 ($2 \times \text{CO}$); $[\alpha]_{\text{D}}^{21} -2.2$ (c. 0.9, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 5.19 (1H, brd, J 8), 4.65 (2H, s), 4.13 (1H, brs), 3.67 (3H, s), 3.65-3.5 (4H, m), 2.7-2.5 (2H, m), 1.43 (9H, s), 0.93 (2H, t, J 8.5),

² $\text{TMSCHN}_2/\text{MeOH:PhMe}$ was used to esterify the carboxylic acid.

0.02 (9H, s); δ_{C} (100 MHz; CDCl_3) 172.0, 155.3, 95.3, 79.6, 69.1, 65.5, 51.8, 47.4, 36.2, 28.5, 18.2, -1.3; (Found: MH^+ , 364.2154. $\text{C}_{16}\text{H}_{33}\text{NO}_6\text{Si}$ requires MH , 364.2155 (0.4 ppm)).

Colorless oil; ν_{max} (film)/ cm^{-1} 3360 (NH), 2954, 2897 (CH), 1716 (CO); $[\alpha]_{\text{D}}^{21}$ -11.6 (*c.* 1.0, CHCl_3); δ_{H} (400 MHz, CDCl_3) 4.7-4.65 (3H, m), 3.88 (1H, brs), 3.61 (2H, m), 3.55-3.4 (2H, m), 1.43 (9H, s), 0.94 (2H, m), 0.85 (2H, d, J 7), 0.04 (9H, s), 0.02 (9H, s); δ_{C} (100 MHz; CDCl_3) 155.0, 95.1, 79.0, 72.5, 65.2, 47.7, 28.5, 20.6, 18.1, -1.0, -1.4; (Found: MH^+ , 378.2497. $\text{C}_{17}\text{H}_{39}\text{NO}_4\text{Si}_2$ requires MH , 378.2496 (-0.4 ppm)).

General SEM Deprotection Reaction-shown for synthesis of compound **30**

0.1M HCl^3 in methanol (2 ml, ~6 ml/mmol) was added to SEM-ether **29** (84 mg, 0.28 mmol). The subsequent reaction mixture was stirred at room temperature for 3h. Excess triethylamine (~0.1 ml) was then added and the volatiles were removed under reduced pressure to deliver a crude mixture, this was then purified by flash column chromatography on silica gel, eluting with diethyl ether to give the *alcohol* **30** (34 mg, 70%) as a colorless oil; ν_{max} (film)/ cm^{-1} 3361 (OH), 2978, 2936 (CH), 1690 (CO); $[\alpha]_{\text{D}}^{21}$ -12.6 (*c.* 1.1, CHCl_3); δ_{H} (400 MHz, CDCl_3) 4.71 (1H, brs), 3.68 (1H, brs), 3.55 (1H, m), 3.43 (1H, dd, J 6.0, 10.5), 3.00 (1H, brs), 1.38 (9H, s), 1.07 (2H, m); δ_{C} (100 MHz; CDCl_3) 156.6, 79.8, 67.4, 48.6, 28.5, 17.2 (t, J 19, CH_2D); (Found: MH^+ , 177.1353. $\text{C}_8\text{H}_{17}\text{NDO}_3$ requires MH , 177.1350 (2.2 ppm)).

³ This was prepared by addition of acetyl chloride (10 mmol) to methanol (100 ml) cooled with a water-bath.

Similar procedures were employed to prepare the following compounds:

Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3348 (OH), 2933, 2861 (CH), 1689 (CO); $[\alpha]_{\text{D}}^{21} -5.8$ (c. 0.8, CHCl_3); $\delta_{\text{H}}(270 \text{ MHz}, \text{CDCl}_3)$ 5.31 (1H, brs), 3.83 (1H, brm), 3.66 (1H, dd, J 5, 11), 3.58 (1H, dd, J 4, 11), 1.7-1.35 (21H, m); $\delta_{\text{C}}(67.5 \text{ MHz}; \text{CDCl}_3)$ 156.2, 79.8, 71.1, 66.4, 49.2, 43.1, 38.0, 37.8, 28.4, 25.6, 22.2, 22.1; (Found: MH^+ , 274.2016. $\text{C}_{14}\text{H}_{27}\text{NO}_4$ requires MH , 274.2018 (0.8 ppm)).

Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3339 (OH), 2980, 2935 (CH), 1682 (CO); $[\alpha]_{\text{D}}^{21} -0.5$ (c. 0.4, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 5.22 (1H, brs), 4.27 (1H, brs), 3.81 (1H, brm), 3.51 (2H, brm), 3.45 (1H, brs), 2.15-2.05 (2H, m), 1.89 (1H, dd, J 9, 14), 1.85-1.7 (2H, m), 1.58 (1H, m), 1.42 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 156.8, 80.0, 73.9, 65.7, 49.4, 41.2, 36.2, 35.8, 28.4, 12.7; (Found: MH^+ , 246.1703. $\text{C}_{12}\text{H}_{23}\text{NO}_4$ requires MH , 246.1705 (0.9 ppm)).

Major diastereomer. Colorless oil; $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 7.4-7.2 (5H, m), 5.18 (1H, brd, J 7.5), 4.73 (1H, brd, J 8.5), 4.11 (1H, brs), 3.95 (1H, brm), 3.72 (1H, brd, J 8.5), 3.63 (1H, dd, J 4, 10.5), 2.60 (1H, brs), 1.82 (2H, m), 1.46 (9H, s); $\delta_{\text{C}}(100 \text{ MHz}; \text{CDCl}_3)$ 157.3, 144.1, 128.4, 127.3, 125.5, 80.2, 70.4, 65.4, 49.7, 42.3, 28.3.

Colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3256 (OH), 2954 (CH), 1695 (CO); $[\alpha]_{\text{D}}^{21} -14.6$ (c. 0.8, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz}, \text{CDCl}_3)$ 4.51 (1H, brs), 3.79 (1H, brs), 3.63 (1H, brd, J 9), 3.44

(1H, dd, *J* 6.5, 10.5), 2.47 (1H, brs), 1.44 (9H, s), 0.74 (2H, m), 0.05 (9H, s); δ_c (100 MHz; CDCl₃) 156.6, 80.0, 69.1, 50.6, 28.8, 20.1, -0.72; (Found: MH⁺, 248.1679. C₁₁H₂₅NO₃Si requires *MH*, 248.1682 (1.2 ppm)).

General Oxidation Method A-shown for synthesis of compound **13**

PDC (413 mg, 1.1) was added to a stirred solution of diol **12e** (50 mg, 0.18 mmol) in anhydrous dimethylformamide (1.5 ml). The reaction mixture was stirred for 6 h, then water (5 ml) was added and the mixture was extracted with dichloromethane (2 × 15 ml). The combined extracts were then washed with water (2 × 20 ml) and brine (20 ml), then dried (MgSO₄) and evaporated under reduced pressure. The crude residue was purified on silica gel, eluting with petrol(40-60):Et₂O 1:1, delivering the lactone **13** (34 mg, 69%) as a white foam; $\nu_{\max}$ (film)/cm⁻¹ 3365 (NH), 2937, 2863 (CH), 1778, 1697 (2 × CO); $[\alpha]_D^{21} +17.7$ (c. 1.1, CHCl₃); δ_H (400 MHz, CDCl₃) 5.05 (1H, brs), 4.51 (1H, brs), 2.73 (1H, brt, *J* 10), 1.8-1.4 (20H, m); δ_c (100 MHz; CDCl₃) 175.0, 155.8, 84.9, 80.8, 51.0, 41.4, 38.6, 36.1, 28.6, 25.2, 22.9, 22.8; (Found: MNH₄⁺, 287.1977. C₁₄H₂₃NO₄ requires *MNH*₄, 287.1971 (-2.1 ppm)).

A similar procedure was employed to prepare the following compound:

Method A, amorphous solid; $\nu_{\max}$ (film)/cm⁻¹ 3321 (NH), 2976, 2945 (CH), 1761, 1712 (2 × CO); $[\alpha]_D^{21} +1.2$ (c. 1.1, CHCl₃); δ_H (400 MHz, CDCl₃) 5.06 (1H, brs), 4.41 (1H, brm), 2.95 (1H, brm), 2.63 (1H, q, *J* 10), 2.36 (1H, q, *J* 10), 2.25-1.95 (3H, m), 1.88 (1H, m), 1.67 (1H, m), 1.45 (9H, s); δ_c (100 MHz; CDCl₃) 174.8, 155.8, 83.2, 80.9, 50.9, 41.5, 34.6, 28.7, 28.3, 12.7. (Found: MH⁺, 242.1398. C₁₂H₁₉NO₄ requires *MH*, 242.1392 (-2.3 ppm)).

General Oxidation Method B-shown for synthesis of compound **28**

PDC (433 mg, 1.2) was added to a stirred solution of alcohol **27** (71 mg, 0.29 mmol) in anhydrous dimethylformamide (3 ml). The reaction mixture was stirred for 18 h, then quenched with water (5 ml) and extracted with EtOAc (2 × 20 ml). The combined extracts were then washed with water (2 × 20 ml) and brine (20 ml), then dried (MgSO₄) and evaporated under reduced pressure to deliver the crude carboxylic acid. This was then dissolved in MeOH:toluene (3 ml, 1:1) and treated with TMSCHN₂ (0.29 ml, 0.58 mmol, 2.0M solution in hexanes) at 0°C. The reaction was stirred for 1h then all volatiles were removed under reduced pressure to deliver a crude mixture, which was then purified by flash column chromatography on silica gel, eluting with Et₂O:petrol(40-60) 1:2 to give the protected *amino acid* **28** (54 mg, 68%) as a colourless waxy solid, m.p. 40-42°C; $\nu_{\text{max}}(\text{film})/\text{cm}^{-1}$ 3373 (NH), 2977, 2955 (CH), 1745, 1715 (2 × CO); $[\alpha]_{\text{D}}^{21} +10.8$ (c. 1.2, CHCl₃); $\delta_{\text{H}}(400 \text{ MHz, CDCl}_3)$ 4.84 (1H, brs), 4.28 (1H, brm), 3.67 (3H, s), 1.37 (9H, s), 1.05 (1H, dd, *J* 6, 9.5), 0.88 (1H, dd, *J* 9.5, 14.5), 0.00 (9H, s); $\delta_{\text{H}}(400 \text{ MHz, CDCl}_3)$ 174.3, 154.9, 79.8, 52.1, 50.6, 28.4, 21.4, -1.3; (Found: MH⁺, 276.1633. C₁₂H₂₅NO₄Si requires *MH*, 276.1631 (-0.6 ppm)).

A similar procedure was employed to prepare the following compound:

Method B, colorless oil; $\nu_{\text{max}}(\text{film})/\text{cm}^{-1}$ 3383 (NH), 2979, 2932 (CH), 1748, 1714 (2 × CO); $[\alpha]_{\text{D}}^{21} +59.9$ (c. 1.1, DCM); ¹H NMR spectrum identical to literature.⁴ *m/z* 308 (10, M⁺), 252 (40, M-*t*Bu), 208 (100, M-Boc).

⁴ Jackson, R. F. W.; Turner, D.; Block, M. H. *J. Chem. Soc., Perkin Trans. I* **1997**, 865-870.

General Oxidation Method C-shown for synthesis of compound 31

Ruthenium chloride (4 mg, 21 μ mol, 5 mol%) was added to a stirred suspension of alcohol **30** (75 mg, 0.43 mmol) and sodium periodate (273 mg, 1.28 mmol) in $\text{CH}_3\text{CN}:\text{H}_2\text{O}:\text{CCl}_4$ (3 ml, 2:3:2) at room temperature. The reaction mixture went dark after 30 min. Stirring was continued for 12 h in total then the reaction was diluted with a saturated solution of ammonium chloride (10 ml) and EtOAc (10 ml). After separation the aqueous was extracted again with EtOAc (2×10 ml). The combined organic extracts were dried (MgSO_4) and evaporated under reduced pressure to deliver the crude carboxylic acid. This was then dissolved in MeOH:toluene (2 ml, 1:1) and treated with TMSCHN_2 (0.42 ml, 0.85 mmol, 2.0M solution in hexanes) at 0°C . The reaction was stirred for 1h then all volatiles were removed under reduced pressure to deliver a crude mixture, which was then purified by flash column chromatography on silica gel, eluting with $\text{Et}_2\text{O}:\text{petrol}(40-60)$ 1:2 to give the protected *amino acid* **31** (53 mg, 61%) as a colorless oil; $\nu_{\text{max}}(\text{film})/\text{cm}^{-1}$ 3382 (NH), 2980 (CH), 1748, 1716 ($2 \times \text{CO}$); $[\alpha]_{\text{D}}^{21} -43.5$ (c. 1.4, CHCl_3); $\delta_{\text{H}}(400 \text{ MHz, CDCl}_3)$ 5.03 (1H, brs), 4.31 (1H, brm), 3.74 (3H, s), 1.44 (9H, s), 1.36 (2H, brd, J 7.5); $\delta_{\text{C}}(100 \text{ MHz; CDCl}_3)$ 174.0, 155.2, 80.0, 52.3, 49.2, 28.5, 18.6 (t, J 20, CH_2D); (Found: MH^+ , 205.1296. $\text{C}_9\text{H}_{17}\text{NDO}_4$ requires MH , 205.1299 (1.3 ppm)).

General Deprotection Reaction-shown for synthesis of compound 34

The protected amino acid **31** (52 mg, 0.26 mmol) was heated at 70°C with 4.5M HCl (2 ml) for 3h. After cooling, the aqueous phase was diluted with water (10 ml) and washed with diethyl ether (10 ml). The aqueous phase was then evaporated under reduced pressure delivering a solid which was triturated with diethyl ether. After decanting and drying, an off-white wax was obtained (31 mg, 95%). $[\alpha]_{\text{D}}^{21} +11.7$ (c. 1.2, 6M HCl); $\delta_{\text{H}}(400 \text{ MHz, D}_2\text{O})$ 4.09 (1H, t, J 7.5), 1.50 (2H, d, J 7.5); $\delta_{\text{C}}(100$

MHz; D₂O) 172.8, 48.7, 15.0 (t, *J* 20, CH₂D); (Found: *M*-Cl⁺, 91.0620. C₃H₇NCIO₂D requires *M*-Cl, 91.0618 (1.9 ppm)).

Similar procedures were employed to prepare the following compounds:

White powder, m.p. ~200°C decomposed (lit.⁵ m.p. 207-209°C decomposed); $[\alpha]_D^{21}$ +35.6 (*c.* 0.104, 6*M* HCl), lit.⁵ $[\alpha]_D$ +34.7 (*c.* 0.098, 6*M* HCl); ¹H NMR spectrum identical to literature.⁵

White Solid, m.p. ~180°C decomposed; $[\alpha]_D^{21}$ +35.2 (*c.* 1.1, 1*M* HCl), lit.⁶ $[\alpha]_D$ +31 (*c.* 0.51, 4*M* HCl); δ_H (400 MHz, D₂O) 4.01 (1H, dd, *J* 5.5, 10.5), 1.22 (1H, dd, *J* 10.5, 14.5), 1.16 (1H, dd, *J* 5.5, 14.5), 0.08 (9H, s); δ_C (100 MHz; D₂O) 17.3, 51.7, 19.3, -2.3; *m/z* (ESI) 176 (100%, M⁺), 162 (35%, M-Cl).

Synthesis of 2-amino-3-boronopropionic acid (+)-38

(a) *n*-BuLi (0.48 ml, 0.95 mmol, 1.97*M* solution in hexanes) was added dropwise to a stirred solution of chloride **10d** (291 mg, 0.89 mmol) in THF (5 ml) at -78°C under an atmosphere of argon. Stirring was continued for 15 min, followed by the dropwise addition of LiNp (5.7 ml, 2.83 mmol, 0.5*M* solution in THF) over 5 min. The dark solution was stirred at -78°C for 2 h. Subsequently triisopropylborate (0.5 ml, 2.15 mmol) was added dropwise to the solution. The reaction mixture was kept cold (*ca.* -78°C to -40°C) overnight then quenched with a saturated solution of ammonium chloride (1 ml), then allowed to warm to rt. To the mixture was then added an

⁵ Lin, W.; He, Z.; Zhang, H.; Zhang, X.; Mi, A.; Jiang, Y. *Synthesis* **2001**, 1007-1009.

⁶ Walkup, R. D.; Cole, D. C.; Whittlesey, B. R. *J. Org. Chem.* **1995**, 60, 2630-2634.

aqueous solution of 0.5M HCl (5 ml, saturated with NaCl), and stirring was continued for 30 min. The solution was then extracted with petrol(40-60) (3 × 20 ml). The combined organics were then extracted with 2M NaOH (3 × 15 ml). These extracts were combined and acidified (~pH 1-2) with 4.5M HCl, then extracted again with Et₂O (3 × 30 ml). The final combined organics were dried (MgSO₄), filtered and evaporated under reduced pressure to give the *boronic acid* **35** (238 mg, 80%) as a light yellow oil. δ_{H} (400 MHz, *d*⁶ Acetone) 6.95-6.8 (1H, br), 6.55-6.45 (1H, br), 5.88 (1H, brs), 4.63 (2H, s), 3.88 (1H, brm), 3.63 (2H, m), 3.53 (1H, dd, *J* 5.5, 9.5), 3.47 (1H, dd, *J* 5.5, 9.5), 1.5-1.35 (9H, several singlets), 1.07 (1H, dd, *J* 6, 16), 1.00 (2H, m), 0.05-0.00 (9H, several singlets).

(b) (–)-Pinane diol (47 mg, 0.28 mmol) was added to a stirred solution of **35** (88 mg, 0.25 mmol) and powdered molecular sieves (~100 mg) in dry dichloromethane (3 ml) under an atmosphere of argon. The reaction mixture was stirred for 18 h then filtered and evaporated. The subsequent residue was purified on silica gel, eluting with petrol(40-60):Et₂O 3:1, delivering *boronate ester* **36** (113 mg, 95%) as a colorless oil; ν_{max} (film)/cm^{–1} 2926 (CH), 1718 (CO); $[\alpha]_{\text{D}}^{21}$ –200 (c. 0.1, CHCl₃); δ_{H} (400 MHz, CDCl₃) 5.06 (1H, brd, *J* 7), 4.65 (2H, s), 4.25 (1H, dd, *J* 2, 9), 4.01 (1H, brs), 3.60 (2H, t, *J* 8.5), 3.53 (2H, d, *J* 4.5), 2.32 (1H, m), 2.20 (1H, m), 2.03 (1H, t, *J* 5), 1.9-1.8 (2H, m), 1.43 (9H, s), 1.37 (3H, s), 1.28 (3H, s), 1.2-1.05 (2H, m), 0.93 (2H, t, *J* 8.5), 0.9-0.8 (4H, m), 0.02 (9H, s); δ_{C} (100 MHz; CDCl₃) 155.4, 95.1, 85.8, 77.8, 71.5, 65.3, 51.4, 39.6, 38.3, 36.2, 35.6, 29.2, 28.7, 28.6, 27.2, 26.6, 24.2, 22.8, 18.2, –1.3; (Found: MH⁺, 484.3271. C₂₄H₄₆NBO₆Si requires *MH*, 484.3266 (–1.1 ppm)).

(c) 0.1M HCl in methanol (2 ml, ~6 ml/mmol) was added to SEM-ether **29** (99 mg, 0.21 mmol). The subsequent reaction mixture was stirred at room temperature for 3h. Excess triethylamine (~0.1 ml) was then added and the volatiles were removed under reduced pressure to deliver a crude mixture, which was then purified by flash column chromatography on silica gel, eluting with petrol(40-60):Et₂O 1:2 to give starting material (4 mg, 4%) and the *alcohol* **37** (49 mg, 68%) as a colorless oil; $\nu_{\max}(\text{film})/\text{cm}^{-1}$ 3420 (OH), 2975, 2929, 2872 (CH), 1694 (CO); $[\alpha]_{\text{D}}^{21}$ -148 (c. 0.9, CHCl₃); $\delta_{\text{H}}(400 \text{ MHz, CDCl}_3)$ 5.01 (1H, brs), 4.24 (1H, d, *J* 2, 9), 3.89 (1H, brs), 3.60 (1H, brd, *J* 10.5), 3.51 (1H, m), 2.31 (1H, m), 2.19 (1H, m), 2.02 (1H, m), 1.95-1.75 (2H, m), 1.41 (9H, s), 1.35 (3H, s), 1.26 (3H, s), 1.25 (1H, m), 1.15-1.0 (2H, m), 0.81 (3H, s); (Found: MH^+ , 354.2452. C₁₈H₃₂NBO₅ requires *MH*, 354.2452 (-0.2 ppm)).

(d) Dess-Martin periodinane (78 mg, 0.18 mmol) was added to a stirred solution of **37** (54 mg, 0.15mmol) in dichloromethane (2 ml) under an atmosphere of argon. After 2 h 1M sodium sulfite (5 ml) was added and the mixture was extracted with Et₂O (2 × 10 ml). The combined organic extracts were washed with 1M sodium hydrogen carbonate (210 ml), dried (Na₂SO₄) and evaporated. The crude mixture was then dissolved in *t*-butanol (3 ml) and treated with a solution of 2-methyl-2-butene (1.5 ml, 3.1 mmol, 2M in THF). A solution of NaH₂PO₄ (167 mg, 1.07 mmol) and NaClO₂ (125 mg, 1.38 mmol) in water (2 ml) was added and the reaction was stirred for 1 h. The *t*-butanol was evaporated *in vacuo* and the residue was partitioned between 0.5M HCl (5 ml) and EtOAc (10 ml). After separation and further extraction (EtOAc 2 × 5

ml), the combined extracts were dried (MgSO₄) and evaporated. The subsequent residue was heated at 70°C with 4.5M HCl (3 ml) for 3 h. After cooling the aqueous solution was washed with dichloromethane (2 × 2 ml). The aqueous solution was then evaporated *in vacuo*. The crude residue was purified on silica gel, eluting with 2:1 EtOH:14M NH₃, delivering the 2-amino-3-boronopropionic acid **38** (15 mg, 74%) as a white solid. After trituration with acetone the product was obtained as a white powder, m.p. >290°C decomposed. δ_{H} (400 MHz, D₂O) 3.94 (1H, brt, *J* 9.5), 1.08 (1H, brdd, *J* 9.5, 13), 0.79 (1H, brdd, *J* 10.5, 12); δ_{C} (100 MHz; D₂O) 178.0, 53.4, 20.4 (br); δ^{11}_{B} (160 MHz; D₂O) 16.38; *m/z* ESI (–ve) 160 ([M–H⁺][–], 75%), 343 ((2[M–H⁺]+Na⁺)[–], 100%). HCl salt⁷ $[\alpha]_{\text{D}}^{21}$ +33.0 (*c.* 0.2, 4.5M HCl); δ_{H} (400 MHz, D₂O) 4.15 (1H, dd, *J* 7, 10), 1.35 (1H, dd, *J* 7, 15), 1.19 (1H, dd, *J* 10, 15); δ_{C} (125 MHz; D₂O) 175.3, 52.4, 18.6 (br); δ^{11}_{B} (160 MHz; D₂O) 24.85.

⁷ The $[\alpha]_{\text{D}}$ was obtained as the HCl salt, obtained by treatment with 10% HCl followed by evaporation.