Highly Selective Supramolecular Catalyzed Allylic Alcohol Isomerization

Dennis H. Leung, Robert G. Bergman, Kenneth N. Raymond

Department of Chemistry, University of California, Berkeley, CA 94720-1460

Supporting Information

Experimental

General. Unless otherwise noted, reactions and manipulations were carried out under inert atmosphere using standard Schlenk and vacuum techniques or in a Vacuum Atmospheres inert atmosphere glove box (N₂) at ambient temperature. ¹H, ¹³C{¹H}, ³¹P{¹H}, and ¹⁹F spectra were recorded at 500, 125, 203, and 376 respectively on a Bruker DRX-500 or AVB-400 spectrometer. All NMR chemical shifts are reported as δ in parts per million (ppm). ¹H and ¹³C{¹H} NMR spectra are reported relative to residual solvent. ³¹P{¹H} NMR spectra are reported relative to trimethyl phosphate as the external standard. ¹⁹F NMR spectra are reported relative to CFCI₃ as the external standard. Coupling constants are reported in Hz. Mass spectrometric (MS) analyses were obtained at the University of California, Berkeley Mass Spectrometry Facility on VT ProSpec, ZAB2-EQ and 70-FE mass spectrometers. All FAB MS data were acquired from samples in a methylene chloride/3-nitrobenzyl alcohol matrix. High resolution TOF electrospray mass spectra (TOF MS ES) were recorded at the Waters facility in Dublin, CA, on a Waters QTOF API mass spectrometer equipped with a Z-spray source. Elemental analyses (EA) were performed by the Microanalytical Laboratory in the College of Chemistry at the University of California, Berkeley. Since an indeterminate

amount of solvent remains fixed to the exterior of the cluster, suitable elemental analyses of the host-guest complexes could not be obtained.

All reagents were used as received from commercial suppliers unless otherwise noted. Anhydrous solvents were passed through a column of activated alumina (type A2, size 12 × 32, Purify) under nitrogen pressure and sparged with nitrogen before use.¹ Deuterated solvents were purchased from Cambridge Isotope Laboratories and were either vacuum transferred from CaH₂ or degassed by three freeze-pump-thaw cycles and placed over activated 4 Å molecular sieves under inert atmosphere. Water and deuterated water were sparged with nitrogen for thirty minutes to remove oxygen and stored under inert atmosphere. Trimethylphosphine (Aldrich) was vacuum-transferred from sodium metal prior to use. Reactions with trimethylphosphine involved condensation of a measured pressure of gas from a bulb of known volume into a reaction vessel at -196 °C. Known-volume bulb vacuum transfers were accomplished with a MKS Baratron gauge attached to a high-vacuum line.

The compounds H₄L and Na₁₂[Ga₄L₆] were prepared using procedure previously developed by Raymond and coworkers.² The complexes [Rh(COD)₂][BF₄] and [Rh(NBD)₂][BF₄] were prepared as previously reported.³ For experiments with gaseous reagents, NMR tubes fitted with Teflon stopcocks (J. Young tubes) were used.

[(PMe₃)₂Rh(COD)][BF₄] (1). A 100-mL Schlenk flask in the glove box was charged with a magnetic stir bar and a solution of [Rh(COD)₂][BF₄] (100 mg, 0.25 mmol) dissolved in 10 mL CH₂Cl₂. The Schlenk flask was removed from the glove box and degassed with three freeze-pump-thaw cycles on a vacuum line. Using a known volume bulb attached to the vacuum line, a known amount of PMe₃ (0.50 mmol) was condensed

into the flask at -196 °C. The solution immediately turned orange and was stirred for 1 hour. The flask was brought into the glove box and the orange solution was filtered through Celite. The solvent was removed *in vacuo* to leave an orange solid. The product was washed with Et₂O, dried under vacuum, and stored in the glove box under N₂. Yield: 86 mg, 76%. ¹H NMR (CD₂Cl₂): δ 4.91 (br, 4H, alkenyl CH), 2.36 (br, 8H, CH₂), 1.43 (d, *J*_{P-H} = 5.6 Hz, 18H, P(CH₃)₃). ¹³C{¹H} NMR (CD₂Cl₂): δ 97.0 (s, alkenyl C), 31.1 (s, CH₂), -15.8 (m, P(CH₃)₃). ³¹P{¹H} NMR (CD₂Cl₂): δ -15.1 (d, *J*_{Rh-P} = 142.6 Hz, P(CH₃)₃). ¹⁹F NMR (CD₂Cl₂): δ -151.5 (s, BF₄). Anal. Cald for C₁₄H₃₀P₂RhBF₄: C, 37.36; H, 6.72. Found: C, 37.13; H, 6.92.

[(dmpe)Rh(COD)][BF₄] (2). In the glove box, a 20-mL scintillation vial was charged with a magnetic stir bar and a solution of [Rh(COD)₂][BF₄] (100 mg, 0.25 mmol) dissolved in 10 mL CH₂Cl₂. A stoichiometric amount of dmpe (0.25 mmol) was added via a gas-tight syringe and the solution was stirred for 1 hour. The resulting orange solution was filtered through Celite and the solvent was removed *in vacuo* to leave an orange solid. The product was washed with Et₂O, dried under vacuum, and stored in the glove box under N₂. Yield: 91 mg, 81%. ¹H NMR (CD₂Cl₂): δ 5.12 (br, 4H, alkenyl CH), 2.45 (m, 4H, CH₂), 2.31 (m, 4H, CH₂), 1.84 (d, *J*_{P-H} = 18.4 Hz, 4H, P(CH₂)), 1.50 (d, *J*_{P-H} = 8.8 Hz, 12H, P(CH₃)). ¹³C{¹H} NMR (CD₂Cl₂): δ 98.8 (m, alkenyl C), 30.6 (s, CH₂), 27.5 (m, P(CH₂)), 13.5 (m, P(CH₃)₃). ³¹P{¹H} NMR (CD₂Cl₂): δ 35.8 (d, *J*_{Rh-P} = 145.8 Hz, *dmpe*). ¹⁹F NMR (CD₂Cl₂): δ -151.5 (s, BF₄). Anal. Cald for C₁₄H₂₈P₂RhBF₄: C, 37.53; H, 6.30. Found: C, 37.92; H, 6.30.

[(PMe₃)₂Rh(NBD)][BF₄] (3). A 100-mL Schlenk flask in the glove box was charged with a magnetic stir bar and a solution of [Rh(NBD)₂][BF₄] (100 mg, 0.27 mmol)

dissolved in 10 mL CH₂Cl₂. The Schlenk flask was removed from the glove box and degassed with three freeze-pump-thaw cycles on a vacuum line. Using a known volume bulb attached to the vacuum line, a known amount of PMe₃ (0.54 mmol) was condensed into the flask at -196 °C. The solution immediately turned orange and was stirred for 1 hour. The flask was brought into the glove box and the orange solution was filtered through Celite. The solvent was removed *in vacuo* to leave an orange solid. The product was washed with Et₂O, dried under vacuum, and stored in the glove box under N₂. Yield: 109 mg, 93%. ¹H NMR (CD₂Cl₂): δ 5.00 (br, 4H, alkenyl CH), 4.08 (s, 2H, CH), 1.74 (s, 2H, CH₂), 1.29 (m, 18H, P(CH₃)). ¹³C{¹H} NMR (CD₂Cl₂): δ 82.9 (m, alkenyl C), 69.3 (m, CH), 54.9 (s, CH₂), 14.8 (m, P(CH₃)). ³¹P{¹H} NMR (CD₂Cl₂): δ -13.7 (d, *J*_{Rh-P} = 153.9 Hz, P(CH₃)). ¹⁹F NMR (CD₂Cl₂): δ -151.5 (s, BF₄). Anal. Calcd for C₁₃H₂₆P₂RhBF₄: C, 35.98; H, 6.04. Found: C, 35.84; H, 5.76.

[(dmpe)Rh(NBD)][BF₄] (4). In the glove box, a 20-mL scintillation vial was charged with a magnetic stir bar and a solution of [Rh(NBD)₂][BF₄] (100 mg, 0.27 mmol) dissolved in 10 mL CH₂Cl₂. A stoichiometric amount of dmpe (0.27 mmol) was added via a gas-tight syringe and the solution was stirred for 1 hour. The resulting orange solution was filtered through Celite and the solvent was removed *in vacuo* to leave an orange solid. The product was washed with Et₂O, dried under vacuum, and stored in the glove box under N₂. Yield: 106 mg, 91%. ¹H NMR (CD₂Cl₂): δ 5.45 (m, 4H, alkenyl CH), 4.17 (s, 2H, CH), 1.82 (s, 2H, CH₂), 1.78 (d, *J*_{P-H} = 17.6 Hz, 4H, CH₂), 1.40 (d, *J*_{P-H} = 12.0 Hz, 12H, P(CH₃)). ¹³C{¹H} NMR (CD₂Cl₂): δ 88.9 (m, alkenyl C), 71.5 (m, CH), 56.8 (s, CH₂), 27.9 (m, P(CH₂)), 12.6 (m, P(CH₃)). ³¹P{¹H} NMR (CD₂Cl₂): δ -13.7 (d,

$J_{\text{Rh-P}} = 153.9 \text{ Hz}$, $P(\text{CH}_3)$). ^{19}F NMR (CD_2Cl_2): δ -151.5 (s, BF_4). Anal. Calcd for $\text{C}_{13}\text{H}_{24}\text{P}_2\text{RhBF}_4$: C, 36.15; H, 5.60. Found: C, 36.80; H, 5.86.

$[(\text{PEt}_3)_2\text{Rh}(\text{COD})][\text{BF}_4]$ (5). In the glove box, a 20-mL scintillation vial was charged with a magnetic stir bar and a solution of $[\text{Rh}(\text{COD})_2][\text{BF}_4]$ (100 mg, 0.25 mmol) dissolved in 10 mL CH_2Cl_2 . Two equivalents of PEt_3 (0.50 mmol) were added via a gas-tight syringe and the solution was stirred for 1 hour. The resulting orange solution was filtered through Celite and the solvent was removed *in vacuo* to leave an orange solid. The product was washed with Et_2O , dried under vacuum, and stored in the glove box under N_2 . Yield: 114 mg, 85%. ^1H NMR (CD_2Cl_2): δ 4.97 (br, 4H, alkenyl CH), 2.32 (br, 8H, CH_2), 1.73 (m, 12H, $\text{P}(\text{CH}_2\text{CH}_3)$), 1.17 (m, 18H, $\text{P}(\text{CH}_2\text{CH}_3)$). $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2): δ 95.6 (m, alkenyl C), 31.1 (s, CH_2), 15.6 (m, $\text{P}(\text{CH}_2\text{CH}_3)$), 8.7 (s, $\text{P}(\text{CH}_2\text{CH}_3)$). $^{31}\text{P}\{^1\text{H}\}$ NMR (CD_2Cl_2): δ 12.1 (d, $J_{\text{Rh-P}} = 140.9 \text{ Hz}$, PEt_3). ^{19}F NMR (CD_2Cl_2): δ -151.5 (s, BF_4). Anal. Calcd for $\text{C}_{20}\text{H}_{42}\text{P}_2\text{RhBF}_4$: C, 44.97; H, 7.92. Found: C, 44.67; H, 7.88.

$(\text{PMe}_3)_2\text{Rh}(\text{OD}_2)[\text{BF}_4]$ (6). In the glove box, a J. Young NMR tube was charged with a solution of **1** (3.0 mg, 6.7×10^{-3} mmol) in 0.5 mL D_2O . The tube was capped, brought out, and degassed via three freeze-pump-thaw cycles on a vacuum line. An excess of H_2 (1 atm) was added and the solution was shaken at room temperature for 2 hours. The color slowly turned from orange to dark yellow. The solution was then degassed via three freeze-pump-thaw cycles to remove excess H_2 . This reactive compound was prepared *in situ* immediately before addition of substrates and was not isolated. A broad resonance for cyclooctane is observed between 1-2 ppm in the ^1H

NMR spectrum. ^1H NMR (D_2O): δ 1.65 (d, $J_{\text{P-H}} = 11.6$ Hz, 18H, $\text{P}(\text{CH}_3)_3$). $^{31}\text{P}\{^1\text{H}\}$ NMR (D_2O): δ 20.3 (d, $J_{\text{Rh-P}} = 154.3$ Hz, $\text{P}(\text{CH}_3)_3$). ^{19}F NMR (D_2O): δ -151.5 (s, BF_4).

General Procedure for Preparation of Rhodium Ga_4L_6 Host-Guest Assemblies. In the glove box, an NMR tube was charged with a solution of the rhodium species (~ 2.3 mg, 5×10^{-3} mmol) dissolved in 0.5 mL D_2O . A stoichiometric amount of $\text{Na}_{12}[\text{Ga}_4\text{L}_6]$ (~ 18 mg, 5×10^{-3} mmol) was added as a solid to this mixture to form a bright yellow/orange solution. The NMR tube was then removed from the glove box and flame-sealed under static vacuum on a vacuum line. The host-guest complex can be isolated by removal of solvent *in vacuo*.

$\text{Na}_{11}[\mathbf{1} \subset \text{Ga}_4\text{L}_6]$. This compound was prepared according to the general procedure outlined above. ^1H NMR (D_2O): δ 8.15 (d, $J = 7.5$ Hz, 12H, *aryl*), 7.63 (d, $J = 8.5$ Hz, 12H, *aryl*), 7.40 (d, $J = 8.0$ Hz, 12H, *aryl*), 7.22 (d, $J = 8.0$ Hz, 4H, *aryl*), 7.06 (d, $J = 7.0$ Hz, 4H, *aryl*), 6.88 (d, $J = 7.5$ Hz, 4H, *aryl*), 6.78 (d, $J = 7.0$ Hz, 12H, *aryl*), 6.64 (t, $J = 8.0$ Hz, 12H, *aryl*), 1.54 (br, 2H, COD), 1.46 (br, 2H, COD), 1.32 (br, 2H, COD), 1.08 (br, 2H, COD), 0.71 (br, 2H, COD), 0.35 (br, 2H, COD), -0.70 (br, 18H, $\text{P}(\text{CH}_3)_3$). $^{13}\text{C}\{^1\text{H}\}$ DEPT 135 NMR (CD_2Cl_2): δ 127.0 (s, CH *aryl*), 119.2 (s, CH *aryl*), 118.7 (s, CH *aryl*), 116.1 (s, CH *aryl*), 115.7 (s, CH *aryl*), 115.4 (s, CH *aryl*), 96.2 (m, COD CH), 94.3 (m, COD CH), 67.2 (s, COD CH_2), 33.4 (s, COD CH_2), 26.3 (s, COD CH_2), 13.5 (m, $\text{P}(\text{CH}_3)_3$). $^{31}\text{P}\{^1\text{H}\}$ NMR (D_2O): δ -17.1 (d, $J_{\text{Rh-P}} = 142.6$ Hz, $\text{P}(\text{CH}_3)_3$). ^{19}F NMR (D_2O): δ -151.5 (s, BF_4). TOF MS ES(-) ($\text{H}_2\text{O}/\text{MeOH}$, 50:50), $\diamond = [\text{Ga}_4\text{L}_6]^{12-}$, calcd. (found), m/z : 1092.04 (1092.04) [$\diamond + 8\text{Na}^+ + \text{Rh}(\text{PMe}_3)_2^+$] $^{3-}$, 1084.75 (1084.72) [$\diamond + 7\text{Na}^+ + \text{H}^+ + \text{Rh}(\text{PMe}_3)_2^+$] $^{3-}$, 1077.39 (1077.39) [$\diamond + 6\text{Na}^+ + 2\text{H}^+ + \text{Rh}(\text{PMe}_3)_2^+$] $^{3-}$, 1070.06 (1070.07) [$\diamond + 5\text{Na}^+ + 3\text{H}^+ + \text{Rh}(\text{PMe}_3)_2^+$] $^{3-}$, 1062.74 (1062.73) [$\diamond + 4\text{Na}^+ + 4\text{H}^+ + \text{Rh}(\text{PMe}_3)_2^+$] $^{3-}$,

Na₁₁[2 ⊂ Ga₄L₆]. This compound was prepared according to the general procedure outlined above. ¹H NMR (D₂O): δ 8.22 (d, *J* = 10.0 Hz, 12H, *aryl*), 7.75 (d, *J* = 8.5 Hz, 12H, *aryl*), 7.46 (dd, *J* = 8.5 Hz, 1.5 Hz, 12H, *aryl*), 7.43 (d, *J* = 8.0 Hz, 4H, *aryl*), 7.43 (d, *J* = 8.0 Hz, 4H, *aryl*), 7.28 (d, *J* = 8.0 Hz, 4H, *aryl*), 7.12 (d, *J* = 7.5 Hz, 4H, *aryl*), 6.95 (t, *J* = 8.0 Hz, 12H, *aryl*), 6.84 (dd, *J* = 7.5 Hz, 1.5 Hz, 12H, *aryl*), 6.70 (t, *J* = 7.5 Hz, 12H, *aryl*), 1.74 (br, 4H, COD), 1.06 (br, 2H, COD), 0.91 (br, 2H, COD), 0.81 (br, 2H, COD), 0.47 (m, 2H, COD), -0.62 (d, *J* = 7.5 Hz, 6H, P(CH₃)₂), -0.86 (d, *J* = 8.5 Hz, 6H, P(CH₃)₂), -1.20 (m, 4H, P(CH₂)). ¹³C{¹H} DEPT 135 NMR (CD₂Cl₂): δ 126.6 (s, CH *aryl*), 119.8 (s, CH *aryl*), 118.7 (s, CH *aryl*), 116.1 (s, CH *aryl*), 115.7 (s, CH *aryl*), 115.4 (s, CH *aryl*), 96.1 (m, COD CH), 95.4 (m, COD CH), 67.2 (s, COD CH₂), 31.2 (s, COD CH₂), 27.7 (s, COD CH₂), 11.1 (m, P(CH₃)₃). ³¹P{¹H} NMR (D₂O): δ 34.5 (d, *J*_{Rh-P} = 145.8 Hz, P(CH₃)). ¹⁹F NMR (D₂O): δ -151.5 (s, BF₄).

Na₁₁[3 ⊂ Ga₄L₆]. This compound was prepared according to the general procedure outlined above. ¹H NMR (D₂O): δ 8.13 (d, *J* = 7.6 Hz, 12H, *aryl*), 7.70 (d, *J* = 8.4 Hz, 12H, *aryl*), 7.35 (dd, *J* = 8.0 Hz, 1.6 Hz, 12H, *aryl*), 7.20 (d, *J* = 8.0 Hz, 4H, *aryl*), 6.94 (t, *J* = 8.0 Hz, 4H, *aryl*), 6.86 (dd, *J* = 8.0 Hz, 1.5 Hz, 4H, *aryl*), 6.76 (dd, *J* = 7.2 Hz, 2.0 Hz, 4H, *aryl*), 6.63 (m, 12H, *aryl*), 2.30 (br, 2H, NBD), 1.94 (br, 2H, NBD), 0.11 (br, 2H, NBD), -0.57 (br, 2H, NBD), -1.35 (m, 18H, NBD). ¹⁹F NMR (D₂O): δ -151.5 (s, BF₄).

Na₁₁[4 ⊂ Ga₄L₆]. This compound was prepared according to the general procedure outlined above. ¹H NMR (D₂O): δ 8.11 (d, *J* = 7.6 Hz, 12H, *aryl*), 7.74 (d, *J* = 8.4 Hz, 12H, *aryl*), 7.36 (dd, *J* = 8.2 Hz, 1.6 Hz, 12H, *aryl*), 7.20 (d, *J* = 8.4 Hz, 4H, *aryl*), 6.98 (t, *J* = 8.0 Hz, 4H, *aryl*), 6.86 (dd, *J* = 7.2 Hz, 4H, *aryl*), 6.76 (dd, *J* = 7.2 Hz,

1.6 Hz, 4H, *aryl*), 6.62 (m, 12H, *aryl*), 2.71 (br, 2H, NBD), 2.42 (br, 2H, NBD), 1.94 (br, 2H, NBD), 0.29 (br, 2H, NBD), -0.79 (d, $J_{\text{P-H}} = 7.6$ Hz, 6H, $\text{P}(\text{CH}_3)_2$), -1.14 (d, $J_{\text{P-H}} = 8.4$ Hz, 6H, $\text{P}(\text{CH}_3)_2$), -1.51 (m, 4H, $\text{P}(\text{CH}_2)_2$). $^{31}\text{P}\{^1\text{H}\}$ NMR (D_2O): δ 32.6 (d, $J_{\text{Rh-P}} = 154.9$ Hz, $\text{P}(\text{CH}_3)_2$). ^{19}F NMR (D_2O): δ -151.5 (s, BF_4).

$\text{Na}_{11}[\mathbf{6} \subset \mathbf{Ga}_4\mathbf{L}_6]$. A J. Young NMR tube was charged with $\text{Na}_{11}[\mathbf{1} \subset \mathbf{Ga}_4\mathbf{L}_6]$ dissolved in D_2O prepared as above. The solution was degassed via three freeze-pump-thaw cycles. An excess of H_2 (1 atm) was added and the solution was allowed to sit at room temperature for 2 hours. The color slowly turned from light orange to dark red. The solution was then degassed via three freeze-pump-thaw cycles to remove excess H_2 . This reactive compound was prepared in situ immediately before addition of substrates and was not isolated. A broad resonance for cyclooctane is observed between 1-2 ppm in the ^1H NMR spectrum. ^1H NMR (D_2O): δ 8.13 (m, *aryl*), 7.82 (d, $J = 8.0$ Hz, *aryl*), 7.61 (d, $J = 8.0$ Hz, *aryl*), 7.39 (d, $J = 8.0$ Hz, *aryl*), 7.30 (d, $J = 8.0$ Hz, *aryl*), 7.20 (d, $J = 8.0$ Hz, *aryl*), 7.03 (t, $J = 8.0$ Hz, *aryl*), 6.86 (d, $J = 7.2$ Hz, *aryl*), 6.76 (t, $J = 7.2$ Hz, *aryl*), 6.60 (m, *aryl*), -1.97 (br m, 18H, $\text{P}(\text{CH}_3)_3$). $^{31}\text{P}\{^1\text{H}\}$ NMR (D_2O): δ -6.5 (d, $J_{\text{Rh-P}} = 197.6$ Hz, $\text{P}(\text{CH}_3)_3$). ^{19}F NMR (D_2O): δ -151.5 (s, BF_4).

General Procedure for Allylic Alcohol Isomerization Reactions. A J. Young NMR tube was charged with either **6** or $\text{Na}_{11}[\mathbf{6} \subset \mathbf{Ga}_4\mathbf{L}_6]$ prepared as above with 2 μL of dioxane as an internal standard. This was brought into the glove box and 10 equivalents of the appropriate allylic alcohol substrate were added directly via a gas tight syringe. An initial time point of the reaction was then obtained immediately by NMR spectroscopy. The reaction progress was monitored by NMR spectroscopy using dioxane as the

reference standard. The solution was allowed to stand at room temperature until the reaction was complete (0.5 h).

Addition of Crotyl Alcohol to 6. A J. Young NMR tube containing **6** dissolved in D₂O was prepared as above. This was brought into the glove box and 10 equivalents of crotyl alcohol were added via gas tight syringe. The solution was allowed to stand at room temperature for 30 min. No isomerization product was observed by NMR spectroscopy. Coordination of crotyl alcohol was difficult to discern in the ¹H NMR spectrum, but is suggested by shifts upfield in the ³¹P NMR spectrum. The two new upfield doublets in the ³¹P NMR spectrum may be due to mono- and bis-coordination of excess crotyl alcohol or multiple binding geometries. This compound was prepared in situ was not isolated. ³¹P{¹H} NMR (CD₂Cl₂): δ -10.3 (d, *J*_{Rh-P} = 110.6 Hz, *P*(CH₃) A), -15.2 (d, *J*_{Rh-P} = 141.2 Hz, *P*(CH₃) B). ¹⁹F NMR (CD₂Cl₂): δ -151.5 (s, BF₄).

References

- (1) Alaimo, P. J.; Peters, D. W.; Arnold, J.; Bergman, R. G. *J. Chem. Educ.* **2001**, 78, 64.
- (2) Caulder, D. L.; Bruckner, C.; Powers, R. E.; Konig, S.; Parac, T. N.; Leary, J. A.; Raymond, K. N. *J. Am. Chem. Soc.* **2001**, 123, 8893-8938.
- (3) Schenck, T. G.; Downes, J. M.; Milne, C. R. C.; Mackenzie, P. B.; Boucher, H.; Whelan, J.; Bosnich, B. *Inorg. Chem.* **1985**, 24, 2334-2337.

Figure S-1. ^1H NMR spectrum of $\text{Na}_{11}[\mathbf{1} \subset \text{Ga}_4\text{L}_6]$ in D_2O .

Figure S-2. ^1H NMR spectrum of $\text{Na}_{11}[\mathbf{2} \subset \text{Ga}_4\text{L}_6]$ in D_2O .

Figure S-3. ^1H NMR spectrum of $\text{Na}_{11}[\mathbf{3} \subset \text{Ga}_4\text{L}_6]$ in D_2O .

Figure S-4. ^1H NMR spectrum of $\text{Na}_{11}[\mathbf{4} \subset \text{Ga}_4\text{L}_6]$ in D_2O .

Figure S-5. ^1H NMR spectrum of **6** in D_2O .

Figure S-6. $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **6** in D_2O .

Figure S-7. ^1H NMR spectrum of $\text{Na}_{11}[\mathbf{6} \subset \text{Ga}_4\text{L}_6]$ in D_2O .

Figure S-8. $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum **6** in the presence of crotyl alcohol in D_2O .