Supplementary Material

Highly Stereoselective Synthesis of (E)-, (Z)- α -Fluoro- α,β -unsaturated esters and (E)-, (Z)- α -Fluoro- α,β -unsaturated Amides from 1-Bromo-1-fluoroalkenes via Palladium-Catalyzed Carbonylation Reactions

Jianjun Xu and Donald J. Burton*

Department of Chemistry, University of Iowa, Iowa City, Iowa 52242

donald-burton@uiowa.edu

Supporting Information Available: Experimental procedures for the synthesis of **Z-6e**, **Z-6f**, **Z-8 (b-g)**, **E-8b**, **E-8g**, **E-8h** and their characterization by ^{19}F , ^1H , ^{13}C NMR, GC-MS, and HRMS; Copies of ^1H and ^{13}C NMR spectra of compounds **Z-6 (a-g)**, **E-6a**, **E-6b**, **E-6c**, **E-6g**, **E-6h**, **Z-8 (a-g)**, **E-8a**, **E-8b**, **E-8g**, **E-8h**; NMR experiment on the existence of intramolecular F...HN coupling in (Z)- and (E)- α -fluoro- α,β -unsaturated amides. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Table of Contents

General Methods.....	S4
Materials.....	S4
Synthesis of (Z)- <i>n</i> -Butyl 2-fluoro-3-(4-fluorophenyl)propenoate (Z-6e).....	S4
Synthesis of (Z)- <i>n</i> -Butyl 2-fluoro-3-(3-nitrophenyl)propenoate (Z-6f).....	S5
General procedures for the preparation of (Z)- α -fluoro- α,β -unsaturated amides Z-8.....	S6
Synthesis of (Z)-3-(2-Chlorophenyl)-2-fluoro- <i>N</i> -phenylacrylamide (Z-8b).....	S7
Synthesis of (Z)-2-Fluoro-3-(4-methoxyphenyl)- <i>N</i> -phenylacrylamide (Z-8c).....	S7
Synthesis of (Z)-3-(4-Chlorophenyl)-2-fluoro- <i>N</i> -phenylacrylamide (Z-8d).....	S8
Synthesis of (Z)-2-Fluoro-3-(4-fluorophenyl)- <i>N</i> -phenylacrylamide (Z-8e).....	S9
Synthesis of (Z)-2-Fluoro-3-(3-nitrophenyl)- <i>N</i> -phenylacrylamide (Z-8f).....	S9
Synthesis of (Z)-2-Fluoro-4-phenylpent-2-enanilide (Z-8g).....	S10
General procedures for the preparation of (E)- α -fluoro- α,β -unsaturated amides E-8....	S11
Synthesis of (E)-3-(2-Chlorophenyl)-2-fluoro- <i>N</i> -phenylacrylamide (E-8b).....	S12
Synthesis of (E)-2-Fluoro-4-phenylpent-2-enanilide (E-8g).....	S12
Synthesis of (E)-2-Fluoro-3-(1-naphthyl)- <i>N</i> -phenylacrylamide (E-8h).....	S13
¹ H and ¹³ C NMR spectra of Z-6, E-6, Z-8 and E-8.....	S15
¹ H and ¹³ C NMR spectra of Z-6a.....	S15
¹ H and ¹³ C NMR spectra of Z-6b.....	S16
¹ H and ¹³ C NMR spectra of Z-6c.....	S17
¹ H NMR spectra of Z-6d.....	S18
¹³ C NMR spectra of Z-6d.....	S20
¹ H NMR spectra of Z-6e.....	S21

^{13}C NMR spectra of Z-6e.....	S23
^1H and ^{13}C NMR spectra of Z-6f.....	S24
^1H and ^{13}C NMR spectra of Z-6g.....	S25
^1H and ^{13}C NMR spectra of E-6a.....	S26
^1H and ^{13}C NMR spectra of E-6b.....	S27
^1H and ^{13}C NMR spectra of E-6c.....	S28
^1H and ^{13}C NMR spectra of E-6g.....	S29
^1H and ^{13}C NMR spectra of E-6h.....	S30
^1H and ^{13}C NMR spectra of Z-8a.....	S31
^1H and ^{13}C NMR spectra of Z-8b.....	S32
^1H and ^{13}C NMR spectra of Z-8c.....	S33
^1H NMR spectra of Z-8d.....	S34
^{13}C NMR spectra of Z-8d.....	S36
^1H NMR spectra of Z-8e.....	S37
^{13}C NMR spectra of Z-8e.....	S39
^1H and ^{13}C NMR spectra of Z-8f.....	S40
^1H and ^{13}C NMR spectra of Z-8g.....	S41
^1H and ^{13}C NMR spectra of E-8a.....	S42
^1H and ^{13}C NMR spectra of E-8b.....	S43
^1H and ^{13}C NMR spectra of E-8g.....	S44
^1H and ^{13}C NMR spectra of E-8h.....	S45
NMR experiment on the existence of intramolecular F...HN coupling in (Z)- and (E)- α -fluoro- α,β -unsaturated amides.....	S46

Supplemental Experimental Section

General Methods. The ^1H , ^{19}F , ^{13}C spectra of the final products were obtained on a 300 MHz NMR spectrometer (TMS, CFCl_3 or CDCl_3 as internal references respectively). GC-MS analyses were performed at 70 eV in the electron-impact mode. Melting points are uncorrected.

Materials. DMF was dried by stirring overnight over CaH_2 , then distilled at reduced pressure. NBu_3 was dried over NaH before distillation. $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$ was prepared by Negishi's procedure.⁴⁸ 1-Bromo-1-fluoroalkenes ($E/Z \approx 1:1$) **1** were prepared from the corresponding aldehydes and CFBr_3 , PPh_3 as described in the literature³⁶ except that THF was used as the solvent.

(Z)-*n*-Butyl 2-fluoro-3-(4-fluorophenyl)propenoate (Z-6e). Similarly, a mixture of 1-bromo-1-fluoro-2-(4-fluorophenyl)ethylene (0.11 g, 0.5 mmol, $E/Z = 87:13$), NBu_3 (0.10 g, 0.57 mmol), $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$ (0.01 g, 0.02 mmol) in *n*-butanol (5 mL) was reacted with CO at room temperature for 144 hours. When the reaction was completed, ^{19}F NMR analysis of the mixture showed that the Z/E ratio of the crude product was 96:4. Silica gel column chromatography (ethyl acetate: hexanes = 5:95, $R_f = 0.32$) followed by recrystallization from hexanes gave white crystals, 0.09 g, yield 75 % (conversion 86 % based on the consumed (E)-1-bromo-1-fluoro-2-(4-fluorophenyl)ethylene), $Z/E = 100:0$. ^{19}F NMR (CDCl_3) δ -110.2 (m, 1 F), -126.8 (d, $^3J_{\text{FH}(\text{trans})} = 34.1$ Hz, 1 F) ppm; ^1H NMR (CDCl_3) δ 7.64 (dd, $J = 8.6$ Hz, $J = 5.6$ Hz, 2 H), 7.09 (t, $J = 8.6$ Hz, 2 H), 6.88 (d, $^3J_{\text{HF}(\text{trans})} = 34.8$ Hz, 1 H), 4.29 (t, $J = 6.6$ Hz, 2 H), 1.73 (pentet, $J = 7.1$ Hz, 2 H), 1.45

(sextet, $J = 7.4$ Hz, 2 H), 0.98 (t, $J = 7.5$ Hz, 3 H) ppm; ^{13}C NMR (CDCl_3) δ 163.2 (dd, $^1J_{\text{CF}} = 251.5$ Hz, $J = 3.5$ Hz), 161.4 (d, $J = 34.0$ Hz), 146.7 (d, $J = 263.6$ Hz), 132.2 (t, $J = 8.3$ Hz), 127.4 (t, $J = 3.5$ Hz), 116.3 (d, $J = 5.0$ Hz), 116.0 (d, $^2J_{\text{CF}} = 22.3$ Hz), 65.8, 30.6, 19.1, 13.7 ppm. GC-MS m/z (relative intensity): 241 ($\text{M}^+ + 1$, 11), 240 (M^+ , 58), 220 (9), 185 (37), 184 (100), 167 (61), 164 (68), 147 (34), 139 (49), 138 (40), 136 (41), 120 (26), 119 (65), 109 (17), 99 (43); HRMS calculated 240.0962 for $\text{C}_{13}\text{H}_{14}\text{O}_2\text{F}_2$, observed 240.0955.

(Z)-*n*-Butyl 2-fluoro-3-(3-nitrophenyl)propenoate (Z-6f). Similarly, a mixture of 1-bromo-1-fluoro-2-(3-nitrophenyl)ethylene (0.30 g, 1.23 mmol, E/Z = 81: 19), NBu_3 (0.24 g, 1.3 mmol), $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$ (0.03 g, 0.04 mmol) in *n*-butanol (4 mL) was reacted with CO at room temperature for 96 hours. ^{19}F NMR analysis of the mixture showed that the Z/E ratio of the crude product was 97: 3. Silica gel column chromatography (ethyl acetate: hexanes = 10: 90, $R_f = 0.33$) followed by recrystallization from hexanes gave white crystals, mp: 63-64 °C, 0.22 g, yield 67 % (conversion 83 % based on the consumed (E)-1-bromo-1-fluoro-2-(3-nitrophenyl)ethylene), Z/E = 100: 0. ^{19}F NMR (CDCl_3) δ -121.3 (d, $^3J_{\text{FH}(\text{trans})} = 33.5$ Hz, 1 F) ppm; ^1H NMR (CDCl_3) δ 8.48 (m, 1 H), 8.23 (dd, $J = 8.2$ Hz, $J = 1.9$ Hz, 1 H), 7.97 (d, $J = 8.0$ Hz, 1 H), 7.61 (t, $J = 8.0$ Hz, 1 H), 6.97 (d, $^3J_{\text{HF}(\text{trans})} = 33.5$ Hz, 1 H), 4.33 (t, $J = 6.7$ Hz, 2 H), 1.75 (pentet, $J = 7.1$ Hz, 2 H), 1.46 (sextet, $J = 7.4$ Hz, 2 H), 0.99 (t, $J = 7.3$ Hz, 3 H) ppm; ^{13}C NMR (CDCl_3) δ 160.4 (d, $J = 34.0$ Hz), 148.3 (d, $^1J_{\text{CF}} = 274.0$ Hz), 148.2, 135.4 (d, $J = 8.6$ Hz), 132.5 (d, $J = 5.1$ Hz), 129.7, 124.4 (d, $J = 8.7$ Hz), 123.8, 114.7 (d, $^2J_{\text{CF}} = 4.1$ Hz), 65.9, 30.3, 18.9, 13.4 ppm; GC-MS m/z (relative intensity): 268 ($\text{M}^+ + 1$, 3), 267 (M^+ , 22), 247 (7), 212 (40), 211 (80), 195 (47), 194 (68), 165 (28), 164 (28), 148 (16), 147 (20), 120 (70), 101 (22), 94 (18), 57

(20), 56 (100); HRMS: MI peak was not detected; elemental analysis: theoretical C 58.43, H 5.24, N 5.24, F 7.12, observed C 58.63, H 5.33, N 5.20, F 6.97.

General procedures for the preparation of (Z)- α -fluoro- α,β -unsaturated amides

Z-8. An oven-dried 120 mL Hastelloy Parr pressure reactor equipped with a stirring bar was charged with 0.07 g (0.10 mmol) of $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$, 0.36 g (3.8 mmol) of freshly distilled aniline, and 4.0 mL of tri-*n*-butylamine. 3.0 Mmol of high E/Z ratio 1-bromo-1-fluoroalkene was added and the pressure reactor was closed (*caution: All reactions should be carried out behind a safety shield in a well-ventilated hood*). The pressure reactor was pressurized to 160 psi with carbon monoxide, and the pressure was released. This process was repeated four times to rid the system of air. Finally this pressure reactor was pressurized to 160 psi and was allowed to stir at room temperature (for 1-bromo-1-fluoroalkenes, R= aryl) or in an oil bath at 50 °C (for 1-bromo-1-fluoroalkenes, R= alkyl). The consumption of carbon monoxide could be roughly determined by the reading on the gauge. When the reaction was completed, the reactor was allowed to cool to room temperature (if necessary) and the pressure was carefully released. The reaction mixture was transferred to a separatory funnel containing 100 mL of ethyl acetate. The organic layer was washed with aqueous 10 % hydrochloric acid (3 X 15 mL), saturated sodium bicarbonate (2 X 15 mL) and brine (2 X 10 mL). The combined organic layers were dried over anhydrous MgSO_4 , concentrated by rotary evaporation, and the crude product was further purified by silica gel column chromatography. Recrystallization from hexanes gave 100 % pure (Z)- α -fluoro- α,β -unsaturated amides.

(Z)-3-(2-Chlorophenyl)-2-fluoro-N-phenylacrylamide (Z-8b). Similarly, a mixture of 1-bromo-1-fluoro-2-(2-chlorophenyl)ethylene (0.71 g, 3.0 mmol, E/Z = 82: 18), aniline (0.34 g, 3.65 mmol), $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$ (0.07 g, 0.10 mmol) in tributylamine (7 mL) was reacted with CO at room temperature for 148 hours. ^{19}F NMR analysis of the mixture showed that the Z/E ratio of the crude product was 89: 11. Silica gel column (ethyl acetate: hexanes = 15: 85, R_f = 0.32) followed by recrystallization from hexanes/ether gave white crystals, mp: 127-128 °C, 0.61 g, yield 74 % (90 % conversion based on the consumed (E)-1-bromo-1-fluoro-2-(2-chlorophenyl)ethylene), Z/E = 100: 0. ^{19}F NMR (CDCl_3) δ -129.0 ppm (dd, $^3J_{\text{FH}(\text{trans})}$ = 37.8 Hz, $J_{\text{F}\cdots\text{HN}}$ = 5.6 Hz, 1 F) ppm; ^1H NMR (CDCl_3) δ 8.02 (broad s, 1 H), 7.85-7.90 (m, 1 H), 7.65 (d, J = 8.0 Hz, 2 H), 7.52 (d, $^3J_{\text{FH}(\text{trans})}$ = 38.5 Hz, 1 H), 7.44-7.47 (m, 1 H), 7.39 (t, J = 7.8 Hz, 2 H), 7.31 (m, 2 H), 7.19 (t, J = 7.0 Hz, 1 H) ppm; ^{13}C NMR (CDCl_3) δ 157.9 (d, $^2J_{\text{CF}}$ = 26.6 Hz), 150.9 (d, $^1J_{\text{CF}}$ = 279.4 Hz), 136.9, 134.9, 131.1 (d, J = 11.9 Hz), 130.5, 130.1, 129.5 (d, J = 3.7 Hz), 129.3, 127.1, 125.4, 120.4, 110.6 (d, $^2J_{\text{CF}}$ = 3.5 Hz) ppm; GC-MS m/z (relative intensity): 277 ($\text{M}^+ + 2$, 4), 275 (M^+ , 12), 241 (17), 240 (100), 183 (17), 120 (49); HRMS calculated 275.0513 for $\text{C}_{15}\text{H}_{11}\text{F}^{35}\text{ClON}$, observed 275.0501.

(Z)-2-Fluoro-3-(4-methoxyphenyl)-N-phenylacrylamide (Z-8c). Similarly, a mixture of 1-bromo-1-fluoro-2-(4-methoxyphenyl)ethylene (0.46 g, 2.0 mmol, E/ Z = 81: 19), aniline (0.23 g, 2.43 mmol), $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$ (0.05 g, 0.06 mmol) in tributylamine (7 mL) was reacted with CO at room temperature for 336 hours. ^{19}F NMR analysis of the mixture showed that the Z/E ratio of the crude product was 96: 4. Silica gel column chromatography (ethyl acetate: hexanes = 15: 85, R_f = 0.25) followed by recrystallization from hexanes/ ether gave white crystals, mp: 156-157 °C, 0.38 g, yield 70 % (conversion

87 % based on the consumed (E)-1-bromo-1-fluoro-2-(4-methoxyphenyl)ethylene), Z/E = 100: 0. ^{19}F NMR (CDCl_3) δ -132.9 (d, $^3J_{\text{FH}(\text{trans})} = 39.3$ Hz, 1 F) ppm; ^1H NMR (CDCl_3) δ 7.99 (broad, 1 H), 7.59-7.65 (m, 4 H), 7.38 (tm, $J = 7.9$ Hz, 2 H), 7.17 (tm, $J = 7.4$ Hz, 1 H), 7.02 (d, $^3J_{\text{HF}(\text{trans})} = 39.9$ Hz, 1 H), 6.94 (d, $J = 8.9$ Hz, 2 H), 3.85 (s, 3 H) ppm; ^{13}C NMR (CDCl_3) δ 160.7 (d, $J = 2.6$ Hz), 158.7 (d, $^2J_{\text{CF}} = 27.7$ Hz), 148.9 (d, $^1J_{\text{CF}} = 271.6$ Hz), 137.2, 132.1 (d, $J = 6.7$ Hz), 129.3, 125.1, 124.1 (d, $J = 2.4$ Hz), 120.4, 114.6, 114.5, 55.5 ppm; GC-MS m/z (relative intensity): 272 ($\text{M}^+ + 1$, 11), 271 (M^+ , 65), 270 (34), 179 (100), 159 (21), 151 (23), 136 (34), 108 (21); HRMS calculated 271.1009 for $\text{C}_{16}\text{H}_{14}\text{FO}_2\text{N}$, observed 271.0996.

(Z)-3-(4-Chlorophenyl)-2-fluoro-N-phenylacrylamide (Z-8d). Similarly, a mixture of 1-bromo-1-fluoro-2-(4-chlorophenyl)ethylene (0.12 g, 0.5 mmol, E/ Z = 88: 12), aniline (0.06 g, 0.66 mmol), $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$ (0.01 g, 0.02 mmol) in tributylamine (5 mL) was reacted with CO at room temperature for 94 hours. ^{19}F NMR analysis of the mixture showed that the Z/E ratio of the crude product was 95: 5. Silica gel column chromatography (ethyl acetate: hexanes = 10: 90, $R_f = 0.36$) followed by recrystallization from hexanes gave white crystals, mp 182- 183 °C, 0.06 g, yield 44 % (conversion 50 % based on the consumed (E)-1-bromo-1-fluoro-2-(4-chlorophenyl)ethylene), Z/E = 100: 0. ^{19}F NMR ($\text{CD}_3\text{C}(\text{O})\text{CD}_3$) δ -126.2 (dd, $^3J_{\text{FH}(\text{trans})} = 35.2$ Hz, $J_{\text{F} \dots \text{HN}} = 30.9$ Hz, 1 F) ppm; ^1H NMR ($\text{CD}_3\text{C}(\text{O})\text{CD}_3$) δ 9.62 (broad, 1 H), 7.82 (dm, $J = 8.3$ Hz, 2 H), 7.73 (dm, $J = 7.1$ Hz, 2 H), 7.49 (dm, $J = 8.5$ Hz, 2 H), 7.36 (tm, $J = 7.2$ Hz, 2 H), 7.14 (tm, $J = 6.9$ Hz, 1 H), 7.02 (dm, $^3J_{\text{HF}(\text{trans})} = 37.6$ Hz, 1 H) ppm; ^{13}C NMR ($\text{CD}_3\text{C}(\text{O})\text{CD}_3$) δ 158.7 (d, $J = 28.5$ Hz), 151.8 (d, $^1J_{\text{CF}} = 279.8$ Hz), 139.0, 135.4 (d, $J = 3.6$ Hz), 132.4 (d, $J = 6.5$ Hz), 131.2 (d, $J = 1.9$ Hz), 129.7 (d, $J = 20.5$ Hz), 125.3, 121.3, 121.2, 113.1 (d, $^2J_{\text{CF}} = 3.7$

Hz) ppm; GC-MS m/z (relative intensity): 278 (4), 277 ($M^+ + 2$, 25), 276 (26), 275 (M^+ , 74), 274 (48), 255 (8), 185 (31), 183 (92), 163 (21), 155 (25), 135 (14), 120 (100), 99 (17), 93 (17); HRMS calculated 275.0513 for $C_{15}H_{11}NOF^{35}Cl$, observed 275.0518.

(Z)-2-Fluoro-3-(4-fluorophenyl)-N-phenylacrylamide (Z-8e). Similarly, a mixture of 1-bromo-1-fluoro-2-(4-fluorophenyl)ethylene (0.11 g, 0.5 mmol, E/Z = 87: 13), aniline (0.06 g, 0.65 mmol), $Cl_2Pd(PPh_3)_2$ (0.01 g, 0.02 mmol) in tributylamine (4 mL) was reacted with CO at room temperature for 140 hours. ^{19}F NMR analysis of the mixture showed that the Z/E ratio of the crude product was 95: 5. Silica gel column chromatography (ethyl acetate: hexanes = 10: 90, R_f = 0.25) followed by recrystallization from hexanes gave white crystals, mp 174- 175 °C, 0.10 g, yield 77 % (conversion 89 % based on the consumed (E)-1-bromo-1-fluoro-2-(4-fluorophenyl)ethylene), Z/E = 100: 0. ^{19}F NMR ($CDCl_3$) δ -110.5 (m, 1 F), -131.0 (dd, $^3J_{FH(trans)} = 38.8$ Hz, $J = 3.7$ Hz, 1 F) ppm; 1H NMR ($CDCl_3$) δ 8.02 (broad, 1 H), 7.62- 7.67 (m, 4 H), 7.38 (tm, $J = 7.9$ Hz, 2 H), 7.18 (tm, $J = 7.3$ Hz, 1 H), 7.11 (tm, $J = 8.7$ Hz, 2 H), 7.04 (d, $J = 40.4$ Hz, 1 H) ppm; ^{13}C NMR ($CDCl_3$) δ 163.1 (dd, $^1J_{CF} = 251.1$ Hz, $J = 3.5$ Hz), 158.1 (d, $J = 27.3$ Hz), 149.5 (d, $^1J_{CF} = 276.0$ Hz), 136.8, 132.2 (t, $J = 8.3$ Hz), 129.2, 127.4 (t, $J = 3.2$ Hz), 125.1, 120.2, 116.0 (d, $^2J_{CF} = 21.9$ Hz), 113.5 (d, $J = 4.2$ Hz) ppm; GC-MS m/z (relative intensity) 260 ($M^+ + 1$, 19), 259 (M^+ , 90); 258 (61), 239 (11), 238 (7), 168 (12), 167 (100), 164 (9), 147 (33), 139 (66), 120 (11), 119 (81), 109 (8), 99 (24), 93 (14); HRMS calculated 259.0809 for $C_{15}H_{11}ONF_2$, observed 259.0796.

(Z)-2-Fluoro-3-(3-nitrophenyl)-N-phenylacrylamide (Z-8f). Similarly, a mixture of 1-bromo-1-fluoro-2-(3-nitrophenyl)ethylene (0.25 g, 1.0 mmol, E/ Z = 81: 19), aniline (0.11 g, 1.2 mmol), $Cl_2Pd(PPh_3)_2$ (0.02 g, 0.03 mmol) in tributylamine (5 mL) was

reacted with CO at room temperature for 110 hours. ^{19}F NMR analysis of the mixture showed that the Z/E ratio of the crude product was 98: 2. Silica gel column chromatography (ethyl acetate: hexanes = 25: 75, R_f = 0.33) followed by recrystallization from hexanes/ ether gave white crystals, mp: 180-182 °C, 0.13 g, yield 46 % (conversion 57 % based on the consumed (E)-1-bromo-1-fluoro-2-(3-nitrophenyl)ethylene), Z/E = 100: 0. ^{19}F NMR (CDCl_3) δ -126.1 (dd, $^3J_{\text{FH}(\text{trans})}$ = 37.5 Hz, $J_{\text{F}\dots\text{HN}}$ = 4.7 Hz, 1 F) ppm; ^1H NMR (CDCl_3) δ 8.56 (m, 1 H), 8.23 (dm, J = 8.3 Hz, 1 H), 8.07 (broad, 1 H), 7.92 (d, J = 7.7 Hz, 1 H), 7.59- 7.66(m, 3 H), 7.40 (tm, J = 7.9 Hz, 2 H), 7.21 (tm, J = 7.5 Hz, 1 H), 7.14 (d, $^3J_{\text{HF}(\text{trans})}$ = 37.8 Hz, 1 H) ppm; ^{13}C NMR (CDCl_3) δ 157.3 (d, J = 27.0 Hz), 151.3 (d, $^1J_{\text{CF}}$ = 281.1 Hz), 148.5, 136.5, 135.7 (d, J = 6.4 Hz), 132.7 (d, J = 2.1 Hz), 129.8, 129.2, 125.4, 124.5 (d, J = 8.7 Hz), 123.9, 120.3, 112.2 (d, $^2J_{\text{CF}}$ = 3.0 Hz) ppm; GC-MS m/z (relative intensity): 287 ($\text{M}^+ + 1$, 10), 286 (M^+ , 100), 269 (10), 239 (10), 238 (6), 194 (31), 164 (18), 148 (18), 147 (22), 136 (17), 120 (61), 94 (16), 93 (19), 77 (17), 65 (19); HRMS calculated 286.0754 for $\text{C}_{15}\text{H}_{11}\text{N}_2\text{O}_2\text{F}$, observed 286.0749.

(Z)-2-Fluoro-4-phenylpent-2-enanilide (Z-8g). Similarly, a mixture of 1-bromo-1-fluoro-3-methyl-3-phenylpropene (0.69 g, 3.0 mmol, E/Z = 83:17), aniline (0.28 g, 3.0 mmol), $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$ (0.07 g, 0.10 mmol) in tributylamine (5 mL) was reacted with CO at 50 °C for 117 hours. ^{19}F NMR analysis of the mixture showed that the Z/E ratio of the crude product was 97: 3. Silica gel column chromatography (ethyl acetate: hexanes = 10: 90, R_f = 0.25) followed by recrystallization from hexanes gave white crystals, mp 95- 96 °C, 0.44 g, 55 % yield (64 % conversion based on the consumed (E)-1-bromo-1-fluoro-3-methyl-3-phenylpropene), Z/E = 100: 0. ^{19}F NMR (CDCl_3) δ -133.6 (dd, $^3J_{\text{FH}(\text{trans})}$ = 35.5 Hz, $J_{\text{F}\dots\text{HN}}$ = 5.1 Hz, 1 F) ppm; ^1H NMR (CDCl_3) δ 7.87 (broad s, 1 H), 7.57 (dm, J = 8.5

Hz, 2 H), 7.23-7.38 (m, 7 H), 7.15 (tt, $J = 7.5$ Hz, $J = 1.1$ Hz, 1 H), 6.41 (dd, $^3J_{\text{FH(trans)}} = 36.4$ Hz, $J = 10.0$ Hz, 1 H), 4.03 (dq, $J = 10.3$ Hz, $J = 7.0$ Hz, 1 H), 1.47 (d, $J = 7.0$ Hz, 3 H) ppm; ^{13}C NMR (CDCl_3) δ 158.0 (d, $^2J_{\text{CF}} = 30.2$ Hz), 149.4 (d, $^1J_{\text{CF}} = 266.3$ Hz), 143.8, 136.9, 129.1, 128.8, 127.0, 126.9, 125.1, 121.5 (d, $^2J_{\text{CF}} = 11.1$ Hz), 120.4, 35.2, 21.5 ppm; GC-MS m/z (relative intensity): 270 ($\text{M}^+ + 1$, 19), 269 (M^+ , 100), 254 (5), 249 (10), 234 (11), 216 (14), 206 (17), 166 (18), 157 (47), 156 (77), 135 (20), 133 (44), 130 (84), 129 (90), 128 (42), 120 (25), 115 (20), 105 (23), 93 (35), 91 (19), 77 (55); HRMS calculated 269.1216 for $\text{C}_{17}\text{H}_{16}\text{FON}$, observed 269.1201.

General procedures for the preparation of (E)- α -fluoro- α,β -unsaturated amides

E-8: An oven-dried 120 mL Hastelloy Parr pressure reactor equipped with a stirring bar was charged with 0.03 g (0.04 mmol) of $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$, 0.14 g (1.5 mmol) of freshly distilled aniline, and 4.0 mL of tri-*n*-butylamine. A mixture of (Z)-1-bromo-1-fluoroalkene (1.0 mmol) and the reduced products was added and the pressure reactor was closed (*caution: All reactions should be carried out behind a safety shield in a well-ventilated hood*). The pressure reactor was pressurized to 160 psi with carbon monoxide, and the pressure was released. This process was repeated four times to rid the system of air. Finally the pressure reactor was pressurized to 160 psi and was allowed to stir in an oil bath at 70 °C. The consumption of carbon monoxide could be roughly determined by the reading on the gauge. When the reaction was completed, the reactor was allowed to cool to room temperature and the pressure was carefully released. The reaction mixture was transferred to a separatory funnel containing 100 mL of ethyl acetate. The organic layer was washed with aqueous 10 % hydrochloric acid (3 X 15 mL), saturated sodium

bicarbonate (2 X 15 mL) and brine (2 X 10 mL). The combined organic layers were dried over anhydrous MgSO_4 , concentrated by rotary evaporation, and the crude product was further purified by silica gel column chromatography. Recrystallization from hexanes gave 100 % pure (Z)- α -fluoro- α,β -unsaturated amides.

(E)-3-(2-Chlorophenyl)-2-fluoro-N-phenylacrylamide (E-8b). Similarly, a mixture of (Z)-1-bromo-1-fluoro-2-(2-chlorophenyl)ethylene (contains (Z)-1-bromo-1-fluoro-2-(2-chlorophenyl)ethylene 0.09 g, 0.4 mmol, E/Z = 0: 100) and the reduced products, aniline (0.06 g, 0.6 mmol), $\text{Cl}_2\text{Pd}(\text{PPh}_3)_2$ (0.01 g, 0.02 mmol) in tributylamine (4 mL) was reacted with CO at 70 °C for 140 hours. Silica gel column chromatography (ethyl acetate: hexanes = 10: 90, R_f = 0.31) followed by recrystallization from hexanes gave white crystals, 0.03 g, 27 % yield, Z/E = 0: 100. ^{19}F NMR (CDCl_3) δ -117.5 (dd, $^3J_{\text{FH(cis)}}$ = 24.1 Hz, $J_{\text{F...HN}}$ = 5.3 Hz) ppm; ^1H NMR (CDCl_3) δ 7.93 (broad s, 1 H), 7.67- 7.70 (m, 1 H), 7.52 (dm, J = 7.7 Hz, 2 H), 7.39- 7.42 (m, 1 H), 7.26- 7.35 (m, 4 H), 7.14 (tm, J = 6.4 Hz, 1 H), 6.96 (d, $^3J_{\text{HF(cis)}}$ = 23.4 Hz, 1 H) ppm; ^{13}C NMR (CDCl_3) δ 157.0 (d, J = 30.6 Hz), 149.6 (d, $^1J_{\text{CF}}$ = 263.4 Hz), 136.5, 133.6 (d, J = 3.1 Hz), 131.9, 130.0, 129.7 (d, J = 10.5 Hz), 129.1, 126.2, 125.2, 120.3, 116.8 (d, $^2J_{\text{CF}}$ = 28.5 Hz) ppm. GC-MS m/z (relative intensity): 277 ($\text{M}^+ + 2$, 0.13), 275 (M^+ , 0.35), 241 (37), 240 (100), 220 (3), 185 (3), 183 (10), 165 (4), 163 (6), 155 (10), 137 (4), 135 (11), 121 (6), 120 (51), 119 (8), 109 (6), 99 (11), 93 (7), 77 (11), 65 (12); HRMS calculated 275.0513 for $\text{C}_{15}\text{H}_{11}\text{ONF}^{35}\text{Cl}$, observed 275.0503.

(E)-2-Fluoro-4-phenylpent-2-enanilide (E-8g). Similarly, a mixture of (Z)-1-bromo-1-fluoro-4-phenyl-1-butene (contains (Z)-1-bromo-1-fluoro-4-phenyl-1-butene 0.23 g, 1.0 mmol, E/Z = 0: 100) and the reduced products, aniline (0.14 g, 1.5 mmol),

Cl₂Pd(PPh₃)₂ (0.03 g, 0.04 mmol) in tributylamine (4 mL) was reacted with CO at 70 °C for 109 hours. Silica gel column chromatography (ethyl acetate: hexanes = 7: 93, *R_f* = 0.34) followed by recrystallization from hexanes gave white crystals, 0.20 g, 74 % yield, Z/E = 0: 100. ¹⁹F NMR (CDCl₃) δ -124.7 (d, ³J_{FH(cis)} = 24.5 Hz, 1 F) ppm; ¹H NMR (CDCl₃) δ 7.95 (broad d, J = 3.3 Hz, 1 H), 7.55 (dm, J = 7.6 Hz, 2 H), 7.26- 7.34 (m, 6 H), 7.16- 7.22 (m, 1 H), 7.12 (tm, J = 7.5 Hz, 1 H), 5.95 (dd, ³J_{HF(cis)} = 24.7 Hz, J = 10.9 Hz, 1 H), 4.98 (dq, J = 10.8 Hz, J = 7.0 Hz, J = 3.0 Hz, 1 H), 1.43 (dd, J = 7.0 Hz, J = 0.9 Hz, 3 H) ppm; ¹³C NMR (CDCl₃) δ 158.3 (d, J = 31.0 Hz), 147.4 (d, ¹J_{CF} = 258.7 Hz), 144.5 (d, J = 2.8 Hz), 136.6, 128.9, 128.5, 126.8, 126.5, 126.1 (d, ²J_{CF} = 15.9 Hz), 124.9, 120.2, 34.7 (d, J = 6.2 Hz), 21.4 ppm. GC-MS *m/z* (relative intensity): 270 (M⁺ + 1, 8), 269 (M⁺, 54), 249 (48), 157 (45), 156 (100), 133 (46), 130 (77), 129 (84), 128 (45), 127 (20), 115 (20), 93 (37), 77 (47); HRMS calculated 269.1216 for C₁₇H₁₆FON, observed 269.1206.

(E)-2-Fluoro-3-(1-naphthyl)-N-phenylacrylamide (E-8h). Similarly, a mixture of (Z)-1-bromo-1-fluoro-2-(1-naphthyl)ethylene (contains (Z)-1-bromo-1-fluoro-2-(1-naphthyl)ethylene 0.25 g, 1.0 mmol, E/Z = 0: 100) and the reduced products, aniline (0.14 g, 1.5 mmol), Cl₂Pd(PPh₃)₂ (0.03 g, 0.04 mmol) in tributylamine (4 mL) was reacted with CO at 70 °C for 175 hours. Silica gel column chromatography (ethyl acetate: hexanes = 15: 85, *R_f* = 0.36) followed by recrystallization from hexanes gave white crystals, mp 122-123 °C, 0.12 g, 42 % yield, Z/E = 0: 100. ¹⁹F NMR (CDCl₃) δ -116.6 (broad d, ³J_{FH(cis)} = 21.4 Hz, 1 F) ppm; ¹H NMR (CDCl₃) δ 7.87- 7.95 (m, 3 H), 7.78 (broad s, 1 H), 7.66 (d, J = 7.1 Hz, 1 H), 7.46- 7.56 (m, 3 H), 7.36 (s, 2 H), 7.30 (d, ³J_{HF(cis)} = 20.5 Hz, 1 H), 7.22- 7.28 (m, 2 H), 7.08 (t, J = 7.3 Hz, 1 H) ppm; ¹³C NMR

(CDCl₃) δ 157.1 (d, J = 31.4 Hz), 150.1 (d, J = 261.2 Hz), 136.6, 133.4, 131.3 (d, J = 2.6 Hz), 129.3, 129.0, 128.7, 127.9, 126.7, 126.2, 125.3, 124.9, 124.3, 120.0, 116.8 (d, $^2J_{\text{CF}}$ = 24.9 Hz) ppm; GC-MS m/z (relative intensity): 292 ($M^+ + 1$, 15), 291 (M^+ , 73), 290 (66), 271 (11), 270 (19), 199 (64), 179 (78), 171 (28), 170 (97), 151 (100), 93 (38); HRMS calculated 291.1059 for C₁₉H₁₄NOF, observed 291.1066.

^1H and ^{13}C NMR spectra of Z-6, E-6, Z-8 and E-8:

Z-6a:

Z-6b:

Z-6c:

Z-6d:

Z-6e:

Z-6f:

Z-6g:

E-6a:

E-6b:

E-6c:

E-6g:

E-6h:

Z-8a:

Z-8b:

Z-8c:

Z-8d:

Z-8e:

Z-8f.

Z-8g:

E-8a:

E-8b:

E-8g:

E-8h:

NMR experiment on the existence of intramolecular F...HN coupling in (Z)- and (E)- α -fluoro- α,β -unsaturated amides. (Z)-2-Fluoro-3,*N*-diphenylacrylamide **Z-8a** (0.01 g), which was chosen as a representative of (Z)- and (E)- α -fluoro- α,β -unsaturated amides, was dissolved in 0.4 mL of CDCl₃ in a NMR tube. ¹⁹F NMR spectrum of the NMR solution showed a typical dd (double (³J_{FH(trans)} = 39.2 Hz) of doublet (J_{F...HN} ~ 3 Hz)) pattern at -130.0 ppm; ¹⁹F {¹H} spectrum showed a singlet peak at -130.0 ppm; One drop of D₂O was added to the NMR solution and the solution was shaken several times. ¹⁹F NMR of the resulted solution showed a d (double (³J_{FH(trans)} = 39.5 Hz), sharp peaks) pattern at -130.0 ppm.