Improved Asymmetric S_NAr Reaction of β -Dicarbonyl Compounds Catalyzed by Quaternary Ammonium Salts Derived from Cinchona Alkaloids

*Sara Kobbelgaard, Marco Bella and Karl Anker Jørgensen**

Danish National Research Foundation: Center for Catalysis, Department of Chemistry,
Aarhus University, DK-8000 Aarhus C, Denmark.

kaj@chem.au.dk

Supporting Information

Contents

General Methods	S3
Materials	S3
General Procedure for the Enantioselective S _N Ar Reaction of Aromatic Fluorides 2a-d with Nucleophile 1a-e Catalyzed by 4a-t	S3
Characterization Data for the Products 3a-e , 5e	S4-5
General Procedure for Opening of α,α -Disubstituted Aryl- β -Ketoesters with Various Nucleophiles to 7a,b,d	S6
Characterization Data for the Products 7a,b,d	S6-7
General Procedure for Diastereoselective Reduction of Optically Active α,α -Disubstituted Aryl- β -Ketoester (3a) to β -Hydroxyester (9a)	S7
Characterization Data for the Products 9a	S7-8

Preparation of Catalyst 4	S8
Characterization Data for the Products 4d,i,k,l,r	S8-10
Preparation of Catalyst 4s	S10
Characterization Data for the Product 4s	S11
References	S11
¹ H and ¹³ C NMR spectra of 3a-h, 5e, 7a,b,d, 9a, 4d,i,k,l,r,s	S12-S30

General methods. The ^1H NMR and ^{13}C NMR spectra were recorded at 400 Mhz and 100 Mhz respectively. The chemical shifts are reported in ppm downfield to TMS ($\delta = 0$) for ^1H NMR and for ^{13}C NMR relative to the central CDCl_3 resonance ($\delta = 77.0$). Flash chromatography (FC) was carried out using Merck silica gel 60 (230-400 mesh). Optical rotations were measured on a Perkin-Elmer 241 polarimeter. The enantiomeric excess (ee) of the products was determined by HPLC using a Daicel Chiralpak AS and Chiralcel OD columns (wavelength = 254 nm) with *i*-PrOH/hexane as the eluent.

Materials. Analytical grade solvents were used as received. All commercially available reagents were used as received. Compounds **1a**, **1c** and **2a-d** are commercially available. Those compounds were prepared according to standard literature procedures: **1b** (deprotonation of corresponding *N*-Boc 2-pyrrolidones and reaction with ethyl chloroformate in THF), **1c**,¹ **1d**,² **1f**,³ **3k**.⁴

General Procedure for the Enantioselective $\text{S}_{\text{N}}\text{Ar}$ Reaction of Aromatic Fluorides **2a-d with Nucleophile **1a-e** Catalyzed by **4a-t**.** In a test tube the catalyst **4** (0.03 mmol, 15 mol%) was dissolved in CH_2Cl_2 (0.8 mL) nucleophile **1a-f** (0.20 mmol, 1.00 eq.), aromatic fluoride **2a-f** (0.25 mmol, 1.25 eq.) and PhMe (3.2 mL) was added. Another test tube contained KOH (150 mg). The test tubes were cooled down to the indicated temperature, and KOH was added to the reaction mixture. The test tube was closed by a rubber septa, no inert atmosphere was used. The reaction mixture was stirred at the indicated temperature until no starting nucleophile could be detected by TLC. The crude mixture was directly purified by FC to afford pure products **3a-j**. The enantiomeric excess was determined by HPLC.

3a: Yield: 89%; ^1H NMR δ (CDCl_3) 8.78 (d, 1H, J 2.4 Hz), 8.34 (dd, 1H, J 8.8 Hz, J 2.0 Hz), 7.41 (d, 1H, J 8.8 Hz), 4.12 (q, 1H, J 7.2 Hz), 4.11 (q, 1H, J 7.2 Hz), 3.2-3.1 (m, 1H), 2.8-2.6 (m, 1H), 2.6-2.2 (m, 3H), 2.1-1.9 (m, 1H), 1.14 (t, 3H, J 7.2 Hz). ^{13}C NMR δ (CDCl_3) 211.3, 168.1, 149.4, 147.2, 140.5, 131.5, 127.4, 121.5, 65.4, 63.0, 38.9, 37.0, 19.6, 14.1. HRMS calc. $\text{C}_{14}\text{H}_{14}\text{N}_2\text{NaO}_7$: 345.0699; found: 345.0699. $[\alpha]_{\text{D}}^{\text{rt}} = -40$ (c 4.05, CDCl_3 , 56% ee (optical rotation performed from another reaction than reported in Table 1). The ee was determined by HPLC using Chiralpak AS column (hexane/*i*-PrOH 80:20); flow rate 1.0 mL/min; $\tau_{\text{major}} = 12.5$ min; $\tau_{\text{minor}} = 15.4$ min.

3b: Yield: 96%; ^1H NMR δ (CDCl_3) 8.50 (s, 1H), 7.60 (s, 1H), 4.14 (q, 2H, J 7.2 Hz), 3.2-3.1 (m, 1H), 2.8-2.6 (m, 1H), 2.6-2.2 (m, 3H), 2.1-1.9 (m, 1H), 1.16 (t, 3H, J 7.2 Hz). ^{13}C NMR δ (CDCl_3) 210.8, 168.2, 148.5, 148.2, 139.1, 137.2, 123.7, 120.6, 65.1, 63.3, 39.0, 37.0, 19.7, 14.3. HRMS calc. $\text{C}_{14}\text{H}_{13}\text{BrN}_2\text{NaO}_7$: 422.9804; found: 422.9803. $[\alpha]_{\text{D}}^{\text{rt}} = -80$ (c 7.67, CDCl_3 , 92% ee). The ee was determined by HPLC using Chiralpak AS column (hexane/*i*-PrOH 90:10); flow rate 1.0 mL/min; $\tau_{\text{major}} = 16.1$ min; $\tau_{\text{minor}} = 18.4$ min.

3c: Yield 16%; ^1H NMR δ (CDCl_3) 8.91 (d, 1H, J 2.4 Hz), 8.57 (dd, 1H, J 8.6 Hz, J 2.3 Hz), 8.21 (d, 1H, J 8.6 Hz), 2.9-2.6 (m, 4H), 2.5-2.4 (m, 2H). ^{13}C NMR (CDCl_3) 203.4, 148.1, 147.4, 135.6, 134.2, 128.1, 122.1, 116.7, 56.5, 37.9, 36.3, 20.2. $[\alpha]_{\text{D}}^{\text{rt}} = +138$ (c 0.61, CH_2Cl_2 , 60% ee). The ee was determined by HPLC using Chiralcel OD column (hexane/*i*-PrOH 80:20); flow rate 1.0 mL/min; $\tau_{\text{major}} = 33.1$ min; $\tau_{\text{minor}} = 38.4$ min.

3d: Yield 36%; ^1H NMR δ (CDCl_3) 8.75 (d, 1H, J 2.8 Hz), 8.40 (dd, 1H, J 9.3 Hz, J 2.7 Hz), 7.78 (d, 1H, J 8.3 Hz), 7.56 (d, 1H, J 9.3 Hz), 7.53 (bs, 1H), 7.49 (d, 1H, J 8.3 Hz), 3.95 (d, 1H, J 17.5 Hz), 3.80 (s, 3H), 3.61 (d, 1H, J 17.6 Hz). ^{13}C NMR (CDCl_3) 198.3, 167.5, 154.6, 149.4, 147.1, 144.0, 141.1, 132.9, 131.3, 129.6, 127.7, 126.9, 126.6, 121.3, 62.5, 53.8, 41.7. HRMS calc. $\text{C}_{17}\text{H}_{11}\text{ClN}_2\text{NaO}_7$: 413.0152; found: 413.0157. $[\alpha]_{\text{D}}^{25} = +53$ (c 1.60, CH_2Cl_2 , 30% ee). The ee was determined by HPLC using Chiralcel OD column (hexane/*i*-PrOH 80:20); flow rate 1.0 mL/min; $\tau_{\text{major}} = 28.8$ min; $\tau_{\text{minor}} = 47.8$ min.

3e: Yield 20%; ^1H NMR δ (CDCl_3) 8.85 (d, 1H, J 2.4 Hz), 8.43 (dd, 1H, J 8.6 Hz, J 2.4 Hz), 7.60 (d, 1H, J 8.7 Hz), 4.2-4.1 (m, 2H), 3.11 (dt, 1H, J 13.9, J 6.5 Hz), 2.83 (dd, 1H, J 13.0 Hz, J 2.8 Hz), 2.6-2.5 (m, 1H), 2.4-2.3 (m, 1H), 2.2-2.1 (m, 1H), 2.0-1.9 (m, 2H), 1.9-1.8 (m, 1H), 1.19 (tri, 3H, J 7.1 Hz). ^{13}C NMR (CDCl_3) 203.2, 168.6, 149.5, 146.9, 140.1, 130.4, 127.0, 120.8, 65.7, 62.6, 40.6, 37.8, 26.1, 22.0, 13.8. HRMS calc. $\text{C}_{15}\text{H}_{16}\text{N}_2\text{NaO}_7$: 359.0850; found: 359.0850. $[\alpha]_{\text{D}}^{25} = -41$ (c 1.42, CH_2Cl_2 , 38% ee). The ee was determined by HPLC using Chiralpak AS column (hexane/*i*-PrOH 90:10); flow rate 1.0 mL/min; $\tau_{\text{major}} = 17.5$ min; $\tau_{\text{minor}} = 19.7$ min.

5e: Yield 16%; ^1H NMR δ (CDCl_3) 8.69 (d, 1H, J 2.7 Hz), 8.22 (dd, 1H, J 2.8 Hz, J 9.2 Hz), 6.97 (d, 1H, J 9.2 Hz), 3.88 (q, 2H, J 7.1 Hz), 2.4-2.3 (m, 2H), 2.2-2.1 (m, 2H), 1.7-1.6 (m, 2H), 1.6-1.5 (m, 2H), 0.87 (t, 3H, J 7.1 Hz). ^{13}C NMR (CDCl_3) 165.1, 155.1, 155.0, 128.8, 122.2, 120.7, 116.6, 60.8, 29.7, 27.8, 25.4, 22.4, 21.6, 13.8. HRMS calc. $\text{C}_{14}\text{H}_{16}\text{N}_2\text{NaO}_7$: 359.0855; found: 359.0849.

General Procedure for Opening of α,α -Disubstituted Aryl- β -Ketoesters with Various Nucleophiles to **7a,b,d.** The β -ketoester (50 mg (**3a**), 0.155 mmol, 1.0 eq.), nucleophile (0.165 mmol, 1.1 eq.) and solvent (2 mL) are combined in a test tube with a magnetic stir bar and maintained at the indicated temperature, until no β -ketoester could be detected by TLC. The crude mixture was directly purified by FC to afford pure products **7a,b,d**.

7a: Yield 84%; ^1H NMR δ (CDCl_3) 8.72 (d, 1H, J 2.1 Hz), 8.41 (dd, 1H, J 8.6 Hz, J 2.0 Hz), 7.78 (d, 1H, J 8.6 Hz), 7.21 (bs, 1H), 4.23 (t, 1H, J 7.3 Hz), 4.2-4.1 (m, 2H), 3.64 (bs, 2H), 2.3-2.1 (m, 3H), 2.0-1.9 (m, 1H), 1.8-1.5 (m, 2H), 1.71 (t, 3H, J 7.1 Hz). ^{13}C NMR (CDCl_3) 172.8, 171.1, 149.3, 146.7, 139.9, 131.5, 127.1, 120.2, 61.8, 46.3, 33.4, 32.0, 23.1, 13.9. HRMS calc. $\text{C}_{14}\text{H}_{18}\text{N}_4\text{NaO}_7$: 377.1073; found: 377.1068.

7b: Yield 49%; ^1H NMR δ (CDCl_3) 8.73 (d, 1H, J 2.4 Hz), 8.43 (dd, 1H, J 8.7 Hz, J 2.4 Hz), 7.81 (d, 1H, J 8.7 Hz), 4.25 (tri, 1H, J 7.3 Hz), 4.2-4.1 (m, 4H), 2.34 (dt, 2H, J 7.2 Hz, J 3.0 Hz), 2.3-2.2 (m, 1H), 2.0-1.9 (m, 1H), 1.7-1.5 (m, 2H), 1.24 (t, 3H, J 7.14 Hz), 1.20 (t, 3H, J 7.12 Hz). ^{13}C NMR (CDCl_3) 172.8, 171.1, 149.5, 146.8, 140.0, 131.5, 127.0, 120.2, 61.8, 60.5, 46.3, 33.5, 32.0, 22.8, 14.2, 14.0. HRMS calc. $\text{C}_{16}\text{H}_{20}\text{N}_2\text{NaO}_8$: 391.1117; found: 391.1116.

7d: Conversion 65% determined by ^1H NMR; ^1H NMR δ (CDCl_3) 8.73 (d, 1H, J 2.3 Hz), 8.41 (dd, 1H, J 8.6 Hz, J 1.7 Hz), 7.78 (d, 1H, J 8.7 Hz), 7.4-7.2 (m, 5H), 5.74 (d, 1H, J 7.0 Hz), 5.11 (quin, 1H, J 6.9 Hz), 4.24 (t, 1H, J 7.5 Hz), 4.2-4.1 (m, 2H), 2.3-2.1 (m, 3H), 2.0-1.8 (m, 1H), 1.8-1.6 (m, 2H), 1.48 (d, 3H, J 6.9 Hz), 1.19 (t, 3H, J 7.1 Hz).

^1H NMR δ (CDCl_3) 8.74 (d, 1H, J 2.3 Hz), 8.40 (dd, 1H, J 8.7 Hz, J 1.9 Hz), 7.75 (d, 1H, J 8.7 Hz), 7.4-7.2 (m, 5H), 5.72 (d, 1H, J 7.0 Hz), 5.11 (quin, 1H, J 6.9 Hz), 4.24 (t, 1H, J 7.5 Hz), 4.2-4.1 (m, 2H), 2.3-2.1 (m, 3H), 2.0-1.8 (m, 1H), 1.8-1.6 (m, 2H), 1.48 (d, 3H, J 6.9 Hz), 1.18 (t, 3H, J 7.2 Hz) ^{13}C NMR (CDCl_3) 171.2, 170.8, 146.7, 144.1, 143.1, 140.0, 131.6, 128.7 (2C), 127.4, 127.1, 126.2, 126.1, 120.2, 61.8, 48.8, 46.3, 35.7, 32.0, 23.3, 21.7, 14.0. HRMS calc. $\text{C}_{22}\text{H}_{25}\text{N}_3\text{NaO}_7$: 486.1590; found: 486.1588. $[\alpha]_D^{rt} = -29$ (c 1.38, CH_2Cl_2 , 38% ee).

General Procedure for Diastereoselective Reduction of Optically Active α,α -Disubstituted Aryl- β -Ketoester (3a) to β -Hydroxyester (9a). To a solution of β -ketoester (**3a**) (113 mg, 0.35 mmol, 1.00 eq.) in THF (5 mL) was added 3 eq. of BH_3 -DMS complex (0.105 mL, 1.05 mmol, 3.00 eq.). After being stirred at room temperature for 7 d, to the reaction mixture was added dropwise 1 mL of 1M HCl and the solution extracted with Et_2O . The organic layer was dried over MgSO_4 , filtered, and evaporated. The crude reaction mixture was purified by FC (Et_2O /pentane 2:1) to give the optically active β -hydroxyester **9a**. The enantiomeric excess was determined by HPLC.

9a: Yield 32%; ^1H NMR δ (CDCl_3) 8.70 (d, 1H, J 2.1 Hz), 8.44 (dd, 1H, J 8.8 Hz, J 1.5 Hz), 8.01 (d, 1H, J 8.8 Hz), 4.55 (t, 1H, J 6.9 Hz), 4.20 (dq, 2H, J 7.1 Hz, J 2.9 Hz), 3.12 (bs, 1H), 2.79 (ddd, 1H, J 13.6 Hz, J 9.9 Hz, J 6.5 Hz), 2.1-2.0 (m, 1H), 2.0-1.8 (m, 3H), 1.7-1.6 (m, 1H), 1.19 (t, 3H, J 7.1 Hz). ^{13}C NMR (CDCl_3) 172.6, 149.1, 146.6, 143.9, 131.0, 126.9, 120.6, 79.7, 62.0, 59.7, 33.5, 32.3, 19.7, 13.9. HRMS calc. $\text{C}_{14}\text{H}_{16}\text{N}_2\text{NaO}_7$: 347.0855; found: 347.0858. $[\alpha]_D^{rt} = +57$ (c 2.47, CH_2Cl_2 , 31% ee). The ee was determined by HPLC using Chiralpak AS column (hexane/*i*-PrOH 80:20); flow rate 1.0 mL/min; $\tau_{\text{major}} = 6.9$ min; $\tau_{\text{minor}} = 8.0$ min.

Stereochemistry at C2 was determined by NOE difference between H2 and the aromatic protons.

Preparation of Catalyst 4:

To a solution in CH_2Cl_2 of commercially available *N*-benzyl cinchonidium chloride (420 mg, 1 mmole) were added benzoyl chloride (1.5 mmol, 1.5 eq.), and Et_3N (200 mg, 2 mmol). The reaction was stirred at rt for 24 h, additional benzoyl chloride (70 mg, 0.5 mmol, 0.5 eq.) was added and the reaction stirred for additional 12 h. Water (15 mL) was added, the mixture was debated, the organic layer dried on MgSO_4 , concentrated *in vacuo* and the product purified by FC (EtOAc, then EtOAc/MeOH 10:1).

4d: Yield: 74%; ^1H NMR δ (CDCl_3) 9.03 (d, 1H, J 8.3 Hz), 8.67 (d, 1H, J 4.5 Hz), 8.22 (d, 2H, J 7.3 Hz), 8.15 (d, 1H, J 8.3 Hz), 8.05 (t, 1H, J 7.7), 7.9-7.8 (m, 3H), 7.78 (t, 1H, J 7.5 Hz), 7.63 (t, 2H, J 7.8 Hz), 7.51 (d, 1H, J 4.6 Hz), 7.5-7.4 (m, 3H), 6.55 (d, 1H, J 11.7 Hz), 5.84 (ddd, 1H, J 17.2 Hz, J 10.4 Hz, J 6.3 Hz), 5.84 (t, 1H, J 11.0 Hz), 5.42 (d, 1H, J 10.5 Hz), 5.21 (d, 1H, J 17.2 Hz), 5.15 (t, 1H, J 10.5 Hz), 4.32 (d, 1H, J 11.7 Hz), 3.8-3.7 (m, 2H), 3.1-3.0 (m, 1H), 3.0-2.9 (m, 1H), 2.7-2.5 (m, 2H), 2.2-2.0 (m, 2H) 1.9-1.7 (m, 1H), 1.5-1.4 (m, 1H). ^{13}C NMR δ (CDCl_3) 164.1, 149.2, 148.1, 140.0, 135.8, 134.9, 133.9 (2C), 130.6, 130.5, 129.8, 129.6 (2C), 129.4 (2C), 129.1 (3C), 127.9, 126.5, 124.3, 124.0, 118.5, 118.1, 68.4, 65.2, 63.3, 60.0, 51.5, 37.5, 26.8, 25.4, 23.2. HRMS calc. $\text{C}_{33}\text{H}_{33}\text{N}_2\text{O}_2$: 489.2542; found: 489.2541. $[\alpha]_D^{25} = -5$ (c 7.00, CDCl_3).

4i: ^1H NMR δ (CDCl_3) 8.82 (d, 1H, J 4.5 Hz), 8.61 (d, 1H, J 8.4 Hz), 8.10 (d, 2H, J 8.2 Hz), 8.07 (d, 1H, J 8.6 Hz), 7.88 (t, 1H, J 7.3 Hz), 7.8-7.7 (m, 4H), 7.50 (d, 1H, J 4.5 Hz), 7.5-7.3 (m, 5H), 6.38 (d, 1H, J 12.0 Hz), 5.70 (ddd, 1H, J 17.2 Hz, J 10.5 Hz, J 6.6 Hz), 5.33 (d, 1H, J 17.2 Hz), 5.1-5.0 (m, 2H), 5.00 (d, 1H, J 10.5 Hz), 4.64 (d, 1H, J 12.0 Hz), 3.8-3.7 (m, 1H), 3.5-3.4 (m, 1H), 2.75 (q, 2H, J 7.6 Hz), 2.7-2.4 (m, 2H), 2.4-2.2 (m, 2H), 2.2-2.1 (m, 1H), 2.0-1.9 (m, 1H), 1.8-1.7 (m, 1H), 1.27 (t, 3H, J 7.6 Hz). ^{13}C NMR (CDCl_3) 164.1, 152.3, 149.2, 148.1, 140.1, 135.9, 133.9 (2C), 130.6, 130.5, 129.8 (2C), 129.8, 129.1 (2C), 129.1, 128.9 (2C), 126.6, 125.3, 124.3, 124.0, 118.5, 118.1, 68.1, 65.3, 63.4, 60.0, 51.5, 37.5, 29.0, 26.8, 25.4, 23.2, 15.1. $[\alpha]_{\text{D}}^{25} = +2$ (c 1.98, CH_2Cl_2).

4k: ^1H NMR δ (CDCl_3) 8.83 (d, 1H, J 4.4 Hz), 8.69 (d, 1H, J 8.2 Hz), 8.15 (d, 2H, J 8.7 Hz), 8.08 (d, 1H, J 8.3 Hz), 7.89 (d, 1H, J 7.6 Hz), 7.8-7.7 (m, 4H), 7.50 (d, 1H, J 4.3 Hz), 7.5-7.3 (m, 3H), 7.06 (d, 2H, J 8.7 Hz), 6.42 (d, 1H, J 11.8 Hz), 5.70 (ddd, 1H, J 17.1 Hz, J 10.5 Hz, J 6.8 Hz), 5.35 (d, 1H, J 17.1 Hz), 5.2-5.1 (m, 1H), 5.1-5.0 (m, 1H), 5.00 (d, 1H, J 10.6 Hz), 4.66 (d, 1H, J 11.8 Hz), 3.91 (s, 3H), 3.8-3.7 (m, 1H), 3.5-3.4 (m, 2H), 2.7-2.6 (m, 1H), 2.4-2.2 (m, 2H), 2.14 (s, 1H), 2.0-1.9 (m, 1H), 1.8-1.7 (m, 1H). ^{13}C NMR (CDCl_3) 164.8, 163.8, 149.2, 148.1, 140.2, 135.9, 134.0 (2C), 131.9 (2C), 130.6, 130.5, 129.8, 129.2, 129.1, 126.7, 124.4, 124.1, 120.0, 118.6, 118.2, 114.7 (3C), 68.0, 65.2, 63.3, 60.0, 55.7, 51.5, 37.6, 26.9, 25.5, 23.3. $[\alpha]_{\text{D}}^{25} = +6$ (c 4.69, CH_2Cl_2).

4l: ^1H NMR δ (CDCl_3) 8.90 (d, 1H, J 4.5 Hz), 8.72 (d, 1H, J 8.7 Hz), 8.11 (d, 1H, J 8.4 Hz), 8.06 (d, 1H, J 8.1 Hz), 7.90 (t, 1H, J 7.3 Hz), 7.84 (d, 2H, J 6.6 Hz), 7.8-7.7 (m, 2H), 7.63 (d, 1H, J 4.5 Hz), 7.60 (d, 1H, J 3.7 Hz), 7.5-7.3 (m, 4H), 6.46 (d, 1H, J 11.8 Hz), 5.72 (ddd, 1H, J 17.0

Hz, J 10.5 Hz, J 6.1 Hz), 5.37 (d, 1H, J 17.3 Hz), 5.19 (d, 1H, J 12.9 Hz), 5.1-4.9 (m, 2H), 4.75 (s, 1H), 4.64 (d, 1H, J 11.9 Hz), 4.0-3.8 (m, 1H), 3.5-3.4 (m, 1H), 3.36 (t, 1H, J 11.7 Hz), 2.7-2.6 (m, 1H), 2.4-2.1 (m, 3H), 1.9-1.8 (m, 1H), 1.8-1.6 (m, 1H). ^{13}C NMR (CDCl_3) 163.7, 149.2, 148.2, 139.7, 135.9, 134.5, 134.1 (2C), 133.3, 132.6, 131.9, 130.6 (2C), 129.9, 129.2 (3C), 127.8, 127.7, 126.7, 124.4, 124.0, 118.6, 118.5, 70.4, 65.1, 63.2, 60.0, 51.1, 37.7, 26.9, 25.2, 23.4. $[\alpha]_D^{25} = -42$ (c 3.00, CH_2Cl_2).

4r: ^1H NMR δ (CDCl_3) 8.49 (d, 1H, J 3.7 Hz), 8.28 (d, 1H, J 8.7 Hz), 8.00 (d, 1H, J 8.8 Hz), 7.9-7.8 (m, 2H), 7.74 (dd, 2H, J 7.9 Hz, J 1.2 Hz), 7.68 (t, 1H, J 7.8 Hz), 7.61 (d, 2H, J 7.7 Hz), 7.5-7.4 (m, 2H), 7.4-7.3 (m, 1H), 7.3-7.2 (m, 3H), 7.2-7.0 (m, 2H), 6.00 (d, 1H, J 11.7 Hz), 5.77 (ddd, 1H, J 17.1 Hz, J 10.4 Hz, J 7.0 Hz), 5.22 (d, 1H, J 17.3 Hz), 5.03 (d, 1H, J 10.5 Hz), 5.0-4.9 (m, 1H), 4.7-4.5 (m, 1H), 4.2-4.1 (m, 1H), 3.5-3.4 (m, 1H), 3.3-3.2 (m, 1H), 2.6-2.5 (m, 1H), 2.3-2.0 (m, 3H), 2.0-1.9 (m, 1H), 1.9-1.8 (m, 1H). ^{13}C NMR (CDCl_3) 146.2, 136.9, 135.9, 135.0, 134.2 (2C), 130.7, 129.7, 129.4 (2C), 129.2 (2C), 127.8 (4C), 126.6, 126.0 (3C), 122.8, 119.2, 118.6, 74.5, 65.7, 63.2, 60.3, 51.0, 37.8, 29.2, 27.1, 25.0. $[\alpha]_D^{25} = -34$ (c 2.80, CH_2Cl_2).

Preparation of Catalyst 4s:

To a solution in CH_2Cl_2 of commercially available *N*-benzyl cinchonidium chloride (210 mg, 0.5 mmol) were added **2a** (0.63 mmol, 1.25 eq.), and KOH (56 mg, 1 mmol, 2.0 eq.). The reaction was stirred at rt for 3 h. The mixture was diluted with water (15 mL) and extracted with CH_2Cl_2 3 times. The combined organic extracts were dried over MgSO_4 , filtrated and concentrated. The mixturer was the poured onto Et_2O (15 mL). The solution was filtered and the resulting yellow solid was collected as the desired product **4s**.

4s: ^1H NMR δ (CDCl_3) 8.82 (d, 1H, J 4.5 Hz), 8.72 (d, 1H, J 2.6 Hz), 8.7-8.6 (m, 1H), 8.11 (d, 1H, J 8.4 Hz), 8.0-7.8 (m, 1H), 7.75 (dd, 1H, J 9.7 Hz, J 2.9 Hz), 7.68 (t, 1H, J 7.7 Hz), 7.59 (d, 2H, J 6.9 Hz), 7.52 (t, 1H, J 7.6 Hz), 7.5-7.3 (m, 3H), 7.10 (d, 1H, J 8.3 Hz), 6.37 (d, 1H, J 9.7 Hz), 5.85 (d, 1H, J 12.1 Hz), 5.69 (ddd, 1H, J 16.9 Hz, J 10.5 Hz, J 6.5 Hz) 5.5-5.4 (m, 1H), 5.29 (d, 1H, J 17.0 Hz) 5.08 (d, 1H, J 10.3 Hz), 4.7-4.5 (m, 1H), 4.5-4.3 (m, 1H), 4.1-4.0 (m, 1H), 3.6-3.4 (m, 1H), 3.34 (t, 1H, J 11.6 Hz), 2.7-2.5 (m, 2H), 2.4-2.3 (m, 1H), 2.25 (s, 1H), 2.0-1.9 (m, 1H), 1.9-1.8 (m, 1H). ^{13}C NMR (CDCl_3) 170.9, 152.6, 149.6, 148.6, 141.1, 138.8, 137.7, 136.1, 135.4, 133.8, 130.7, 130.2, 129.3 (2C), 128.7, 128.4, 126.7 (2C), 125.8, 124.1, 122.6, 121.9, 118.3, 117.5, 67.6, 64.5, 60.6, 50.6, 37.9, 26.7, 24.8, 22.4. $[\alpha]_D^{25} = +28$ (c 0.47, CH_2Cl_2).

References

1. Dehli, J. R.; Gotor, V. *J. Org. Chem.* **2002**, *67*, 6816.
2. McCann, S. F.; Annis, G. D.; Shapiro, R.; Piotrowski, D.; W. Lahm, G. P.; Long, J. K.; Lee, K. C.; Hughes, M. M.; Myers, B. J.; Griswold, S. M.; Reeves, B. M.; March, R. W.; Sharpe, P. L.; Lowder P.; Barnette, W. E.; Wing, K. D. *Pest. Manag. Sci.* **2001**, *57*, 153.
3. Albarella, J. P. *J. Org. Chem.* **1977**, *42*, 2009.
4. Barton, D. H. R.; Blazejewski, J. C.; Charpiot, B.; Finet, J. P.; Motherwell, W. B.; Papoula, M. T. B.; Stanforth S. P. *J. Chem. Soc. Perkin Trans. I*, **1985**, 2667.

3b

3d

3f

3g

3h

7a

7b

7d

4d

4i

4k

4l