InCl₃-Catalyzed Regio- and Stereoselective Thiolytic of α,β -Epoxycarboxylic Acids in Water

Francesco Fringuelli, Ferdinando Pizzo,* Simone Tortoioli and Luigi Vaccaro*

CEMIN - Dipartimento di Chimica, Laboratorio di Chimica Organica, Università di Perugia, Via Elce di Sotto, 8 I-06123, Perugia, Italia

Fax: (+39) 075 5855560

E-mail: luigi@unipg.it; pizzo@unipg.it

CONTENTS

page S2 Experimental Section

page S3-S17 Full characterization data (¹H NMR, ¹³C NMR, IR, GC-MS, R_f).

Experimental Section

All chemicals were purchased and used without any further purification. In this work we have used de-ionized water. GC analyses were performed with an SPB-5 fused silica capillary column (30 m, 0.25 mm diameter), an “on column” injector system, a FID detector and hydrogen as the carrier gas. GC-MS analyses were carried out with 70 eV electron energy. All ^1H NMR and ^{13}C NMR spectra were recorded at 200 MHz or 400 MHz, and at 50.3 or 100.6 MHz respectively, using a convenient deuterated solvent (reported in the characterization charts) and the residual peak as internal standard, or TMS in the case of CDCl_3 . The pH values of the aqueous reaction medium were controlled using a pH-meter apparatus equipped with a combined refillable pH electrode and with an Autocompensating Temperature Control (ATC) probe. All melting points are uncorrected. Thin Layer Chromatography analyses were performed on silica gel on aluminum plates and UV and/or KMnO_4 or phosphomolybdic acid were used as revealing agents. Column chromatography were performed by using silica gel 230-400 mesh and eluting as reported in the following characterization data. Compounds **3a**, **c-e**, **4a-e**, and **5a-c**, **e** are new compounds and characterization data (^1H NMR, ^{13}C NMR, IR, GC-EIMS, R_f) are listed below.

The relative stereochemistry of all the products was assigned on the basis of the H-2/H-3 J^4 coupling values, and on the H-2/H-3 ^1H NMR and C-2/C-3 ^{13}C NMR ppm values. The assignment was in agreement with the data available in literature for related compounds.¹

¹ (a) Chong, J. M.; Sharpless, K. B. *J. Org. Chem.* **1985**, *50*, 1560-1563. (b) Woydowski, K.; Fleischhauer, J.; Shiffer, J.; Liebsher, J. *J. Chem. Soc., Perkin Trans. 1* **1999**, 149-153. (c) Fringuelli, F.; Pizzo, F.; Vaccaro, L. *J. Org. Chem.* **2001**, *66*, 3354-3358. (d) Amantini, D.; Fringuelli, F.; Pizzo, F.; Vaccaro, L. *J. Org. Chem.* **2001**, *66*, 4463-4467. (e) Tosaki, S.; Tsuji, R.; Ohshima, T.; Shibasaki, M. *J. Am. Chem. Soc.* **2005**, *127*, 2147-2155.

Typical uncatalyzed procedure at pH 9.0

To a 30 °C thermostated mixture of thiol **2a** (1.05 mmol) in water (2 mL) α,β -epoxy acid (1.0 mmol) was added. The pH was adjusted to pH 9.0 by adding NaOH 5 M aq. sol. The pH was kept constant for the entire reaction time by adding a 10% aqueous H_2SO_4 solution. The reaction was monitored by GLC analyses on the corresponding methyl ester derivative, and when the starting material was completely consumed, the resulting mixture was cooled at 0 °C, acidified to pH 1.0 by adding some drops of 10% aqueous H_2SO_4 and extracted by EtOAc (3 x 3 mL). The organic layer was evaporated under reduced pressure to furnish the almost pure the β -hydroxy sulfide products. Mixture of products were separated by silica gel column chromatography (see below).

Typical procedure of the Lewis acid-catalyzed thiolysis at pH 4.0

In a flask thermostated at 30 °C, thiol (1.05 mmol) and 1.0 mL of a 0.1 M aqueous solution of the Lewis acid (0.1 mmol) were added to 0.7 mL of water. By addition of little amounts of NaOH 0.5 M the pH was brought to the desiderated value. The α,β -epoxy acid (1.0 mmol) was then added under stirring and the pH was kept constant at pH 4.0 by adding a 5 M NaOH solution. When the reaction was complete, the resulting mixture was cooled at 0 °C, acidified to pH 1.0 by adding some drops of 10% aqueous H_2SO_4 and extracted with EtOAc (3 x 2 mL). The combined organic layers were dried over Na_2SO_4 and evaporated under reduced pressure to give the corresponding β -hydroxy sulfides that were purified by recrystallization or by column chromatography.

Re-use of the catalyst. After extraction, the mother liquors coming from InCl_3 -catalyzed thiolysis at pH 4.0 can be reused by adding phenylthiol (**1a**) and an α,β -epoxycarboxylic acid and after adjusting the pH to 4.0. No significant decrease in the catalytic efficiency was observed after 5 consecutive runs.

trans-2-hydroxy-1-(phenylthio)cyclohexene-1-carboxylic acid (**3a**)

Characterized by using a mixture with its regioisomer **4a**; ^1H NMR (CDCl_3 , 200 MHz) δ = 1.34-2.17 (m, 8H), 3.88 (dd, J = 3.6, 9.0 Hz, 1H), 7.18-7.55 (m, 5H); ^{13}C NMR (CDCl_3 , 50.3 MHz) δ = 22.2, 22.7, 23.0, 24.8, 59.2, 73.9, 128.9, 129.7, 129.8, 137.5, 174.7; GC-EIMS on methyl ester derivative (266) m/z = 266 (M^+ , 80), 234 (35), 207 (25), 179 (30), 110 (100), 69 (40).

Rel(2S,3S)-3-hydroxy-2-(phenylthio)hexanoic acid (**3c**)

solid; purified by recrystallization from EtOAc; 85% yield; m. p. = 107-109 °C; ^1H NMR (CDCl_3 , 400 MHz) δ = 0.94 (t, J = 7.3 Hz, 3H), 1.36-1.44 (m, 1H), 1.51-1.61 (m, 2H), 1.79-1.84 (m, 1H), 3.65 (d, J = 6.7 Hz, 1H), 3.96 (ddd, J = 3.5, 6.7, 8.7 Hz, 1H), 4.76 (br s, 1H), 7.31-7.33 (m, 3H), 7.49-7.52 (m, 2H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ = 13.8, 18.7, 36.2, 56.3, 71.5, 128.4, 129.2, 132.7, 132.9, 175.5; IR (CHCl_3): ν = 2977 (s), 2936 (m), 2401 (s), 1711 (s), 1522 (s), 1477 (m), 1423 (s), 1333 (w), 1219 (s) cm^{-1} ; GC-EIMS on methyl ester derivative (254) m/z = 254 (M^+ , 1), 182 (74), 150 (18), 123 (100), 109 (14); anal. calcd. for $\text{C}_{12}\text{H}_{16}\text{O}_3\text{S}$ (240.3): C 59.97; H 6.71; found: C 60.23; H 6.79.

Rel(2S,3S)-3-hydroxy-2-(phenylthio)butyric acid (3d)

solid; purified by recrystallization with 65/35-petroleum ether/EtOAc; 92% yield; m. p. = 83-85 °C; ^1H NMR (CDCl_3 , 400 MHz) δ = 1.44 (d, J = 6.3 Hz, 3H), 3.59 (d, J = 7.2 Hz, 1H), 4.14 (apparent quintuplet, J = 6.6 Hz, 1H), 7.32-7.35 (m, 3H), 7.50-7.52 (m, 2H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ = 20.4, 57.6, 67.9, 128.4, 129.2, 132.5, 132.9, 176.2; IR (CHCl_3): ν = 3620 (m), 3432 (s), 3028 (s), 1726 (s), 1392 (s), 1224 (s), 1047 (s), 878 (m) cm^{-1} ; GC-EIMS on methyl ester derivative (226.3) m/z = 226 (M^+ , 4), 182 (86), 150 (24), 123 (100), 109 (21), 77 (15), 45 (17); anal. calcd. for $\text{C}_{10}\text{H}_{12}\text{O}_3\text{S}$ (212.3): C 56.58; H 5.70; found: C 56.89; H 5.77.

Rel(2S,3S)-3-hydroxy-2-methyl-2-(phenylthio)pentanoic acid (3e)

Characterized by using a chromatographically enriched mixture; eluent: 70/28/2, petroleum ether/EtOAc/AcOH; 95% overall yield of **3e** and **4e**; ^1H NMR (CDCl_3 , 400 MHz) δ = 1.05 (t, J = 7.3 Hz, 3H), 1.36 (s, 3H), 1.39-1.45 (m, 1H), 1.99-2.09 (m, 1H), 3.78 (d, J = 1.1, 9.9 Hz, 1H), 7.30-7.34 (m, 2H), 7.46-7.50 (m, 3H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ = 11.1, 17.0, 23.7, 58.8, 75.8, 129.0, 129.6, 129.8, 137.0, 178.1; GC-EIMS on methyl ester derivative (254.3) m/z = 254 (M^+ , 7), 195 (11), 151 (100), 123 (22), 110 (4), 43 (24).

trans-1-hydroxy-2-(phenylthio)cyclohexene-1-carboxylic acid (4a)

solid; purified by recrystallization from *n*-hexane; 96% yield; m. p. = 134-135 °C; ^1H NMR (CDCl_3 , 200 MHz) δ = 1.18-1.43 (m, 1H), 1.60-1.75 (m, 4H), 2.00-2.12 (m, 3H), 3.18 (dd, J = 4.0, 10.0 Hz, 1H), 4.94 (br s, 2H), 7.20-7.30 (m, 3H), 7.44-7.49 (m, 2H); ^{13}C NMR (CDCl_3 , 50.3 MHz) δ = 24.7, 29.1, 30.6, 35.5, 56.5, 77.9, 127.2, 129.0, 132.1, 135.9, 176.6; IR (CHCl_3): ν = 3538 (s), 3053 (m), 2936 (s), 2616 (m), 1692 (s), 1438 (s), 1304 (w),

1051 (m), 910 (w) cm^{-1} ; GC-EIMS on methyl ester derivative (266) $m/z = 266$ (M^+ , 85), 248 (30), 207 (35), 157 (20), 110 (100), 79 (35); anal. calcd. for $\text{C}_{13}\text{H}_{16}\text{O}_3\text{S}$ (275): C 61.88; H 6.39; found: C 61.93; H 6.41.

Rel(2S,3R)-2-hydroxy-3-phenyl-3-(phenylthio)propanoic acid (4b)

solid; purified by recrystallization with 85/15-petroleum ether/EtOAc; 95% yield; m. p. = 129-130 $^{\circ}\text{C}$; ^1H NMR (CD_3COCD_3 , 400 MHz) $\delta = 4.62$ (d, $J = 4.1$ Hz, 1H), 4.75 (d, $J = 4.1$ Hz, 1H), 7.22-7.30 (m, 6H), 7.40-7.48 (m, 4H); ^{13}C NMR (CD_3COCD_3 , 100.6 MHz) $\delta = 56.7$, 73.5, 128.1, 128.4, 128.7, 129.8, 130.3, 132.6, 135.8, 138.6, 173.2; IR (CHCl_3): $\nu = 3516$ (m), 3014 (s), 2584 (m), 1730 (s), 1481 (w), 1276 (m), 1101 (s), 913 (w) cm^{-1} ; GC-EIMS on methyl ester derivative (288.3) $m/z = 288$ (M^+ , 1), 199 (100), 165 (8), 119 (10), 91 (36), 77 (4); anal. calcd. for $\text{C}_{15}\text{H}_{14}\text{O}_3\text{S}$ (274.3): C 65.67; H 5.14; found: C 65.77; H 5.13.

Rel(2s,3R)-2-hydroxy-3-(phenylthio)hexanoic acid (4c)

solid; purified by recrystallization with 9/1-petroleum ether/EtOAc; 96% yield; m. p. = 93-94 $^{\circ}\text{C}$; ^1H NMR (CDCl_3 , 400 MHz) $\delta = 0.94$ (t, $J = 7.2$ Hz, 3H), 1.49-1.69 (m, 6H), 3.58 (ddd, $J = 2.9$, 4.5, 9.4 Hz, 1H), 4.31 (d, $J = 2.8$ Hz, 1H), 7.28-7.34 (m, 3H), 7.46-7.48 (m, 2H); ^{13}C NMR (CDCl_3 , 100.6 MHz) $\delta = 13.7$, 20.6, 31.1, 52.9, 71.8, 127.7, 129.3, 132.2, 133.5, 175.7; IR (CHCl_3): $\nu = 3621$ (w), 3025 (s), 2400 (m), 1764 (w), 1522 (m), 1424 (m), 1226 (s), 1046 (m), 929 (m) cm^{-1} ; GC-EIMS on methyl ester derivative (254) $m/z = 254$ (M^+ , 18), 165 (80), 123 (100), 77 (7), 55 (11); anal. calcd. for $\text{C}_{12}\text{H}_{16}\text{O}_3\text{S}$ (240.3): C 59.97; H 6.71; found: C 60.13; H 6.77.

Rel(2S,3R)-2-hydroxy-3-(phenylthio)butyric acid (4d)

oil; purified by chromatography on silica gel, eluent: 63/35/2 petroleum ether/EtOAc/AcOH; 94% yield; ^1H NMR (CDCl_3 , 400 MHz) $\delta = 1.28$ (d, $J = 7.0$ Hz, 3H), 3.72 (dd, $J = 2.7$, 7.0, 14.1 Hz, 1H), 4.31 (d, $J = 2.7$ Hz, 1H), 6.54 (br s, 2H), 7.25-7.33 (m, 3H), 7.44-7.48 (m, 2H); ^{13}C NMR (CDCl_3 , 100.6 MHz) $\delta = 20.4$, 57.6, 67.9, 128.4, 129.2, 132.5, 132.9, 176.2; IR (CHCl_3): $\nu = 2987$ (m), 2933 (w), 1731 (s), 1602 (w), 1480 (m), 1376 (m), 1252 (s); cm^{-1} ; GC-EIMS on methyl ester derivative (226.3) $m/z = 226$ (M^+ , 17), 167 (4), 149 (8), 137 (100), 109 (29), 77 (7); anal. calcd. for $\text{C}_{10}\text{H}_{12}\text{O}_3\text{S}$ (212.3): C 56.58; H 5.70; found: C 56.89; H 5.85.

Rel(2S,3R)-2-hydroxy-2-methyl-3-(phenylthio)pentanoic acid (4e)

solid; purified by recrystallization with 85/15-petroleum ether/EtOAc; 95% yield; m. p. = 94-96 °C; ^1H NMR (CDCl_3 , 400 MHz) δ = 1.18 (t, J = 7.2 Hz, 3H), 1.57 (s, 3H), 1.60-1.78 (m, 2H), 3.24 (dd, J = 2.8, 11.1 Hz, 1H), 5.44 (br s, 1H), 7.20-7.31 (m, 3H), 7.47-7.48 (m, 2H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ = 12.7, 24.9, 25.0, 61.5, 78.5, 126.9, 129.1, 131.2, 136.7, 178.7; IR (CHCl_3): ν = 3516 (m), 2980 (s), 2582 (w), 1723 (s), 1584 (w), 1454 (w), 1169 (m), 1042 (w), 881 (w) cm^{-1} ; GC-EIMS on methyl ester derivative (288.3) m/z = 254 (M^+ , 9), 195 (9), 151 (100), 123 (29), 109 (22), 73 (9); anal. calcd. for $\text{C}_{12}\text{H}_{16}\text{O}_3\text{S}$ (240.3): C 59.97; H 6.71; found: C 60.12; H 6.50.

***trans*-2-(allylthio)-1-hydroxycyclohexanecarboxylic acid (5a)**

oil; chromatography on silica gel, eluent: 70/28/2-petroleum ether/EtOAc/AcOH; 65% yield; ^1H NMR (CDCl_3 , 400 MHz) δ = 1.31-1.42 (m, 1H), 1.64-1.67 (m, 4H), 2.05-2.12 (m, 3H), 2.68 (dd, J = 4.7, 10.6 Hz, 1H), 3.19 (dd, J = 6.8, 13.6 Hz, 1H), 3.31 (dd, J = 7.6, 13.6 Hz, 1H), 5.08-5.16 (m, 2H), 5.74-5.84 (m, 1H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ = 21.8, 24.7, 30.5, 36.2, 50.5, 60.5, 77.5, 117.5, 134.4, 179.0; IR (CHCl_3): ν = 2940 (s), 2860 (m), 1714 (s), 1636 (w), 1448 (m), 1331 (w), 1230 (s) cm^{-1} ; GC-EIMS on methylester derivative (230) m/z = 230 (M^+ , 95), 171 (71), 141 (51), 129 (68), 101 (100), 79 (43), 67 (60); anal. calcd. for $\text{C}_{14}\text{H}_{16}\text{O}_3\text{S}$ (216.3): C 55.53; H 7.46; found: C 55.43; H 7.30.

Rel(2S,3R)-3-(allylthio)-2-hydroxy-3-phenylpropanoic acid (5b)

oil; chromatography on silica gel, eluent: 70/28/2-petroleum ether/EtOAc/AcOH; 90% yield; ^1H NMR (CDCl_3 , 400 MHz) δ = 2.95 (dd, J = 8.0, 13.9 Hz, 1H), 3.10-3.15 (m, 1H), 4.26 (d, J = 3.6 Hz, 1H), 4.61 (d, J = 3.6 Hz, 1H), 5.00-5.14 (m, 2H), 5.74-5.78 (m, 1H), 7.27-7.31 (m, 3H), 7.33-7.38 (m, 2H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ = 34.4, 51.2, 73.0, 117.9, 128.1, 128.4, 129.1, 133.6, 136.2; GC-EIMS on methylester derivative: decomposition; anal. calcd. for $\text{C}_{12}\text{H}_{14}\text{O}_3\text{S}$ (238.3): C 60.48; H 5.92; found: C 60.45; H 5.90.

Rel(2S,3R)-3-(allylthio)-2-hydroxy-3-hexanoic acid (5c)

oil; chromatography on silica gel, eluent: 70/28/2-petroleum ether/EtOAc/AcOH; 86% yield; ^1H NMR (CDCl_3 , 400 MHz) δ = 0.90 (t, J = 7.0 Hz, 3H), 1.20-1.65 (m, 4H), 2.97-3.08 (m, 1H), 3.10-3.30 (m, 2H), 4.26 (d, J = 1.1 Hz, 1H), 5.05-5.25 (m, 2H), 5.70-5.90 (m, 1H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ = 13.6, 20.3, 31.8, 34.7, 47.9, 72.9, 117.6, 134.0, 176.1; GC-EIMS on methylester derivative (218) m/z = 129 (100), 95 (13), 87 (77), 55 (20); anal. calcd. for $\text{C}_9\text{H}_{16}\text{O}_3\text{S}$ (204,3): C, 52.91; H, 7.89; found: C 52.81; H 7.80.

Rel(2S,3R)-3-(allylthio)-2-hydroxy-2-methyl-3-pentanoic acid (5e)

oil; chromatography on silica gel, eluent: 70/28/2-petroleum ether/EtOAc/AcOH; 85% yield; ^1H NMR (CDCl_3 , 400 MHz) δ = 1.06 (t, J = 7.3 Hz, 3H), 1.40-1.60 (m, 2H), 1.56 (s, 3H), 2.55-2.65 (m, 2H), 3.04-3.25 (m, 1H), 5.05-5.15 (m, 2H), 5.73-5.85 (m, 1H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ = 12.4, 24.5, 24.7, 36.5, 55.8, 78.2, 117.5, 134.4, 179.7; GC-EIMS on methylester derivative (218) m/z = 218 (M^+ 3), 115 (100), 81 (26), 73 (43), 43 (34); anal. calcd. for $\text{C}_9\text{H}_{16}\text{O}_3\text{S}$ (204,3): C, 52.91; H, 7.89; found: C 52.96; H 7.86.