Supporting Information

Indirect Cation Pool Method. Rapid Generation of Alkoxy-carbenium Ion Pools from Thioacetals

Seiji Suga, Kouichi Matsumoto, Koji Ueoka, and Jun-ichi Yoshida*

*Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering,
Kyoto University, Nishikyo-ku, Kyoto 615-8510, Japan.*

General Remarks.

GC analysis was performed on a gas chromatograph (SHIMADZU GC-14B) equipped with a flame ionization detector using a fused silica capillary. ^1H and ^{13}C NMR spectra were recorded in CDCl_3 on a Varian Gemini 2000 (^1H 300 MHz, ^{13}C 75 MHz), Varian MERCURY plus-400 (^1H 400 MHz, ^{13}C 100 MHz), or JEOL ECA-600P (^1H 600 MHz, ^{13}C 150 MHz) spectrometer with Me_4Si as an internal standard unless otherwise noted. Mass spectra were obtained on JEOL JMS SX-102A mass spectrometer. CSI (cold-spray ionization) mass spectra were recorded on a JEOL JMS-T100CSK spectrometer. IR spectra were measured with a SHIMADZU FTIR 1600 spectrometer. Thin-layer chromatography (TLC) was carried out by using Merck precoated silica gel F_{254} plates (thickness 0.25 mm). Flash chromatography was carried out on a column of silica gel (Kanto Chem. Co., Silica Gel N, spherical, neutral, 40-100 μm). Gel permeation chromatography (GPC) was carried out on Japan Analytical Industry LC-908 equipped with JAIGEL-1H and 2H using CHCl_3 as eluent. All reactions were carried out under Ar atmosphere unless otherwise noted.

Materials.

Dichloromethane was washed with water, distilled from P_2O_5 , redistilled from dried K_2CO_3 to remove a trace amount of acid, and stored over molecular sieves 4A. Dry THF was used as obtained (Kanto Chemical Co., Inc.). ArSSAr ($\text{Ar} = p\text{-FC}_6\text{H}_4$)¹ was prepared according to the procedures in the literatures.

Thioacetals.

1-Methoxy-1-phenylthiononane,² 2-phenylthiotetrahydropyran,³ and 2-phenylthiotetrahydrofuran³ were prepared according to the literature procedures.

(Methoxy(phenylthio))methylbenzene. To a toluene solution of benzaldehyde dimethyl acetal (786.8 mg, 5.17 mmol) and thiophenol (612.0 mg, 5.55 mmol), was added $\text{BF}_3\cdot\text{OEt}_2$ (736.0 mg, 5.19 mmol) at -78°C . The solution was stirred for 2 h. Dry pyridine (2.4 ml) was added. The mixture was diluted with ether and was washed with 1 M NaOH solution and water. After drying over Na_2SO_4 and removal of solvent, colorless oil thus obtained was purified by flash chromatography (hexane/EtOAc 50:1 to 10:1) to

obtain the title compound (757.8 mg, 3.29 mmol, 64%).⁴

1-Methoxy-1-(4-chlorophenylthio)nonane. To a solution of 4-chlorobenzenethiol (5.19 g, 36.0 mmol) in tetrahydrofuran (THF) (64 ml), was added triethylamine (7.2 ml, 52 mmol) and chloromethyl methyl ether (3.3 ml, 43 mmol) at 0 °C. After being stirred at room temperature for 2 h, the reaction mixture was partitioned between saturated aqueous NH₄Cl and ether. The organic phase was separated, washed with saturated aqueous NaHCO₃, and was dried over MgSO₄. The solvent was removed under reduced pressure and the residue was purified by distillation (130 °C, 14 mmHg), to obtain (4-chlorophenylthio)methyl methyl ether (5.76 g, 30.5 mmol).

To a solution of (4-chlorophenylthio)methyl methyl ether (1.51 g, 8.0 mmol) in THF (20 ml), was added butyllithium (1.54 M in hexane, 6.4 ml, 10 mmol) at -78 °C. The mixture was stirred at -45 °C for 3 h and was cooled to -78 °C. 1-Iodooctane (2.62 g, 10.9 mmol) was added, and the mixture was stirred at -78 °C for 20 min. Saturated aqueous NH₄Cl was added. The organic materials were extracted with ether, and the organic extracts were washed with brine and dried over MgSO₄. After removal of the solvent, the residue was purified by flash chromatography (hexane) to obtain the title compound (1.60 g, 5.3 mmol, 66 %): TLC *R_f* 0.73 (hexane/EtOAc 20:1); ¹H NMR (400 MHz, CDCl₃) δ 0.87 (t, *J* = 6.8 Hz, 3H), 1.21-1.28 (m, 10H), 1.40-1.43 (m, 2H), 1.36-1.44 (m, 2H), 1.62-1.78 (m, 2H), 3.46 (s, 3H), 4.57 (t, *J* = 6.8 Hz, 1H), 7.23-7.26 (m, 2H), 7.36-7.40 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 14.2, 22.8, 26.3, 29.2, 29.3, 29.5, 31.9, 35.6, 55.4, 90.9, 128.7, 131.6, 133.5, 134.6; IR (neat) 2953, 1476, 1130, 1092, 623 cm⁻¹; LRMS (EI) *m/z* 300 (M⁺), 269 (M⁺-OCH₃), 187 (M⁺-C₈H₁₇), 157 (M⁺-SC₆H₄Cl); HRMS (EI) calcd for C₁₆H₂₅OClS: 300.1315, found: 300.1313.

¹H NMR spectrum of 1-methoxy-1-(4-chlorophenylthio)nonane.

¹³C NMR spectrum of 1-methoxy-1-(4-chlorophenylthio)nonane.

1-Methoxy-1-(4-methylphenylthio)nonane. To a solution of 4-methylbenzenethiol (6.13 g, 49.4 mmol) in THF (100 ml), was added triethylamine (10 ml, 73 mmol) and chloromethyl methyl ether (4.69 g, 58.3 mmol) at 0 °C. After being stirred at room temperature for 2 h, the reaction mixture was partitioned between saturated aqueous NH_4Cl and ether. The organic phase was separated, washed with saturated aqueous NaHCO_3 , and was dried over MgSO_4 . The solvent was removed under reduced pressure and the residue was purified by distillation to obtain (4-methylphenylthio)methyl methyl ether (6.71 g, 39.9 mmol).

To a solution of (4-methylphenylthio)methyl methyl ether (3.21 g, 19.1 mmol) in THF (45 ml), was added butyllithium (1.54 M in hexane, 15 ml, 23.1 mmol) at -78 °C. The mixture was stirred at -45 °C for 3 h and cooled to -78 °C. 1-Iodooctane (5.90 g, 24.6 mmol) was added, and the mixture was stirred at -78 °C for 30 min. Saturated aqueous NH_4Cl was added. The organic materials were extracted with ether, and the organic extracts were washed with brine and dried over MgSO_4 . After removal of the solvent, the residue was purified by flash chromatography (hexane/EtOAc 50:1) to obtain the title compound (4.43 g, 15.8 mmol, 83 %): TLC R_f 0.42 (hexane/EtOAc 20:1); ^1H NMR (300 MHz, CDCl_3) δ 0.87 (t, J = 6.6 Hz, 3H), 1.22-1.34 (m, 10H), 1.36-1.52 (m, 2H), 1.64-1.80 (m, 2H), 2.32 (s, 3H), 3.46 (s, 3H), 4.54 (t, J = 6.6 Hz, 1H), 7.09 (m, 2H), 7.35 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 14.1, 21.1, 22.6, 26.2, 29.1, 29.2, 29.4, 31.8, 35.6, 55.4, 91.1, 129.4, 129.4, 134.0, 137.5; IR (neat) 2924, 1466, 1088, 644 cm^{-1} ; LRMS (EI) m/z 280 (M^+), 157 ($\text{M}^+ - \text{SC}_6\text{H}_4\text{CH}_3$); HRMS (EI) calcd for $\text{C}_{17}\text{H}_{28}\text{OS}$: 280.1861, found: 280.1859.

¹H NMR spectrum of 1-methoxy-1-(4-methylphenylthio)nonane.

¹³C NMR spectrum of 1-methoxy-1-(4-methylphenylthio)nonane.

Electrochemical Generation and Accumulation of $\text{ArS}(\text{ArSSAr})^+$.

Typical Procedure and CSI-MS Analysis. The anodic oxidation was carried out in an H-type divided cell (4G glass filter) equipped with a carbon felt anode (Nippon Carbon JF-20-P7, ca. 320 mg, dried at 250 °C/1 mmHg for 1 h before use) and a platinum plate cathode (40 mm x 20 mm). In the anodic chamber was placed a solution of ArSSAr ($\text{Ar} = p\text{-FC}_6\text{H}_4$) (101.9 mg, 0.401 mmol) in 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2$ (8.0 mL). In the cathodic chamber were placed 0.3 M $\text{Bu}_4\text{NBF}_4/\text{CH}_2\text{Cl}_2$ (8.0 mL) and trifluoromethanesulfonic acid (41.0 mg, 0.273 mmol). The constant current electrolysis (8 mA) was carried out at -78 °C with magnetic stirring until 0.67 F/mol of electricity was consumed. The anodic solution was analyzed by CSI-MS (spray temperature; 0 °C): HRMS (CSI) m/z calcd for $\text{C}_{18}\text{H}_{12}\text{F}_3\text{S}_3$ ($\text{ArS}(\text{ArSSAr})^+$ ($\text{Ar} = p\text{-FC}_6\text{H}_4$), M^+): 318.0047, found: 318.0087, m/z calcd for $\text{C}_{34}\text{H}_{48}\text{BF}_7\text{S}_3$ ($[\text{Bu}_4\text{NBF}_4]\text{ArS}(\text{ArSSAr})^+$ ($\text{Ar} = p\text{-FC}_6\text{H}_4$), M^+) 710.2925, found: 710.2945.

CSI-MS of $\text{ArS}(\text{ArSSAr})^+$ ($\text{Ar} = p\text{-FC}_6\text{H}_4$) (spray temperature; 0 °C).

NMR Analysis of Electrochemically Generated $\text{ArS}(\text{ArSSAr})^+$ ($\text{Ar} = p\text{-FC}_6\text{H}_4$). The anodic oxidation was carried out in a divided cell equipped with a carbon felt anode and a platinum plate cathode. In the anodic chamber were placed a solution of $\text{ArS}(\text{ArSSAr})^+$ ($\text{Ar} = p\text{-FC}_6\text{H}_4$) (52.9 mg, 0.208 mmol) in 4.0 mL of 0.3 M Bu_4NBF_4 in $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1). In the cathodic chamber were placed trifluoromethanesulfonic acid (21.0 mg, 0.140 mmol) and 0.3 M Bu_4NBF_4 in $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1) (4.0 mL). The constant current electrolysis (4.0 mA) was carried out at -78 °C with magnetic stirring. After 0.67 F/mol of electricity was consumed, the reaction mixture of the anodic chamber was transferred to a 5 mm ϕ NMR tube with a septum cap under Ar atmosphere at -78 °C. The NMR measurement was carried out at -80 °C. Chemical shifts are reported using the methylene signal of CH_2Cl_2 at δ 5.32 (^1H NMR) as a standard. The huge signal coming from CH_2Cl_2 is reduced by usual pulse techniques.⁵ ^1H NMR (600 MHz, $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1)) δ 7.22-7.32 (t, $J = 8.6$ Hz, 6H), 7.44-7.74 (br s, 6H).

¹H NMR spectrum of electrochemically generated ArS(ArSSAr)⁺ (Ar = *p*-FC₆H₄)
(-80 °C).

NMR Analysis of $\text{ArS}(\text{ArSSAr})^+$ ($\text{Ar} = p\text{-FC}_6\text{H}_4$) Generated by the Reaction of ArSSAr and SbCl_5 . $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1) (0.6 mL) and ArSSAr ($\text{Ar} = p\text{-FC}_6\text{H}_4$) (30.7 mg, 0.121 mmol) were added to a 5 mm ϕ NMR tube with a septum cap under Ar atmosphere at -78°C . Then SbCl_5 (35.3 mg, 0.118 mmol) was added and the NMR tube was shaken at the same temperature. The NMR measurement was carried out at -80°C . Chemical shifts are reported using the methylene signal of CH_2Cl_2 at δ 5.32 (^1H NMR) as a standard. The huge signal coming from CH_2Cl_2 is reduced by usual pulse techniques: ^1H NMR (600 MHz, $\text{CH}_2\text{Cl}_2/\text{CD}_2\text{Cl}_2$ (10:1)) δ 7.22-7.31 (t, $J = 8.6$ Hz, 6H), 7.47-7.58 (br t, 6H).

¹H NMR spectrum of ArS(ArSSAr)⁺ (Ar = *p*-FC₆H₄) generated by the reaction of ArSSAr and SbCl₅ (-80 °C).

Generation of Alkoxycarbenium Ions.

Typical Procedure. The anodic oxidation was carried out in an H-type divided cell (4G glass filter) equipped with a carbon felt anode and a platinum plate cathode (40 mm x 20 mm). In the anodic chamber was placed a solution of ArSSAr (Ar = *p*-FC₆H₄) (101.9 mg, 0.401 mmol,) in 0.3 M Bu₄NBF₄/CH₂Cl₂ (8.0 mL). In the cathodic chamber were placed 0.3 M Bu₄NBF₄/CH₂Cl₂ (8.0 mL) and trifluoromethanesulfonic acid (41.0 mg, 0.273 mmol). The constant current electrolysis (8 mA) was carried out at -78 °C with magnetic stirring until 0.67 F/mol of electricity was consumed. To the anodic chamber containing electrogenerated ArS(ArSSAr)⁺, was added 1-methoxy-1-phenylthiononane (53.3 mg, 0.200 mmol) and the mixture was stirred for 5 min at -78 °C. The solution thus obtained was used for the subsequent reaction.

NMR Analysis of Alkoxycarbenium Ion (2). The anodic oxidation was carried out in a divided cell equipped with a carbon felt anode and a platinum plate cathode. In the anodic chamber were placed a solution of ArSSAr (Ar = *p*-FC₆H₄) (81.9 mg, 0.322 mmol) in 4.0 mL of 0.3 M Bu₄NBF₄/CH₂Cl₂/CD₂Cl₂ (10:1). In the cathodic chamber were placed trifluoromethanesulfonic acid (32.4 mg, 0.216 mmol) and 0.3 M Bu₄NBF₄ in CH₂Cl₂/CD₂Cl₂ (10:1) (4.0 mL). The constant current electrolysis (4.0 mA) was carried out

^{13}C NMR spectrum of alkoxy-carbenium ion (**2**) ($-80\text{ }^{\circ}\text{C}$).

Reactions of Alkoxy-carbenium Ion with Carbon Nucleophiles.

4-Methoxy-dodec-1-ene (Typical Procedure). The electrolysis of ArSSAr (Ar = *p*-FC₆H₄) (102 mg, 0.401 mmol) was carried out as described above. To the ArS(ArSSAr)⁺ thus generated in the anodic chamber, was added 1-methoxy-1-phenylthiononane (52.7 mg, 0.198 mmol) at $-78\text{ }^{\circ}\text{C}$ and the reaction mixture was stirred for 5 min. Then allyltrimethylsilane (56.6 mg, 0.495 mmol) was added and the resulting solution was stirred at $-78\text{ }^{\circ}\text{C}$ for 15 min. The reaction was quenched with Et₃N. The solvent was removed under reduced pressure and the residue was quickly filtered through a short column (2 x 3 cm) of silica gel to remove Bu₄NBF₄. The silica gel was washed with ether (150 mL). The GC analysis of the combined filtrate indicated that the title compound was formed in 89 % yield (GC *t*_R 7.1 min, column, CBP-1; 0.22 mm ϕ x 0.25 μm x 25 m; initial oven temperature, $100\text{ }^{\circ}\text{C}$; rate of temperature increase, $10\text{ }^{\circ}\text{C}/\text{min}$). The isolated product was identified by the comparison of its spectral data with those of an authentic sample.⁶

The other products, 3-(1-methoxynonyl)cyclohexene (a mixture of two diastereomers; ca. 5:4 by ^{13}C NMR),⁶ 3-methoxy-1-phenylundecan-1-one,⁶ 3-methoxy-2,2-dimethylundecanoic acid methyl ester,⁶ 2-(cyclohex-2-enyl)tetrahydropyran (a mixture of two diastereomers; ca. 7:1 by GC

analysis),⁶ and 2-(cyclohex-2-enyl)tetrahydrofuran (a mixture of two diastereomers; ca. 9:1 by GC analysis),⁶ and (1-methoxy-but-3-enyl)benzene⁷ were identified by the comparison of their spectral data with those of authentic samples.

4-Fluorophenyl(phenyl)disulfide. The title compound was obtained as a byproduct in the reactions described above: ¹H NMR (300 MHz, CDCl₃) δ 6.94-7.04 (m, 2H), 7.20-7.35 (m, 3H), 7.40-7.52 (m, 4H); ¹³C NMR (75 MHz, CDCl₃) δ 116.2 (*J* = 22.8 Hz), 127.4, 128.0, 129.1, 130.6 (*J* = 8.0 Hz), 132.3 (*J* = 3.4 Hz), 136.8, 162.4 (*J* = 247.1 Hz); IR (neat) 3061, 1879, 1590, 1487, 1478, 1439, cm⁻¹; LRMS (EI) *m/z* 236 (M⁺), 127 (M⁺-C₆H₅S), 109 (M⁺-C₆H₄FS); HRMS (EI) calcd for C₁₂H₉FS₂ (M⁺): 236.01297, found: 236.0132.

¹H NMR spectrum of 4-fluorophenyl(phenyl)disulfide.

¹³C NMR spectrum of 4-fluorophenyl(phenyl)disulfide.

Evaluation of Reaction Time of Alkoxycarbenium Ion **2** with Allyltrimethylsilane.

The electrolysis of ArSSAr (Ar = *p*-FC₆H₄) (103.1 mg, 0.405 mmol) was carried out as described above. To the ArS(ArSSAr)⁺ thus generated in the anodic chamber, was added 1-methoxy-1-phenylthiononane (54.3 mg, 0.204 mmol) at -78 °C, and the reaction mixture was stirred for 5 min. Then, allyltrimethylsilane (46.5 mg, 0.407 mmol) was added and the resulting solution was stirred at -78 °C for 1 min, followed by the treatment with 1-(trimethylsiloxy)cyclohexene (70.0 mg, 0.411 mmol) at -78 °C for 10 min. The reaction was quenched with Et₃N. The solvent was removed under reduced pressure and the residue was quickly filtered through a short column (2 x 3 cm) of silica gel to remove Bu₄NBF₄. The silica gel was washed with ether (150 mL). The GC analysis of the combined filtrate indicated that the 4-methoxy-dodec-1-ene was formed in 81 % yield (GC *t*_R 7.09 min, column, CBP-1; 0.22 mm ϕ x 0.25 μ m x 25 m; initial oven temperature, 100 °C; rate of temperature increase, 10 °C/min). Only a trace amount of the product derived from 1-(trimethylsiloxy)cyclohexene was obtained, indicating that the reaction of **2** with allyltrimethylsilane completed within 1 min at -78 °C.

References

- (1) Hirano, M.; Yakabe, S.; Monobe, H.; Morimoto, T. *J. Chem. Reserch (S)* **1998**, 472.
- (2) Yoshida, J.; Sugawara, M.; Tatsumi, M.; Kise, N. *J. Org. Chem.* **1998**, 63, 5950.
- (3) Braga, A. L.; Sliveria, C. C.; Dornelles, L.; Zeni, G.; Galarza, F. A. D.; Wessjohann, L. A. *Syn. Commun.* **1995**, 25, 3155.
- (4) Sato, T.; Otera, J.; Nozaki, H. *Tetrahedron* **1989**, 45, 1209.
- (5) Guéron, M.; Plateau, P.; Decorps, M. *Prog. Nucl. Magn. Reson. Spectrosc.* **1991**, 23, 135.
- (6) Suga, S.; Suzuki, S.; Yamamoto, A.; Yoshida, J. *J. Am. Chem. Soc.* **2000**, 122, 10244.
- (7) Okajima, M.; Suga, S.; Itami, K.; Yoshida, J. *J. Am. Chem. Soc.* **2005**, 127, 6930.