Indium-Mediated Reaction of Trialkylsilyl Propargyl Bromide with Aldehydes: Highly Regioselective Synthesis of Allenic and Homopropargylic Alcohols

Man-Jing Lin and Teck-Peng Loh*

*Department of Chemistry, National University of Singapore, 3 Science Drive 3,
Singapore 117543*

Supporting information:

Experimental Section

General: THF was distilled from sodium/benzophenone. Hexane and dichloromethane were distilled from CaH₂. TLC was carried out with pre-coated Merck 60 F₂₅₄ plates. Silica gel 60 (Merck, 400-630 mesh) was used for column chromatography. ¹H and ¹³C NMR spectra were taken in CDCl₃ on Bruker DPX300 and referenced to internal tetramethylsilane. MS (EI) spectra were recorded on a Hewlett-Packard 5890A gas chromatogram. IR spectra were recorded on a Bio-RAD FTS 165 FT-IR Spectrometer.

Indium-Mediated Propargylation of Aldehydes; General Procedure:

Trimethylsilyl propargyl bromide (191 mg, 1.0 mmol, 2 equiv) was added to a mixture of the indium powder (114 mg, 1.0 mmol, 2 equiv) and indium tribromide (18 mg, 0.05 mmol, 0.1 equiv) in 1 mL anhydrous THF at room temperature with stirring under an atmosphere of dry nitrogen. After 15 min, nonyl aldehyde (86 µL, 0.5 mmol, 1 equiv) was added at room temperature. The mixture was refluxed overnight, and finally quenched with 10 mL of 1M HCl solution. The aqueous layer was extracted with ethyl acetate (10 mL × 3). The combined organic extracts were washed with brine, dried over anhydrous magnesium sulfate, concentrated under vacuum, and purified by flash silica gel column chromatography to provide 121 mg (95% yield) of 1-Trimethylsilyl-dodec-1-yn-4-ol (5e) as a clear oil.

1-Phenyl-6-trimethylsilyl-hex-5-yn-3-ol (5a)

Yield%: 85%

R_f 0.40 (hexane : ethyl acetate 4:1)

¹H NMR: (300 MHz, CDCl₃)

δ 7.31-7.26 (m, 2H, phenyl), 7.22-7.20 (m, 3H, phenyl), 3.76 (m, 1H, **CHOH**), 2.73 (m, 2H, Ph**CH**₂), 2.48 (dd, *J* = 4.86, 16.71 Hz, 1H, C≡C**HH**), 2.38 (dd, *J* = 6.63, 16.71 Hz, 1H, C≡C**HH**), 1.85 (m, 2H, PhCH₂**CH**₂), 0.16 (s, 9H, Si(**CH**₃)₃) ppm

¹³C NMR: (75 MHz, CDCl₃)

δ 141.7, 128.5, 128.4, 125.9, 103.0 (C≡CSi), 87.8 (**CHOH**), 69.1 (C≡CSi), 37.8 (CH₂**CHOH**), 31.8 (PhCH₂), 29.0 (CH₂C≡C), 0.11 (Si(**CH**₃)₃) ppm

FTIR: 3434, 3020, 2961, 2170, 1718, 1492, 1454, 1411, 1251, 1216, 1040, 845, 758, 700, 669 cm⁻¹

EIHRMS: Calcd for C₁₅H₂₂OSi : 246.1440, found : 246.1396.

1-Phenyl-6-trimethylsilanyl-hex-1-en-5-yn-3-ol (5b)

Yield%: 89%

R_f 0.40 (hexane : ethyl acetate 4:1)

¹H NMR: (300 MHz, CDCl₃)

δ 7.41-7.25 (m, 5H, Phenyl), 6.67 (d, *J* = 16.0 Hz, 1H, Ph**CH**), 6.27 (dd, *J* = 5.91, 9.75 Hz, 1H, PhCH=**CH**), 4.45 (m, 1H, **CHOH**), 2.62 (dd, *J* = 4.5, 16.71 Hz, 1H, C≡C**HH**), 2.36 (dd, *J* = 6.27, 16.71 Hz, 1H, C≡C**HH**), 2.25 (br, 1H, **OH**), 0.176 (s, 9H, Si(**CH**₃)₃) ppm

¹³C NMR: (75 MHz, CDCl₃)

δ 136.5, 131.1, 130.2, 128.6, 127.8 (Ph**CH**), 126.6 (PhCH=**CH**), 102.5 (C≡CSi), 88.1 (**CHOH**), 70.7 (C≡CSi), 29.3 (CH₂**CHOH**), 0.06 (Si(**CH**₃)₃) ppm

FTIR: 3366, 3027, 2958, 2900, 2358, 2176, 1933, 1600, 1493, 1449, 1250, 1100, 1029, 966, 843, 748, 693 cm⁻¹

EIHRMS: Calcd for C₁₅H₂₀OSi : 244.1283, found : 244.1278.

1-Cyclohexyl-4-trimethylsilanyl-but-3-yn-ol (5c)

Yield%: 90%

R_f 0.48 (hexane : ethyl acetate 4:1)

¹H NMR: (300 MHz, CDCl₃)

δ 3.43 (m, 1H, **CHOH**), 2.43 (dd, *J* = 3.6, 16.71 Hz, 1H, **C≡CCHH**), 2.32 (dd, *J* = 7.65, 16.71 Hz, 1H, **C≡CCHH**), 2.12 (br, 1H, **OH**), 1.21-0.96 (m, 11H, c-C₆**H**₁₁), 0.117 (s, 9H, Si(**CH**₃)₃) ppm

¹³C NMR: (75 MHz, CDCl₃)

δ 103.8 (**C≡CSi**), 87.3 (**CHOH**), 73.8 (**C≡CSi**), 42.6 (**CHCHOH**), 28.9, 28.1 (**CH₂CHOH**), 26.4, 26.1, 26.0 (2C, overlap), 0.01 (Si(**CH**₃)₃) ppm

FTIR: 3402, 2928, 2851, 2663, 2174, 1713, 1450, 1422, 1349, 1249, 1198, 1103, 1086, 1015, 842, 760, 698, 651 cm⁻¹

EIHRMS: Calcd for C₁₃H₂₄OSi : 224.1596, found : 224.1594.

1-Phenyl-4-trimethylsilanyl-but-3-yn-1-ol (5d)

Yield%: 92%

R_f 0.38 (hexane : ethyl acetate 4:1)

¹H NMR: (300 MHz, CDCl₃)

δ 7.38-7.26 (m, 5H, phenyl), 4.84 (t, J = 6.27 Hz, 1H, PhCHOH), 2.65 (d, J = 6.27 Hz, 2H, CH₂C≡CSi), 2.45 (brd, 1H, CHOH), 0.17(s, 9H, Si(CH₃)₃) ppm

¹³C NMR: (75 MHz, CDCl₃)

δ 142.5, 128.9, 127.8, 125.7, 103.0, 87.8 (CH₂C≡CSi), 72.3 (CHOH), 31.0 (CHCH₂C), 0.06 (Si(CH₃)₃) ppm

FTIR: (neat) ν 3396, 3033, 2960, 2902, 2177, 1724, 1604, 1494, 1454, 1250, 1202, 1039, 846, 759, 700 cm⁻¹

EIHRMS: Calcd for C₁₃H₁₈OSi : 218.1127, found: 218.1125

1-Trimethylsilanyl-dodec-1-yn-4-ol (5e)

Yield%: 95%

R_f 0.45 (hexane : ethyl acetate 4:1)

¹H NMR: (300 MHz, CDCl₃)

δ 3.71 (m, 1H, CHOH), 2.45 (dd, J = 4.89, 16.71 Hz, 1H, C≡CCHH), 2.33 (dd, J = 6.96, 16.71 Hz, 1H, C≡CCHH), 1.96 (br, 1H, OH), 1.50 (m, 2H, CH₂CHOH), 1.27 (m, 12H, (CH₂)₆), 0.87 (t, J = 5.9 Hz, 3H, CH₃), 0.15 (s, 9H, Si(CH₃)₃) ppm

¹³C NMR: (75 MHz, CDCl₃)

δ 103.4 (C≡CSi), 86.9 (CHOH), 70.0 (C≡CSi), 36.2 (CH₂CHOH), 31.9, 29.5, 29.5, 29.3, 28.9, 25.6, 22.7 (CH₂CH₃), 14.1 (CH₃), 0.07 (Si(CH₃)₃) ppm

FTIR: 3365, 2957, 2926, 2855, 2176, 1465, 1250, 1084, 1005, 842, 760, 698, 648 cm⁻¹

EIHRMS: Calcd for C₁₅H₃₀OSi : 254.2066, found : 254.2067.

1-Triisopropylsilanyl-dodec-1-yn-4-ol (9a)

Yield%: 58%

R_f 0.52 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 3.72 (m, 1H, CHOH), 2.50 (dd, $J = 4.86, 16.71$ Hz, 1H, $\text{C}\equiv\text{CCHH}$), 2.39 (dd, $J = 6.6, 16.71$ Hz, 1H, $\text{C}\equiv\text{CCHH}$), 1.27 (m, 14H, $(\text{CH}_2)_7$), 1.07 (m, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$), 0.88 (t, $J = 6.96$ Hz, 3H, CH_3) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 104.8 ($\text{C}\equiv\text{CSi}$), 83.6 (CHOH), 70.1 ($\text{C}\equiv\text{CSi}$), 36.2, 31.9, 29.6, 29.5, 29.3, 28.9 ($\text{CH}_2\text{C}\equiv\text{C}$), 25.6, 22.7, 18.6 ($\text{SiCH}(\text{CH}_3)_2$), 14.1 (CH_3), 11.2 ($\text{SiCH}(\text{CH}_3)_2$) ppm

FTIR: 3429, 2927, 2856, 2177, 1712, 1648, 1466, 1378, 1250, 1033, 843, 760 cm^{-1} .

EIHRMS: Calcd for $\text{C}_{21}\text{H}_{42}\text{OSi}$: 338.3055, found : 338.3017.

1-(*tert*-Butyl-diphenyl-silanyl)-dodec-1-yn-4-ol (9f)

Yield%: 35%

R_f 0.49 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 7.79 (m, 4H, Phenyl), 7.37 (m, 6H, Phenyl), 3.85 (m, 1H, CHOH), 2.65 (dd, $J = 5.22, 17.07$ Hz, 1H, $\text{C}\equiv\text{CCHH}$), 2.55 (dd, $J = 6.6, 17.07$ Hz, 1H, $\text{C}\equiv\text{CCHH}$), 1.28 (m, 14H, $(\text{CH}_2)_7$), 1.09 (s, 9H, $\text{Si}(\text{C}(\text{CH}_3)_3)_3$), 0.88 (t, $J = 6.96$ Hz, 3H, CH_3) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 135.5, 135.5, 133.5, 133.5, 129.5, 129.5, 127.7, 127.7, 107.6 ($\text{C}\equiv\text{CSi}$), 82.8 (CHOH), 70.2 ($\text{C}\equiv\text{CSi}$), 36.4, 31.9, 29.6, 29.5, 29.3, 29.2 ($\text{CH}_2\text{C}\equiv\text{C}$), 27.1 ($\text{SiC}(\text{CH}_3)_3$), 25.6, 22.7, 18.5 ($\text{SiC}(\text{CH}_3)_3$), 14.1 (CH_3) ppm

FTIR: 3359, 2959, 2928, 2856, 2175, 1428, 1378, 1109, 820 cm^{-1} .

ESIHRMS: Calcd for $\text{C}_{28}\text{H}_{40}\text{OSiNa}$: 443.2746, found : 443.2751.

20-(1'-Hydroxy-4'-Trimethylsilanyl-3'-butyn-1'-yl)-3-oxopregn-4-ene (5f)

Yield%: 65%

R_f 0.50 (hexane : ethyl acetate 1:1)

^1H NMR: (300 MHz, CDCl_3)

δ 5.72 (s, 1H, 4- CH), 3.87 (m, 1H, 22- CH), 1.18 (s, 3H, 19- CH_3), 0.91 (d, $J = 6.63$ Hz, 3H, 21- CH_3), 0.73 (s, 3H, 18- CH_3), 0.16 (s, 9H, $\text{Si}(\text{CH}_3)_3$) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 199.6 (3- C), 171.5 (5- C), 123.8 (4- CH), 104.2 ($\text{C}\equiv\text{CSi}$), 87.3 (22- CH), 71.8 ($\text{C}\equiv\text{CSi}$), 55.8, 53.8, 52.6, 42.4, 39.6, 39.4, 38.6, 35.7, 34.0, 32.9, 32.0, 29.7, 27.7, 26.9, 24.1, 21.1, 17.4, 15.3, 11.8, 11.5, 0.08 ($\text{Si}(\text{CH}_3)_3$) ppm

FTIR: 3445, 3019, 2937, 1659, 1455, 1215, 1024, 844, 756, 669 cm^{-1}

EIHRMS: Calcd for $\text{C}_{28}\text{H}_{44}\text{O}_2\text{Si}$: 440.3111, found : 440.3102

20-(1'-Hydroxy-3'-butyn-1'-yl)-3-oxopregn-4-ene (10)

Yield%: 90%

R_f 0.41 (hexane : ethyl acetate 1:1)

^1H NMR: (300 MHz, CDCl_3)

δ 5.73 (s, 1H, 4-CH), 3.90 (m, 1H, 22-CH), 1.19 (s, 3H, 19-CH₃), 0.92 (d, $J = 6.63$ Hz, 3H, 21-CH₃), 0.73 (s, 3H, 18-CH₃) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 199.6 (3-C), 171.5 (5-C), 123.8 (4-C), 81.7 (C $\equiv$ CSi), 71.8 (22-CH), 70.4 (C $\equiv$ CSi), 55.8, 53.8, 52.5, 42.4, 39.7, 39.6, 38.6, 35.7, 34.0, 33.0, 32.0, 29.7, 27.7, 25.5, 24.1, 21.1, 17.4, 11.9, 11.4 ppm

FTIR: 3435, 3017, 2941, 2360, 1659, 1444, 1216, 756, 668 cm^{-1}

EIHRMS: Calcd for $\text{C}_{25}\text{H}_{36}\text{O}_2$: 368.2715, found : 368.2713.

20-(1-Hydroxy-3-buten1-yl)-3-oxopregn-4-ene (11)

Yield%: 98%

R_f 0.51 (hexane : ethyl acetate 2:1)

^1H NMR: (300 MHz, CDCl_3)

δ 5.87-5.76 (m, 1H, $\text{CH}=\text{CH}_2$), 5.73 (br, 1H, 4- CH), 5.13 (d, $J = 7.65$ Hz, 1H, $\text{CH}=\text{CH}_2$), 5.09 (br, 1H, $\text{CH}=\text{CH}_2$), 3.71 (dd, $J = 4.89, 8.36$ Hz, 1H, 22- CH), 1.18 (s, 3H, 19- CH_3), 0.93 (d, $J = 5.91$ Hz, 3H, 21- CH_3), 0.72 (s, 3H, 18- CH_3) ppm

^{13}C NMR: (75 MHz, CDCl_3)

199.6 (3- C), 171.5 (5- C), 135.7 ($\text{CH}_2=\text{CH}$), 123.8 (4- CH), 117.5 ($\text{CH}_2=\text{CH}$), 72.4 (22- CH), 55.8, 53.8, 52.6, 42.4, 40.2, 40.1, 39.7, 38.6, 35.7 (2C, overlap), 34.0, 32.9, 32.0, 27.7, 24.1, 21.1, 17.4, 11.9, 11.7 ppm

FTIR: 3368, 2939, 1674, 1432, 1229, 886 cm^{-1} .

EIHRMS: Calcd for $\text{C}_{25}\text{H}_{38}\text{O}_2$: 370.2888, found : 370.2872.

Indium-Mediated Allenylation of Aldehydes; General Procedure:

To a suspension of indium powder (57 mg, 0.5 mmol, 2 equiv) in $\text{H}_2\text{O}/\text{THF}$ (0.5 mL, 5:1) was added nonyl aldehyde (43 μL , 0.25 mmol, 1 equiv) and then triisopropylsilyl propargyl bromide (69 mg, 2 equiv). The mixture was vigorously stirred at room temperature for 10 hours. Standard workup and purified by flash silica gel column chromatography to provide 44 mg (52% yield) 3-Triisopropylsilyl-dodeca-1,2-dien-4-ol (**8a**).

3-Triisopropylsilyl-dodeca-1,2-dien-4-ol (**8a**)

Yield%: 52%

R_f 0.55 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 4.56 (m, 2H, $\text{C}=\text{C}=\text{CH}_2$), 4.01 (m, 1H, CHOH), 1.27 (m, 14H, $(\text{CH}_2)_7$), 1.09 (m, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$), .088 (t, $J = 6.45$ Hz, 3H, CH_3) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 209.6 ($\text{C}=\text{C}=\text{CH}_2$), 96.9 ($\text{C}=\text{C}=\text{CH}_2$), 72.4 (CHOH), 69.8 ($\text{C}=\text{C}=\text{CH}_2$), 38.3, 31.9, 29.6, 29.5, 29.3, 26.2, 22.7, 18.7 ($\text{CH}(\text{CH}_3)_2$), 14.1 (CH_3), 11.7 ($\text{SiCH}(\text{CH}_3)_2$) ppm

FTIR: 3468, 2927, 2866, 1922, 1465, 1067, 1008, 883, 811, 761 cm^{-1} .

EIHRMS: Calcd for $\text{C}_{21}\text{H}_{42}\text{OSi}$: 338.3005 , found : 338.3004.

1-Phenyl-4-triisopropylsilanyl-hexa-4,5-dien-3-ol (8b)

Yield%: 71%

R_f 0.54 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 7.32-7.16 (m, 5H, Phenyl), 4.61 (m, 2H, $\text{C}=\text{C}=\text{CH}_2$), 4.06 (brs, 1H, CHOH), 2.78 (m, 2H, CH_2Ar), 1.90 (m, 2H, $\text{CH}_2\text{CH}_2\text{Ar}$), 1.08 (m, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 209.6 ($\text{C}=\text{C}=\text{CH}_2$), 142.1, 128.5, 128.3, 125.8, 96.7 ($\text{C}=\text{C}=\text{CH}_2$), 72.6 (CHOH), 69.1 ($\text{C}=\text{C}=\text{CH}_2$), 40.1 ($\text{CH}_2\text{CH}_2\text{Ar}$), 32.6 (CH_2Ar), 18.6 (CH_3), 11.6 ($\text{SiCH}(\text{CH}_3)_2$) ppm

FTIR: 3449, 2944, 2866, 1922, 1717, 1450, 883, 809, 748, 699 cm^{-1} .

EIHRMS: Calcd for $\text{C}_{21}\text{H}_{34}\text{OSi}$: 330.2379, found : 330.2390.

1-Phenyl-4-triisopropylsilanyl-hexa-1,4,5-trien-3-ol (8c)

Yield%: 46%

R_f 0.54 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 7.40-7.21 (m, 5H, Phenyl), 6.58 (d, J = 16.02 Hz, 1H, PhCH=CH), 6.30 (dd, J = 6.69, 15.69 Hz, 1H, PhCH=CH), 4.71 (m, 1H, CHOH), 4.66 (m, 2H, C=C=CH₂), 1.08 (m, 21H, SiCH(CH₃)₃) ppm

^{13}C NMR: (75 MHz, CDCl₃)

δ 210.1 (C=C=CH₂), 131.8, 130.3, 129.7, 128.6, 127.7, 126.6, 96.3 (C=C=CH₂), 73.0 (CHOH), 70.9 (C=C=CH₂), 18.7 (CH₃), 11.7 (SiCH(CH₃)₂)₃ ppm

FTIR: 3626, 3020, 2394, 1657, 1218, 925, 776, 743 cm⁻¹.

EIHRMS: Calcd for C₂₁H₃₂OSi : 328.2222, found : 328.2213.

1-Cyclohexyl-2-triisopropylsilanyl-buta-2,3-dien-1-ol (8d)

Yield%: 61%

R_f 0.52 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl₃)

δ 4.55 (m, 2H, C=C=CH₂), 3.83 (t, J = 5.58 Hz, 1H, CHOH), 1.27-0.92 (m, 11H, c-C₆H₁₁), 1.09 (m, 21H, Si(CH(CH₃)₂)₃) ppm

^{13}C NMR: (75 MHz, CDCl₃)

δ 209.4 (C=C=CH₂), 95.4 (C=C=CH₂), 74.0 (CHOH), 72.2 (C=C=CH₂), 43.5 (CHCHOH), 31.1, 26.7, 26.5, 26.5, 26.1, 18.7 (CH₃), 11.8 (Si(CH(CH₃)₂)₃) ppm

FTIR: 3429, 2930, 2866, 1928, 1637, 1459, 1217, 1077, 1002, 873, 758 cm⁻¹.

EIHRMS: Calcd for C₁₉H₃₆OSi : 308.2535, found : 308.2533.

1-Phenyl-2-triisopropylsilanyl-buta-2,3-dien-1-ol (8e)

Yield%: 59%

R_f 0.55 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 7.41-7.25 (m, 5H, Phenyl), 5.16 (s, 1H, CHOH), 4.64 (m, 2H, $\text{C}=\text{C}=\text{CH}_2$), 1.08 (m, 21H, $\text{Si}(\text{CH}(\text{CH}_3)_2)_3$) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 209.8 ($\text{C}=\text{C}=\text{CH}_2$), 143.4, 128.3, 127.8, 127.2, 97.4 ($\text{C}=\text{C}=\text{CH}_2$), 73.2 (CHOH), 72.5 ($\text{C}=\text{C}=\text{CH}_2$), 18.6 (CH_3), 11.7 ($\text{SiCH}(\text{CH}_3)_2$) ppm

FTIR: 3440, 2945, 2866, 1925, 1459, 1192, 1018, 883, 817, 760, 700 cm^{-1} .

EIHRMS: Calcd for $\text{C}_{19}\text{H}_{30}\text{OSi}$: 302.2066, found : 302.2071.

3-(*tert*-Butyl-diphenyl-silanyl)-dodeca-1,2-dien-4-ol (8f)

Yield%: 45%

R_f 0.55 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 7.72-7.67 (m, 4H, Phenyl), 7.47-7.35 (m, 6H, Phenyl), 4.80 (m, 2H, $\text{C}=\text{C}=\text{CH}_2$), 3.83 (br s, 1H, CHOH), 1.38-.098 (m, 14H, $(\text{CH}_2)_7$), 1.14 (s, 9H, $\text{SiC}(\text{CH}_3)_3$), 0.87 (t, $J = 6.96$ Hz, 3H, CH_3) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 210.0 (C=C=CH₂), 136.3, 136.3, 133.7, 133.7, 129.5, 129.5, 127.8, 127.7, 96.1 (C=C=CH₂), 74.1 (CHOH), 70.3 (C=C=CH₂), 37.5, 31.8, 29.4, 29.2, 29.2, 28.1 (SiC(CH₃)₃), 25.8, 22.7, 19.5 (SiC(CH₃)₃), 14.1 (CH₂CH₃) ppm

FTIR: 3582, 3081, 2928, 2857, 1925, 1463, 1428, 110, 998, 820, 758, 739, 701 cm⁻¹.

EIHRMS: Calcd for C₂₈H₄₀OSi : 420.2848, found : 420.2846.

4-(*tert*-Butyl-diphenyl-silanyl)-1-phenyl-hexa-4,5-dien-3-ol (8g)

Yield%: 50%

R_f 0.39 (hexane : ethyl acetate 4:1)

¹H NMR: (300 MHz, CDCl₃)

δ 7.72-7.67 (m, 4H, Phenyl), 7.44-7.35 (m, 6H, Phenyl), 7.26-7.12 (m, 3H, Phenyl), 6.97-6.95 (m, 2H, Phenyl), 4.83 (m, 2H, C=C=CH₂), 3.90 (br s, 1H, CHOH), 2.67-2.59 (m, 1H, PhCH), 2.42-2.30 (m, 1H, PhCH), 1.70-1.62 (m, 2H, PhCH₂CH₂), 1.10 (s, 9H, SiC(CH₃)₃) ppm

¹³C NMR: (75 MHz, CDCl₃)

δ 210.0 (C=C=CH₂), 142.0, 136.2, 136.2, 133.6, 133.6, 129.6, 129.5, 128.4, 128.2, 127.9, 127.8, 125.6, 96.0 (C=C=CH₂), 74.3 (CHOH), 69.8 (C=C=CH₂), 39.2 (PhCH₂CH₂), 32.4 (PhCH₂) 28.1 (SiC(CH₃)₃), 19.5 (SiC(CH₃)₃) ppm

FTIR: 3563, 3455, 3071, 2932, 2859, 1924, 1428, 1105, 820, 757, 701, 609 cm⁻¹.

EIHRMS: Calcd for C₂₈H₃₂OSi : 412.2222, found : 412.2236.

4-(*tert*-Butyl-diphenyl-silanyl)-1-phenyl-hexa-1,4,5-trien-3-ol (8h)

Yield%: 52%

R_f 0.42 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 7.74-7.68 (m, 4H, Phenyl), 7.46-7.26 (m, 6H, Phenyl), 7.24-7.13 (m, 3H, Phenyl), 7.10 (m, 2H, Phenyl), 6.07-5.96 (m, 2H, PhCH=CH), 4.88 (m, 2H, C=C=CH_2), 4.52 (br s, 1H, CHOH), 1.14 (s, 9H, $\text{SiC(CH}_3)_3$) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 210.7 (C=C=CH_2), 136.8, 136.5, 136.4, 133.5, 133.3, 130.7, 129.9, 129.7, 129.5, 128.3, 127.8, 127.7, 127.4, 126.5, 95.5 (C=C=CH_2), 74.7 (CHOH), 71.4 (C=C=CH_2), 28.1 ($\text{SiC(CH}_3)_3$), 19.5 ($\text{SiC(CH}_3)_3$) ppm

FTIR: 3583, 3226, 2931, 2861, 1629, 1106, 751, 698 cm^{-1} .

EIHRMS: Calcd for $\text{C}_{28}\text{H}_{30}\text{OSi}$: 410.2066, found : 410.2054.

2-(*tert*-Butyl-diphenyl-silanyl)-1-cyclohexyl-buta-2,3-dien-1-ol (8i)

Yield%: 56%

R_f 0.55 (hexane : ethyl acetate 4:1)

^1H NMR: (300 MHz, CDCl_3)

δ 7.70-7.65 (m, 4H, Phenyl), 7.44-7.33 (m, 6H, Phenyl), 4.79 (m, 2H, C=C=CH_2), 3.66-3.63 (m, 1H, CHOH), 1.26-0.88 (m, 11H, $\text{c-C}_6\text{H}_{11}$), 1.14 (s, 9H, $\text{SiC(CH}_3)_3$) ppm

^{13}C NMR: (75 MHz, CDCl_3)

δ 209.8 (C=C=CH₂), 136.3, 136.3, 133.8, 133.8, 129.5, 129.5, 127.7, 127.7, 94.8 (C=C=CH₂), 74.4 (CHOH), 74.0 (C=C=CH₂), 42.6, 30.8, 28.2 (SiC(CH₃)₃), 26.4, 26.3, 26.0, 25.7, 19.6 (SiC(CH₃)₃) ppm

FTIR: 3435, 2993, 2935, 2860, 1924, 1634, 1319, 1105, 998, 832, 754, 701 cm⁻¹.

EIHRMS: Calcd for C₂₆H₃₄OSi : 390.2379, found : 390.2366.

2-(*tert*-Butyl-diphenyl-silanyl)-1-phenyl-buta-2,3-dien-1-ol (8j)

Yield%: 55%

R_f 0.40 (hexane : ethyl acetate 4:1)

¹H NMR: (300 MHz, CDCl₃)

δ 7.54-7.51 (m, 2H, Phenyl), 7.42-7.38 (m, 2H, Phenyl), 7.35-7.23 (m, 6H, Phenyl), 7.14-7.12 (m, 3H, Phenyl), 7.00-6.97 (m, 2H, Phenyl), 4.95 (m, 1H, CHOH), 4.81 (m, 2H, C=C=CH₂), 1.10 (s, 9H, SiC(CH₃)₃) ppm

¹³C NMR: (75 MHz, CDCl₃)

δ 211.1 (C=C=CH₂), 136.3, 136.3, 133.6, 133.6, 133.1, 129.5, 129.5, 129.3, 127.8, 127.8, 127.6, 127.2, 96.5 (C=C=CH₂), 74.5 (CHOH), 73.0 (C=C=CH₂), 28.1 (SiC(CH₃)₃), 19.6 (SiC(CH₃)₃) ppm

FTIR: 3437, 3020, 2923, 2858, 2401, 1637, 1226, 754, 669 cm⁻¹.

EIHRMS: Calcd for C₂₆H₂₈OSi : 384.1909, found : 384.1918.

NMR Studies:

Lewis acid: InBr_3

Lewis acid: InF_3

References

1. Isaac, M. B.; Chan, T. H. *Chem. Commun.* **1995**, 1003.
2. Hammond, G. B.; Wang, Z. G. *J. Org. Chem.* **2000**, 65, 6547.
3. Loh T. P.; Hu, Q. Y.; Chok, Y. K.; Tan, K. T. *Tetrahedron Lett.* **2001**, 42, 9277..