Intermolecular and Intramolecular α -Amidoalkylation Reactions using Bismuth Triflate as Catalyst

by

Frédéric Pin,[†] Sébastien Comesse,[†] Bernard Garrigues,[‡] Štefan Marchalín[‡] and Adam Daich^{†}*

URCOM, EA 3221, UFR des Sciences & Techniques, Université du Havre, 25 rue Philippe Lebon, BP: 540, F-76058 Le Havre Cedex, France; Hétérochimie Fondamentale et Appliquée, UMR-5069 CNRS, Université Paul Sabatier, 118 route de Narbonne, F-31062 Toulouse Cedex 04, France; and Department of Organic Chemistry, Slovak University of Technology, SK-81237 Bratislava, Slovak Republic

adam.daich@univ-lehavre.fr

RECEIVED DATE (to be automatically inserted after your manuscript is accepted if required according to the journal that you are submitting your paper to)

Bismuth(III) triflate mediated intermolecular and intramolecular α -amidoalkylation processes.

Authors may be reached by phone at (+33) 02-32-74-44-03, by fax at (+33) 02-32-74-43-91 or by E.mail at the following address: adam.daich@univ-lehavre.fr

¹H- NMR discussion

(For more details, please see NMR spectra of all products in the supporting Information file).

The structure of all products and intermediates reported herein was confirmed by their ¹H, ¹³C NMR spectra including DEPT programs and elemental analyses. So, the ¹H NMR spectra of allyl derivatives **4a-I** and alkyl ones as **5a** and **6a** recorded in CDCl₃ showed the methylene group of the –N-CH₂– functionalities as an AB system due to the diastereotopic effect with a coupling constant of $J=14.1-15.6$ Hz characteristic of geminate protons. Also, the angular proton CH in **4a-i**, **5a** and **6a** appears as a doublet of doublet centered at $\delta=3.53$ up to 5.07 ppm with $J=3.9-5.0$ Hz and $J=4.7-7.0$ Hz characteristic of the coupling observed in the –CH-CH₂– enchainment installed in the above final allyl and alkyl products by the α -amidoalkylation process. Of particular note, this proton resonated in general up-field compared to the same one in corresponding α -hydroxy lactams **2a-i** and α -acetoxy lactams **3a-i** in which it appeared as a doublet (CH(OH)) at $\delta=5.56$ up to 5.77 ppm with $J=8.8-10.4$ Hz and as a singlet at $\delta=6.86$ up to 7.11 ppm, respectively. In the case of **3i**, this proton appears as

doublet at 6.06 ppm with $J=5.5$ Hz. More interestingly, the ^{13}C NMR spectra of **4a-i** revealed the presence of two sp^2 -hybridized carbons corresponding to $\text{CH}=\text{CH}_2$ fragment of the allyl group and the absence of the quaternary ester carbonyl function ($\text{C}=\text{O}$) and CH_3 of the acyl group, in the starting α -acetoxy lactams **3a-i**, respectively. These values are comparable to that reported in the literature for related allyl products.² In the case of products **5a** and **6a**, an additional acyl and benzoyl carbonyl function ($\text{C}=\text{O}$) instead of $-\text{CH}=\text{CH}_2$ residue were appeared at $\delta=205.8$ ppm and $\delta=197.3$, respectively.

Several key features in the ^1H and ^{13}C NMR spectra of the cyclized compounds **4j-l** and **4p,g** were diagnostic of the structure. For instance, in the ^1H NMR spectra of isoindoloisoquinoline **4j** and isoindolo-[1,3]benzoxazines **4k,l** the methylene protons of the $-\text{N}-\text{CH}_2-$ functionalities, which were an AB system due to the diastereotopic effect in α -acetoxy lactams congeners **3j-l** (with $J=14.5$, 10.5 and 11.5 Hz, respectively), appeared also as an AB system but interestingly with increasing coupling constant values, which are respectively of $J=15.5$, 16.8 and 17.0 Hz.^{1,2} The spectra showed also the disappearance of two signals corresponding to the CH_3 protons of the lateral acetyl group and one aromatic proton in **3j-l**. For pyrroloisoquinolinone **4p**, the ^1H NMR revealed the presence of the $-\text{CH}_2-\text{CH}_2-$ group and absence of one proton in the aromatic region while in the NMR spectra of pyrroloisoquinolinone **4g**, this chain appears as two triplet at $\delta=2.94$ and 3.72 with $J=6.3$ Hz as a consequence of the absence of the stereogenic center in the nitrogen adjacent position. They revealed also the presence of a singlet corresponding to the angular proton in **4j-l** at $\delta=5.69$, 5.96 and 6.14 ppm, shifted downfield compared to those in corresponding α -acetoxy lactams **3j** ($\delta=6.89$ ppm), **3k** ($\delta=7.18$ ppm) and **3l** ($\delta=7.16$ ppm), respectively. It is of particular note that the latter proton appeared as a doublet with trans coupling constant of $J=4.0$ Hz in (1*S*,10*bR*)-1-acetoxy-1,2,3,5,6,10*b*-hexahydro-pyrrolo[2,1-*a*]isoquinolin-3-one (**4p**),³ while in the case of **4q** no expected angular proton at nitrogen α -position appeared. As a consequence of the spontaneous carbon-carbon double bond isomerization in the pyrrolidinone ring leading to the formation of the more stable isomer in which the olefin was in conjugation with the benzene ring, an angular proton at the adjacent position of the lactamic carbonyl function appeared as a doublet of quadruplet at $\delta=3.24$ ppm with $J=7.8$ Hz ($\text{CH}-\text{CH}_3$ coupling) and $J=3.1$ Hz ($\text{CH}-\text{CH}=\text{}$ coupling).

¹ (a) Katritzky, A. R.; Mehta, S.; He, H.-Y. *J. Org. Chem.* **2001**, *66*, 148-152 and references therein. (b) Hucher, N.; Daïch, A.; Decroix, B. *J. Heterocycl. Chem.* **1998**, *35*, 1477-1483.

² See Winn, M.; Zaugg, H. E. *J. Org. Chem.* **1968**, *33*, 3779 and also reference 1a.

³ Lee, Y. S.; Kang, D. W.; Lee, S.-J.; Park, H. *J. Org. Chem.* **1995**, *60*, 7149-7152.

Carbon-13 NMR data proved to be of particular value in the assignment of structure of cyclized compounds as isoindoloisoquinoline **4j**, isoindolo[1,3]benzoxazines **4k,l** and pyrrolo[2,1-*a*]isoquinolones **4p,g**. The most distinguishing feature in the ^{13}C NMR spectra was the regular appearance of one downfield signal at $\delta=168.2$ ppm, 168.2 ppm and 168.6 ppm for **4j,k,l** and $\delta=170.3$ ppm and 180.0 ppm for **4p,g**, which have been attributed to the lactamic carbonyl C=O. From these results, it is apparent that the variation of the central nucleus of the triheterocyclic systems **4j-l** and **4p** led to only small changes in the chemical shifts values for this resonance. In comparison with their α -acetoxy lactams congeners **3j-l** and **3p**, the difference of the chemical shift values are of $\Delta\delta=+0.2$, $+0.4$, $+0.8$ and $+0.2$ ppm. But we noted in this data set for the same carbon in **4g**, a consistent downfield shift of $\Delta\delta=+9.2$ ppm in comparison between the cyclized product **4g** and α -acetoxy lactam **3g**. This is due probably, in the latter case, to the presence of a double bond in conjugation with benzene ring as constraint system. Furthermore, the key feature in the ^{13}C NMR spectra of products **4j-l**, **4p** and **4q** was the respective appearance of thirteen, seven and nine signals in the aromatic region. Moreover, two of these resonances including a carbonyl signal and one aromatic quaternary carbon disappear in the corresponding DEPT program spectra, as a consequence of the intramolecular α -amidoalkylation cyclization. In the case of **4q**, an additional quaternary carbon located at α -position of the lactamic nitrogen atom disappears also after similar NMR measurements.

II- Experimental Section

General remarks. Melting points were taken with a capillary melting point apparatus and are uncorrected. The infrared (IR) absorption spectra were determined as solutions in potassium bromide and are indicated in cm^{-1} . The ^1H and ^{13}C NMR spectra were recorded as solutions in CDCl_3 or $\text{DMSO}-d_6$ at 200 or 300 MHz (^1H) and 50.3 or 75 MHz (^{13}C), respectively, and chemical shifts (δ) are expressed in ppm relative to TMS as internal standard. Thin layer chromatography (TLC) was performed using silica gel analytical plates (F_{254}) of 0.25 mm thickness. The detection on TLC plates was performed by UV light at 254 or 365 nm or using *p*-anisaldehyde revelator. The analytical results of elemental analysis are within 0.4 % of theoretical values and were obtained from INSA institution at Rouen, F-76130 Mt-St-Aignan in France.

6-(4-Methoxybenzyl)-6H-pyrrolo[3,4-b]pyridine-5,7-dione (1g). A mixture of *p*-methoxybenzylamine (1.37 g, 10 mmol), pyridine-2,3-dicarboxylic anhydride (1.49 g, 10 mmol) and three drops of anhydrous triethylamine in toluene (100 mL) was refluxed with a Dean-stark apparatus for 36 hours. The reaction mixture was cooled, then concentrated under reduced pressure. The residue was

dissolved into dichloromethane (20 mL), washed with 5% hydrochloric acid solution then with a 5% sodium hydrogenocarbonate solution. The organic layer was dried over magnesium sulfate, concentrated under *vacuo*, and purified by chromatography on silica gel column using a mixture of cyclohexane/ethyl acetate (3/2) as eluent. This product was isolated as a white solid in 78% yield; R_f 0.56 (cyclohexane/AcOEt 2:3); mp 130 °C; IR (KBr): ν 1781, 1725, 1515, 1466, 1392, 1344, 1249 cm^{-1} ; ^1H NMR (CDCl_3): δ 3.74 (s, 3H, OCH_3), 4.82 (s, 2H, CH_2N), 6.83 (d, 2H, $J = 8.6$ Hz, H_{aro}), 7.36 (d, 2H, $J = 8.6$ Hz, H_{aro}), 7.56 (dd, 1H, $J = 7.8$ and 5.5 Hz, H_{aro}), 8.11 (dd, 1H, $J = 7.8$ and 1.6 Hz, H_{aro}), 8.91 (dd, 1H, $J = 5.5$ and 1.6 Hz, H_{aro}); ^{13}C NMR (CDCl_3): δ 41.5 (CH_2), 55.4 (CH_3), 114.2 (2CH_{aro}), 127.5 (CH_{aro}), 127.6 (C_q), 128.3 (C_q), 129.6 (C_q), 130.5 (2CH_{aro}), 131.4 (CH_{aro}), 155.4 (CH_{aro}), 159.5 (C_q), 166.1 ($\text{C}=\text{O}$), 166.2 ($\text{C}=\text{O}$); Anal. Calcd. for $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_3$ (268.08): C, 67.16; H, 4.51; N, 10.44. Found: C, 67.07; H, 4.42; N, 10.34.

2-(4-Methoxybenzyl)-2H-pyrrolo[3,4-*b*]quinoline-1,3-dione (1h). To a solution of quinoline-2,3-dicarboxylic anhydride (995 mg, 5 mmol) in 50 mL of dry THF was added at room temperature under stirring dropwise *p*-methoxybenzylamine (685 mg, 5 mmol). After 2 h of reaction, 1 equiv. of DCC (1.03 g, 5 mmol) and 10 mol% of DMAP were added and the reaction was heated at reflux for 12 h. The reaction mixture was filtered on short column of a Celite pad and the filtrate was concentrated in *vacuo*. The residue was redissolved in dichloromethane and filtered again using same protocol. The solvent was carefully removed under vacuum to give white crystals, which were purified by flash-chromatography, eluting with cyclohexane/EtOAc 1:3. This led to isolation of the expected imide **1h** as a white solid in 70% yield; R_f 0.47 (cyclohexane/AcOEt 1:1); mp 208 °C; IR (KBr): ν 1779, 1720, 1628, 1615, 1593, 1515, 1467, 1395 cm^{-1} ; ^1H NMR (CDCl_3): δ 3.75 (s, 3H, OCH_3), 4.91 (s, 2H, CH_2N), 6.83 (d, 2H, $J = 8.6$ Hz, H_{aro}), 7.43 (d, 2H, $J = 8.6$ Hz, H_{aro}), 7.73 (dd, 1H, $J = 7.8$ and 7.0 Hz, H_{aro}), 7.91 (dd, 1H, $J = 8.6$ and 7.0 Hz, H_{aro}), 8.02 (d, 1H, $J = 7.8$ Hz, H_{aro}), 8.40 (d, 1H, $J = 8.6$ Hz, H_{aro}), 8.61 (s, 1H, H_{aro}); ^{13}C NMR (CDCl_3): δ 41.8 (CH_2), 55.5 (CH_3), 114.3 (2CH_{aro}), 123.2 (C_q), 128.3 (C_q), 128.9 (C_q), 129.7 (CH_{aro}), 130.1 (CH_{aro}), 130.6 (2CH_{aro}), 131.7 (CH_{aro}), 132.8 (CH_{aro}), 132.9 (CH_{aro}), 150.8 (C_q), 151.0 (C_q), 159.6 (C_q), 165.9 ($\text{C}=\text{O}$), 166.1 ($\text{C}=\text{O}$); Anal. Calcd. for $\text{C}_{19}\text{H}_{14}\text{N}_2\text{O}_3$ (318.1): C, 71.69; H, 4.43; N, 8.80. Found: C, 71.54; H, 4.34; N, 8.71.

General Procedure for the Reduction of Imides 1a-p. NaBH_4 (3 to 6 equiv.) was added by portions over 20 minutes at 0 °C to a stirred solution of the imide **1a-q** (1 equiv.) in MeOH (20 mL per 1 mmol of imide **1**). The reaction mixture was directly concentrated under reduced pressure, then CH_2Cl_2 was added. The organic layer was washed with water, brine, dried over MgSO_4 and the solvent evaporated. The residue was then purified by chromatography on a silica gel column to give

the expected hydroxy lactams **2a-q**. For reduction of imides **1g,h**, 1.2 equiv. NaBH₄ and 1 equiv. of Mg(ClO₄)₂ in a mixture of THF/CHCl₃ (1/1 of V/V) were used.

2-(Furan-2-ylmethyl)-3-hydroxyisoindolin-1-one (2a). This product was isolated as a brown solid; 93% yield; *R_f* 0.24 (CH₂Cl₂); mp 127 °C; IR (KBr): ν 3300 (OH), 1706 cm⁻¹; ¹H NMR (CDCl₃): δ 3.90 (d, 1H, *J* = 9.4 Hz, OH), 4.24 (d, 1H, *J* = 15.6 Hz, CH₂N), 4.60 (d, 1H, *J* = 15.6 Hz, CH₂N), 5.69 (d, 1H, *J* = 9.4 Hz, CH), 6.25 (s, 2H, H_{fur}), 7.28 (s, 1H, H_{fur}), 7.38-7.64 (m, 4H, H_{aro}); ¹³C NMR (CDCl₃): δ 35.6 (CH₂), 81.6 (CH), 108.7 (CH_{aro}), 110.6 (CH_{aro}), 123.6 (CH_{aro}), 123.7 (CH_{aro}), 129.9 (CH_{aro}), 131.3 (C_q), 132.7 (CH_{aro}), 142.7 (CH_{aro}), 144.1 (C_q), 150.3 (C_q), 167.4 (C=O); Anal. Calcd. for C₁₃H₁₁NO₃ (229.07): C, 68.11; H, 4.84; N, 6.11. Found: C, 68.01; H, 4.76; N, 6.05.

3-Hydroxy-2-(4-methoxybenzyl)-isoindolin-1-one (2f). This product was isolated as a white solid; 94% yield; *R_f* 0.33 (cyclohexane/AcOEt 1:1); mp 158 °C; IR (KBr): ν 3691, 3559, 2686, 2411, 1700, 1684, 1613 cm⁻¹; ¹H NMR (CDCl₃): δ 3.35 (d, 1H, *J* = 10.4 Hz, OH), 3.74 (s, 3H, OCH₃), 4.17 (d, 1H, *J* = 14.9 Hz, CH₂N), 4.79 (d, 1H, *J* = 14.9 Hz, CH₂N), 5.56 (d, 1H, *J* = 10.4 Hz, CH), 6.79 (d, 2H, *J* = 8.6 Hz, H_{aro}), 7.22 (d, 2H, *J* = 8.6 Hz, H_{aro}), 7.38-7.55 (m, 3H, H_{aro}), 7.63 (d, 1H, *J* = 7.0 Hz, H_{aro}); ¹³C NMR (CDCl₃): δ 42.2 (CH₂), 55.4 (CH₃), 81.1 (CH), 114.2 (2CH_{aro}), 123.5 (CH_{aro}), 123.6 (CH_{aro}), 129.1 (C_q), 129.8 (CH_{aro}), 130.0 (2CH_{aro}), 131.5 (C_q), 132.5 (CH_{aro}), 144.2 (C_q), 159.2 (C_q), 167.5 (C=O); Anal. Calcd. for C₁₆H₁₅NO₃ (269.11): C, 71.36; H, 5.61; N, 5.20. Found: C, 71.22; H, 5.49; N, 5.15.

7-Hydroxy-6,7-dihydro-6-(4-methoxybenzyl)-pyrrolo[3,4-b]pyridin-5-one (2g). This product was isolated as a white solid; 72% yield; *R_f* 0.27 (cyclohexane/AcOEt 1:4); mp 170 °C; IR (KBr): ν 3164, 1663, 1615, 1587, 1513, 1439, 1422 cm⁻¹; ¹H NMR (DMSO-*d*₆): δ 3.76 (s, 3H, OCH₃), 4.34 (d, 1H, *J* = 14.9 Hz, CH₂N), 4.91 (d, 1H, *J* = 14.9 Hz, CH₂N), 5.62 (d, 1H, *J* = 8.8 Hz, CH), 6.91 (d, 2H, *J* = 8.8 Hz, H_{aro}), 6.99 (d, 1H, *J* = 8.8 Hz, OH), 7.29 (d, 2H, *J* = 8.6 Hz, H_{aro}), 7.59 (dd, 1H, *J* = 7.8 and 4.4 Hz, H_{aro}), 8.13 (dd, 1H, *J* = 7.8 and 1.5 Hz, H_{aro}), 8.80 (dd, 1H, *J* = 4.4 and 1.5 Hz, H_{aro}); ¹³C NMR (DMSO-*d*₆): δ 41.5 (CH₂), 55.1 (CH₃), 80.4 (CH), 113.9 (2CH_{aro}), 124.6 (CH_{aro}), 125.2 (C_q), 129.2 (2CH_{aro}), 129.3 (C_q), 131.1 (CH_{aro}), 152.8 (CH_{aro}), 158.5 (C_q), 163.8 (C_q), 164.5 (C=O); Anal. Calcd. for C₁₅H₁₄N₂O₃ (270.1): C, 66.66; H, 5.22; N, 10.36. Found: C, 66.53; H, 5.09; N, 10.28.

3-Hydroxy-2,3-dihydro-2-(4-methoxybenzyl)-pyrrolo[3,4-b]quinolin-1-one (2h). This product was isolated as a white solid; 77% yield; *R_f* 0.47 (cyclohexane/AcOEt 2:3); mp 211 °C; IR (KBr): ν 3066, 1706, 1634, 1616, 1514, 1428, 1412 cm⁻¹; ¹H NMR (DMSO-*d*₆): δ 3.78 (s, 3H, OCH₃), 4.43 (d, 1H, *J* = 15.1 Hz, CH₂N), 4.98 (d, 1H, *J* = 15.1 Hz, CH₂N), 5.77 (d, 1H, *J* = 9.0 Hz, CH), 6.94 (d, 2H, *J* = 8.4 Hz, H_{aro}), 7.08 (d, 1H, *J* = 9.0 Hz, OH), 7.35 (d, 2H, *J* = 8.4 Hz, H_{aro}), 7.76 (dd, 1H, *J* = 7.8

and 7.0 Hz, H_{aro}), 7.95 (dd, 1H, *J* = 8.4 and 7.0 Hz, H_{aro}), 8.19 (d, 1H, *J* = 7.8 Hz, H_{aro}), 8.26 (d, 1H, *J* = 8.4 Hz, H_{aro}), 8.82 (s, 1H, H_{aro}); ¹³C NMR (DMSO-*d*₆): δ 41.7 (CH₂), 55.0 (CH₃), 80.6 (CH), 113.9 (2CH_{aro}), 123.0 (C_q), 127.3 (CH_{aro}), 127.7 (C_q), 129.0 (CH_{aro}), 129.1 (C_q), 129.2 (2CH_{aro}), 129.8 (CH_{aro}), 131.5 (CH_{aro}), 131.9 (CH_{aro}), 149.2 (C_q), 158.5 (C_q), 162.5 (C_q), 164.3 (C=O); Anal. Calcd. for C₁₉H₁₆N₂O₃ (320.12): C, 71.24; H, 5.03; N, 8.74. Found: C, 71.07; H, 5.00; N, 8.66.

5-Hydroxy-3-methyl-1-phenethyl-1H-pyrrol-2(5H)-one (2qB). To a solution of 3-methyl-1-phenethyl-1H-pyrrole-2,5-dione (**1q**, 1.075 g, 5 mmol) in methanol 20 mL was added in one portion and under stirring cerium(III) chloride heptahydrate (1.86 g, 5 mmol). After 10 min of reaction at room temperature the solution was cooled to 0 °C and 190 mg of sodium borohydride (5 mmol) was added portionwise. After 1 h of reaction at 0 °C and additional 1 h at room temperature, the reaction mixture was quenched with ice water and methanol removed under reduced pressure. The solution was extracted twice with dichloromethane (10 mL) and the combined organic layers washed with water, brine, dried over MgSO₄ and evaporated to dryness to give after chromatography separation, eluting with cyclohexane/AcOEt 1:4), the expected minor hydroxy lactam **2qB** in 28.7% yield (overall yield of 86%). This product was isolated as a white solid; *R*_f 0.25 (cyclohexane/AcOEt 1:4); IR (KBr): ν 3054, 1712, 1645, 1612, 1507, 1433 cm⁻¹; ¹H NMR (CDCl₃): δ 1.85 (s, 3H, CH₃), 2.13 (d, 1H, *J* = 11.3 Hz, OH), 2.88 (dd, 2H, *J* = 7.8 and 7.0 Hz, CH₂), 3.51 (ddd, 1H, *J* = 14.9, 7.8 and 7.0 Hz, NCH₂), 3.76 (ddd, 1H, *J* = 14.9, 7.8 and 7.0 Hz, NCH₂), 4.98 (d, 1H, *J* = 11.3 Hz, CHOH), 6.45 (s, 1H, =CH), 7.16-7.31 (m, 5H, H_{aro}); ¹³C NMR (CDCl₃): δ 11.1 (CH₃), 34.9 (CH₂), 41.2 (CH₂), 82.2 (CH), 126.7 (CH_{aro}), 128.8 (2CH_{aro}), 129.0 (2CH_{aro}), 137.1 (C_q), 138.6 (CH_{aro}), 139.1 (C_q), 170.6 (C=O); Anal. Calcd. for C₁₃H₁₅NO₂ (217.11): C, 71.87; H, 6.96; N, 6.45. Found: C, 71.69; H, 6.87; N, 6.37.