Intramolecular Additions of Alcohols and Carboxylic Acids to Inert Olefins Catalyzed by Silver(I) Triflate

Cai-Guang Yang, Nicholas W. Reich, Zhangjie Shi, Chuan He*

Department of Chemistry, the University of Chicago,

5735 S. Ellis Avenue, Chicago, IL 60617,

E-mail: chuanhe@uchicago.edu

General Methods. Catalytic cyclization reactions were performed in sealed oven-dried glass tubes under an atmosphere of dry nitrogen. Flash column chromatography was performed employing EM Science Geduran silica gel 60 (35-75 μ m), and eluting with n-pentane/ethyl ether. Thin layer chromatography was performed on EM Science silica gel 60 F254 plates (250 μ m). NMR spectra were recorded on operating at Bruker DRX-400/500 spectrometer for ^1H NMR and 100/125 MHz for ^{13}C NMR in CDCl_3 unless noted otherwise and were referenced to tetramethylsilane (TMS) at 0.00 ppm. The chemical shifts are expressed in ppm and coupling constants are given in Hz. Data for ^1H NMR are recorded as follows: chemical shift (ppm), multiplicity (s, singlet; d, doublet; t, triplet; m, multiplet), coupling constant (Hz), integration. Data for ^{13}C NMR are reported in terms of chemical shift (δ , ppm). Low-resolution mass spectra were obtained from the Mass Spectrometry Facility at the University of Chicago. Elemental analyses were performed by Microanalytical Laboratory at University of Illinois at Urbana-Champaign. Silver triflate was purchased from Aldrich and used as received. 1,2-Dichloroethane (DCE) was redistilled over CaH_2 before used. THF was freshly distilled over sodium benzoquinone ketyl under dry nitrogen.

Most products are known compounds, and the references are given. For unknown products, copies of ^1H NMR and ^{13}C NMR spectrum are attached at the end. Elemental analysis for product (Table 2, entry 4) is also obtained.

2,2-Diphenyl-4-hexen-1-ol (Table 2, entry 3).¹ To a solution of methyl diphenylacetate (2.26 g, 10.0 mmol) in THF was added a solution of LDA (1.8 M, 6.8 mL, 12.0 mmol) dropwise at -78 $^{\circ}\text{C}$ and the resulting solution was stirred for additional 20 min. Crotyl bromide (1.45 mL, 12.0 mmol) was added and the reaction mixture was warmed to rt, stirred overnight, quenched with HCl (3 N, 10.0 mL), and extracted with ether (3 $\times$ 50 mL). The combined ether extracts were washed with brine (40 mL), dried (MgSO_4), and concentrated under vacuum. The residue was purified by silica gel chromatography to give methyl 2,2-diphenyl-4-pentenoate (2.24 g, 80%) as a viscous colorless oil.

To a suspension of LiAlH_4 (380 mg, 10.0 mmol) in THF (10 mL) was slowly added a solution of methyl 2,2-diphenyl-4-pentenoate (1.12 g, 5 mmol) at 0 $^{\circ}\text{C}$. The resulting mixture was refluxed for 2 h, and quenched by carefully addition of water (2.5 mL) and aqueous NaOH (15%, 3 mL) at 0 $^{\circ}\text{C}$. The resulting suspension was filtered through Celite and eluted with ether. The ether eluant was dried over MgSO_4 and concentrated under vacuum. The residue was purified by silica gel chromatography to give the title compound (0.91 g, 90%) as colorless oil. For *E*: ^1H NMR (500 MHz) δ 7.28 – 7.25 (m,

4H), 7.20 – 7.15 (m, 6H), 5.48 (m, 1H), 5.04 (m, 1H), 4.09 (d, $J = 6.5$ Hz, 2H), 2.87 (d, $J = 7.0$ Hz, 2H), 1.55 (d, $J = 6.5$ Hz, 3H), 1.27 (t, $J = 7.0$ Hz, 1H); ^{13}C NMR (125 MHz) δ 145.5, 128.6, 128.2, 128.1, 128.0, 126.6, 126.2, 67.9, 51.7, 39.7, 17.9; MS (m/z , EI) 235 $[(\text{M} - \text{OH})^+]$.

2-(1-Naphenyl)-4-methyl-4-penten-1-ol (Table 2, entry 4). The title compound was prepared in 72% yield as a colorless oil from reaction of methyl 1-naphthaleneacetate and 3-bromo-2-methylpropene employing a procedure analogous to the one used to synthesize 2,2-diphenyl-4-hexen-1-ol. ^1H NMR (500 MHz) δ 8.21 (d, $J = 8.5$ Hz, 1H), 7.90 (d, $J = 8.0$ Hz, 1H), 7.77 (d, $J = 8.0$ Hz, 1H), 7.58 – 7.40 (m, 4H), 4.02 (m, 1H), 3.91 (t, $J = 5.5$ Hz, 2H), 2.68 (dd, $J = 14.5$ and 7.5 Hz, 1H), 2.56 (dd, $J = 14.5$ and 7.0 Hz, 1H), 1.79 (s, 3H), 1.65 (t, $J = 5.5$ Hz, 1H); ^{13}C NMR (125 MHz) δ 143.7, 138.1, 134.1, 132.2, 129.0, 127.1, 126.0, 125.5, 125.4, 123.8, 122.8, 112.4, 66.7, 40.7, 39.7, 22.5; MS (m/z , EI) 226 (M^+).

3-Phenyl-4-penten-1-ol (Table 2, entry 5). The title compound was synthesized according to a published procedure.² ^1H NMR (400 MHz) δ 7.34 - 7.30 (m, 2H), 7.23 – 7.20 (m, 3H), 5.99 (ddd, $J = 17.3$ and 10.2 and 7.2 Hz, 1H), 5.10 (m, 1H), 5.06 (m, 1H), 3.63 (m, 2H), 3.48 (q, $J = 7.6$ Hz, 1H), 1.98 (m, 2H), 1.41 (br s, 1H); ^{13}C NMR (100 MHz) δ 143.6, 141.7, 128.5, 127.5, 126.3, 114.3, 60.8, 46.2, 37.8; MS (m/z , EI) 162 (M^+).

2,2-Bis(2-methylallyl)propane-1,3-diol (Table 2, entry 6).¹ The title compound was synthesized in 76% yield as a colorless oil from reaction of dimethyl malonate and 3-bromo-2-methylpropene employing a procedure analogous to the one used to synthesize 2,2-diphenyl-4-hexen-1-ol. ^1H NMR (500 MHz) δ 4.91 (m, 2H), 4.79 (m, 2H), 3.66 (d, $J = 4.7$ Hz, 4H), 2.40 (t, $J = 4.7$ Hz, 2H), 2.14 (s, 4H), 1.82 (s, 6H); ^{13}C NMR (125 MHz) δ 142.8, 115.0, 68.0, 43.0, 40.7, 25.2; MS (m/z , EI) 184 (M^+).

2-(*tert*-Butyldimethylsiloxymethyl)-4-methyl-2-(methylallyl)-4-penten-1-ol (Table 2, entry 7).¹ The title compound was prepared in 83% yield from 2,2-bis(2-methylallyl)propane-1,3-diol as a colorless oil according to a known procedure.³ ¹H NMR (500 MHz) δ 4.89 (m, 2H), 4.74 (m, 2H), 3.59 (m, 4H), 2.69 (t, J = 6.0 Hz, 1H), 2.13 (ABq, J = 13.5 Hz, 4H), 1.79 (s, 6H), 0.90 (s, 9H), 0.06 (s, 6H); ¹³C NMR (125 MHz) δ 142.8, 115.0, 68.8, 68.5, 43.0, 40.6, 25.8, 25.4, 18.1, -5.7.

2-(2-Methylallyl)propane-1,3-diol (Table 2, entry 8). The title compound was synthesized in 68% yield as a colorless oil from reaction of dimethyl malonate and 3-bromo-2-methylpropene employing a procedure analogous to the one used to synthesize 2,2-diphenyl-4-hexen-1-ol. ¹H NMR (400 MHz) δ 4.79 (s, 1H), 4.72 (s, 1H), 3.79-3.75 (m, 2H), 3.65-3.60 (m, 2H), 3.01 (m, 3H), 1.98 (s, 3H), 1.74 (s, 3H); ¹³C NMR (100 MHz) δ 143.3, 112.2, 64.9, 39.5, 36.5, 27.5, 22.1; MS (m/z , EI) 130 (M^+).

2-(Hydroxymethyl)-4-methylpent-4-enyl triisopropylsilane (Table 2, entry 9). The title compound was prepared in 67% isolated yield from 2-(2-methylallyl)propane-1,3-diol as a colorless oil according to a known procedure.³ ¹H NMR (400 MHz) δ 4.78 (m, 1H), 4.71 (m, 1H), 3.90-3.85 (m, 1H), 3.79-3.71 (m, 1H), 3.70-3.63 (m, 2H), 2.58 (br s, 1H), 2.08-1.98 (m, 1H), 1.97-1.95 (m, 2H), 1.74 (s, 3H), 1.10-1.05 (m, 21H); ¹³C NMR (125 MHz) δ 143.4, 112.2, 67.8, 66.9, 39.7, 36.5, 22.3, 18.0, 11.8; MS (m/z , EI) 286 (M^+).

2-(Hydroxymethyl)-4-methylpent-4-enyl acetate (Table 2, entry 10). The title compound was prepared in 54% isolated yield from 2-(2-methylallyl)propane-1,3-diol as pale yellow oil using a procedure adapted from Clarke.⁴ ¹H NMR (500 MHz) δ 4.82 (m, 1H), 4.71 (m, 1H), 4.21-4.18 (m, 1H), 4.09-4.04 (m, 1H), 3.62-3.58 (m, 1H), 3.53-3.47 (m, 1H), 2.34 (br s, 1H), 2.07 (s, 3H), 2.08-2.01 (m, 3H), 1.74 (s, 3H); ¹³C NMR (125 MHz) δ 171.7, 142.8, 112.4, 64.3, 62.2, 37.9, 36.5, 22.0, 20.7; MS (m/z , EI) 172 (M^+).

2-(Hydroxymethyl)-4-methylpent-4-enyl benzoate (Table 2, entry 11). The title compound was prepared in 50% isolated yield from 2-(2-methylallyl)propane-1,3-diol using a procedure adapted from Clarke.⁴ ¹H NMR (400 MHz) δ 7.60-7.57 (m, 2H), 7.56-7.44 (m, 3H), 4.85 (s, 1H), 4.80 (s, 1H), 4.50-4.47 (m, 1H), 4.34-4.3 (m, 1H), 3.69 (m, 1H), 3.61 (m, 1H), 2.70 (br s, 1H), 2.17 (m, 3H), 1.78 (s, 3H); ¹³C NMR (125 MHz) δ 167.1, 142.8, 133.2, 129.9, 129.6, 128.4, 112.8, 64.5, 62.4, 38.4, 36.7, 22.2; MS (m/z, EI) 234 (M⁺).

meso-2-Cyclohexenyl-2-(1-Naphthyl)ethanol (Table 2, entry 12). The title compound was synthesized in 67% yield as a colorless oil from reaction of 1-naphthaleneacetate and 3-bromocyclohexene employing a procedure analogous to the one used to synthesize 2,2-diphenyl-4-hexen-1-ol. ¹H NMR (500 MHz) δ 8.16 (dd, J = 8.0 and 7.5 Hz, 2H), 7.90 (dd, J = 7.5 and 1.5 Hz, 2H), 7.78 (m, 2H), 7.56 – 7.47 (m, 8H), 5.95 – 5.92 (m, 1H), 5.86 – 5.82 (m, 1H), 5.65 – 5.62 (m, 1H), 5.41 (m, 1H), 4.08 – 4.04 (m, 4H), 3.70 (m, 2H), 2.75 (m, 1H), 2.66 (m, 1H), 2.02 (m, 3H), 1.94 (m, 1H), 1.82 (m, 1H), 1.69 (m, 1H), 1.56 (m, 2H), 1.44 (m, 3H), 1.27 (m, 1H); ¹³C NMR (125 MHz) δ 138.1, 137.5, 134.1, 133.0, 130.0, 129.0, 128.9, 128.8, 128.1, 127.1, 127.0, 126.0, 125.5, 125.4, 125.3, 124.1, 123.3, 123.2, 64.6, 64.1, 37.9, 37.5, 27.2, 27.1, 25.4, 25.1, 22.0, 21.1, 20.7; MS (m/z, EI) 252 (M⁺).

2,2-Diphenyl-5-methyl-4-hexen-1-ol (Table 2, entry 13).¹ The title compound was synthesized in 70% yield as a colorless oil from reaction of methyl diphenylacetate and 3,3-dimethylallyl bromide employing a procedure analogous to the one used to synthesize 2,2-diphenyl-4-hexen-1-ol. ¹H NMR (400 MHz) δ 7.33 - 7.29 (m, 4H), 7.24 – 7.19 (m, 6H), 4.86 (m, 1H), 4.13 (d, J = 6.8 Hz, 2H), 2.89 (d, J = 7.2 Hz, 2H), 1.62 (s, 3H), 1.56 (s, 3H), 1.24 (t, J = 6.8 Hz, 1H); ¹³C NMR (100 MHz) δ 145.5, 134.5, 128.2, 128.1, 126.2, 119.7, 68.2, 53.4, 52.2, 34.9, 25.9, 17.8; MS (m/z, EI) 249 [(M - OH)⁺].

2,2-Dimethyl-5-phenyl-4-penten-1-ol (Table 2, entry 14). The title compound was synthesized in 70% yield as a colorless oil from reaction of methyl isobutyrate and cinnamyl bromide employing a procedure analogous to the one used to synthesize 2,2-diphenyl-4-hexen-1-ol. ^1H NMR (500 MHz) δ 7.36 – 7.34 (m, 2H), 7.30 – 7.27 (m, 2H), 7.21 – 7.18 (m, 1H), 6.40 (d, J = 16.0 Hz, 1H), 6.28 – 6.22 (m, 1H), 3.36 (s, 2H), 2.17 (dd, J = 7.5 and 1.0 Hz, 2H), 1.65 (br s, 1H), 0.91 (s, 6H); ^{13}C NMR (125 MHz) δ 137.6, 132.3, 128.5, 127.1, 126.9, 126.0, 71.6, 42.3, 36.1, 23.9.

2-(2-Cyclohexenyl-3-methyl)-2,2-dimethylethanol (6). The title compound was synthesized in 70% yield as a colorless oil from reaction of methyl isobutyrate and 1-methyl-3-bromohexene⁵ employing a procedure analogous to the one used to synthesize 2,2-diphenyl-4-hexen-1-ol. ^1H NMR (500 MHz) δ 5.40 (br s, 1H), 3.47 (dd, J = 10.9 and 5.2 Hz, 1H), 3.38 (dd, J = 10.9 and 6.6 Hz, 1H), 2.07 (m, 1H), 1.90 (m, 1H), 1.84 – 1.79 (m, 2H), 1.74 – 1.71 (m, 1H), 1.66 (s, 3H), 1.48 (m, 1H), 1.36 (t, J = 6.0 Hz, 1H), 1.17 (m, 1H), 0.86 (s, 6H); ^{13}C NMR (125 MHz) δ 135.4, 122.3, 70.2, 41.4, 37.5, 29.9, 24.2, 23.6, 23.0, 22.0, 21.6; MS (m/z , EI) 168 (M^+).

4-Methyl-2-naphthyl-4-pentenoic acid (Table 4, entry 3). A mixture of 4-methyl-2-naphthyl-4-pentenoic acid methyl ester (1.27 g, 5.0 mmol) and NaOH (1.0 g, 25.0 mmol) in 20 mL of methanol and 5 mL of water was refluxed until reaction was completed. The solvent was removed under vacuum and the residue was dissolved into 20 mL of water. The resulting mixture was washed with ether (2×40 mL). The aqueous phase was acidified with HCl solution to pH 1 ~ 2 and extracted with ether (3×50 mL). The combined ether phases were washed with brine, dried over MgSO_4 , and concentrated under vacuum. The residue was subjected to silica gel chromatography to give the title compound (1.08 g, 90%) as a white solid. ^1H NMR (500 MHz) δ 8.21 (d, J = 8.5 Hz, 1H), 7.90 (d, J = 7.5 Hz, 1H), 7.82 (d, J = 8.5 Hz, 1H), 7.59 (m, 2H), 7.53 (m, 1H), 7.48 (t, J = 7.5 Hz, 1H), 4.83 (s, 2H), 4.72 (dd, J = 9.0 and 6.0 Hz, 1H), 3.05 (dd, J = 15.0 and 9.0 Hz, 1H), 2.65 (dd, J = 15.0 and 6.0 Hz, 1H), 1.80 (s, 3H); ^{13}C NMR (125 MHz) δ 179.9, 142.5, 134.3, 134.0, 131.5, 129.0, 128.1, 126.5, 125.7, 125.4, 125.0, 122.9, 112.4, 44.9, 40.5, 22.6; MS (m/z , EI) 240 (M^+).

2,2-Dimethyl-4-hexenoic acid (Table 4, entry 4). The title compound was prepared in 69% yield from reaction of methyl isobutyrate and crotyl bromide followed by hydroxylation employing a procedure analogous to the one used to synthesize 4-methyl-2-naphthyl-4-pentenoic acid. For *E*: ^1H NMR (500 MHz) δ 5.50 (m, 1H), 5.40 (m, 1H), 2.22 (d, $J = 7.3$ Hz, 2H), 1.66 (dd, $J = 5.0$ and 1.0 Hz, 3H), 1.17 (s, 6H); ^{13}C NMR (125 MHz) δ 184.8, 128.8, 126.2, 43.2, 42.4, 24.5, 18.0; For *Z*: ^1H NMR (500 MHz) δ 5.60 (m, 1H), 5.40 (m, 1H), 2.32 (d, $J = 7.8$ Hz, 2H), 1.62 (dt, $J = 5.2$ and 0.8 Hz, 3H), 1.21 (s, 6H); ^{13}C NMR (125 MHz) δ 184.9, 126.9, 125.4, 42.5, 37.1, 24.5, 12.9.

2,2,5-Trimethyl-4-hexenoic acid (Table 4, entry 5).⁶ The title compound was prepared in 62% yield from reaction of methyl isobutyrate and 3,3-dimethylallyl bromide followed by hydroxylation employing a procedure analogous to the one used to synthesize 4-methyl-2-naphthyl-4-pentenoic acid. ^1H NMR (500 MHz) δ 5.12 (m, 1H), 2.25 (d, $J = 7.6$ Hz, 2H), 1.71 (s, 3H), 1.61 (s, 3H), 1.18 (s, 6H); ^{13}C NMR (125 MHz) δ 185.0, 134.5, 119.5, 42.7, 38.4, 26.0, 24.5, 17.9; MS (m/z , EI) 156 (M^+).

3-Phenyl-4-pentenoic acid (Table 4, entry 6). The title compound was synthesized according to a published procedure.⁷ ^1H NMR (400 MHz) δ 7.32 - 7.28 (m, 2H), 7.22 - 7.19 (m, 3H), 5.96 (m, 1H), 5.09 (m, 1H), 5.06 (dt, $J = 5.5$ and 1.3 Hz, 1H), 3.84 (q, $J = 7.7$ Hz, 1H), 2.75 (m, 2H); ^{13}C NMR (100 MHz) δ 178.5, 142.1, 139.9, 128.6, 127.5, 126.8, 115.0, 45.1, 39.9; MS (m/z , EI) 176 (M^+).

Bis(2-methylallyl)malonic acid (Table 4, entry 7).⁸ The title compound was synthesized in 56% yield from reaction of dimethyl malonate and 3-bromo-2-methylpropene followed by hydroxylation employing a procedure analogous to the one used to prepare 4-methyl-2-naphthyl-4-pentenoic acid. ^1H NMR (500 MHz) δ 4.86 (t, $J = 1.5$ Hz, 2H), 4.77 (br s, 2H), 2.76 (s, 4H), 1.71 (s, 6H); ^{13}C NMR (125 MHz) δ 177.7, 139.7, 115.5, 56.7, 45.7, 23.1.

meso-2-Cyclohexenyl-2-naphthyl acetic acid (Table 4, entry 8). The title compound was synthesized in 66% yield from reaction of 1-naphthaleneacetate and 3-bromocyclohexene followed by hydroxylation employing a procedure analogous to the one used to prepare 4-methyl-2-naphthyl-4-pentenoic acid. ^1H NMR (500 MHz) δ 8.22 (d, $J = 8.5$ Hz, 1H), 8.19 (d, $J = 8.5$ Hz, 1H), 7.87 (d, $J = 8.0$ Hz, 2H), 7.80 (d, $J = 8.0$ Hz, 2H), 7.72 (d, $J = 7.0$ Hz, 1H), 7.66 (d, $J = 7.0$ Hz, 1H), 7.55 (m, 2H), 7.51 – 7.46 (m, 4H), 5.86 (s, 2H), 5.57 (m, 1H), 5.10 (m, 1H), 4.28 (d, $J = 11.0$ Hz, 2H), 3.10 (m, 2H), 2.10 (m, 1H), 2.00 (m, 4H), 1.86 (m, 1H), 1.66 – 1.50 (m, 3H), 1.45 – 1.36 (m, 2H), 1.06 – 1.00 (m, 1H); ^{13}C NMR (125 MHz) δ 179.6, 179.4, 134.02, 133.98, 133.3, 133.0, 132.4, 132.3, 129.4, 129.2, 128.99, 128.96, 128.90, 128.03, 127.99, 127.6, 126.43, 126.39, 125.69, 125.61, 125.5, 125.3, 123.3, 123.2, 51.1, 38.3, 38.2, 28.2, 26.0, 25.3, 25.2, 21.2, 20.9.

2,2-Dimethyl-5-hexenoic acid (Table 4, entry 9).⁹ The title compound was synthesized in 64% yield from reaction of methyl isobutyrate and 4-bromobutene followed by hydroxylation employing a procedure analogous to the one used to prepare 4-methyl-2-naphthyl-4-pentenoic acid. ^1H NMR (500 MHz) δ 5.79 (m, 1H), 5.02 (m, 1H), 4.94 (m, 1H), 2.05 (m, 2H), 1.64 (m, 2H), 1.21 (s, 6H); ^{13}C NMR (125 MHz) δ 185.0, 138.2, 114.6, 41.9, 39.5, 29.2, 24.9; MS (m/z, EI) 142 (M^+).

$\alpha,\alpha,3$ -Trimethyl-2-cyclohexene-1-acetic acid (9). The title compound was synthesized in 70% yield as colorless oil from reaction of methyl isobutyrate and 1-methyl-3-bromohexene⁵ followed by hydroxylation employing a procedure analogous to the one used to prepare 4-methyl-2-naphthyl-4-pentenoic acid. ^1H NMR (500 MHz) δ 5.21 (br s, 1H), 2.48 (m, 1H), 1.89 (m, 1H), 1.85 – 1.79 (m, 2H), 1.68 (m, 1H), 1.65 (s, 3H), 1.52 – 1.48 (m, 1H), 1.18 (m, 1H), 1.14 (s, 3H), 1.10 (s, 3H); ^{13}C NMR (125 MHz) δ 185.2, 136.4, 121.5, 45.3, 42.9, 30.0, 24.0, 23.9, 22.6, 21.5.

General procedure for the silver triflate catalyzed intramolecular cyclization: A mixture of hydroxyl-olefin or carboxyl-olefin (1.0 mmol) and silver triflate (13.0 mg,

0.05 mmol) in 2 mL of dry 1,2-dichloroethane was heating at 83 °C for 15 – 36 h in a dry sealed tube under nitrogen atmosphere. The reaction was cooled to rt and the catalyst was removed by filtration through a short pad of silica gel. The ether eluant was then concentrated and the residue was subjected to silica gel chromatography to give the corresponding cyclic ethers or lactones.

2-Methyl-5,5-diphenyltetrahydropyran (Table 2, entry 3).¹ ¹H NMR (500 MHz) δ 7.43 – 7.41 (m, 2H), 7.28 – 7.23 (m, 4H), 7.17 – 7.16 (m, 4H), 4.63 (dd, J = 12.5 and 3.0 Hz, 1H), 3.56 (d, J = 12.5 Hz, 1H), 3.56 (m, 1H), 2.48 – 2.38 (m, 2H), 1.54 (dq, J = 11.0 and 3.5 Hz, 1H), 1.29 – 1.18 (m, 1H), 1.16 (d, J = 6.0 Hz, 3H); ¹³C NMR (125 MHz) δ 146.7, 145.8, 128.9, 128.2, 127.9, 126.9, 126.2, 125.5, 75.0, 73.9, 45.7, 34.7, 29.7, 21.6; MS (m/z, EI) 252 (M⁺).

2-Ethyl-4,4-diphenyltetrahydrofuran (Table 2, Entry 3).¹ ¹H NMR (400 MHz) δ 7.33 – 7.18 (m, 10H), 4.62 (dd, J = 8.7 and 0.8 Hz, 1H), 4.12 (d, J = 8.7 Hz, 1H), 3.96 (m, 1H), 2.60 (ddd, J = 12.0 and 5.7 and 1.0 Hz, 1H), 2.28 (dd, J = 12.0 and 9.6 Hz, 1H), 1.68 (m, 1H), 1.56 (m, 1H), 0.94 (t, J = 7.5 Hz, 3H); ¹³C NMR (100 MHz) δ 146.4, 146.0, 128.3, 128.2, 127.1, 126.3, 126.1, 80.1, 76.7, 55.9, 44.4, 28.9, 10.3; MS (m/z, EI) 252 (M⁺).

2,2-Dimethyl-4-naphthyltetrahydrofuran (Table 2, entry 4). ¹H NMR (400 MHz) δ 8.16 (d, J = 8.5 Hz, 1H), 7.89 (dd, J = 7.9 and 1.5 Hz, 1H), 7.76 (d, J = 7.9 Hz, 1H), 7.58 – 7.44 (m, 4H), 4.42 (dd, J = 8.3 and 7.5 Hz, 1H), 4.30 (m, 1H), 4.02 (dd, J = 8.3 and 7.5 Hz, 1H), 2.36 (dd, J = 12.3 and 7.9 Hz, 1H), 2.10 (dd, J = 12.3 and 9.7 Hz, 1H), 1.44 (s, 6H); ¹³C NMR (100 MHz) δ 137.8, 133.9, 132.0, 128.9, 127.0, 125.9, 125.5, 123.4, 122.7, 81.1, 72.4, 45.8, 41.2, 28.9, 28.4; MS (m/z, EI) 226 (M⁺); Calcd for C₁₆H₁₈O: C, 84.91, H, 8.02; found C, 84.88, H, 7.94.

2-Methyl-3-phenyl-tetrahydrofuran (Table 2, entry 5). *Trans* : *cis* = 2.7 : 1. ¹H NMR (500 MHz) δ 7.33 – 7.15 (m, 5H), [4.19 – 4.13 (m), 4.06 – 4.00 (m), 2H, 1 : 3], 3.89 – 3.83 (m, 1H), [3.32 (m), 2.80 (q, J = 8.9 Hz), 1H, 1 : 3], 2.42 – 2.35 (m, 1H), 2.21 – 2.06 (m, 1H), [1.22 (d, J = 6.0 Hz), 0.84 (d, J = 6.4 Hz), 3H, 3 : 1]; For *trans*: ¹³C NMR (125

MHz) δ 141.4, 128.5, 127.5, 126.5, 82.1, 67.3, 52.9, 35.3, 18.9; For cis: 141.7, 128.3, 128.1, 126.2, 78.1, 66.8, 48.3, 32.7, 16.8; MS (m/z, EI) 162 (M^+).

3,3,8,8-Tetramethyl-2,7-dioxaspiro[4.4]nonane (Table 2, entry 6).¹ ^1H NMR (500 MHz) δ 3.80 (d, J = 8.5 Hz, 2H), 3.61 (d, J = 8.5 Hz, 2H), 1.79 (s, 4H), 1.22 (s, 6H), 1.19 (s, 6H); ^{13}C NMR (125 MHz) δ 80.2, 76.5, 52.8, 49.3, 28.9, 28.4; MS (m/z, EI) 184 (M^+).

(1,1-Dimethylethyl)dimethyl[[tetrahydro-5,5-dimethyl-3-(2-methyl-2-propenyl)-3-furanyl]methyl]silane (Table 2, entry 7).¹ ^1H NMR (500 MHz) δ 4.82 (m, 1H), 4.70 (m, 1H), 3.72 (d, J = 9.0 Hz, 1H), 3.51 (d, J = 9.0 Hz, 1H), 3.46 (d, J = 9.5 Hz, 1H), 3.40 (d, J = 9.5 Hz, 1H), 2.24, 2.16 (ABq, J = 13.4 Hz, 2H), 1.73 (br s, 3H), 1.63, 1.59 (ABq, J = 13.4 Hz, 2H), 1.25 (s, 3H), 1.24 (s, 3H), 0.89 (s, 6H), 0.030 (s, 3H), 0.024 (s, 3H); ^{13}C NMR (125 MHz) δ 143.0, 114.1, 80.5, 73.5, 65.5, 49.8, 46.5, 42.6, 29.4, 28.7, 25.8, 24.4, 18.2, -5.6.

(Tetrahydro-5,5-dimethylfuran-3-yl)methanol (Table 2, entry 8). ^1H NMR (500 MHz) δ 3.97 (dd, J = 9.0 and 7.5 Hz, 1H), 3.67-3.57 (m, 3H), 2.57 (m, 1H), 2.05 (br s, 1H), 1.90 (dd, J = 12.5 and 7.5 Hz, 1H), 1.42 (dd, J = 12.5 and 7.5 Hz, 1H), 1.30 (s, 3H), 1.21 (s, 3H); ^{13}C NMR (125 MHz) δ 81.1, 69.5, 65.0, 42.4, 41.5, 28.7, 27.6; MS (m/z, EI) 130 (M^+).

(Tetrahydro-5,5-dimethylfuran-3-yl)methoxytriisopropylsilane (Table 2, entry 9). ^1H NMR (400 MHz) δ 3.94 (dd, J = 8.8 and 7.6 Hz, 1H), 3.70-3.60 (m, 3H), 2.57 (m, 1H), 1.83 (dd, J = 12.4 and 8.4 Hz, 1H), 1.45 (dd, J = 12.4 and 8.4 Hz, 1H), 1.29 (s, 3H), 1.21

(s, 3H), 1.12-1.03 (m, 21H); ^{13}C NMR (125 MHz) δ 80.8, 69.6, 65.5, 42.9, 41.4, 28.8, 27.9, 18.1, 12.1; MS (m/z, EI) 286 (M^+).

(Tetrahydro-5,5-dimethylfuran-3-yl)methyl acetate (Table 2, entry 10). ^1H NMR (500 MHz) δ 4.11 (dd, J = 11.0 and 6.0 Hz, 1H), 4.00-3.95 (m, 2H), 3.61 (dd, J = 9.0 and 7.0 Hz, 1H), 2.68 (m, 1H), 2.06 (s, 3H), 1.93 (dd, J = 12.5 and 8.0 Hz, 1H), 1.44 (dd, J = 12.5 and 8.0 Hz, 1H), 1.30 (s, 3H), 1.22 (s, 3H); ^{13}C NMR (125 MHz) δ 170.8, 80.6, 69.1, 66.0, 41.4, 38.9, 28.4, 27.5, 20.7; MS (m/z, EI) 172 (M^+).

(Tetrahydro-5,5-dimethylfuran-3-yl)methyl benzoate (Table 2, entry 11). ^1H NMR (500 MHz) δ 8.03-8.00 (m, 2H), 7.57-7.54 (m, 1H), 7.45-7.41 (m, 1H), 4.34 (dd, J = 11.0 and 6.5 Hz, 1H), 4.23 (dd, J = 10.5 and 8.0 Hz, 1H), 4.03 (dd, J = 8.5 and 7.0 Hz, 1H), 3.71 (dd, J = 8.5 and 7.0 Hz, 1H), 2.83 (m, 1H), 1.98 (dd, J = 12.5 and 8.0 Hz, 1H), 1.53 (dd, J = 12.5 and 8.0 Hz, 1H), 1.32 (s, 3H), 1.23 (s, 3H); ^{13}C NMR (125 MHz) δ 166.3, 132.9, 130.0, 129.4, 128.3, 80.8, 69.2, 66.5, 41.6, 39.2, 28.6, 27.5; MS (m/z, EI) 234 (M^+).

endo-cis-3-Naphthyl-octahydrobenzofuran (Table 2, entry 12). ^1H NMR (500 MHz) δ 8.08 (d, J = 8.5 Hz, 1H), 7.88 (d, J = 8.0 Hz, 1H), 7.75 (d, J = 7.5 Hz, 1H), 7.57 – 7.45 (m, 4H), 4.55 (dd, J = 9.0 and 8.0 Hz, 1H), 4.20 (q, J = 4.5 Hz, 1H), 4.17 (dd, J = 9.0 and 4.0 Hz, 1H), 3.99 (dt, J = 8.0 and 4.0 Hz, 1H), 2.29 – 2.25 (m, 1H), 2.00 – 1.91 (m, 2H), 1.70 – 1.66 (m, 1H), 1.63 – 1.54 (m, 3H), 1.46 – 1.43 (m, 1H), 1.29 – 1.26 (m, 1H); ^{13}C NMR (125 MHz) δ 138.8, 134.0, 131.9, 129.0, 126.9, 126.0, 125.5, 125.4, 123.2, 123.0, 76.3, 71.9, 45.6, 44.7, 28.5, 27.6, 23.7, 21.1; MS (m/z, EI) 252 (M^+).

***exo-cis*-3-Naphthyltetrahydrobenzofuran (Table 2, entry 12).** ^1H NMR (500 MHz) δ 8.15 (d, $J = 8.0$ Hz, 1H), 7.87 (d, $J = 8.5$ Hz, 1H), 7.75 (d, $J = 8.5$ Hz, 1H), 7.56 – 7.49 (m, 2H), 7.42 (m, 1H), 7.37 (d, $J = 6.0$ Hz, 1H), 4.60 (dd, $J = 10.0$ and 8.5 Hz, 1H), 4.42 (td, $J = 8.5$ and 3.5 Hz, 1H), 4.36 (q, $J = 3.0$ Hz, 1H), 4.24 (t, $J = 8.5$ Hz, 1H), 2.49 – 2.44 (m, 1H), 2.11 – 2.08 (m, 1H), 1.62 – 1.54 (m, 2H), 1.47 – 1.42 (m, 2H), 1.08 – 0.99 (m, 2H), 0.66 – 0.63 (m, 1H); ^{13}C NMR (125 MHz) δ 133.8, 133.4, 132.5, 128.9, 127.1, 125.9, 125.5, 125.0, 123.6, 123.2, 78.7, 68.2, 45.1, 41.6, 28.8, 24.2, 23.1, 20.4; MS (m/z , EI) 252 (M^+).

2,2-Dimethyl-5,5-diphenyltetrahydropyran (Table 2, entry 13). ^1H NMR (500 MHz) δ 7.35 – 7.34 (m, 4H), 7.31 – 7.28 (m, 4H), 7.21 – 7.18 (m, 2H), 4.09 (s, 2H), 2.46 (t, $J = 6.0$ Hz, 2H), 1.43 (t, $J = 6.0$ Hz, 2H), 1.26 (s, 6H); ^{13}C NMR (125 MHz) δ 146.4, 128.0, 127.8, 125.9, 71.2, 68.9, 45.8, 32.5, 30.7, 26.3; MS (m/z , EI) 266 (M^+).

2-Phenyl-5,5-dimethyltetrahydropyran (Table 2, entry 14). ^1H NMR (500 MHz) δ 7.38 – 7.31 (m, 4H), 7.27 – 7.23 (m, 1H), 4.23 (dd, $J = 11.0$ and 0.8 Hz, 1H), 3.62 (dd, $J = 11.0$ and 0.8 Hz, 1H), 3.34 (d, $J = 11.0$ Hz, 1H), 1.84 – 1.75 (m, 1H), 1.70 – 1.66 (m, 1H), 1.63 – 1.58 (m, 1H), 1.56 – 1.49 (m, 1H), 1.13 (s, 3H), 0.87 (s, 3H); ^{13}C NMR (125 MHz) δ 143.1, 128.2, 127.2, 125.8, 80.2, 78.6, 37.2, 30.4, 29.8, 27.2, 23.4.

***cis,cis*-3,3,7-Trimethyloctahydrobenzofuran (7).** ^1H NMR (500 MHz) δ 4.06 (dd, $J = 3.5$ and 3.0 Hz, 1H), 3.63 (d, $J = 8.0$ Hz, 1H), 3.51 (d, $J = 8.0$ Hz, 1H), 1.74 – 1.71 (m, 1H), 1.61 – 1.53 (m, 2H), 1.48 – 1.44 (m, 1H), 1.41 – 1.38 (m, 1H), 1.19 – 1.11 (m, 2H), 1.13 (s, 3H), 1.09 – 1.02 (m, 1H), 1.03 (d, $J = 7.0$ Hz, 3H), 0.98 (s, 3H); ^{13}C NMR (125 MHz) δ 80.7, 78.3, 48.2, 41.6, 34.6, 28.5, 25.1, 24.4, 21.4, 18.9; MS (m/z , EI) 168 (M^+).

1,4,4-Trimethyl-2-oxabicyclo[3.3.1]nonan (8). ^1H NMR (500 MHz) δ 3.90 (d, J = 12.0 Hz, 1H), 3.31 (d, J = 12.0 Hz, 1H), 2.0 – 1.9 (m, 3H), 1.87 – 1.83 (m, 1H), 1.61 – 1.56 (m, 2H), 1.50 – 1.42 (m, 2H), 1.27 (m, 1H), 1.10 (s, 3H), 1.08 (s, 3H), 0.90 (s, 3H); ^{13}C NMR (125 MHz) δ 74.0, 69.8, 37.7, 36.9, 36.4, 32.8, 30.2, 28.6, 26.9, 24.4, 21.1; MS (m/z, EI) 168 (M^+).

Dihydro-3,3,5-trimethyl-2(3H)-furanone (Table 4, entry 2).¹⁰ mp 49 – 50 °C; ^1H NMR (500 MHz) δ 4.56 (m, 1H), 2.18 (dd, J = 12.7 and 5.8 Hz, 1H), 1.70 (dd, J = 12.7 and 9.9 Hz, 1H), 1.40 (d, J = 6.1 Hz, 3H), 1.27 (s, 3H), 1.25 (s, 3H); ^{13}C NMR (125 MHz) δ 181.9, 73.3, 45.1, 40.8, 25.0, 24.2, 21.0; MS (m/z, EI) 128 (M^+).

Dihydro-5,5-dimethyl-3-(1-naphthyl)-2(3H)-furanone (Table 4, entry 3). mp 112 – 114 °C; ^1H NMR (400 MHz) δ 7.91 – 7.86 (m, 2H), 7.83 – 7.80 (m, 1H), 7.57 – 7.49 (m, 2H), 7.47 – 7.46 (m, 2H), 4.72 (dd, J = 10.5 and 9.6 Hz, 1H), 2.73 (dd, J = 12.9 and 9.6 Hz, 1H), 2.29 (dd, J = 12.9 and 10.5 Hz, 1H), 1.58 (s, 3H), 1.56 (s, 3H); ^{13}C NMR (100 MHz) δ 176.7, 134.0, 133.6, 131.2, 129.2, 128.3, 126.4, 125.8, 125.6, 122.8, 82.5, 44.5, 44.0, 28.9, 27.5; MS (m/z, EI) 240 (M^+).

Dihydro-5-ethyl-3,3-dimethyl-2(3H)-furanone (Table 4, entries 4 and 9).¹⁰ ^1H NMR (500 MHz) δ 4.31 (m, 1H), 2.11 (dd, J = 12.7 and 6.0 Hz, 1H), 1.70 (m, 1H), 1.68 (dd, J = 12.7 and 10.0 Hz, 1H), 1.58 (m, 1H), 1.22 (s, 3H), 1.21 (s, 3H), 0.95 (t, J = 7.4 Hz, 3H); ^{13}C NMR (125 MHz) δ 181.9, 78.2, 42.9, 40.3, 28.5, 24.9, 24.3, 9.4; MS (m/z, EI) 142 (M^+).

Tetrahydro-3,3,6-trimethyl-2H-pyran-2-one (Table 4, entries 4 and 9).¹¹ ^1H NMR (500 MHz) δ 4.37 (m, 1H), 1.79 – 1.75 (m, 1H), 1.72 – 1.56 (m, 3H), 1.28 (d, J = 6.5 Hz, 3H), 1.20 (s, 3H), 1.19 (s, 3H); ^{13}C NMR (125 MHz) δ 177.4, 77.8, 37.5, 34.2, 27.6, 27.5, 21.8; MS (m/z, EI) 142 (M^+).

Tetrahydro-3,3,6,6-tetramethyl-2H-pyran-2-one (Table 4, entry 5).¹² mp 79 - 80 °C; ¹H NMR (500 MHz) δ 1.82 – 1.79 (m, 2H), 1.76 – 1.74 (m, 2H), 1.38 (s, 6H), 1.27 (s, 6H); ¹³C NMR (125 MHz) δ 177.6, 82.4, 37.1, 32.0, 31.5, 28.9, 27.6; MS (m/z, EI) 156 (M⁺).

***trans*-Dihydro-5-methyl-4-phenyl-2(3H)-furanone (Table 4, entry 6).**¹³ ¹H NMR (400 MHz) δ 7.38 (m, 2H), 7.30 (m, 1H), 7.27 (m, 2H), 4.56 (m, 1H), 3.24 (m, 1H), 2.94 (dd, J = 17.6 and 8.5 Hz, 1H), 2.79 (dd, J = 17.6 and 11.1 Hz, 1H), 1.42 (d, J = 6.2 Hz, 3H); ¹³C NMR (100 MHz) δ 175.4, 138.2, 129.0, 127.7, 127.1, 83.0, 49.5, 37.3, 19.1; MS (m/z, EI) 176 (M⁺).

***cis*-Dihydro-5-methyl-4-phenyl-2(3H)-furanone (Table 4, entry 6).**¹⁴ ¹H NMR (400 MHz) δ 7.37 – 7.29 (m, 3H), 7.15 – 7.13 (m, 2H), 4.93 (m, 1H), 3.76 (m, 1H), 2.95 (dd, J = 17.6 and 9.0 Hz, 1H), 2.81 (dd, J = 17.6 and 6.1 Hz, 1H), 1.00 (d, J = 6.5 Hz, 3H); ¹³C NMR (100 MHz) δ 176.6, 137.5, 128.6, 127.6, 127.4, 79.8, 44.6, 34.7, 16.5; MS (m/z, EI) 176 (M⁺).

3,3,8,8-Tetramethyl-2,7-dioxaspiro[4.4]nonane-1,6-dione (Table 4, entry 7).¹⁵ ¹H NMR (400 MHz) δ 2.80 (d, J = 13.4 Hz, 2H), 2.17 (d, J = 13.4 Hz, 2H), 1.59 (s, 6H), 1.45 (s, 6H); ¹³C NMR (100 MHz) δ 174.1, 83.5, 55.0, 47.7, 29.3, 28.4; MS (m/z, EI) 212 (M⁺).

***exo-cis*-Hexahydro-*endo*-3-naphthyl-2(3H)-benzofuranone (Table 4, entry 8).** ¹H NMR (400 MHz) δ 7.94 – 7.88 (m, 2H), 7.82 (d, J = 8.2 Hz, 1H), 7.69 (d, J = 7.1 Hz, 1H), 7.57 – 7.48 (m, 3H), 4.77 (q, J = 4.0 Hz, 1H), 4.76 (d, J = 6.1 Hz, 1H), 2.81 (m, 1H), 2.35 (m, 1H), 1.67 (m, 1H), 1.57 (m, 2H), 1.45 (m, 1H), 1.04 (m, 2H), 0.79 (m, 1H);

^{13}C NMR (100 MHz) δ 176.8, 133.6, 131.8, 129.1, 128.5, 128.0, 127.3, 126.3, 125.6, 125.2, 122.0, 77.0, 50.0, 40.1, 27.6, 24.1, 23.1, 19.7; MS (m/z , EI) 266 (M^+).

endo-cis-Hexahydro-exo-3-naphthyl-2(3H)-benzofuranone (Table 4, entry 8). ^1H NMR (500 MHz) δ 7.88 (dd, $J = 7.7$ and 1.1 Hz, 1H), 7.82 (d, $J = 7.9$ Hz, 1H), 7.80 (d, $J = 7.9$ Hz, 1H), 7.57 – 7.50 (m, 2H), 7.42 (dd, $J = 7.9$ and 7.9 Hz, 1H), 7.28 (d, $J = 7.1$ Hz, 1H), 4.65 (q, $J = 4.6$ Hz, 1H), 4.28 (d, $J = 4.1$ Hz, 1H), 2.52 (m, 1H), 2.10 – 2.01 (m, 2H), 1.74 – 1.51 (m, 5H), 1.36 – 1.30 (m, 1H); ^{13}C NMR (125 MHz) δ 178.3, 134.1, 131.9, 131.4, 129.1, 128.3, 126.5, 125.9, 125.3, 124.9, 122.8, 77.3, 51.8, 43.2, 27.8, 26.9, 22.6, 20.2; MS (m/z , EI) 266 (M^+).

cis,cis-Hexahydro-3,3,7-trimethyl-2(3H)-benzofuranone (10). ^1H NMR (500 MHz) δ 4.48 (dd, $J = 4.0$ and 3.5 Hz, 1H), 1.91 (ddd, $J = 12.0$ and 5.5 and 4.0 Hz, 1H), 1.72 (m, 1H), 1.65 (m, 2H), 1.48 (m, 1H), 1.27 (s, 3H), 1.19 – 1.08 (m, 2H), 1.13 (s, 3H), 1.11 (d, $J = 7.0$ Hz, 3H), 1.06 – 0.99 (m, 1H); ^{13}C NMR (125 MHz) δ 182.2, 79.5, 45.8, 45.4, 34.0, 27.8, 24.3, 23.8, 22.9, 19.1, 18.3; MS (m/z , EI) 182 (M^+).

1,4,4-Trimethyl-2-oxabicyclo[3.3.1]nonan-3-one (11). ^1H NMR (500 MHz) δ 2.13 (dq, $J = 14.0$ and 2.5 Hz, 1H), 1.91 – 1.87 (m, 2H), 1.76 (m, 1H), 1.61 – 1.58 (m, 2H), 1.44 – 1.35 (m, 3H), 1.33 (s, 3H), 1.32 (s, 3H), 1.26 (s, 3H); ^{13}C NMR (125 MHz) δ 179.1, 81.3, 39.4, 38.2, 37.5, 34.9, 29.6, 29.0, 26.9, 23.7, 17.6; MS (m/z , EI) 182 (M^+).

Reference

1. Qian, H.; Han, X.; Widenhoefer, R. A. *J. Am. Chem. Soc.* **2004**, *126*, 9536-9537.
2. Dussalt, P. H.; Zope, U. R. *J. Org. Chem.* **1995**, *60*, 8218-8222.
3. Harnden, M. P.; Jarvest, R. L. *J. Chem. Soc. Perkin Trans. I* **1991**, 2073-2079.
4. Clarke, P. A. *Tetrahedron Lett.* **2002**, *43*, 4761-4763.
5. Breton, G. W.; Shugart, . H.; Hughey, C. A.; Conrad, B. P.; Perala, S. M. *Molecules [online computer file]*, **2001**, *6*, 655-662.

6. Fujita, T.; Watanabe, S.; Suga, K.; Nakayama, H. *Synthesis* **1979**, 310-311.
7. Li, W.; Hanau, C. E.; dAvignon, A.; Moeller, K. D. *J. Org. Chem.* **1995**, *60*, 8155-8170.
8. Bien, S.; Ovadia, D. *J. Chem. Soc. Perkin Trans. I* **1974**, 333-336.
9. Horvat, S.; Karallas, P.; White, J. M. *J. Chem. Soc. Perkin Trans. II* **1998**, 2151-2154.
10. Tiecco, M.; Tingoli, M.; Testaferri, L.; Bartoli, D. *Synth. Commun.* **1989**, *19*, 2817-2824.
11. Souma, Y.; Iyoda, J.; Sano, H. *Bull. Chem. Soc. Jpn.* **1976**, *49*, 3291-3295.
12. Jeanne-Carlier, R.; Bourelle-Wagnier, F. *Tetrahedron Lett.* **1975**, *16*, 1841-1842.
13. Reissig, H.-U.; Angert, H. *J. Org. Chem.* **1993**, *58*, 6280-6285.
14. Brown, E.; Deroye, C.; Touet, J. *Tetrahedron: Asymmetry* **1998**, *9*, 1605-1614.
15. Fristad, W. E.; Hershberger, S. S. *J. Org. Chem.* **1985**, *50*, 1026-1031.

Figure S1 Olefin 1 in CD₂Cl₂ at 298 K.

Figure S2 Olefin **1** with 20 mol % of silver triflate in CD₂Cl₂ at 298 K

Figure S3 Olefin **1** with 50 mol % of silver triflate in CD₂Cl₂ at 298 K.

Figure S4 Olefin **1** with an equimolar of silver triflate in CD₂Cl₂ at 298 K.

Figure S5 Olefin **1** in 1,2-dichloroethane- d_4 at 353 K.

Figure S6 Olefin **1** with 50 mol % Silver triflate in 1,2-dichloroethane- d_4 at 298 K.

Figure S7 Olefin **1** with 50 mol % Silver triflate in 1,2-dichloroethane- d_4 at 353 K.

Figure S8 Olefin **1** with 50 mol % Silver triflate in 1,2-dichloroethane- d_4 at 353 K for 1.5 h.

Figure S9 Olefin **1** with 50 mol % Silver triflate in 1,2-dichloroethane- d_4 at 353 K for 2.5 h.

Figure S10 Olefin **1** with 50 mol % Silver triflate in 1,2-dichloroethane- d_4 at 353 K for 12 h.

Figure S11 Excess amount of silver triflate was added to the reaction mixture at 353 K.

Figure S12 Excess amount of silver triflate was added to the reaction mixture at 353 K after 1.5 h.

Figure S13 Excess amount of silver triflate was added to the reaction mixture at 353 K after 3.5 h. Reaction complete.